UNCLASSIFIED

AD 298 008

Reproduced
by the

ARMED SERVICES TECHNICAL INFORMATION AGENCY
ARLINGTON HALL STATION
ARLINGTON 12, VIRGINIA

Reproduced From
Best Available Copy

UNCLASSIFIED

NOTICE: When government or other drawings, specifications or other data are used for any purpose other than in connection with a definitely related government procurement operation, the U. S. Government thereby incurs no responsibility, nor any obligation whatsoever; and the fact that the Government may have formulated, furnished, or in any way supplied the said drawings, specifications, or other data is not to be regarded by implication or otherwise as in any manner licensing the holder or any other person or corporation, or conveying any rights or permission to manufacture, use or sell any patented invention that may in any way be related thereto.

63-2-5

CATALOGED BY ASTIA
AS AD No. 298008

OEG

This is an interim report of continuing research. It does not necessarily represent the opinion of OEG or the U. S. Navy. It may be modified or withdrawn at any time.

Interim Research Memorandum

OPERATIONS EVALUATION GROUP

Center for Naval Analyses

WASHINGTON 25, D. C.

DISCLAIMER NOTICE



THIS DOCUMENT IS BEST QUALITY AVAILABLE. THE COPY FURNISHED TO DTIC CONTAINED A SIGNIFICANT NUMBER OF PAGES WHICH DO NOT REPRODUCE LEGIBLY.

9 November 1962

INTERIM RESEARCH MEMORANDUM

OPERATIONS EVALUATION GROUP

ESTIMATING THE LATERAL RANGE CURVE FROM OBSERVED DETECTION RANGES

By R.D. Arnold and J. Bram

IRM-27

This is an interim report of continuing research. It does not necessarily represent the opinion of OEG or the U.S. Navy. It may be modified or withdrawn at any time.

ABSTRACT

The problem of estimating the distribution of detection ranges, e.g., of a sonar or radar, in which there is no information on the total number of detection opportunities or the target tracks, has been treated in OEG IRM-7, (reference (1)). A more detailed discussion of the assumptions and derivations is presented here; the result differs slightly from that given in IRM-7.

We are given an observer with a detection device, and a collection of numbers r_i, $i=1, \ldots, N$, representing the ranges at which targets of a certain type have been detected. It is assumed that the detection device has a range r, called the detection range, which varies randomly as time passes, but that it is essentially constant during the passage of any one target. A target is assumed to be detected, at range r, if and only if it passes within a range r of the detection device. The path of each target is a straight line. Only one target at a time passes by.

It is known that there are cases in which targets pass directly by the detector and go undetected. We may think of the detector as "not working" in such cases. It is easy to include this situation in our model; we simply define the detection range r to be $r=-1$ when the device is not working. If γ_0 is the probability that the detection device is not working, then, of course, $r = -1$ with probability γ_0. We cannot estimate γ_0 in the situation described here.

The main thing is the assumption that the detection range against the given type of target is a random variable associated with the detection device; it does not depend on the individual target, or indeed, on the presence or absence of a target. During the total time of the "experiment," N targets were detected, at ranges $r_1, \ldots, r_N$; we don't know how many targets passed by that were not detected.

We shall denote by ρ the lateral range of any target (detected or not); this is the shortest distance from the observer to the straight line path traversed by the target, i.e., the "distance of closest approach." From the definition of the detection range, we see that a target is detected if and only if $0 \le \rho \le r$. Both ρ and r are random variables; ρ is associated with the target, and r with the detection device. It follows that ρ are r are independent random variables.

In the following, r_0 denotes an arbitrary, but fixed, positive number.

We are interested in two functions:

$$f(r_0) = \Pr(r \ge r_0), \tag{1}$$

and

$$L(r_0) = \Pr(\text{a target is detected, given that its lateral range is } r_0). \tag{2}$$

The first function $f(r_o)$ depends only on the detection device, but we can estimate it only through observed detections, i.e., targets are needed.

The probability space that is involved here is evidently the (ρ, r) plane, $\rho \geq o$, $r \geq -1$. The observed ranges $r_1, \ldots, r_N$ of the detected targets are independent samples of a certain probability distribution. This distribution is nothing more nor less than the distribution of detection ranges (of the detection device) given a (target) detection, i.e., the distribution of r given $o \leq \rho \leq r$, if we recall that a target is detected if and only if $o \leq \rho \leq r$.

Regarding (2), we can write

$$L(r_o) = Pr(o \leq \rho \leq r \mid \rho = r_o)$$

$$= Pr(o \leq r_o \leq r \mid \rho = r_o) .$$

But in the last expression, the two events $r_o \leq r$ and $\rho = r_o$ are independent. Hence

$$L(r_o) = Pr(r_o \leq r) = f(r_o) . \tag{3}$$

by (1). This appears in [1] as Theorem 1.

So all we need now is a reasonable estimate for $f(r_o)$. Since we have plenty of samples of the distribution of r given detection, we can write

$$Pr(r_o \leq r < r_o + \Delta r_o \mid \text{detection}) \approx \frac{\Delta n}{N} , \tag{4}$$

in which Δn is the number of observed detections with ranges in the interval $(r_o, r_o + \Delta r_o)$, $r_o \geq o$, and N is the total number of detections. But

$$\begin{aligned} &Pr(r_o \leq r < r_o + \Delta r_o \mid \text{detection}) \\ &= Pr(r_o \leq r < r_o + \Delta r_o \mid o \leq \rho \leq r) \\ &= \frac{Pr(r_o \leq r < r_o + \Delta r_o \text{ and } \rho \leq r)}{Pr(\rho \leq r)} . \end{aligned} \tag{5}$$

Let $\alpha = Pr(\rho \le r) = Pr(\text{detection})$.

Now the numerator of (5) differs only negligibly from

$$Pr(r_o \le r < r_o + \Delta r_o \text{ and } \rho \le r_o)$$

if Δr_o is small, and provided that ρ has a continuous distribution. Since $\{r_o \le r < r_o + \Delta r_o\}$ and $\{\rho \le r_o\}$ are independent events, (4) and (5) now yield

$$\frac{Pr(r_o \le r < r_o + \Delta r).\ Pr(\rho \le r_o)}{\alpha} \approx \frac{\Delta n}{N},$$

or

$$Pr(r_o \le r < r_o + \Delta r_o) \approx \frac{\alpha}{N} \quad \frac{\Delta n}{Pr(\rho \le r_o)}. \tag{6}$$

Now let us suppose that ρ is uniformly distributed in an interval $o \le \rho \le K$. Then (6) becomes

$$\left.\begin{aligned} Pr(r_o \le r < r_o + \Delta r_o) &= \frac{\alpha}{N} \quad \frac{R\,\Delta n}{r_o} \\ &\qquad \text{if } r_o < R. \\ &= \frac{\alpha}{N}\ \Delta n \quad \text{if} \quad r_o \ge R. \end{aligned}\right\} \tag{7}$$

In an unpublished working paper, J. Neuendorffer, who first considered this problem, groups his observed ranges into range bands. While his estimate is basically good, it contains an artificial bias that is introduced by the choice of end points for his range bands.

Here we shall simply take as our estimate for $f(r_o)$ — or rather $1 - f(r_o)$, which is nondecreasing — that discrete distribution which is determined by the right side of (7). That equation will be satisfied if we take the jumps in $1 - f(r)$ — we drop the subscript from r_o — to be located in $r = -1$ and

$r = r_1, \ldots, r_N$, with magnitudes $\gamma_o, \gamma_1, \ldots$ in which

$$\gamma_o = Pr(r = -1), \text{ unknown,}$$

$$\left.\begin{aligned} \gamma_i &= \frac{\alpha}{N} \frac{R}{r_i} \quad \text{if } r_i \le R, \\ \gamma_i &= \frac{\alpha}{N} \quad \text{if } r_i > R. \end{aligned}\right\} \tag{8}$$

By summing for $i = 1, \ldots, N$, we obtain

$$1 - \gamma_o = \frac{\alpha}{N} \left[\sum_{r_i \le R} \frac{R}{r_i} + M \right], \tag{9}$$

in which M is the number of observed r_i larger than R. From (9), we can estimate α/N:

$$\alpha/N = \frac{1 - \gamma_o}{\sum_{r_j \le R} \frac{R}{r_j} + M} \tag{10}$$

and (8) can be written

$$\gamma_o = Pr(r = -1),$$

$$\left.\begin{aligned} \gamma_i &= \frac{R}{r_i} \cdot \frac{1 - \gamma_o}{\sum_{r_j \le R} \frac{R}{r_j} + M} \quad \text{if } r_i \le R \\ \gamma_i &= \frac{1 - \gamma_o}{\sum_{r_j \le R} \frac{R}{r_j} + M} \quad \text{if } r_i > R, \end{aligned}\right\} \tag{11}$$

and finally,

$$f(r) = \sum_{r_i \geq r} \gamma_i = L(r) . \tag{12}$$

This estimate, for $r \geq o$, contains the unknown γ_o in the constant factor $1 - \gamma_o$, which does not affect the shape of the curve.

INTERIM RESEARCH MEMORANDUM
IRM-27

Reference: 1. C. F. Kent, Processing Detection Data Gathered from Targets of Opportunity, OEG IRM-7, 29 Dec 1961