

UNCLASSIFIED

AD 4 2 3 9 9 3

DEFENSE DOCUMENTATION CENTER

FOR

SCIENTIFIC AND TECHNICAL INFORMATION

CAMERON STATION, ALEXANDRIA, VIRGINIA



UNCLASSIFIED


423993

D1-82-0303

"Also available from the author"

BOEING SCIENTIFIC
RESEARCH
LABORATORIES

**Generating a Variable from the Tail
of the Normal Distribution**



G. Marsaglia

Mathematics Research

September 1963

CATALOGUE NO. 2-1-1-1
48 / 100

GENERATING A VARIABLE FROM THE TAIL OF THE NORMAL DISTRIBUTION

by

G. Marsaglia

Mathematical Note No. 322

Mathematics Research Laboratory

BOEING SCIENTIFIC RESEARCH LABORATORIES

September 1963

Very fast procedures for generating normal variables may be based on representing the density function as a mixture, with the dominant terms chosen so that they lead to fast procedures in the computer. See [1] - [3]. There is always the problem of handling the tail of the normal distribution. Variables from the tail are needed so infrequently that convenience is a more important consideration than speed in searching for methods. The following method is very convenient - easy to understand and easy to program. At the same time, it is reasonably fast, requiring a logarithm and a square root.

The idea is to transform the tail of the normal distribution to the unit interval and then use the rejection technique. It goes as follows:


To generate a standard normal variable X , conditioned by $X > a$, i.e., with density $ce^{-.5x^2}$, $x > a$, generate pairs of uniform over (0,1) random variables U_1, U_2 until

$$(1) \quad U_2 < a(a^2 - 2 \ln U_1)^{-\frac{1}{2}}$$

then put $X = (a^2 - 2 \ln U_1)^{\frac{1}{2}}$.

The density of U_1 , given condition (1), is a multiple of $(a^2 - 2 \ln u_1)^{-\frac{1}{2}}$, $0 < u_1 < 1$, hence the density of $X = (a^2 - 2 \ln U_1)^{\frac{1}{2}}$ is a multiple of $e^{-.5x^2}$, $a < x < \infty$.

Graphs of $u_2 = a(a^2 - 2 \ln u_1)^{-\frac{1}{2}}$ for $a = 1$ and 3 are plotted in this figure. When $a = 3$, the probability of event (1) is .88, so that the efficiency of the rejection technique is satisfactorily high.



REFERENCES

- [1] G. Marsaglia, "Expressing a Random Variable in Terms of Uniform Random Variables", Annals Math. Stat. 32, (1961), pp. 894-898.
- [2] G. Marsaglia "Random Variables and Computers", Transactions of the Third Prague Conference, to appear.
- [3] G. Marsaglia, M. D. MacLaren, T. A. Bray, "Fast Procedure for Generating Normal Variables", Comm. of the ACM, to appear.