

UNCLASSIFIED

AD NUMBER
AD813804
NEW LIMITATION CHANGE
TO Approved for public release, distribution unlimited
FROM Distribution authorized to U.S. Gov't. agencies and their contractors; Critical Technology; MAY 1967. Other requests shall be referred to Department of the Army, Fort Detrick, Attn: Technical Releases Branch, Frederick, MD 21701.
AUTHORITY
Army Biological Defense Research Lab ltr dtd 28 Sep 1971

THIS PAGE IS UNCLASSIFIED

AD

AD 813804

TECHNICAL MANUSCRIPT 388

DISULFUR MONOXIDE:
PRODUCTION BY DESULFOVIBRIO

Warren P. Iverson


MAY 1967

DEPARTMENT OF THE ARMY
Fort Detrick
Frederick, Maryland


STATEMENT #2 UNCLASSIFIED

This document is subject to special export controls and each transmittal to foreign governments or foreign nationals may be made only with prior approval of

W.P. Iverson
Frederick, Md.



DISCLAIMER NOTICE



THIS DOCUMENT IS BEST QUALITY AVAILABLE. THE COPY FURNISHED TO DTIC CONTAINED A SIGNIFICANT NUMBER OF PAGES WHICH DO NOT REPRODUCE LEGIBLY.

Reproduction of this publication in whole or in part is prohibited except with permission of the Commanding Officer, Fort Detrick, ATTN: Technical Releases Branch, Technical Information Division, Fort Detrick, Frederick, Maryland, 21701. However, DDC is authorized to reproduce the publication for United States Government purposes.

DDC AVAILABILITY NOTICES

Qualified requesters may obtain copies of this publication from DDC.

Foreign announcement and dissemination of this publication by DDC is not authorized.

Release or announcement to the public is not authorized.

DISPOSITION INSTRUCTIONS

Destroy this publication when it is no longer needed. Do not return it to the originator.

The findings in this publication are not to be construed as an official Department of the Army position, unless so designated by other authorized documents.

DEPARTMENT OF THE ARMY
Fort Detrick
Frederick, Maryland 21701

TECHNICAL MANUSCRIPT 388

DISULFUR MONOXIDE: PRODUCTION
BY DESULFOVIBRIO

Warren P. Iverson

Special Operations Division
COMMODITY DEVELOPMENT AND ENGINEERING LABORATORY

Project 1C522301A061

May 1967

ACKNOWLEDGMENT

The author extends his thanks to A. Macdonald, Jr., and Dr. C.K. Huston, Fort Detrick for the infrared analysis; L. Daasch, CRDL, Edgewood, Maryland, for the mass spectrographic analysis; Dr. J. Ladino, Fort Detrick, for the sulfate analysis; and Dr. R.J. Myers, Univ. of California, Berkeley, for technical advice.

DISULFUR MONOXIDE: PRODUCTION BY DESULFOVIBRIO**ABSTRACT**

Desulfovibrio desulfuricans growing on agar surfaces produces a gas that appears to be identical to "Schenk's sulfur monoxide," later identified as disulfur monoxide. The gas stimulates surface growth of Desulfovibrio on an agar medium and is used by the cells as an electron donor for the reduction of benzyl viologen.

Previous investigations with Desulfovibrio desulfuricans (Mid-Continent strain A) had indicated the presence of a volatile substance or gas that was produced by this organism growing on plates of trypticase soy broth* plus 2% agar.¹ This gas was characterized by its ability to stimulate growth of the same organism on agar plates of A^{PI}** medium² where growth was ordinarily very poor and also by its ability to be used by the organism as an electron donor for the reduction of benzyl viologen (BV). Violet (reduced) areas of the dye were produced in agar [2% agar plus 0.01 M tris(hydroxymethyl)aminomethane - HCl buffer, pH 7.0 ± 1, and 0.01% BV] around heavy concentrations of Desulfovibrio cells when petri plates of the agar were placed in contact with this gas in a helium or nitrogen atmosphere.

Attempts to identify this gas by gas chromatography were unsuccessful. When the volatile material was flushed with helium from a Brewer jar containing inoculated trypticase soy agar plates through a series of dry glass traps (cooled with NaCl-ice, methanol - dry ice, and liquid nitrogen successively), the gas, as indicated by the two biological properties described above, could be concentrated in the liquid nitrogen trap. An odor resembling that of H₂S was detected in the liquid nitrogen trap as well as from the outlet tube from this trap, which was immersed in water. The water produced blackening of lead acetate paper. The odor of H₂S could be detected, however, only infrequently from a Brewer jar containing inoculated plates of trypticase soy broth plus agar. A sample of the gas from the trap in liquid nitrogen was analyzed in the mass spectrometer but no sulfur compounds could be detected. Substances with mass peaks of 44, 28, and 16 corresponding to CO₂, N₂ or CO⁻, and CH₄ were detected. CO₂ and CH₄ have previously been reported as metabolic products of Desulfovibrio.^{3,4} Because the trap, cooled with liquid nitrogen, was evacuated to remove helium before analysis, the compound might have been removed. In an attempt to isolate the sulfide compound,

* Baltimore Biological Laboratories.

** American Petroleum Institute.

the biologically produced gas was passed (using helium as a carrier) through a series of three lead acetate traps (5% solution w/v) and then to the atmosphere by bubbling through water. No darkening in any of the three traps was noticed, but a strip of lead acetate paper placed in the water exposed to the air darkened in a minute or less. An odor resembling that of H_2S was also present. The gas appeared to form a sulfide only when in contact with oxygen.

A small amount of black precipitate was obtained in a week by placing a lead acetate solution (5% w/v) in a petri plate on top of 12 inoculated plates of trypticase soy broth plus 4 ml per liter sodium lactate plus agar in a Brewer jar with an atmosphere of hydrogen with traces of oxygen present. The precipitate was collected and the gas resulting from the addition of concentrated H_2SO_4 to the precipitate in nitrogen atmosphere stimulated growth on API agar (growth and blackening in 17 hours) and acted as a hydrogen donor for BV reduction.

Another sample of the precipitate was similarly obtained and the gas was analyzed in the mass spectrometer. Mass peaks corresponding to H_2S^- , SO_2^- , and SO^- were found.

H_2S had no stimulatory effect on growth and no hydrogen donor ability. SO_2 similarly had no effect in stimulating growth on API agar, but very slight reduction of BV by Desulfovibrio cells did occur. "Sulfur monoxide," now established as S_2O by Meschi and Myers,⁵ was prepared according to the method of Schenk⁶ by passing an electrical discharge (about 5 kv) through SO_2 (generated by adding concentrated H_2SO_4 to anhydrous reagent grade Na_2SO_3) in a straight glass tube (74 cm long; 2.5 cm I.D.) with platinum electrodes sealed at each end at low pressure (2 to 3 mm). The S_2O was flushed with helium into an evacuated Brewer jar with inoculated plates of API agar and BV agar. Sufficient S_2O was produced after repeating the procedure seven or eight times so that growth was visible on the API medium within 17 hours, as well as evidence of reduction of BV.

The infrared (IR) spectra of SO_2 before and after electrical discharge are indicated in Figure 1, A and B. The absorption peaks for SO_2 at $1,155\text{ cm}^{-1}$ and $1,125\text{ cm}^{-1}$ were replaced by a single peak at $1,135\text{ cm}^{-1}$. Peaks similar to those in Figure 1 B were also obtained by passing an electrical discharge through SO_2 in S vapor, using the apparatus described by Jones.⁷ The gas appears to be stable in a dry glass trap, as evidenced by the appearance of the $1,135\text{ cm}^{-1}$ peak, for at least 24 hours. This stability is in agreement with the finding of Cordes and Schenk.⁸





Figure 1. Infrared Absorption Bands. A. IR absorption bands of SO_2 ; path length 10 cm; NaCl windows. B. IR absorption bands of SO_2 after passage of electrical discharge; path length 10 cm; NaCl windows. C. IR absorption band of gas produced by *Desulfovibrio* collected in liquid N_2 trap; path length 10 cm; NaCl windows.

The IR spectrum of the gas produced by Desulfovibrio and collected in the liquid nitrogen-cooled trap is shown in Figure 1 C. A peak at 1,135 cm^{-1} , identical to the peak of the electrically discharged SO_2 , is present. No bands for H_2S or SO_2 were found.

The IR spectra reported by Jones⁷ for S_2O indicates bands at 1,165 cm^{-1} and 679 cm^{-1} . Blukis and Myers,⁹ using frozen films, reported that the 679 cm^{-1} band disappears and the band at 1,165 cm^{-1} decreases in intensity and shifts to 1,300 cm^{-1} after heating the film to 280 K. On further examination it appears that the band found at 1,135 cm^{-1} and the one reported by Myers at 1,130 cm^{-1} are probably identical and are due to a decomposition product of S_2O , the band being caused by an S-O stretch.

Schenk¹⁰ and Murthy¹¹ reported that $\text{S}_2\text{O}_3^{--}$, S^{--} , S were produced by "SO" or S_2O (because the properties previously attributed to the presence of SO in the discharge product are shown to be due to S_2O) when it reacted with alkaline solutions. It was of interest that (0.02%) $\text{Na}_2\text{S}_2\text{O}_3$ incorporated in the API agar also stimulated growth and acted as an electron donor for the reduction of BV.

The possibility is suggested that the organisms may be producing SO as an intermediate in the reduction of sulfate from trypticase soy broth (7 to 8 mg SO_4^{--} as the barium salt per gram dry medium):



SO is quite unstable, decomposing within 3 seconds, and probably forms S_2O immediately according to the following reaction: $3 \text{SO} \longrightarrow \text{S}_2\text{O} + \text{SO}_2$.⁵ SO_2 , however, was not detected in the traps during these observations.

A large number of extracellular, irregularly shaped particles was found among the cells of Desulfovibrio that were reducing BV on agar in the presence of the gas produced by growth of the organism on trypticase soy broth agar plates. Using the technique described by Skerman,¹² in which sulfur is extracted and recrystallized from pyridine, a large number of rhombic crystals were found among the cells after several hours, suggesting that the organisms might have been producing sulfur. It cannot be too strongly stressed that until the complete products of the reaction of the gas with the agar are known, it would be impossible to state whether sulfur is produced by the organisms or by a direct breakdown of S_2O with water.¹¹

It was previously mentioned that the gas has an odor resembling that of H_2S in contact with air. Cordes and Schenk¹³ also observed a similar odor from their "sulfur monoxide." Murthy¹¹ reported S^{--} as a reaction product with 2N NaOH.

It thus appears that the gas S_2O , or its decomposition product, is produced by Desulfovibrio growing on agar plates and that it may be an intermediate product in the reduction of sulfate.

LITERATURE CITED

1. Iverson, W.P. 1966. Growth of Desulfovibrio on the surface of agar media. *Appl. Microbiol.* 14:529-534.
2. Allred, R.C. 1958. Methods used for the counting of sulfate reducing bacteria and for the screening of bacteriocides. *Prod. Mon.* 22:32-34.
3. Postgate, J. 1959. Sulfate reduction by bacteria. *Annu. Rev. Microbiol.* 13:505-520.
4. Sisler, F.D.; Zobell, C.E. 1951. Hydrogen utilization by some marine sulfate-reducing bacteria. *J. Bacteriol.* 62:117-127.
5. Meschi, D.J.; Myers, R.J. 1956. Disulfur monoxide: I. Its identification as the major constituent in Schenk's "sulfur monoxide." *J. Amer. Chem. Soc.* 78:6220-6223.
6. Schenk, P.W. 1933. Über das Schwefelmonoxyd: I. Mitteilung. *Z. Anorg. Allg. Chem.* 211:150-159.
7. Jones, A.V. 1950. Infra-red and ultraviolet spectra of sulphur monoxide. *J. Chem. Phys.* 18:1263-1268.
8. Cordes, H.; Schenk, P.W. 1933. On sulfur monoxide. *Trans. Faraday Soc.* 30:31-32.
9. Blukis, U.; Myers, R.J. 1965. Disulfur monoxide: III. Its infra-red spectrum and thermodynamic functions. *J. Phys. Chem.* 69:1154-1156.
10. Schenk, P.W. 1951. Über das Schwefelmonoxyd: XII. Reaktionen des Schwefelmonoxyds in Wasser. *Z. Anorg. Allg. Chem.* 265:169-179.
11. Murthy, A.R.V. 1952. Studies on the chemical behavior of some compounds of sulfur: I. Production of disulfur monoxide by the combustion of sulfur in oxygen at low pressure. *Proc. Indian Acad. Sci.* 36A:388-401.
12. Skerman, V.B.D. 1959. A guide to the identification of bacteria. The Williams & Wilkins Company, Baltimore, Md. 228 p.
13. Cordes, H.; Schenk, P.W. 1933. Über das Schwefelmonoxyd: II. Darstellung und Eigenschaften von Reinen Schwefelmonoxyd. *Z. Anorg. Allg. Chem.* 214:33-43.

Unclassified

Security Classification

DOCUMENT CONTROL DATA - R&D		
<i>(Security classification of title, body of abstract and indexing annotation must be entered when the overall report is classified)</i>		
1. ORIGINATING ACTIVITY (Corporate author) Department of the Army Fort Detrick, Frederick, Maryland, 21701		2a. REPORT SECURITY CLASSIFICATION Unclassified
		2b. GROUP
3. REPORT TITLE DISULFUR MONOXIDE: PRODUCTION BY <u>DESULFOVIBRIO</u>		
4. DESCRIPTIVE NOTES (Type of report and inclusive dates)		
5. AUTHOR(S) (Last name, first name, initial) Iverson, Warren P.		
6. REPORT DATE May 1967	7a. TOTAL NO. OF PAGES 12	7b. NO. OF REFS 13
8a. CONTRACT OR GRANT NO. a. PROJECT NO. IC522301A061 c. d.	9a. ORIGINATOR'S REPORT NUMBER(S) Technical Manuscript 388	
9b. OTHER REPORT NO(S) (Any other numbers that may be assigned this report)		
10. AVAILABILITY/LIMITATION NOTICES Qualified requesters may obtain copies of this publication from DDC. Foreign announcement and dissemination of this publication by DDC is not authorized. Release or announcement to the public is not authorized.		
11. SUPPLEMENTARY NOTES	12. SPONSORING MILITARY ACTIVITY Department of the Army Fort Detrick, Frederick, Maryland, 21701	
13. ABSTRACT <p><u>Desulfovibrio desulfuricans</u> growing on agar surfaces produces a gas that appears to be identical to "Schenk's sulfur monoxide," later identified as disulfur monoxide. The gas stimulates surface growth of <u>Desulfovibrio</u> on an agar medium and is used by the cells as an electron donor for the reduction of benzyl viologen.</p>		
14. Key Words Disulfides <u>Desulfovibrio desulfuricans</u> Agar Reduction Benzyl viologen		

DD FORM 1473
1 JAN 64

Unclassified

Security Classification