REPORT DOCUMENTATION PAGE

Form Approved
OMB No. 074-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503

1. AGENCY USE ONLY (Leave blank)

2. REPORT
DATE 1996

3. REPORT TYPE AND DATES COVERED
Scientific paper presented 14-19 July 1996

4. TITLE AND SUBTITLE

Laser induced fluorescence spectroscopy for *in-situ* detection of petroleum contamination

5. FUNDING NUMBERS

N/A

6. AUTHOR(S)

D.S. Knowles, S.H. Lieberman, M. Davey, & B. Wingfield

7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES)

Naval Command, Control & Ocean Surveillance Center
53475 Strothe Rd.
San Diego, CA 92152

Computer Sciences Corp.
4050 Hancock Street
San Diego, CA 92110

University of California, San Diego
San Diego, CA

8. PERFORMING ORGANIZATION
REPORT NUMBER

N/A

9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES)

SERDP
901 North Stuart St. Suite 303
Arlington, VA 22203

10. SPONSORING / MONITORING
AGENCY REPORT NUMBER

N/A

11. SUPPLEMENTARY NOTES

Scientific paper presented at the 1996 XX International Quantum Electronics Conference. 14-19 July 1996. This work was supported in part by SERDP. The United States Government has a royalty-free license throughout the world in all copyrightable material contained herein. All other rights are reserved by the copyright owner.

12a. DISTRIBUTION / AVAILABILITY STATEMENT

Approved for public release: distribution is unlimited

12b. DISTRIBUTION CODE

A

13. ABSTRACT (Maximum 200 Words)

A fiber-optic chemical sensor has been developed using laser induced fluorescence (LIF) to provide high resolution, real-time, in-situ field detection of underground petroleum contamination. Recent experiments using a 308 nm XeCl laser source have shown improved performance, particularly at detecting light molecular weight (i.e. aviation) fuels.

19980710 007

14. SUBJECT TERMS

SERDP, Petroleum contamination, Laser induced fluorescence (LIF)

15. NUMBER OF PAGES

4

16. PRICE CODE N/A

17. SECURITY CLASSIFICATION
OF REPORT

unclass.

18. SECURITY
CLASSIFICATION
OF THIS PAGE

unclass.

19. SECURITY CLASSIFICATION
OF ABSTRACT

unclass.

20. LIMITATION OF
ABSTRACT

UL

NSN 7540-01-280-5500

Standard Form 298 (Rev. 2-89)
Prescribed by ANSI Std. Z39-18
298-102

DTIC QUALITY INSPECTED 1

Laser induced fluorescence spectroscopy for *in-situ* detection
of petroleum contamination

59 - 1996

David S. Knowles & Stephen H. Lieberman

Naval Command, Control & Ocean Surveillance Center

53475 Strothe Rd.

San Diego, CA 92152

(619) 553-1281 (phone)

(619) 553-2876 (Fax)

dknowles@nosc.mil

Michele Davey

Computer Sciences Corp.

4045 Hancock St.

San Diego, CA 92110

Blair Wingfield

University of California, San Diego

San Diego, CA

Abstract

A fiber-optic chemical sensor has been developed using laser induced fluorescence (LIF) to provide high resolution, real-time, in-situ field detection of underground petroleum contamination. Recent experiments using a 308 nm XeCl laser source have shown improved performance, particularly at detecting light molecular weight (i.e. aviation) fuels.

Laser induced fluorescence spectroscopy for *in-situ* detection of petroleum contamination

David S. Knowles* & Stephen H. Lieberman
Naval Command, Control & Ocean Surveillance Center
53475 Strothe Rd.
San Diego, CA 92152

Michele Davey
Computer Sciences Corp.
4045 Hancock St.
San Diego, CA 92110

Blair Wingfield
University of California, San Diego
San Diego, CA

Fuel storage tanks, hidden underground at sites such as gasoline stations and airports throughout the world, are leaving an unexpected legacy: the soil beneath the tank is often heavily contaminated with toxic hydrocarbons that may have been leaking for decades.[1] Studies have shown that certain components of fuels are carcinogenic, and while natural bacteria will slowly degrade most hydrocarbons, the carcinogenic compounds can remain in the ground long after the source has been removed.

The need for rapid assessment of such sites has led to the development of spectroscopic systems that are capable of remotely measuring the presence of underground petroleum contamination. [2-4] These systems use laser induced fluorescence (LIF) to detect the presence of petroleum products by exciting fluorescence in a component of most petroleum, the polynuclear aromatic hydrocarbons (PAHs). The PAHs are typically excited in the UV, in our case using a 337 nm nitrogen laser, and the emission is characteristically Stokes shifted towards the visible [5]. Because petroleum compounds are a complex mixture of different PAHs (along with a large range of alkanes, cycloalkanes and other organic compounds), the emission spectrum is a convolution of a large number of individual molecular spectra. While it is not possible to deconvolve the measured spectrum to determine the exact composition of the contaminant, the spectral shape of the

[Knowles, Lieberman, Davey, Wingfield, "Laser-Induced fluorescence spectroscopy..."]

emission does provide information on the class of fuel that may be present underground, and the intensity of the fluorescence is related to the concentration of the contamination.[6]

We have recently begun experiments using a 308 nm XeCl laser in place of the nitrogen laser as the pump source for our LIF petroleum sensor. Roughly speaking, the energy needed for exciting the PAHs is inversely related to the number of rings so the shorter wavelength of the XeCl laser is able to more efficiently excite fluorescence in lighter fuels that contain only one and two ring PAHs, such as jet fuels, which have been difficult to detect using the nitrogen laser. The XeCl source also appears to improve the detection limit compared to the nitrogen laser system, perhaps due to stronger absorption at the shorter wavelengths. The effects of different excitation wavelengths on the measured spectral shapes for various fuel types is still under investigation. We will present the results of field investigations at various contaminated sites, as well as laboratory studies to compare the predicted performance of the XeCl and nitrogen laser based systems.

References

- [1]. General Accounting Office, "Hazardous Materials: Upgrading of underground storage tanks can be improved to avoid costly cleanups", GAO Report GAO/NSIAD-92-117.
- [2]. S.H.Lieberman, G. A. Theriault, S. S. Cooper, P. G. Malone, R.S. Olson, P.W. Lurk, "Rapid, subsurface, in situ field screening of petroleum hydrocarbon contamination using laser induced fluorescence over optical fibers", Proceedings of the Second International Symposium - Field Screening Methods for Hazardous Wastes and Toxic Chemicals, Las Vegas, NV, Feb 1991, pg 57-63.
- [3]. R. Neissner, A. Krupp, "Detection and chemical characterization of polycyclic aromatic hydrocarbon aerosols by means of laser induced fluorescence", Part. Part. Syst. Charact., 8, (1991), 23-28.

[Knowles, Lieberman, Davey, Wingfield, "Laser-Induced fluorescence spectroscopy..."]

[4]. R.W.St. Germain, G. D. Gillispie, "Transportable tunable dye laser for field analysis of aromatic hydrocarbons in groundwater", Proceedings of the Second International Symposium - Field Screening Methods for Hazardous Wastes and Toxic Chemicals, Las Vegas, NV, Feb 1991, pg 789-792.

[5]. J. W. Birks, **Photophysics of Aromatic Molecules**, Wiley - Interscience, London, 1970.

[6]. D.S.Knowles, S.H.Lieberman, "Field results from the SCAPS Laser-Induced Fluorescence (LIF) sensor for in-situ, subsurface detection of petroleum hydrocarbons", Proceedings of the European Symposium on Optics for environmental and Public Safety, Munich, Germany, June 1995.

*presenter/corresponding author