

AD_____

AWARD NUMBER: W81XWH-04-1-0009

TITLE: GKLF as a novel target in selenium chemoprevention of prostate cancer

PRINCIPAL INVESTIGATOR: Yan Dong, Ph.D.

CONTRACTING ORGANIZATION: Health Research Incorporated
Roswell Park Cancer Institute
Buffalo, New York 14263

REPORT DATE: February 2006

TYPE OF REPORT: Annual

PREPARED FOR: U.S. Army Medical Research and Materiel Command
Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Approved for Public Release;
Distribution Unlimited

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

REPORT DOCUMENTATION PAGE

Form Approved
OMB No. 0704-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number. **PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.**

1. REPORT DATE (DD-MM-YYYY) 01-02-2006			2. REPORT TYPE Annual		3. DATES COVERED (From - To) 1 Feb 2005 – 31 Jan 2006	
4. TITLE AND SUBTITLE GKLF as a novel target in selenium chemoprevention of prostate cancer					5a. CONTRACT NUMBER	
					5b. GRANT NUMBER W81XWH-04-1-0009	
					5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S) Yan Dong, Ph.D. E-Mail: yan.dong@roswellpark.org					5d. PROJECT NUMBER	
					5e. TASK NUMBER	
					5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Health Research Incorporated Roswell Park Cancer Institute Buffalo, New York 14263					8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) U.S. Army Medical Research and Materiel Command Fort Detrick, Maryland 21702-5012						
10. SPONSOR/MONITOR'S ACRONYM(S)					11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
13. SUPPLEMENTARY NOTES						
14. ABSTRACT The present study investigated the functional significance of the zinc finger transcription factor gut-enriched krüppel-like factor (GKLF) in mediating selenium action in the androgen receptor (AR)-null PC-3 human prostate cancer cells. We found that over-expression of GKLF enhances selenium inhibition of DNA synthesis and induction of apoptosis. Furthermore, knocking down the expression of GKLF greatly attenuates the growth suppressive and apoptosis inducing activities of selenium. Therefore, our data support an important role of GKLF induction in selenium action in the AR-null prostate cancer cells. However, we found that, in cells expressing a functional AR, the disruption of AR signaling is most likely more important than the induction of GKLF signaling for selenium action. Selenium treatment significantly decreases the expression of AR and AR-regulated genes implicated in prostate carcinogenesis (PSA, KLK2, ABCC4, DHCR24, and GUCY1A3) in five human prostate cancer cell lines irrespective of their AR genotype (wild-type vs. mutant) or sensitivity to androgen-stimulated growth. Transfection of AR in the androgen-dependent LNCaP cells weakens significantly the inhibitory effect of selenium on cell proliferation and AR target gene expression. Since the vast majority of prostate cancers, including those refractory to hormone therapy, express a functional AR, the disruption of AR signaling is probably more important for selenium action and more relevant to selenium chemoprevention of prostate cancer than the induction of GKLF.						
15. SUBJECT TERMS selenium, cancer chemoprevention, prostate cancer, GKLF transcription factor, androgen receptor						
16. SECURITY CLASSIFICATION OF:				17. LIMITATION OF ABSTRACT	18. NUMBER OF PAGES	19a. NAME OF RESPONSIBLE PERSON USAMRMC
a. REPORT U	b. ABSTRACT U	c. THIS PAGE U	19b. TELEPHONE NUMBER (include area code)			

Table of Contents

Cover.....	1
SF 298.....	2
Introduction.....	4
Body.....	4-6
Key Research Accomplishments.....	6-7
Reportable Outcomes.....	7
Conclusions.....	7-8
References.....	8-9
Appendices.....	10-18

Molecular Cancer Therapeutics, 4, 1047-1055, 2005.

A. INTRODUCTION:

This project was designed initially to investigate the role of the zinc finger transcription factor gut-enriched krüppel-like factor (GKLF) in contributing to the molecular effects of selenium in cancer chemoprevention. In the first annual progress report, we described the mechanistic basis for GKLF upregulation by selenium and the effect of GKLF overexpression on the growth of prostate cancer cells. Our data indicated a growth suppressive and pro-apoptotic function of GKLF in the androgen receptor (AR)-null PC-3 cells. However, the LNCaP cells, which contain a functional AR, responded to GKLF overexpression by inducing the expression of AR, and the effect of which predominated, leading to a modest stimulation of cell growth. We also found that selenium is able to markedly suppress AR expression. Exogenous expression of AR attenuated the growth suppressive activity of selenium, although accompanied by a significant increase in GKLF level. The data suggest that disruption of AR signaling is probably more important than the induction of GKLF signaling for selenium action in AR-expressing cells. Therefore, as approved by the DOD Prostate Cancer Research Program, we continued our GKLF study in the AR-null PC-3 cells, but shifted our research focus to selenium suppression of AR signaling in the androgen-responsive LNCaP cells.

B. BODY:

Results for Task 2 (To determine the effect of GKLF overexpression on the growth of prostate cancer cells as well as selenium growth inhibition):

Overexpression of GKLF enhances MSA inhibition of DNA synthesis in PC-3 cells. In an attempt to investigate the biological consequence of GKLF induction by MSA, we transiently transfected PC-3 cells with a GKLF expression construct, pcDNA3.1/His B-GKLF (kindly provided by Dr. Anil K. Rustgi at the University of Pennsylvania) and assessed the effect of GKLF overexpression on MSA growth inhibition. Cell proliferation was quantified with the combined use of the cell viability detection reagent WST-1 (Roche) and a BrdU ELISA system (Roche), which measures BrdU incorporation during DNA synthesis. The BrdU ELISA data were normalized by the WST-1 results to correct for the differences in cell number. As shown in Fig. 1, after exposure to 5 μ M MSA for 16 hr, DNA synthesis was suppressed by 37% in cells transfected with the mock vector. The suppression was significantly greater (50%, $P < 0.05$) in cells overexpressing GKLF.





Fig. 1. Enhanced MSA effect on DNA synthesis suppression by GKLF overexpression as detected by ELISA of BrdU incorporation. Columns, % of untreated control. The results are presented as mean \pm SE ($n = 3$). *, $P < 0.05$, statistically different from mock-transfectants.


Overexpression of GKLF induces apoptosis and enhances the effect of MSA on apoptosis induction in PC-3 cells. We next assessed the effect of GKLF overexpression on

MSA induction of apoptosis by using the Cell Death Detection ELISA^{PLUS} kit (Roche), which quantitatively detects apoptotic nucleosomes. As shown in Fig. 2, exogenous expression of GKLf in PC-3 cells led to a significant increase of apoptosis. In addition, the effect of MSA on apoptosis induction was greatly enhanced by GKLf overexpression ($P < 0.01$ compared with MSA-treated mock-transfectants).



Results for Task 3 (To assess the effect of GKLf gene knockdown on the growth inhibitory and apoptosis inducing actions of selenium):

GKLf knockdown weakens the growth suppressive activity of MSA in PC-3 cells. In order to further establish the role of GKLf induction in mediating MSA action, we adopted the small interference RNA (siRNA) technique to knock down the expression of GKLf. The predesigned StealthTM Select RNAi for GKLf from Invitrogen was used in this study. The siRNA at a concentration of 40 nM was transiently transfected into PC-3 cells using the Lipofectamine 2000 reagent (Invitrogen). As shown in Fig.3, the GKLf siRNA transfection not only markedly repressed basal GKLf mRNA expression (~90%), but also prevented the induction of this gene by MSA.



The response of the GKLf knockdown cells to MSA-mediated DNA synthesis inhibition was subsequently examined. The BrdU ELISA assay was performed at 16 hr after 10 μ M MSA treatment, and the data are presented in Fig. 4. MSA treatment inhibited DNA synthesis by 73% in the scrambled siRNA-transfected cells, as opposed to 50% in the GKLf siRNA-transfected cells ($P < 0.05$). Thus, GKLf knockdown was able to significantly mitigate the growth suppressive activity of MSA.




Fig. 4. Weakened inhibitory effect of MSA on DNA synthesis by GKLf knockdown as detected by ELISA of BrdU incorporation. Columns, % of untreated control. The results are presented as mean \pm SE ($n = 3$). *, $P < 0.05$, statistically different from scrambled siRNA-transfected samples.

GKLf knockdown attenuates the apoptosis induction activity of MSA in PC-3 cells.

MSA-induced apoptosis was also examined in the GKLf knockdown cells. Without MSA treatment, no difference was observed in apoptosis between transfection with scrambled siRNA or GKLf siRNA (data not shown). In cells treated with 10 μ M MSA for 16 hr, MSA-mediated apoptosis induction was greatly attenuated by GKLf gene knockdown ($P < 0.01$). Together with the BrdU data, our results demonstrated an important role of GKLf induction in mediating the growth inhibitory and apoptosis inducing actions of MSA.




Fig. 5. Attenuated apoptosis inducing activity of MSA by GKLf knockdown as detected by ELISA of DNA fragmentation. The results are expressed as increase of apoptosis compared with untreated control. *, $P < 0.01$, statistically different from scrambled siRNA-transfected samples.

Results for Task 4 (To investigate the effect of androgen receptor overexpression on the growth inhibitory and gene-expression modulating activities selenium):

Please see the attached article (Dong *et al.*, Molecular Cancer Therapeutics, 4, 1047-1055, 2005) in Appendix for detailed description of the specific aspects of the research pertinent to this task.

C. KEY RESEARCH ACCOMPLISHMENTS:

- Overexpression of GKLf enhances selenium inhibition of DNA synthesis in the AR-null PC-3 cells.
- Overexpression of GKLf induces apoptosis and enhances the effect of selenium on apoptosis induction in PC-3 cells.
- GKLf knockdown weakens the DNA synthesis suppressive activity of selenium in PC-3 cells.

- GKLf knockdown attenuates the apoptosis induction activity of selenium in PC-3 cells.
- Selenium treatment significantly decreases the expression of androgen receptor (AR) and prostate specific antigen (PSA) in five human prostate cancer cell lines (LNCaP, LAPC-4, CWR22Rv1, LNCaP-C81 and LNCaP-LN3), irrespective of their AR genotype (wild-type vs. mutant) or sensitivity to androgen-stimulated growth.
- Selenium suppression of AR *trans*-activation is accounted for primarily by the reduction of AR protein level.
- Selenium inhibition of five AR-regulated genes implicated in prostate carcinogenesis (PSA, KLK2, ABCC4, DHCR24, and GUCY1A3) is significantly attenuated by AR overexpression.
- Transfection of AR in LNCaP cells weakens significantly the inhibitory effect of selenium on cell growth and proliferation.

D. REPORTABLE OUTCOMES:

- ***Publication:***

Dong, Y., Zhang, H., Gao, A.C., Marshall, J.R., and Ip, C. (2005) Androgen receptor signaling intensity is a key factor in determining the sensitivity of prostate cancer cells to selenium inhibition of growth and cancer-specific biomarkers. *Molecular Cancer Therapeutics*, 4, 1047-1055.

- ***Presentation:***

Roswell Park Cancer Institute Chemotherapy/Chemoprevention Research Round, January, 2006, Oral Presentation, "Androgen-Signaling Suppression in Selenium Chemoprevention of Prostate Cancer".

- ***Funding applied:***

American Cancer Society Research Scholar Grant (Dong, PI), submitted in October, 2005.

E. CONCLUSIONS:

The results from the current study indicate an important role of GKLf induction in mediating the growth suppressive and apoptosis induction activities of selenium in the AR-null PC-3 cells. However, in cells expressing a functional AR, the disruption of AR signaling is most likely more important than the induction of GKLf signaling for selenium action. The vast majority of prostate cancers express a functional AR. Although GKLf has growth suppressive activity in the AR-null cells, such activity might be overshadowed by AR signaling in AR-expressing cells. Therefore, it would be imperative to continue our research effort on the study of selenium suppression of AR signaling.

Almost all patients with advanced prostate cancer respond initially to treatments that interfere with the AR signaling process. However, these treatments often fail after prolonged use

and recurrence becomes a major clinical issue (1). The development of hormone refractory prostate cancer is not associated with loss of AR (2, 3). Instead, the accumulation of several molecular alterations frequently leads to a lower threshold requirement of androgens for the proliferation and survival of prostate cancer cells. Amplification and/or overexpression of AR can hyper-sensitize cells to sub-physiological levels of androgens (4-7). A recent report by Chen *et al.* (4) claimed that increased AR expression is both necessary and sufficient to convert prostate cancer growth from androgen-dependent to -independent, and that AR antagonists may display agonistic activity in cells with elevated AR expression. In addition, AR gene mutations could result in a promiscuous receptor with a broad ligand-binding and *trans*-activation spectrum (8). A selenium intervention strategy aimed at diminishing the expression of AR could be helpful not only for reducing incident prostate cancer, but also for preventing relapses after endocrine therapy. In addition, the fact that selenium suppresses AR signaling provides a sound rationale for using selenium in combination with an anti-androgen as a new modality for not only the prevention but also the treatment of prostate cancer.

F. REFERENCES:

- 1 Koivisto,P., Kolmer,M., Visakorpi,T. and Kallioniemi,O.P. Androgen receptor gene and hormonal therapy failure of prostate cancer, *Am.J.Pathol.*, 152: 1-9, 1998.
- 2 Hobisch,A., Culig,Z., Radmayr,C., Bartsch,G., Klocker,H. and Hittmair,A. Androgen receptor status of lymph node metastases from prostate cancer, *Prostate*, 28: 129-135, 1996.
- 3 Sadi,M.V., Walsh,P.C. and Barrack,E.R. Immunohistochemical study of androgen receptors in metastatic prostate cancer. Comparison of receptor content and response to hormonal therapy, *Cancer*, 67: 3057-3064, 1991.
- 4 Chen,C.D., Welsbie,D.S., Tran,C., Baek,S.H., Chen,R., Vessella,R., Rosenfeld,M.G. and Sawyers,C.L. Molecular determinants of resistance to antiandrogen therapy, *Nat.Med.*, 10: 33-39, 2004.
- 5 Koivisto,P., Kononen,J., Palmberg,C., Tammela,T., Hyytinen,E., Isola,J., Trapman,J., Cleutjens,K., Noordzij,A., Visakorpi,T. and Kallioniemi,O.P. Androgen receptor gene

amplification: a possible molecular mechanism for androgen deprivation therapy failure in prostate cancer, *Cancer Res.*, 57: 314-319, 1997.

- 6 Kokontis,J., Takakura,K., Hay,N. and Liao,S. Increased androgen receptor activity and altered c-myc expression in prostate cancer cells after long-term androgen deprivation, *Cancer Res.*, 54: 1566-1573, 1994.
- 7 Visakorpi,T., Hyytinen,E., Koivisto,P., Tanner,M., Keinanen,R., Palmberg,C., Palotie,A., Tammela,T., Isola,J. and Kallioniemi,O.P. In vivo amplification of the androgen receptor gene and progression of human prostate cancer, *Nat.Genet.*, 9: 401-406, 1995.
- 8 Zhao,X.Y., Malloy,P.J., Krishnan,A.V., Swami,S., Navone,N.M., Peehl,D.M. and Feldman,D. Glucocorticoids can promote androgen-independent growth of prostate cancer cells through a mutated androgen receptor, *Nat.Med.*, 6: 703-706, 2000.

Androgen receptor signaling intensity is a key factor in determining the sensitivity of prostate cancer cells to selenium inhibition of growth and cancer-specific biomarkers

Yan Dong,¹ Haitao Zhang,¹ Allen C. Gao,² James R. Marshall,¹ and Clement Ip¹

¹Division of Cancer Prevention and Population Sciences and
²Departments of Medicine, Pharmacology and Therapeutics,
Roswell Park Cancer Institute, Buffalo, New York

Abstract

Our previous report showed that methylseleninic acid (MSA) significantly decreases the expression of androgen receptor and prostate-specific antigen (PSA) in LNCaP cells. The present study extended the above observations by showing the universality of this phenomenon and that the inhibitory effect of MSA on prostate cancer cell growth and cancer-specific biomarkers is mediated through androgen receptor down-regulation. First, MSA decreases the expression of androgen receptor and PSA in five human prostate cancer cell lines (LNCaP, LAPC-4, CWR22Rv1, LNCaP-C81, and LNCaP-LN3), irrespective of their androgen receptor genotype (wild type versus mutant) or sensitivity to androgen-stimulated growth. Second, by using the ARE-luciferase reporter gene assay, we found that MSA suppression of androgen receptor transactivation is accounted for primarily by the reduction of androgen receptor protein level. Third, MSA inhibition of five androgen receptor-regulated genes implicated in prostate carcinogenesis (*PSA*, *KLK2*, *ABCC4*, *DHCR24*, and *GUCY1A3*) is significantly attenuated by androgen receptor overexpression. Fourth, transfection of androgen receptor in LNCaP cells weakened noticeably the inhibitory effect of MSA on cell growth and proliferation. Androgen receptor signaling has been documented extensively to play an important role in the development of both androgen-dependent and -independent prostate

cancer. Our finding that MSA reduces androgen receptor availability by blocking androgen receptor transcription provides justification for a mechanism-driven intervention strategy in using selenium to control prostate cancer progression. [Mol Cancer Ther 2005;4(7):1047–55]

Introduction

Prostate cancer is the second most common cancer and the second leading cause of cancer death in men in the United States. Androgen plays an important role not only in maintaining the function of the prostate but also in promoting the development of prostate cancer (1). Androgen binds to the androgen receptor, which subsequently translocates to the nucleus and interacts with specific androgen-responsive elements (ARE) on the promoters of target genes. The interaction leads to the activation or repression of genes involved in the proliferation and differentiation of the prostate cells (2). Prostate-specific antigen (PSA) and kallikrein 2 (KLK2) are two well-accepted targets of androgen receptor. PSA, also known as kallikrein 3, is an established serum marker for the diagnosis and prognosis of prostate cancer. Although KLK2 is not as widely used as PSA, it is increasingly recognized to provide added information to disease staging (3, 4).

The randomized, placebo-controlled Nutritional Prevention of Cancer trial showed that selenium supplementation reduced the incidence of prostate cancer by 50% (5, 6). This trial was designed initially to assess the effect of selenium on nonmelanoma skin cancer. Because men accounted for a sizable proportion of the cohort (974 of a total of 1,312), there was sufficient power to analyze the changes in prostate cancer risk. When the prostate cancer data were further stratified, there was evidence of a greater reduction in risk from selenium supplementation among men who had low baseline plasma PSA levels (6). Early-stage prostate cancer is mostly responsive to androgen stimulation. The inference that the protective effect of selenium might be more pronounced in early-stage prostate cancer, as reflected by low PSA secretion, lends credence to the idea that selenium might affect androgen signaling.

Recently, we reported that a selenium metabolite, in the form of methylseleninic acid (MSA), greatly down-regulates the expression of androgen receptor and PSA in the androgen-responsive LNCaP human prostate cancer cells (7, 8). The suppression of androgen receptor signaling occurs well before any significant growth inhibition, which is accompanied by correlative changes in numerous cell

Received 4/25/05; accepted 5/11/05.

Grant support: Department of Defense Postdoctoral Fellowship Award and New Investigator Award W81XWH-04-1-0009 (Y. Dong); Roswell Park Alliance Foundation (C. Ip); National Cancer Institute grant CA91990 (C. Ip); and National Cancer Institute, Cancer Center Support Grant P30 CA16056 (Roswell Park Cancer Institute).

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked advertisement in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

Requests for reprints: Yan Dong, Department of Cancer Chemoprevention, Roswell Park Cancer Institute, Elm and Carlton Streets, Buffalo, NY 14263. Phone: 716-845-1583. E-mail: yan.dong@roswellpark.org

Copyright © 2005 American Association for Cancer Research.

cycle and apoptosis regulatory molecules (9–13). Androgen receptor signaling involves multiple steps, the receptor itself is just one of many effectors that participate in the process. For example, heat shock proteins are known to modulate the stability of androgen receptor as well as its affinity to androgen (14, 15). The transactivating activity of androgen receptor can be affected markedly by a large number of coactivators and corepressors (16). Our microarray analysis suggests that MSA alters the expression of several heat shock proteins, coactivators, and corepressors of the superfamily of steroid hormone receptors (17). In view of these confounding effects, the present study was designed to determine the role of androgen receptor down-regulation per se in MSA interference of androgen receptor signaling. Our approach was to use the ARE-luciferase reporter gene assay to find out the extent to which selenium suppression of androgen receptor transactivation could be reversed when the luciferase activity is normalized based on androgen receptor protein level. We also investigated whether androgen receptor transfection might attenuate selenium-mediated down-regulation of five androgen receptor targets: *PSA*, *KLK2*, ATP-binding cassette C4 (*ABCC4*, also known as *MRP4*), 24-dehydrocholesterol reductase (*DHCR24*, also known as *seladin-1*), and soluble guanylate cyclase 1 α 3 (*GUCY1A3*). These five androgen-inducible genes were selected based on the criteria that they are consistently overexpressed in prostate cancer compared with normal prostate tissue (18). Finally, in an effort to evaluate the biological significance of the selenium-androgen receptor signaling axis, we investigated whether androgen receptor overexpression might block the growth inhibitory effect of selenium.

Materials and Methods

Selenium Reagent, Prostate Cancer Cell Lines, 3-(4,5-dimethylthiazol-2-yl)-2,5-Diphenyltetrazolium Bromide Assay, and Bromodeoxyuridine-Labeling Analysis

MSA was synthesized as previously described (19). The LNCaP and CWR22Rv1 human prostate cancer cell lines were obtained from American Type Culture Collection (Manassas, VA). The LAPC-4 cell line was provided by Dr. Charles L. Sawyers at the University of California at Los Angeles Jonsson Comprehensive Cancer Center. The two androgen-unresponsive LNCaP sublines, LNCaP-LN3 and LNCaP-C81, were obtained from Dr. Curtis A. Pettaway (University of Texas M.D. Anderson Cancer Center) and Dr. Ming-Fong Lin (University of Nebraska Medical Center), respectively. The cells were cultured in RPMI 1640 supplemented with 10% fetal bovine serum, 100 unit/mL penicillin, 100 μ g/mL streptomycin, and 2 mmol/L glutamine. The 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) assay for cell growth and the bromodeoxyuridine (BrdUrd) labeling for DNA synthesis were done as described in our previous publication (17).

Transient Transfection of Androgen Receptor

The procedure was carried out using the Lipofect-AMINE Plus reagent (Invitrogen, Carlsbad, CA) per instruction of the manufacturer. At 24 hours before transfection, cells were plated in growth medium without antibiotics at a density to reach 90% to 95% confluency at transfection. The pSG5hAR androgen receptor expression vector (20) or the pSG5 mock plasmid (Stratagene, La Jolla, CA) was introduced into LNCaP cells with or without the cotransfection of the pEGFP-F membrane-GFP-encoding construct (BD Biosciences, San Jose, CA). The purpose of the green fluorescent protein (GFP) was to enable us to enrich for the subset of positively androgen receptor-transfected cells. During cotransfection, the two plasmids were added at 1:1 molar ratio. The amount of DNA transfected was 12 μ g per 10-cm culture dish. The DNA/liposome mixture was removed at 3 hours after transfection. For the MTT assay, the cells were trypsinized 16 hours later and plated in triplicate onto a 96-well plate. Cells were allowed to recover for an additional 24 hours before exposure to 10 μ mol/L MSA. The MTT assay was conducted at 48 hours post-MSA treatment. For the BrdUrd-labeling analysis, the cells were subjected to MSA treatment at 24 hours posttransfection and labeled with BrdUrd after 24 hours of MSA treatment.

Reporter Gene Assay

The ARE-luciferase reporter plasmid, containing three repeats of the ARE region ligated in tandem to the luciferase reporter (20), was transiently transfected into cells at a concentration of 9 μ g per 10-cm culture dish. After incubating with the transfection mixture for 3 hours, the cells were trypsinized, resuspended in medium containing charcoal-stripped serum and 10 nmol/L dihydrotestosterone (Sigma, St. Louis, MO), and plated in triplicate onto 6-well plates. Cells were allowed to recover for an additional 24 hours before exposure to 10 μ mol/L MSA. After 6 or 16 hours of MSA treatment, cells were lysed in reporter lysis buffer (Promega, Madison WI), and the luciferase activity was assayed using the Luciferase Assay System (Promega). Protein concentration in cell extracts was determined by the bicinchoninic acid protein assay kit (Pierce, Rockford, IL). Luciferase activities were normalized by the protein concentration of the sample. The transfection experiments were repeated thrice.

Western Blot Analysis

Details of the procedure for Western blot analysis were described previously (17). Immunoreactive bands were quantitated by volume densitometry and normalized to glyceraldehyde-3-phosphate dehydrogenase. The following monoclonal antibodies were used in this study (source): anti-glyceraldehyde-3-phosphate dehydrogenase (Chemicon, Temecula, CA), anti-androgen receptor (BD Biosciences), and anti-PSA (Lab Vision, Fremont, CA).

Real-time Reverse Transcription-PCR

Real-time reverse transcription-PCR analysis was done as described previously (21). The PCR primers and

Taqman probes for β -actin, androgen receptor, PSA, KLK2, ABCC4, DHCR24, and GUCY1A3 were Assays-on-Demand products from Applied Biosystems (Foster City, CA). The PCR conditions were as follows: an initial incubation at 50°C for 2 minutes, then a denaturation at 95°C for 10 minutes followed by 40 cycles of 95°C for 15 seconds and 60°C for 1 minute. The relative quantitation of gene expression was done using the comparative C_T ($\Delta\Delta C_T$) method (22).

Androgen Receptor mRNA Stability Assay

Actinomycin D (5 μ g/mL) was added to the cultures to stop new RNA synthesis at the time of MSA treatment, and androgen receptor mRNA levels were measured by real-time reverse transcription-PCR at hourly intervals for the next 6 hours. The turnover of androgen receptor mRNA was determined by comparing mRNA levels over time in cells treated with or without MSA.

Statistical Analysis

The Student's two-tailed *t* test was used to determine significant differences between treatment and control values, and $P < 0.05$ was considered statistically significant.

Results

MSA Depresses Androgen Receptor Transcription

Figure 1 shows the effect of MSA on androgen receptor transcript and protein levels as well as androgen receptor mRNA stability in LNCaP cells. The decrease in androgen receptor transcript, as determined by real-time reverse transcription-PCR, occurred very quickly (Fig. 1A). On the average, there was about a 50% reduction in the first three hours after treatment with 10 μ mol/L MSA; by 6 hours, the magnitude of inhibition rose to 80%. At the protein level, there was no change in androgen receptor in the first two hours (Fig. 1B). A modest decrease began to appear at 3 hours, and the inhibition became very pronounced at 6 hours (Fig. 1B). The observation is consistent with the time-dependent sequence of reduced mRNA leading to decreased protein expression. To determine whether the down-regulation of androgen receptor mRNA was due to decreased transcription or increased mRNA degradation, we did an mRNA stability assay under the condition in which new RNA synthesis was blocked. Actinomycin D was added to the culture at the time of MSA treatment, and androgen receptor mRNA levels were followed in a 6-hour time course experiment. Because actinomycin D could be cytotoxic, we also monitored cell growth for up to 8 hours and did not observe cell death or significant growth inhibition during this period. Our results showed that treatment with MSA actually increased the stability of androgen receptor mRNA (Fig. 1C). This observation rules out increased mRNA degradation as a contributing factor. Therefore, the decrease in androgen receptor mRNA level by MSA is likely to be accounted for by a vigorous block of androgen receptor transcription.

We next examined the effect of MSA on the expression of androgen receptor and PSA in four additional human

prostate cancer cell lines: LAPC-4, CWR22Rv1, LNCaP-C81, and LNCaP-LN3. The LAPC-4 cells are androgen responsive and express a wild-type androgen receptor (23), as opposed to LNCaP cells that are also androgen responsive but express a mutant, although functional, androgen receptor. The other three cell lines are all androgen-unresponsive and express a mutant but functional androgen receptor (24–27). As shown in Fig. 2 (left), MSA decreased androgen receptor and PSA transcript levels progressively as a function of time in all four cell lines examined. The reduction in androgen receptor and PSA proteins (right) paralleled the drop in the transcripts. In LAPC-4, CWR22Rv1, and LNCaP-C81 cells, a decrease in PSA transcript was already detectable as early as 3 hours, at a time when there was no apparent loss of the androgen receptor protein. The data suggest that MSA disrupts androgen receptor signaling through additional mechanism(s) beyond reducing the availability of the androgen receptor protein.




Figure 1. Effect of MSA on androgen receptor (AR) expression in LNCaP cells. **A**, inhibition of androgen receptor mRNA level as determined by real-time RT-PCR. **B**, inhibition of androgen receptor protein level as determined by Western blot analysis. **C**, androgen receptor mRNA stability in the presence or absence of MSA. Bars, SE. *, $P < 0.05$, statistically different compared with untreated control.




Figure 2. Effect of MSA on androgen receptor (AR) and PSA expression in LAPC-4, CWR22Rv1, LNCaP-C81, and LNCaP-LN3 cells. *Left*, mRNA levels as determined by real-time RT-PCR; *right*, protein levels as determined by Western blot analysis. *Columns*, % inhibition; *bars*, SE. *, $P < 0.05$, statistically different compared with untreated control.

MSA-Mediated Androgen Receptor Down-Regulation Leads to a Reduction of Androgen Receptor Transactivating Activity

The transactivation of androgen receptor is an indicator of androgen receptor signaling and can be quantified readily by a reporter gene assay. To investigate whether the reduced availability of androgen receptor by MSA is a major factor in modulating androgen receptor transcriptional activity, we transiently transfected LNCaP cells with the ARE-luciferase reporter plasmid and normalized the luciferase activity based on the level of the androgen receptor protein. This normalization step eliminates the level of androgen receptor expression as a determinant of androgen receptor transactivation. The luciferase reporter assay was carried out at

6 and 16 hours after treatment with 10 $\mu\text{mol/L}$ MSA. At these two time points, androgen receptor protein level was inhibited by 60% and 77%, respectively (Fig. 3B, *inset*). As can be seen in Fig. 3A, without normalizing for the difference in androgen receptor protein level between the MSA-treated and -untreated samples, the ARE-promoter activity was decreased by 65% or 75%, respectively, after 6 or 16 hours of MSA treatment. However, after normalization, the ARE-promoter activity was inhibited by a meager 15% at the 6-hour time point, and the inhibition disappeared completely at 16 hours (Fig. 3B). These findings suggest that the reduced availability of the androgen receptor protein is the major factor in contributing to MSA disruption of androgen receptor signaling.

Overexpression of Androgen Receptor Attenuates the Effect of MSA on the Down-Regulation of Androgen Receptor – Regulated Genes

To delineate the role of low androgen receptor abundance as a cause of reduced PSA expression by selenium, we transiently transfected LNCaP cells with a wild-type androgen receptor construct and determined the response of PSA to MSA. After 3 hours of MSA exposure, PSA transcript was depressed by about 75% in the mock-transfected cells but only by about 45% in the androgen receptor-transfected cells (Fig. 4A). Based on our routine experience of a 40% transfection efficiency as determined by GFP cotransfection analysis (described below), we believe that the inhibitory effect of MSA on PSA mRNA might have been reversed completely in positive androgen receptor transfectants. Our conclusion was derived from the following theoretical calculation: 40% of $(1 - x) + 60%$ of $(1 - a) = 1 - b$, where $x = %$ inhibition in positive androgen receptor transfectants, $a = 75%$ inhibition in mock transfectants, and $b = 45%$ inhibition in the mixed population of androgen receptor-transfected cells. Solving for x in the above equation gave a value of 0% inhibition. In




Figure 3. Effect of MSA on ARE-promoter activity before (**A**) and after (**B**) normalizing by androgen receptor (AR) protein level. **B**, representative Western blot analysis of androgen receptor protein level in the cell extracts (*inset*). Androgen receptor protein level was depressed by 60% or 77% at 6 or 16 h, respectively. *Columns*, % inhibition; *bars*, SE. *, $P < 0.05$, statistically different compared with untreated control.




Figure 4. Effect of androgen receptor (AR) overexpression on MSA down-regulation of PSA. **A**, PSA mRNA expression as determined by real-time RT-PCR, in androgen receptor- or mock-transfected LNCaP cells treated with MSA. *Columns*, % inhibition. *, $P < 0.05$, statistically different compared with the value from the mock transfectant. **B**, PSA Western blot analysis in androgen receptor- or mock-transfected LNCaP cells treated with MSA.

other words, there was no inhibition of PSA expression by MSA in the positive androgen receptor transfectants (i.e., complete reversal). The difference between the mock- and androgen receptor-transfected cells, although still apparent, was not as great at 4 and 6 hours compared with that at 3 hours. The fact that a robust androgen receptor presence was not sufficient to completely counteract the suppressive effect of MSA on the transcription of PSA at the later time points suggests that there could be a delay in the recruitment of additional mechanisms by which MSA might diminish androgen receptor signaling. We also studied the protein level of PSA by Western blotting. The Western analysis was done at 24 hours after MSA treatment. As shown in Fig. 4B, PSA protein was depressed by about 70% in the mock-transfected cells but only by about 40% in the androgen receptor-transfected cells. The protein levels of androgen receptor in the mock- and androgen receptor-transfected cells are also shown in Fig. 4B for confirmation purposes.

Using a bioinformatic data mining approach, we recently identified five additional androgen-inducible genes that are expressed at a higher level in prostate cancer compared

with normal prostate tissue; furthermore, their expression is repressed by MSA (18). These genes are *KLK2*, *ABCC4* (also known as *MRP4*), *DHCR24* (also known as *seladin-1*), *GUCY1A3*, and long-chain fatty acid CoA ligase 3 (*FACL3*). MSA down-regulation of their expression only occurs in LNCaP cells but not in the androgen-unresponsive PC-3 cells that express an extremely low level of androgen receptor (18). To verify that the decreased expression of these genes is a direct consequence of MSA suppression of androgen receptor signaling, we applied the same androgen receptor overexpression protocol as described above and used real-time reverse transcription-PCR to quantitate their transcript levels. The *FACL3* gene was not included in this study as no Assays-on-Demand primers and probes are available for this gene. The results are shown in Fig. 5. Androgen receptor transfection significantly muted the inhibition of gene expression by MSA. In general, the difference in % inhibition between the mock and androgen receptor transfectants was greatest at 3 hours and narrowed gradually with time. The overall pattern was very similar for *KLK2*, *ABCC4*, *DHCR24*, and *GUCY1A3*. The data thus show a key role of androgen receptor down-regulation in mediating the inhibitory effects of MSA on the expression of androgen receptor-regulated genes.

Overexpression of Androgen Receptor Interferes with MSA-Mediated Growth Inhibition

In an effort to evaluate the biological significance of MSA suppression of androgen receptor signaling, we transiently transfected LNCaP cells with a wild-type androgen receptor and assessed the response of the androgen

receptor-overexpressing cells to MSA-induced growth inhibition. The MTT assay was conducted at 48 hours post-MSA, and the data are presented in Fig. 6A. In the absence of MSA, cell growth was not altered by the transfection of androgen receptor (data not shown), indicating that the endogenous level of androgen receptor is not a limiting factor for the growth of these cells. MSA treatment inhibited growth by 40% in the mock transfectants, as opposed to 27% in the androgen receptor transfectants. The difference is statistically significant ($P = 0.003$). Thus, androgen receptor overexpression was able to weaken the growth suppressive activity of MSA. One reason that the difference was seemingly compressed was due to the fact that only a fraction of cells was successfully transfected, and in this study, cell growth was assessed using the whole cell population. To address the last problem, we cotransfected cells with the androgen receptor expression vector and a membrane-GFP-encoding construct. The cells were then subjected to BrdUrd labeling, and the data were analyzed by gating just the GFP-positive cells. As shown in Fig. 6B, after selecting for the subset of GFP-positive cells, we found that MSA inhibited DNA synthesis by a very modest 16% in the androgen receptor transfectants, as opposed to 72% in the mock transfectants. Because the GFP and androgen receptor cDNAs are not located in the same plasmid construct, it is possible that not all the cells positive for GFP are also positive for the transfected androgen receptor. Thus, our selection process only led to an enrichment, rather than an exclusive selection, of double-positive cells. Therefore, the difference




Figure 5. Effect of androgen receptor (*AR*) overexpression on MSA down-regulation of *KLK2*, *ABCC4*, *DHCR24*, and *GUCY1A3* mRNA expression as determined by real-time RT-PCR, in androgen receptor- or mock-transfected LNCaP cells treated with MSA. Columns, % inhibition. *, $P < 0.05$, statistically different compared with the value from the mock transfectant.




Figure 6. Effect of androgen receptor (AR) overexpression on MSA inhibition of cell growth. **A**, MTT cell growth assay in androgen receptor- or mock-transfected LNCaP cells treated with MSA. Western blot confirmation of androgen receptor protein level (inset). **B**, BrdUrd labeling of selected GFP-positive or nonselected androgen receptor-transfected LNCaP cells treated with MSA. Columns, % inhibition compared with untreated control. *, $P < 0.05$, statistically different from mock transfectant. **, $P < 0.05$, statistically different from mock transfectant and nonselected androgen receptor transfectant.

between the mock transfectants and the androgen receptor transfectants might have been even more pronounced if all the cells used in the experiment were successfully transfected with androgen receptor. Figure 6B also shows that when we did the BrdUrd labeling experiment with the nonenriched androgen receptor-transfected cells, the inhibition by MSA was about 45%, a value half-way between that achieved by the mock transfectants and the enriched androgen receptor transfectants.

Discussion

Our previous report showed that selenium significantly decreases the expression and the transactivating activity of androgen receptor in LNCaP cells (21). The present study extended the above observations by showing the universality of this phenomenon and a key role of androgen receptor down-regulation in mediating the inhibitory effects of selenium on prostate cancer cell growth and the expression of cancer-specific biomarkers. First, selenium decreases the expression of androgen receptor and PSA in five human prostate cancer cell lines, irrespective of their androgen receptor genotype (wild type versus mutant) or sensitivity to androgen-stimulated growth. Second, a reporter gene assay with the ARE-luciferase construct indicated that depletion of the androgen receptor protein

is a major factor for selenium depression of androgen receptor transactivating activity. Third, overexpression of androgen receptor greatly weakens the inhibitory effects of selenium on prostate cancer cell proliferation as well as the expression of five androgen receptor-regulated genes implicated in prostate carcinogenesis: *PSA*, *KLK2*, *ABCC4*, *DHCR24*, and *GUCY1A3*. These findings, however, do not necessarily exclude additional mechanisms by which selenium diminishes androgen receptor signaling (e.g., via modulation of ligand binding, androgen receptor dimerization, nuclear translocation, and the interaction of androgen receptor with its coregulators). In fact, our previous report provided some evidence that selenium is able to inhibit the binding of androgen receptor to the ARE in the absence of a drop in the androgen receptor level (21).

A selenium intervention strategy aimed at diminishing the expression of androgen receptor could be helpful not only for reducing prostate cancer incidence but also for preventing relapses after endocrine therapy. Almost all patients with advanced prostate cancer respond initially to treatments that interfere with the androgen receptor-signaling process. However, these treatments often fail after prolonged use and recurrence becomes a major clinical issue (28). The development of hormone refractory prostate cancer is not associated with loss of androgen receptor (29, 30). Instead, the appearance of several molecular alterations frequently leads to a lower threshold requirement of androgens for the proliferation and survival of prostate cancer cells. Androgen receptor gene mutations could result in a promiscuous receptor with a broad ligand-binding and transactivation spectrum (31). Amplification and/or overexpression of androgen receptor may hypersensitize cells to subphysiologic levels of androgens (32–35). A recent report by Chen et al. (35) claimed that increased androgen receptor expression is both necessary and sufficient to convert prostate cancer growth from androgen-dependent to -independent and that androgen receptor antagonists may display agonistic activity in cells with elevated androgen receptor expression. On the other hand, several studies showed that knocking down the expression of androgen receptor inhibits the growth of prostate cancer cells, both *in vitro* and *in vivo*, and induces apoptosis (36–39). Because selenium blocks the transcription of androgen receptor (see Fig. 1), this treatment modality may prove to be effective in prostate cancer intervention.

The down-regulation of androgen receptor targets by selenium has important clinical implication. We have studied *PSA*, *KLK2*, *ABCC4*, *DHCR24*, and *GUCY1A3*. All these genes are expressed at a higher level in prostate cancer compared with normal prostate tissue (18). *PSA* and *KLK2* are prostate-specific differentiation markers. They belong to the serine protease family and are both secretory proteins. *PSA* is the most useful serum marker for the diagnosis and prognosis of prostate cancer. The combined use of *PSA* and *KLK2* has been shown to improve the specificity of biochemical detection of prostate cancer (40–44) and the accuracy in predicting tumor grade and

stage (3, 4). ABCC4 (also known as MRP4) is a member of the multidrug resistance-associated protein family of transporters. Overexpression of ABCC4 in neuroblastoma is associated with poor prognosis and resistance to the topoisomerase I poison irinotecan and its active metabolite SN-38 (45). Thus, the down-regulation of MRP4 by selenium might represent a potential mechanism by which selenium enhances the therapeutic efficacy of a number of anticancer drugs, including irinotecan (46). DHCR24 (also known as seladin-1) is an antiapoptotic protein, it inhibits the activity of caspase 3 (47). The overexpression of this gene has also been reported in adrenocortical adenoma cells compared with adjacent nontumor cells (48). GUCY1A3 catalyzes the conversion of GTP to the second messenger cyclic guanosine 3',5'-monophosphate, which regulates the activity of protein kinases, phosphodiesterases, and ion channels (49). Future selenium intervention trial may consider monitoring androgen receptor, PSA, KLK2, ABCC4, DHCR24, and GUCY1A3 in biopsied prostate samples, to obtain a more comprehensive picture of an individual's responsiveness to selenium. Recent data also showed that cellular PSA is more sensitive than secretory PSA to selenium intervention (50). This is one more reason why it is preferable to do the analysis in biopsied prostate tissue.

Acknowledgments

We thank Hai Wang, Dorothy Donovan, Tamora Loftus, Todd Parsons, and Cathy Russin for their excellent technical assistance.

References

- Culig Z, Hobisch A, Hittmair A, et al. Expression, structure, and function of androgen receptor in advanced prostatic carcinoma. *Prostate* 1998;35:63–70.
- Koivisto P, Kolmer M, Visakorpi T, Kallioniemi OP. Androgen receptor gene and hormonal therapy failure of prostate cancer. *Am J Pathol* 1998;152:1–9.
- Haese A, Becker C, Noldus J, et al. Human glandular kallikrein 2: a potential serum marker for predicting the organ confined versus non-organ confined growth of prostate cancer. *J Urol* 2000;163:1491–7.
- Recker F, Kwiatkowski MK, Piironen T, et al. Human glandular kallikrein as a tool to improve discrimination of poorly differentiated and non-organ confined prostate cancer compared with prostate-specific antigen. *Urology* 2000;55:481–5.
- Clark LC, Combs GF Jr, Turnbull BW, et al. Effects of selenium supplementation for cancer prevention in patients with carcinoma of the skin. A randomized controlled trial. Nutritional Prevention of Cancer Study Group. *JAMA* 1996;276:1957–63.
- Duffield-Lillico AJ, Reid ME, Turnbull BW, et al. Baseline characteristics and the effect of selenium supplementation on cancer incidence in a randomized clinical trial: a summary report of the Nutritional Prevention of Cancer Trial. *Cancer Epidemiol Biomarkers Prev* 2002;11:630–9.
- Dong Y, Lee SO, Zhang H, et al. Prostate specific antigen expression is down-regulated by selenium through disruption of androgen receptor signaling. *Cancer Res* 2004;64:19–22.
- Cho SD, Jiang C, Malewicz B, et al. Methyl selenium metabolites decrease prostate specific antigen expression by inducing protein degradation and suppressing androgen-stimulated transcription. *Mol Cancer Ther* 2004;3:605–12.
- Dong Y, Zhang H, Hawthorn L, Ganther HE, Ip C. Delineation of the molecular basis for selenium-induced growth arrest in human prostate cancer cells by oligonucleotide array. *Cancer Res* 2003;63:52–9.
- Jiang C, Wang Z, Ganther H, Lu J. Caspases as key executors of methyl selenium-induced apoptosis (anoikis) of DU-145 prostate cancer cells. *Cancer Res* 2001;61:3062–70.
- Jiang C, Wang Z, Ganther H, Lu J. Distinct effects of methylseleninic acid versus selenite on apoptosis, cell cycle, and protein kinase pathways in DU145 human prostate cancer cells. *Mol Cancer Ther* 2002;1:1059–66.
- Wang Z, Jiang C, Lu J. Induction of caspase-mediated apoptosis and cell-cycle G₁ arrest by selenium metabolite methylselenol. *Mol Carcinog* 2002;34:113–20.
- Zu K, Ip C. Synergy between selenium and vitamin E in apoptosis induction is associated with activation of distinctive initiator caspases in human prostate cancer cells. *Cancer Res* 2003;63:6988–95.
- Neckers L, Ivy SP. Heat shock protein 90. *Curr Opin Oncol* 2003;15:419–24.
- Georget V, Terouanne N, Nicolas J-C, Sultan C. Mechanism of antiandrogen action: key role of Hsp90 in conformational change and transcriptional activity of the androgen receptor. *Biochemistry* 2002;41:11824–31.
- Heinlein CA, Chang C. Androgen receptor (AR) coregulators: an overview. *Endocr Rev* 2002;23:175–200.
- Dong Y, Zhang H, Hawthorn L, Ganther HE, Ip C. Delineation of the molecular basis for selenium-induced growth arrest in human prostate cancer cells by oligonucleotide array. *Cancer Res* 2003;63:52–9.
- Zhang H, Dong Y, Zhao H, et al. Microarray data mining for potential selenium targets in chemoprevention of prostate cancer. *Cancer Genomics and Proteomics* 2005;2:97–114.
- Ip C, Thompson HJ, Zhu Z, Ganther HE. *In vitro* and *in vivo* studies of methylseleninic acid: evidence that a monomethylated selenium metabolite is critical for cancer chemoprevention. *Cancer Res* 2000;60:2882–6.
- Yeh S, Chang C. Cloning and characterization of a specific co-activator, ARA70, for the androgen receptor in human prostate cells. *Proc Natl Acad Sci U S A* 1996;93:5517–21.
- Dong Y, Lee SO, Zhang H, et al. Prostate specific antigen (PSA) expression is down-regulated by selenium through disruption of androgen receptor signaling. *Cancer Res* 2004;64:19–22.
- Livak LJ, Schmittgen TD. Analysis of relative gene expression data using real-time quantitative PCR and the 2^{-ΔΔCT} method. *Methods* 2001;25:402–8.
- Klein KA, Reiter RE, Redula J, et al. Progression of metastatic human prostate cancer to androgen independence in immunodeficient SCID mice. *Nat Med* 1997;3:402–8.
- Igawa T, Lin FF, Lee MS, et al. Establishment and characterization of androgen-independent human prostate cancer LNCaP cell model. *Prostate* 2002;50:222–35.
- Nakashiro K, Hara S, Shinohara Y, et al. Phenotypic switch from paracrine to autocrine role of hepatocyte growth factor in an androgen-independent human prostatic carcinoma cell line, CWR22R. *Am J Pathol* 2004;165:533–40.
- Pettaway CA, Pathak S, Greene G, et al. Selection of highly metastatic variants of different human prostatic carcinomas using orthotopic implantation in nude mice. *Clin Cancer Res* 1996;2:1627–36.
- Veldscholte J, Berrevoets CA, Ris-Stalpers C, et al. The androgen receptor in LNCaP cells contains a mutation in the ligand binding domain which affects steroid binding characteristics and response to antiandrogens. *J Steroid Biochem Mol Biol* 1992;41:665–9.
- Koivisto P, Kolmer M, Visakorpi T, Kallioniemi OP. Androgen receptor gene and hormonal therapy failure of prostate cancer. *Am J Pathol* 1998;152:1–9.
- Hobisch A, Culig Z, Radmayr C, et al. Androgen receptor status of lymph node metastases from prostate cancer. *Prostate* 1996;28:129–35.
- Sadi MV, Walsh PC, Barrack ER. Immunohistochemical study of androgen receptors in metastatic prostate cancer-comparison of receptor content and response to hormonal therapy. *Cancer* 1991;67:3057–64.
- Zhao XY, Malloy PJ, Krishnan AV, et al. Glucocorticoids can promote androgen-independent growth of prostate cancer cells through a mutated androgen receptor. *Nat Med* 2000;6:703–6.
- Koivisto P, Kononen J, Palmberg C, et al. Androgen receptor gene amplification: a possible molecular mechanism for androgen deprivation therapy failure in prostate cancer. *Cancer Res* 1997;57:314–9.

33. Kokontis J, Takakura K, Hay N, Liao S. Increased androgen receptor activity and altered c-myc expression in prostate cancer cells after long-term androgen deprivation. *Cancer Res* 1994;54:1566–73.
34. Visakorpi T, Hyytinen E, Koivisto P, et al. *In vivo* amplification of the androgen receptor gene and progression of human prostate cancer. *Nat Genet* 1995;9:401–6.
35. Chen CD, Welsbie DS, Tran C, et al. Molecular determinants of resistance to antiandrogen therapy. *Nat Med* 2004;10:33–9.
36. Eder IE, Culig Z, Ramoner R, et al. Inhibition of LncAP prostate cancer cells by means of androgen receptor antisense oligonucleotides. *Cancer Gene Ther* 2000;7:997–1007.
37. Eder IE, Hoffmann J, Rogatsch H, et al. Inhibition of LNCaP prostate tumor growth *in vivo* by an antisense oligonucleotide directed against the human androgen receptor. *Cancer Gene Ther* 2002;9:117–25.
38. Liao X, Tang S, Thrasher JB, Griebeling TL, Li B. Small-interfering RNA-induced androgen receptor silencing leads to apoptotic cell death in prostate cancer. *Mol Cancer Ther* 2005;4:505–15.
39. Zegarra-Moro OL, Schmidt LJ, Huang H, Tindall DJ. Disruption of androgen receptor function inhibits proliferation of androgen-refractory prostate cancer cells. *Cancer Res* 2002;62:1008–13.
40. Hecht SS. Metabolic activation and detoxification of tobacco-specific nitrosoamines: a model for cancer prevention strategies. *Drug Metab Rev* 1994;26:373–90.
41. Magklara A, Scorilas A, Catalona WJ, Diamandis EP. The combination of human glandular kallikrein and free prostate-specific antigen (PSA) enhances discrimination between prostate cancer and benign prostatic hyperplasia in patients with moderately increased total PSA. *Clin Chem* 1999;45:1960–6.
42. Nam RK, Diamandis EP, Toi A, et al. Serum human glandular kallikrein-2 protease levels predict the presence of prostate cancer among men with elevated prostate-specific antigen. *J Clin Oncol* 2000;18:1036–42.
43. Partin AW, Catalona WJ, Finlay JA, et al. Use of human glandular kallikrein 2 for the detection of prostate cancer: preliminary analysis. *Urology* 1999;54:839–45.
44. Recker F, Kwiatkowski MK, Piironen T, et al. The importance of human glandular kallikrein and its correlation with different prostate specific antigen serum forms in the detection of prostate carcinoma. *Cancer* 1998;83:2540–7.
45. Norris MD, Smith J, Tanabe K, et al. Expression of multidrug transporter MRP4/ABCC4 is a marker of poor prognosis in neuroblastoma and confers resistance to irinotecan *in vitro*. *Mol Cancer Ther* 2005;4:547–53.
46. Cao S, Durrani FA, Rustum YM. Selective modulation of the therapeutic efficacy of anticancer drugs by selenium containing compounds against human tumor xenografts. *Clin Cancer Res* 2004;10:2561–9.
47. Greeve I, Hermans-Borgmeyer I, Brellinger C, et al. The human DIMINUTO/DWARF1 homolog seladin-1 confers resistance to Alzheimer's disease-associated neurodegeneration and oxidative stress. *J Neurosci* 2000;20:7345–52.
48. Sarkar D, Imai T, Kambe F, et al. The human homolog of Diminuto/Dwarf1 gene (hDiminuto): a novel ACTH-responsive gene overexpressed in benign cortisol-producing adrenocortical adenomas. *J Clin Endocrinol Metab* 2001;86:5130–7.
49. Zabel U, Weeger M, La M, Schmidt HH. Human soluble guanylate cyclase: functional expression and revised isoenzyme family. *Biochem J* 1998;335:51–7.
50. Cho SD, Jiang C, Malewicz B, et al. Methyl selenium metabolites decrease prostate-specific antigen expression by inducing protein degradation and suppressing androgen-stimulated transcription. *Mol Cancer Ther* 2004;3:605–11.