AD _____

Award Number: W81XWH-09-1-0437

TITLE: Treatment of Traumatic Brain Injury by Localized Application of Sub-atmospheric Pressure to the Site of Cortical Impact

PRINCIPAL INVESTIGATOR: Michael Morykwas, Ph.D.

CONTRACTING ORGANIZATION: Wake Forest University Health Sciences
Winston-Salem, NC 27157

REPORT DATE: July 201G

TYPE OF REPORT: Annual

PREPARED FOR: U.S. Army Medical Research and Materiel Command
Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Approved for Public Release;
Distribution Unlimited

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

REPORT DOCUMENTATION PAGE			Form Approved OMB No. 0704-0188		
Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number. PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.					
1. REPORT DATE July 201G		2. REPORT TYPE Annual		3. DATES COVERED 1 July 201F – 30 June 201G	
4. TITLE AND SUBTITLE Treatment of Traumatic Brain Injury by Localilzed Application of Sub-atmospheric Pressure to the Site of Cortical Impact				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER W81XWH-09-1-0437	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S) Michael Morykwas, Ph.D. E-Mail: "o o qt {ny cB y cngj gcmj Qf w				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Wake Forest University Health Sciences Winston-Salem, NC 27157				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) U.S. Army Medical Research and Materiel Command Fort Detrick, Maryland 21702-5012				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION / AVAILABILITY STATEMENT Approved for Public Release; Distribution Unlimited					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT Gcej "o clqt"y ct"vopf u"q"j cxg"e"-uki pcwtg"lplwt {ø"y kj "tewo cve"dtclp"lplwt { "VDK"cuuqekcvgf "y kj "y g"Kcs"y ct " "Qr gtcvqp"KcsKHggf go "KKcpf "Qr gtcvqp"Gpf wt lpi "Hggf go +f wg"q"y g"j ki j "lpekf gpeg"qh'r gtupppgnlplwtgf "d { "KGF " "to r tqxkugf "gzmqukxg"fxlegu+0"Vj g"ncti g"i { tgeprj cke"dtclp"qh'uy lpg"ku'uko kret"q"j wo cpu."y wu"e"uy lpg"o qf gn'qh"e" eqpvtqmgf "eqtveclko r cev"EEKlplwt { +tgcvgf "d { "y g"eqpvtqmgf "cr r rdecvqp"qh'uw d/cvo qur j gtke"r tguuwtg"y cu"go r mq { gf 0" Y qtnleqo r rvgf "lp"l gct"3"uj qy gf "y cv322"o o "J i "xcewwo "cr r rdecvqp"q"y g"ukg"qh"e"lqecndtclp"lplwt { "f getgcugf "y g" xqnwo g"qh"j go qttj ci g"cpf "lpwtgf "ctgc0"Y qtnleqo r rvgf "lp"l gct"4"uj qy gf "y cv322"o o "J i "xcewwo "pggf gf "q"dg"cr r rkgf " q"y g"ukg"qh"y g"lqecndlplwt { "hqt"e"o lolo wo "qh"7"f c { u"q"r tggpvrvg"lpetgcugu"lp"lvtc"etcpkcnr tguuwtg"cpf "tguwo r vqp"qp" y g"lplwt { "r tqegu0"lp"l gct"5."y qtnleqo r rvgf "uj qy gf "y cv"y cv"5"j qwt"f gnc { "dgy ggp"lplwt { "cpf "lpklcvqp"qh"xcewwo " vtgcvo gpv"y cu"cu"ghhecekqwu"cu"lo o gf lcvg"vtgcvo gpv."cpf "y cv"e"8"j qwt"f gnc { "dgy ggp"lplwt { "cpf "lpklcvqp"qh"xcewwo " vtgcvo gpv"y cu"uwn"ghhecekqwu0"Vj ku"cmqy u"ht"e"tgcruk"e"y qtnlpi "y lpf qy "hqt"uweeguuhw"cr r rdecvqp"qh'eqpvtqmgf ." qecrk gf "xcewwo "q"y g"ukg"qh"e"lqecndVKD					
15. SUBJECT TERMS Vtewo cve"dtclp"lplwt { ".uw d/cvo qur j gtke"r tguuwtg"vtgcvo gpv"xcewwo ."o gej cplecn"kuuwtg"tguuuekcvqp					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT	18. NUMBER OF PAGES	19a. NAME OF RESPONSIBLE PERSON USAMRMC
a. REPORT U	b. ABSTRACT U	c. THIS PAGE U			19b. TELEPHONE NUMBER (include area code)
			UU	11	

Table of Contents

	<u>Page</u>
Introduction.....	5
Body.....	5
Key Research Accomplishments.....	11
Reportable Outcomes.....	11
Conclusion.....	11
References.....	12
Appendices.....	12

Introduction

Each major war tends to have a ‘signature injury’, with traumatic brain injury (TBI) associated with the Iraq war (Operation Iraqi Freedom II and Operation Enduring Freedom) due to the high incidence of personnel injured by IED (improvised explosive devices). Our previous study in a rat model showed that Mechanical Tissue Resuscitation (MTR – the controlled application of vacuum) to the cerebral cortex following a controlled cortical impact (CCI) injury reduces brain edema and the extent of injury, modulates metabolites in injured neuronal tissues, preserves neuronal tissue, and improves functional recovery. The large gyrencephalic brain of swine is similar to humans, thus a swine model of CCI injury and MTR treatment was evaluated for future human clinical applications. Year 3 goals were to determine the length of time between the initial injury event and efficacious application of vacuum.

Body

Year 3 goals stated in the Statement of Work included determination of the length of time between creation of a focal TBI and then efficacious application of vacuum application to the site of the focal injury. Injuries were created and animals were treated with 100 mm Hg vacuum immediately (based on Year 1 and 2 studies), or after a 3 or 6 hour delay.

During Year 3 the MRI scanner used for imaging during Years 1 and 2 was removed and replaced with a new, more powerful scanner. This resulted in a hiatus in the project due to the inability to scan.

A total of thirty two (32) female domestic swine (22-33 kg) were procured (including those treated immediately (0 hour delay) in previous years). For MTR treatment, a sterile vacuum dressing was placed in the bony defect and 100 mm Hg was applied continuously either immediately post injury, or after either a 3 or 6 hour delay. Vacuum was applied for a total of 5 days. Intracranial pressure was monitored by telemetry up until the time of MR imaging. Five days post surgery, all animals were analyzed by MRI (GE Signa EchoSpeed 1.5-T scanner). Parameters analyzed included: MR sagittal T1 imaging; coronal T2 imaging; coronal MPGR (Multi-Planer Gradient Echo); Axial T2* Contrast Enhanced Perfusion (0.2 ml/Kg Magnevist contrast by power injection). All animals were euthanized 10 days post injury and perfused with 4% para-formaldehyde through the ascending aorta 8 days post-injury. The brain was removed and postfixed in the same fixative solution overnight at 4°C. After a PBS rinse, the brains were placed in 30% sucrose at 4°C before they were snap-frozen in O.C.T and stored at -80°C. Coronal sections of the injured area were cut into 20 µm thick sections using a cryostat, mounted, and kept frozen until use. Sections were collected every 0.5 mm through all injured area over a total distance of 2 cm. Histological staining and analysis is being completed.

Figure 1. Representative histologic cross sections of brains for control (non-treated animals), immediate treatment, a 3 hour delay between injury creation and vacuum application, or a 6 hour delay between injury creation and vacuum application. The large area of hemorrhage and necrosis can be seen in the non-treated brain. Much smaller areas are present in the immediate (0 hour delay) and the 3 hour delay animal. A small area of hemorrhage and damage is present in the 6 hour delay brain.

Figure 2. MRI T2 analysis of hyperintense areas of the image of the injured areas showed areas of increased water content (swelling), with largest volume in the non-treated animals. The volume of the area of injury for animals treated immediately was not significantly ($p < 0.05$) different that the volume for animals treated after a 3 hour delay before vacuum initiation. The volume of the injured area for animals treated following a 6 hour delay between injury and vacuum application was not significantly smaller than the non-treated animals. The volume for the 6 hour delay animals was also not significantly larger than the areas of injury for the 0 hour or 3 hour delay animals.

Figure 3. Gradient echo MR imaging was used to measure the volume of the hemorrhage in the animals. All treated animals had a significantly smaller volume of hemorrhage than the non-treated animals. The volume of the hemorrhage for the 0 hour delay and the 6 hour delay were similar, with the same degree of significance ($p<0.05$). The volume for the 3 hour delay was very significantly smaller ($p<0.01$) than the non-treated volume. The volumes of hemorrhage for the treated animals were not significantly different from each other.

Total Intracranial hemorrhage volume measured in gradient echo MRI images in traumatic brain injured pigs.

Pigs were treated under negative pressure 100 mmHg for 5 days with different application start times.

* Indicates $P<0.05$; ** Indicates $P<0.01$

Figure 4. Histologic cross sections were imaged and the volume of the area of necrosis was determined. The volume of the area of necrosis for all treated animals was significantly ($p<0.05$) smaller than the area of necrosis for non-treated animals. The volume of area of necrosis for animals in the 3 hour and 6 hour delay groups was very significantly ($p<0.01$) than the area for the non-treated animals. The area of necrosis between the 0 hour delay and the 3 or 6 hour delay was not significantly different.

Total necrotic volume in series of histological sections of traumatic brain injured pigs. Pigs were treated under negative pressure 100 mmHg for 5 days with different application start times.

* indicates $P<0.05$; ** indicates $P<0.01$

Table 1. Quantitative Table with data and sample sizes for the data presented graphically above. The sample size for histologic determination of the necrotic brain volume for the 0 hour delay group was 7 animals. All other groups contained 8 animals.

Damaged Volumes (mm ³)	Injured Only Mean±SD	No Delay Treated Mean±SD	3-hrs Delay Treated Mean±SD	6-hrs Delay Treated Mean±SD
Injured Brain Tissue volume (T2 MR)	7900.72±2248.78 N=8	5621.47±1435.46 N=8	5004.30±2202.92 N=8	6919.37±2038.23 N=8
Intracranial Hemorrhage Volume (GE MRI)	553.18±283.40 N=8	232.69±194.47 N=8	102.94±154.47 N=8	175.59±229.87 N=8
Brain Necrotic Volume (H&E Staining)	985.81±395.64 N=8	388.21±486.26 N=7	414.91±341.93 N=8	338.21±195.69 N=8

Mean damaged brain volume measurements in MR images and H&E staining in traumatic brain injured pigs. Pigs were treated under negative pressure 100 mmHg for 5 days with different application start times.

Trends apparent from the MRI data show that a 3 hour delay provides for a better result than animals treated after a 6 hour delay in vacuum application. The area of injury as determined from the T2 imaging analysis exhibited a significantly smaller area of injury in animals treated after a 3 hour delay compared to animals treated after a 6 hour delay. Animals treated after a 6 hour delay still exhibited a smaller area of injury and a smaller hemorrhage volume compared to non-treated animals.

Histologic analysis of the area of necrosis shows that animals treated after the 6 hour delay had a non-statistically significant smaller area of necrosis than animals treated after a 3 hour delay. All treated animals had a statistically smaller area of necrosis than non-treated animals.

The data shows that animals treated with 100 mm Hg vacuum applied continuously for 5 days and after a 3 hour delay between injury and vacuum application exhibited as good of a result as those animals treated immediately post injury. Animals treated after a 6 hour delay between injury and vacuum application still exhibited a favorable result, although it may not have been as efficacious as the 3 hour delay group.

Key Research Accomplishments

- Determination that application of 100 mm Hg sub-atmospheric pressure for 5 days to the site of cortical injury following a 3 or 6 hour delay between injury and application of the vacuum was still efficacious. Animals in the 3 hour delay group showed a trend for a more favorable result compared to animals in the 6 hour delay group.

Reportable Outcomes

Presentation:

Zheng Z, Argenta L, Morykwas M. Mechanical Tissue Resuscitation treatment reduces injured brain tissue swellings and intracerebral hemorrhages in a pig traumatic brain injury model (delay treatment). (Abstract) Congress of Neurological Surgeons 2012 Annual Meeting, October 4 - 10, 2012

Accepted for publication (not yet published):

Johnston ME, Zheng Z, Maldjian JA, Whitlow CT, Morykwas MJ, Jung Y: Cerebral Blood Flow Quantification in Swine using Pseudo-Continuous Arterial Spin Labeling, Journal of Magnetic Resonance Imaging.

Conclusion

This portion of the study demonstrates that the use of mechanical tissue resuscitation (MTR) treatment of a continuous vacuum of 100 mm Hg for 5 days was as efficacious after a 3 or 6 hour delay between injury creation as compared to animals treated immediately after injury creation (0 hour delay).

The expectation of rapid translation of the technique to humans is still anticipated with interest from industry for commercialization of the product and technique, although no formal licensing discussions have begun.

References

N.A.

Appendices

N.A.