

UNCLASSIFIED

AD 91694

Armed Services Technical Information Agency

Reproduced by

DOCUMENT SERVICE CENTER

KNOTT BUILDING, DAYTON, 2, OHIO

This document is the property of the United States Government. It is furnished for the duration of the contract and shall be returned when no longer required, or upon recall by ASTIA to the following address: Armed Services Technical Information Agency, Document Service Center, Knott Building, Dayton 2, Ohio.

NOTICE: WHEN GOVERNMENT OR OTHER DRAWINGS, SPECIFICATIONS OR OTHER DATA ARE USED FOR ANY PURPOSE OTHER THAN IN CONNECTION WITH A DEFINITELY RELATED GOVERNMENT PROCUREMENT OPERATION, THE U. S. GOVERNMENT THEREBY INCURS NO RESPONSIBILITY, NOR ANY OBLIGATION WHATSOEVER; AND THE FACT THAT THE GOVERNMENT MAY HAVE FORMULATED, FURNISHED, OR IN ANY WAY SUPPLIED THE SAID DRAWINGS, SPECIFICATIONS, OR OTHER DATA IS NOT TO BE REGARDED BY IMPLICATION OR OTHERWISE AS IN ANY MANNER LICENSING THE HOLDER OR ANY OTHER PERSON OR CORPORATION, OR CONVEYING ANY RIGHTS OR PERMISSION TO MANUFACTURE, USE OR SELL ANY PATENTED INVENTION THAT MAY IN ANY WAY BE RELATED THERETO.

UNCLASSIFIED

12 March 1956

COSMIC RAY INTENSITY AT HIGH ALTITUDES

on February 23, 1956*

By

J. A. Van Allen and C. E. McIlwain

Department of Physics
State University of Iowa
Iowa City, Iowa

*Assisted by National Science Foundation and by joint
program of Office of Naval Research and Atomic Energy
Commission.

A balloon flight of a single Geiger tube was made¹ from Iowa City ($\lambda = 52^\circ\text{N}$) during the period

1. The authors are indebted to Dr. J. A. Simpson for informing us of the fact that a great cosmic ray storm was in progress. The onset of the storm occurred at 03^h49^m G.M.T. February 23.

19^h33^m to 21^h12^m Greenwich Mean Time on February 23, 1956. The apparatus consisted of a single Victoreen 1B85 Geiger tube of calibrated effective length and a radio transmitter, all housed in a pressure-tight aluminum shell of wall thickness 0.032". Identical apparatus has been flown by this laboratory on previous occasions by means of balloons and rockets. The axis of the tube was nearly vertical. The geomagnetic latitude of equipment while at the highest altitude was estimated to be 53°N .

The raw counting rate of the Geiger tube during the ascent of the balloon is shown as a function of time in Figure 1. The measured effective length of the tube was 6.8 ± 0.2 cm. and the diameter was 1.90 ± 0.02 cm. The maximum observed value of the counting rate was 53.0 ± 0.3 counts/second during the period 20^h55^m to 21^h08^m G.M.T.

Correcting for dead time and intrinsic efficiency ($\epsilon = 0.98$) and dividing by the appropriate geometric factor of the counter² we find

2. E. C. Ray, M. S. Thesis, State University of Iowa
(June 1953).

$$(1) \quad \bar{J} = 0.76 \pm 0.03 \text{ (cm}^2 \text{ sec sterad)}^{-1}$$

for the angular average value, over the upper hemisphere, of the unidirectional intensity of all charged particles, the average being taken in the usual manner.³ An equivalent

3. L. H. Meredith, J. A. Van Allen, and M. B. Gottlieb,
Phys. Rev. 99, 198 (1955).

statement of the result is

$$(2) \quad J_0 = 4.8 \pm 0.2 \text{ (cm}^2 \text{ sec)}^{-1}$$

where J_0 is the flux through a sphere of unit cross-section.

During the period 1950-1955, this laboratory has made a large number of balloon and rocket flights of single Geiger tubes and Geiger tube telescopes at geomagnetic latitudes ranging from 41°N to 89°N. A summary of pertinent

on an atmospheric pressure basis assuming ^{various} constant rates of ascent in m/sec, an assumption concerning balloon performance which is well grounded in our experience with flights of clusters of rubber balloons of the type used. The results of this process are shown in Figure 2.

From this plot it appears likely that the actual rate of rise was about 4 m/sec and that the previously quoted intensities (1) and (2) likely refer to the vicinity of the Pfozzer-Regener maximum.

We are indebted particularly to Drs. F. B. McDonald and K. A. Anderson and to Messrs. G. Ludwig and W. R. Webber for assistance.

Figure Captions

Figure 1 Observed counting rate of single Geiger tube as a function of time during the ascent of the balloon. Statistical standard errors are shown.

Figure 2 Replots of observed counting rate data of Figure 1 (assuming various constant rates of rise of balloon) showing comparison with normal situation at $\lambda = 53^\circ$ (solid curve).




FIG. 1

COUNTS PER SECOND




FIG. 2