

UNCLASSIFIED

AD NUMBER

AD313998

CLASSIFICATION CHANGES

TO: unclassified

FROM: confidential

LIMITATION CHANGES

TO:
Approved for public release, distribution unlimited

FROM:
Distribution authorized to U.S. Gov't. agencies and their contractors; Administrative/Operational Use; DEC 1959. Other requests shall be referred to Redstone Arsenal, Huntsville, AL.

AUTHORITY

31 Dec 1971, DoDD 5200.10; RSIC D/A ltr, 16 Nov 1972

THIS PAGE IS UNCLASSIFIED

AD

313 998

Reproduced by

Armed Services Technical Information Agency

ARLINGTON HALL STATION; ARLINGTON 12 VIRGINIA

NOTICE: WHEN GOVERNMENT OR OTHER DRAWINGS, SPECIFICATIONS OR OTHER DATA ARE USED FOR ANY PURPOSE OTHER THAN IN CONNECTION WITH A DEFINITELY RELATED GOVERNMENT PROCUREMENT OPERATION, THE U. S. GOVERNMENT THEREBY INCURS NO RESPONSIBILITY, NOR ANY OBLIGATION WHATSOEVER; AND THE FACT THAT THE GOVERNMENT MAY HAVE FORMULATED, FURNISHED, OR IN ANY WAY SUPPLIED THE SAID DRAWINGS, SPECIFICATIONS, OR OTHER DATA IS NOT TO BE REGARDED BY IMPLICATION OR OTHERWISE AS IN ANY MANNER LICENSING THE HOLDER OR ANY OTHER PERSON OR CORPORATION, OR CONVEYING ANY RIGHTS OR PERMISSION TO MANUFACTURE, USE OR SELL ANY PATENTED INVENTION THAT MAY IN ANY WAY BE RELATED THERETO.

Best Available Copy

CONFIDENTIAL

S-22 671

Copy No. _____

Dec. 15, 1959

ROHM & HAAS COMPANY 10

REDSTONE ARSENAL RESEARCH DIVISION
HUNTSVILLE, ALABAMA




AD No. 313 998

ASTIA FILE COPY

TRAFLUOROHYDRAZINE

FC



A NEW HIGH ENERGY LIQUID

OXIDIZER

FILE COPY
Returned to
ASTIA
ARLINGTON HALL STATION
ARLINGTON 12, VIRGINIA
ASTIA 11543

ASTIA
 RECEIVED
 DEC 21 1959
 RECEIVED
 TIPDR A

This document contains information affecting the national defense of the United States within the meaning of the Espionage Laws, Title 18, U. S. C., Sections 793 and 794. The transmission or the revelation of its contents in any manner to an unauthorized person is prohibited by law.

CONFIDENTIAL

Best Available Copy

**Best
Available
Copy**

C O N F I D E N T I A L

ROHM & HAAS COMPANY

REDSTONE ARSENAL RESEARCH DIVISION
HUNTSVILLE, ALABAMA

REPORT NO. 3.22

TETRAFLUOROHYDRAZINE
(A new, high energy, liquid oxidizer)


By

CHARLES B. COLBURN

and

KENNETH A. WILDE

Approved:



Robert W. Walker
Head, Physical & Polymer
Chemistry Section



Allen R. Deschere
General Manager

December 15, 1959

ARMY ORDNANCE CORPS
Project Number TU2-10
RESEARCH ON ROCKET PROPELLANTS AND ROCKET MOTORS
Contract No. DA 01-021-ORDR15

Best Available Copy C O N F I D E N T I A L

C O N F I D E N T I A L

ROHM & HAAS COMPANY

REDSTONE ARSENAL RESEARCH DIVISION
HUNTSVILLE, ALABAMA

ABSTRACT

✓
Tetrafluorohydrazine gives the following theoretical equilibrium impulses at 1000 psi → 1 atmos. with the following fuels:

H₂ - 357 sec.; N₂H₄ - 333; NH₃ - 325; UMDH - 316; Hidyne - 314;
JP-4 -297.

The physical and chemical properties of tetrafluorohydrazine indicate that it would be much easier to handle as a liquid oxidizer than are oxygen and fluorine.

The preparation, chemical and physical properties of tetrafluorohydrazine are briefly reviewed.

↑

C O N F I D E N T I A L

CONFIDENTIAL

TETRAFLUOROHYDRAZINE (A new, high energy, liquid oxidizer)

The new heat of formation of tetrafluorohydrazine ($\Delta H_{25}^{\circ}[\text{N}_2\text{F}_4] = -2.0 \pm 2.5 \text{ kcal./mole}$)¹ which has been determined at the National Bureau of Standards indicates that tetrafluorohydrazine is a much more powerful oxidizer than previously believed. The calculated specific impulses (Table I) would indicate that tetrafluorohydrazine lies between fluorine and oxygen in its oxidative power. However, its physical and chemical properties are much superior to those of fluorine and oxygen (Table II).

It is immediately apparent from these tables that tetrafluorohydrazine exceeds liquid oxygen with all fuels listed (except hydrogen) and approaches the performance of fluorine as a high energy oxidizer. Because of its properties outlined in Table II it is much more readily handled than are fluorine and oxygen. Because of its low freezing point it would be easier to work with than is N_2O_4 .

Preparation

The preparation of tetrafluorohydrazine was first reported by this Division². Subsequent reports in this series have covered the development of satisfactory methods of preparation of tetrafluorohydrazine.

Tetrafluorohydrazine is prepared by the reaction at elevated temperatures of nitrogen trifluoride with various fluorine acceptors such as copper, arsenic, bismuth, antimony, glass^{2,3} and carbon⁴. The

¹ G. T. Armstrong, S. Marantz and C. F. Coyle NBS Report No. 6584 October, 1959.

² Rohm & Haas Company Quarterly Progress Report on Physical Chemistry, No. P-57-4, March, 1957.

³ C. B. Colburn, A. Kennedy JACS 80 5004 (1958).

⁴ Stauffer Chemical Company Report No. SCC-26-QFR-6 December, 1958.

CONFIDENTIAL

CONFIDENTIAL

-2-

Table I

Specific Impulse¹ of Various Fuel Oxidizer Mixtures

Fuel	Oxidizer					
	N ₂ F ₄		N ₂ F ₄ ² N ₂ O ₄	O ₂	F ₂	N ₂ O ₄
Hydrogen	Eq. ³	357		391 ⁵	410 ⁵	
	Froz. ⁴	326		388 ⁵	398 ⁵	
Hydrazine	Eq.	333		313 ⁵	363 ⁵	292 ⁵
	Froz.	311		301 ⁵	334 ⁵	283 ⁵
Ammonia	Eq.	325		294 ⁵	357 ⁵	
	Froz.	306		285 ⁵	330 ⁵	
UDMH	Eq.	316		310 ⁵		285 ⁵
	Froz.	286		295 ⁵		274 ⁵
Hidyne ⁶	Eq.	314		306 ⁵		282 ⁵
	Froz.	285		291 ⁵		271 ⁵
JP-4 (CH _{1.93})	Eq.	297	314	300 ^{5,7}		276 ^{5,7}
	Froz.	277	285	286 ^{5,7}		263 ^{5,7}

AD - 313 996

¹ Calculated using ideal assumptions, expansion from 1000 psi to 1 atmos. See Rohm & Haas Company Quarterly Progress Report on Physical Chemistry, No. P-59-16, September 1959 for method of calculation.

² For carbon containing fuels it is interesting to note that a mixture of N₂F₄ and N₂O₄ will give a higher specific impulse than either constituent alone in much the same fashion as do mixtures of F₂ and O₂.

³ Equilibrium flow.

⁴ Frozen flow.

⁵ These values taken from Rocketdyne Publication 505-X (Revised 1 January, 1959) "Rocket Engine Propellants".

⁶ 60 wt. % UDMH 40% diethylene triamine.

⁷ These numbers are for RP-1 which has essentially the same composition as JP-4.

CONFIDENTIAL

-3-

Table II

Physical Properties of N₂F₄, F₂, O₂, N₂O₄


	<u>N₂F₄</u>	<u>O₂³</u>	<u>F₂³</u>	<u>N₂O₄³</u>
Boiling Point °C.	-73 ¹	-183	-187	21.3
Melting Point °C.	-163 (approx.)	-218.4	-223	-9.3
Critical Temperature °C.	36 ¹			
Critical Pressure Atmos.	77 ¹			
Density gr/cc.	1.5 ² at -100°C.	1.14 ⁻¹⁸³	1.108 ⁻¹⁸⁷	1.491 ⁰

¹ C. B. Colburn, A. Kennedy JACS 80 5004 (1958).

² Private communication Charles Mader to W. D. Niederhauser, 15 November, 1957.

³ These values are handbook values.

reaction can be represented as follows:



The loss of hydrogen by difluoroamine¹ on contact with various solids has also been reported to give tetrafluorohydrazine². Electrical discharge through nitrogen trifluoride has also been reported as yielding tetrafluorohydrazine and difluorodiazine³.

By far the most efficient method for the preparation of tetrafluorohydrazine is the reaction of nitrogen trifluoride with carbon⁴. This process has been perfected⁵ and scaled up to a small pilot plant operation⁶. The yield of tetrafluorohydrazine from the reaction of nitrogen trifluoride with carbon is 77% at 440°C. with a residence time

¹ A. Kennedy, C. B. Colburn, JACS 81 2906 (1959).

² E. A. Lawton & J. Q. Weber JACS 81 4755 (1959).

³ J. W. Frazer, I. Inorg. Z Nuclear Chem. 11 166-167 (1959).

⁴ Stauffer Chemical Company Report No. SCC-26-QPR-6 December, 1958.

⁵ Rohm & Haas Company Quarterly Progress Report on Physical Chemistry, No. P-59-9, June, 1959.

⁶ Ibid., Quarterly Progress Report on Solid Propellant Processing, No. P-59-8 June, 1959.

CONFIDENTIAL

of approximately six seconds. The yield is between 62-71% in the reaction between nitrogen trifluoride and copper at 375°C. and with a residence time of thirteen minutes. These reactions take place in flow systems and hence the process can be made continuous.

Summary of Chemical and Physical Properties of Tetrafluorohydrazine

Heat of formation	= -2.0 + 2.5 kcal. /mole
Boiling point	- 73°C.
Freezing point	-162°C. (approximately)
Critical temperature	36°C.
Critical pressure	77 atmospheres (estimated)
Vapor pressure equation	
	$\log P_{(\text{mm})} = -692/T + 6.33$
Density	1.5 gm. /cc.
Heat of vaporization	3170 cal. /mole

Mass Cracking Pattern of N₂F₄

(Obtained on a Consolidated Model 620 Mass Spectrometer)

m/e	Ion	Pattern
52	NF ₂ ⁺	90.6%
33	NF ⁺	100.0%
28	N ₂ ⁺	7.7%
19	F ⁺	4.7%
14	N [±]	8.5%

Sensitivity div. /m 19.0

Instrument sensitivity 100 div. /m for m/e 43 for n butane.

Infrared Spectrum of Tetrafluorohydrazine

(See Fig. 1 Rohm & Haas Company QPR Physical Chemistry Report, P-57-10, page 1).

CONFIDENTIAL

-5-

N. M. R. Spectrum of Tetrafluorohydrazine

The F^{19} nuclear magnetic resonance spectrum of tetrafluorohydrazine consisted of a single broad unresolved band at a field of approximately 75 p. p. m. lower than that of the F^{19} nuclei of trifluoroacetic acid. Resolution of the expected triplet was not observed.

Tetrafluorohydrazine does not react with glass, stainless steel, or copper at room temperature.

Summary

Tetrafluorohydrazine is a powerful new oxidizing agent which lies between liquid oxygen and liquid fluorine in its performance behaviour with conventional fuels. Its physical and chemical properties indicate that it would be much easier to work with in liquid motors than fluorine and oxygen. Tetrafluorohydrazine can be prepared with relative ease.

Acknowledgement

The authors wish to acknowledge the enthusiastic support and guidance of Dr. W. D. Niederhauser in this work.

Best Available Copy

CONFIDENTIAL