

AFOSR 4772

63-4.

408388

AD

AF EOAR GRANT 62 - 120

TR

30 April 1963

CATALOGED BY DDC
AS AD No.

TECHNICAL REPORT

STATISTICAL MODEL APPLIED TO THE REGION
OF THE ν_3 FUNDAMENTAL OF CO_2 AT $1200^\circ K$

U.P. OPPENHEIM

TECHNION - ISRAEL INSTITUTE OF
TECHNOLOGY
DEPARTMENT OF PHYSICS
HAIFA, ISRAEL

The research reported in this document has been sponsored by
the

OFFICE OF SCIENTIFIC RESEARCH, OAR

through the European Office, Aerospace Research, United States
Air Force

408 388

DDC
JUN 19 1963

Abstract

A new method is developed in order to correlate the observed emissivity of a statistical band with the experimental parameters of the gas (pressure, optical path, etc.). Use is made of curves of growth for every frequency in the band and the method is applied to the region of the 4.3μ bands of CO_2 at a temperature of 1200° K. Experimental results for CO_2 are described, which were obtained by heating the gas in cells of different length in an electrical furnace. Good quantitative agreement is found with experimental results of other workers. It is shown that the statistical model predicts the emissivity correctly over wide ranges of pressure and optical path.

Report

The work done under the Grant was reported in a paper which was published in the Journal of the Optical Society of America, Number 3, pp. 344 - 350 (1963), under the title:

"Statistical model applied to the region of the ν_3 fundamental of CO_2 at 1200°K ", by U.P. Oppenheim and Y. Ben-Aryeh.

No other publications resulted from the present work.

TECHNION

HAIFA, ISRAEL

DEPT PHYSICS

30 April 1963

AF EOAR 62-120

TR

PHYSICS

TECHNION

HAIFA, ISRAEL

DEPT PHYSICS

30 April 1963

AF EOAR 62-120

TR

PHYSICS

STATISTICAL MODEL APPLIED TO THE REGION OF THE

²₃ FUNDAMENTAL OF CO₂ AT 1200° K

U.P. OPPENHEIM

ABSTRACT: A new method is developed in order to correlate the observed emissivity of a statistical band with the experimental parameters of the gas. The method is applied to the region of the 4.3 μ bands of CO₂ at 1200° K.

STATISTICAL MODEL APPLIED TO THE REGION OF THE

²₃ FUNDAMENTAL OF CO₂ AT 1200° K

U.P. OPPENHEIM

ABSTRACT: A new method is developed in order to correlate the observed emissivity of a statistical band with the experimental parameters of the gas. The method is applied to the region of the 4.3 μ bands of CO₂ at 1200° K.

TECHNION

HAIFA, ISRAEL

DEPT PHYSICS

30 April 1963

AF EOAR 62-120

TR

PHYSICS

TECHNION

HAIFA, ISRAEL

DEPT PHYSICS

30 April 1963

AF EOAR 62-120

TR

PHYSICS

STATISTICAL MODEL APPLIED TO THE REGION OF THE

²₃ FUNDAMENTAL OF CO₂ AT 1200° K

U.P. OPPENHEIM

ABSTRACT: A new method is developed in order to correlate the observed emissivity of a statistical band with the experimental parameters of the gas. The method is applied to the region of the 4.3 μ bands of CO₂ at 1200° K.

STATISTICAL MODEL APPLIED TO THE REGION OF THE

²₃ FUNDAMENTAL OF CO₂ AT 1200° K

U.P. OPPENHEIM

ABSTRACT: A new method is developed in order to correlate the observed emissivity of a statistical band with the experimental parameters of the gas. The method is applied to the region of the 4.3 μ bands of CO₂ at 1200° K.