# THE UNIVERSITY OF MICHIGAN

03105-38-7

DDC

LUIL 27 195

IJOIN B

May 1965

COLLEGE OF LITERATURE, SCIENCE, AND THE ARTS DEPARTMENT OF COMMUNICATION SCIENCES

Technical Report

Transparent Categories and Categories of Transition Systems

YEHOSHAFAT GIVE'ON

# AD466445

Under Contract with: Department of the Navy Office of Naval Research Contract No. Nonr-1224(21)

Washington, D. C.

Administered through:

OFFICE OF RESEARCH ADMINISTRATION . ANN A BOR

**BEST AVAILABLE COPY** 

NOTICE: When government or other drawings, specifications or other data are used for any purpose other than in connection with a definitely related government procurement operation, the U. S. Government thereby incurs no responsibility, nor any obligation whatsoever; and the fact that the Government may have formulated, furnished, or in any way supplied the said drawings, specifications, or other data is not to be regarded by implication or otherwise as in any manner licensing the holder or any other person or corporation, or conveying any rights or permission to manufacture, use or sell any patented invention that may in any way be related thereto. THE UNIVERSITY OF MICHIGAN COLLEGE OF LITERATURE, SCIENCE, AND THE ARTS Department of Communication Sciences

1880 (A)

1

Technical Report

#### TRANSPARENT CATEGORIES AND CATEGORIES OF TRANSITION SYSTEMS

Yehoshafat Give'on

ORA Project 03105

under contract with:

DEPARTMENT OF THE NAVY OFFICE OF NAVAL RESEARCH CONTRACT NO. Nonr-1224(21) WASHINGTON, D.C.

administered through:

OFFICE OF RESEARCH ADMINISTRATION AN

ANN ARBOR

May 1965

#### RESEARCH PROGRESS REPORT

Title: "Transparent Categories and Categories of Transition Systems" Y. Give'on, University of Michigan Technical Report 03105-38-T, May 1965; Nonr 1224(21).

Background: The Logic of Computers Group of the Communication Sciences Department of The University of Michigan is investigating the application of logic and mathematics to the design of computing automata. The application of the techniques and concepts of abstract algebra to automata forms a part of this investigation.

<u>Condensed Report Contents</u>: Several recent results in automata theory give evidence of the importance of homomorphisms in the study of transition systems and automata. It is natural therefore to inquire how much information can be retrieved from the algebra of homomorphism compositions with respect to transition systems. The natural mathematical framework for the discussion of this problem is categorical algebra.

We define a category  $\mathbf{A}_{W}$  of the transition systems with input W, where W is any arbitrary fixed monoid, and with arbitrary sets of states. A preliminary study of  $\mathbf{A}_{W}$  (Give'on 1964) shows that one can reconstruct the internal structure of any transition system from the way homomorphisms (i.e., the morphisms of  $\mathbf{A}_{W}$ ) behave around it.

In this paper we show that  $\mathbf{A}_W$  has a generator,  $\mathbf{M}_W$  (which is W operating on itself as a transition system) and that there exists a functor Nor :  $\mathbf{A}_W \longrightarrow \mathbf{A}_W$  naturally equivalent to the identity functor of  $\mathbf{A}_W$  which factors through  $\operatorname{Hom}_{\mathbf{A}_W}(\mathbf{M}_W)$ .

A general exposition of the nature of properties which are retrievable from the "morphism-behavior" in an arbitrary category is presented so that it provides a rigorous general basis for studying "retrievable" properties and categories in which every structural property of objects and morphisms is "retrievable." Finally, we prove that for a very broad class of input monoids, which includes all the types of input-monoids encountered in automata theory, the categories  $\mathbf{A}_W$  are <u>transparent</u>. That is, anything which can be said about the structure of transition systems with input W, can be said by referring to their homomorphisms only. In particular, all the automorphisms of  $\mathbf{A}_W$ , for this type of W, are naturally equivalent to the identity functor of  $\mathbf{A}_W$ . Π

For further information: The complete report is available in the major Navy technical libraries and can be obtained from the Defense Documentation Center. A few copies are available for distribution by the author.

#### 1. INTRODUCTION

Several recent results in automata theory (in particular, Hartmanis & Stearns 1964, Zeiger 1964) give evidence of the importance of homomorphisms in the study of transition systems and automata. It is natural therefore to inquire how much information can be retrieved from the algebra of homomorphism compositions with respect to transition systems. The natural mathematical framework for the discussion of this problem is categorical algebra.

We define a category  $\mathbf{A}_W$  of the transition systems with input W, where W is any arbitrary fixed monoid, and with arbitrary sets of states. A preliminary study of  $\mathbf{A}_W$  (Give'on 1964) shows that one can reconstruct the internal structure of any transition system from the way homomorphisms (i.e., the morphisms of  $\mathbf{A}_W$ ) behave around it.

In this paper we show that  $\mathbf{A}_W$  has a generator,  $\mathbf{M}_W$  (which is W operating on itself as a transition system) and that there exists a functor Mor :  $\mathbf{A}_W \longrightarrow \mathbf{A}_W$  naturally equivalent to the identity functor of  $\mathbf{A}_W$  which factors through Hom  $\mathbf{A}_W$  ( $\mathbf{M}_W$ ,-).

-1-

\$

A general exposition of the nature of properties which are retrievable from the "morphism-behavior" in an arbitrary category is presented so that it provides a rigorous general basis for studying "retrievable" properties and categories in which every structural property of objects and morphisms is "retrievable." E

 $\prod$ 

Finally, we prove that for a very broad class of input monoids, which includes all the types of input-monoids encountered in automata theory, the catesories  $R_W$  are <u>transparent</u>. That is, anything which can be said about the structure of transition systems with input W, can be said by referring to their homomorphisms only. In particular, all the automorphisms of  $A_W$ , for this type of W, are naturally equivalent to the identity functor of  $A_W$ .

Some elementary acquaintance with categorical algebra is needed. In particular, we shall make use of the following notions:

(i) Category, it objects and its morphisms.

(ii) Epic, monic, and invertible morphisms versus surjective, injective, and bijective functions.

-2-

(iii) Initial and terminal objects.

(iv) <u>Functors</u>, <u>natural transformations</u> and <u>natural equivalences</u> of functors.

• •

¥.....

(v) <u>Embedding</u> functors, <u>automorphism</u> functors, and <u>adjoint</u> functors.

The reader who is not familiar with these notions is referred to the literature (Kan 1958, Freyd 1964, and MacLane 1965). Additional issues of categorical algebra with reference to automata theory are discussed in (Give'on 1965).

### 2. CATEGORIES OF TRANSITION SYSTEMS

2.1 Let W be a fixed monoid, We denote by  ${\bf A}_{\rm W}$  the category specified as follows.

E

The objects of  $\mathbf{A}_W$  are <u>transition-systems</u> with input W. That is, systems of the form

$$A = (S(A) \times W \xrightarrow{\lambda} S(A))$$

where:

(i) S(A) is any set, the <u>set of states</u> of A; (ii)  $\lambda_{A} : S(A) \times \mathbb{N} \longrightarrow S(A)$  is a function, the <u>transition</u>

function of A, with the following properties (we write s.o) for  $\lambda_A(s,\omega)$ :

(iii)  $s \cdot l_{W} = s$  for all  $s \in S(A)$ , where  $l_{W}$  is the identity

element of W;

(iv)  $s \cdot (\omega_1 \omega_2) = (\mathbf{s} \cdot \omega_1) \cdot \omega_2$  for all  $s \in S(A)$  and  $a \parallel \omega_1 \cdot \omega_2 \in W$ .

The morphisms of  $\mathbf{A}_{ij}$  are of the form

$$\Lambda \xrightarrow{t} B$$

-4-

where  $f : S(A) \longrightarrow S(B)$  is a function satisfying  $f(s \cdot \omega) = f(s) \cdot \omega$  for all seS(A) and all  $\omega \in W$ . (Note that  $s \cdot \omega$  on the left hand of this equation refers to the transition function of A, while  $f(s) \cdot \omega$  refers to the transition function of B.)

The composition of the morphisms of  $A_W$  is determined in an obvious manner by the composition of the functions which underly the morphisms. That is,  $(C \xrightarrow{g} D)(A \xrightarrow{f} B)$  is defined <u>only when B = C</u> and then it is equal to  $A \xrightarrow{gf} D$ .

2.2 As in many other "natural" categories of mathematical systems, we have a <u>forgetful</u> functor  $S : A_{W} \longrightarrow S$  from  $A_{W}$  to S the category of sets where S(A) is the set of states of A and  $S(A \xrightarrow{f} b) = (f : S(A) \longrightarrow S(B))$ .

Note that  $\mathbf{A}_{_{\mathrm{N}}}$  contains, among its objects, an <u>empty</u> object to be denoted by  $\phi_{_{\mathrm{A}}}$ . Here we adopt the useful convention that for any set T there exists a unique function which is injective (i.e., one-one into) from  $\phi$ , the empty set, into T. Thus, the transition function of  $\phi_{\mathrm{lv}}$  is this "empty" function :  $\phi \ge 1 \longrightarrow \phi$  ( $\phi \ge 1 = \phi$ ) and for any object  $\Lambda$  of  $\mathbf{A}_{\mathrm{lv}}$ there exists a unique morphism  $\phi_{\mathrm{lv}} = \frac{\sigma_{_{\mathrm{A}}}}{\longrightarrow} \Lambda$  which is determined by the "empty"

-5-

 $\phi = S(\phi_{k}) \longrightarrow S(A).$ 

The forgetful functor  $S : A_W \longrightarrow S$  has an adjoint (cf. Kan 1958, MacLane 1965), the functor  $Fr : S \longrightarrow A_W$ , which assigns to each set T, an object Fr(T) of  $A_W$  which is free on T <u>c</u> S(Fr(T)).

The functor  $Fr : S \longrightarrow A_W$  can be specified as follows. For any set T, Fr(T) is the transition system defined by:

$$S(Fr(T)) = T \times W$$
,

$$(t, \omega_1) \cdot \omega_2 = (t, \omega_1 \omega_2)$$

6

For any function  $f: T_1 \longrightarrow T_2$  there exists a unique morphism

 $Fr(T_1) \xrightarrow{Fr(f)} Fr(T_2) \text{ such that for any te } T_1 : [Fr(f)](t, 1_W) = (f(t), 1_W).$ Hence Fr(T) is "free on  $T \ge \{1_W\}$ ". We identify the elements of  $T \ge \{1_W\}$ with the elements of  $T : t \equiv (t, 1_W).$ 

If  $T_1$  and  $T_2$  are sets which have the same cardinality, then  $Fr(T_1)$  and  $Fr(T_2)$  are isomorphic (i.e., there exists an invertible morphism  $Fr(T_1) \longrightarrow Fr(T_2)$  of  $A_W$ . In particular, if T is a single-element set then we denote Fr(T) by  $M_W$ .

-6-

 $M_W$  serves a very important role in  $A_W$  as we shall see later. Note that  $M_W$  may be defined as W operating on itself. That is,

> $S(M_W) = W$ ,  $\omega_1 \cdot \omega_2 = \omega_1 \omega_2$ .

2.3  $\mathbb{A}_{W}$  shares with the obelian "natural" categories, e.g., of groups or of modules, (cf. Fréyd 1964) the property that the monic (respectively, the epic, and the invertible) morphisms are precisely those morphisms of  $\mathbb{A}_{W}$  whose underlying functions are injective (respectively, surjective and bijective). The arguments that establish these facts are similar to the arguments employed in the category of groups for the same end.

The existence of the forgetful functor  $S : A_W \longrightarrow S$  implies that a morphism  $A \xrightarrow{f} B$  is invertible in  $A_W$  iff f is bijective. Since S is an embedding functor, every morphism of  $A_W$  whose underlying function is injective (respectively, surjective) must be monic (respectively, epic).

In order to prove the converse (for monic and epic morphisms of  $\mathbf{A}_{u}$ ) we need some additional observations about  $\mathbf{A}_{u}$ . These observations

-7-

3Ľ

will be incorporated in the proofs of the following lemmata.

2.3.1 <u>LENMA</u>: If  $A \xrightarrow{9} B$  is an epic morphism of  $\mathbf{A}_{W}$  then  $e : S(A) \longrightarrow S(B)$  is surjective.

I

Π

<u>Proof:</u> The image of  $e : S(A) \longrightarrow S(B)$  is a subset e(S(A)) of S(B) such that for any  $\omega \in W$  and any  $s \in e(S(A))$ ,  $s \cdot \omega \in e(S(A))$ . Hence e(S(A)) is a transition system e(A) which is a sub-system of B.

We define a new object B/e(A) of  $A_{u}$  by:

 $S(B/e(A)) = (S(B) - e(S(A))) \cup \{s_{*}\}$  where  $s_{*} \in S(B)$ ;

the transition function of B/e(A) is the same as of B except for the cases where  $s \cdot \omega c e(S(A))$ ; in these cases we set  $s \cdot \omega = s_{\star}$ , and for all  $\omega c w$  we set  $s_{\star} \cdot \omega = s_{\star}$ .

Obviously, B/e(A) is formed from B by contracting e(S(A)) to a single state s<sub>\*</sub>. This contraction takes the form of a canonical morphism  $B \xrightarrow{q_e} B/e(A)$ , where  $q_e : S(B) \longrightarrow S(B/e(A))$  is identical on S(B) - e(S(A)) and it maps all of e(S(A)) onto s<sub>\*</sub>.

-8-

In addition to  $B \xrightarrow{q_e} B/e(A)$ , we have another morphism  $B \xrightarrow{Z} B/e(A)$  which maps all of e(S(A)) onto  $s_*$ . Clearly e is surjective iff  $q_e = z$ .

Obviously  $ze = q_e$ , since both map all of S(A) onto s<sub>\*</sub>. But e is epic and therefore  $ze = q_e$  implies  $q_e = z_e$ .

2.3.2 <u>LENMA</u>: If  $A \xrightarrow{j} B$  is a monic morphism of  $A_W$  then  $j : S(A) \longrightarrow S(B)$  is injective.

<u>Proof:</u> Assume that for  $s_1, s_2 \in S(A)$  we have  $j(s_1) = j(s_2)$ .

We define two morphisms  $\mathbb{M}_{W} \xrightarrow{f_{1}} \mathbb{A}$  and  $\mathbb{M}_{W} \xrightarrow{f_{2}} \mathbb{A}$  by  $f_{1}(\mathbb{I}_{W}) = s_{1}$ , and  $f_{2}(\mathbb{I}_{W}) = s_{2}$ . Obviously,  $jf_{1} = jf_{2}$ , and since j is monic, it follows that  $f_{1} = f_{2}$ ; i.e.,  $s_{1} = s_{2}$ .

2.4 For any object A of  $A_W$  any any subset T <u>c</u> S(A), we define A(T),

-9-

the subsystem of A generated by T, as follows:

 $S(A(T)) = T \cdot W = \{t \cdot \omega : t_E T \text{ and } \omega_E W \},$ 

 $(\mathbf{t} \cdot \boldsymbol{\omega}_1) \cdot \boldsymbol{\omega}_2 = \mathbf{t} \cdot (\boldsymbol{\omega}_1 \boldsymbol{\omega}_2).$ 

A subset T of S(A) is said to generate A iff A(T) = A; i.e., iff T·W = S(A). In particular A is said to be monogenic iff A is generated by a single-element subset of S(A). N

0

Π

For example,  $M_W$  is monogenic since  $\{1_W\}$  generates  $M_W$  (Obviously for any  $\omega \epsilon W$  :  $1_W \cdot \omega = \omega$ ). More generally,  $M_W$  is generated by  $\{u\}$  iff there exists  $v \epsilon W$  such that  $uv = 1_W$ .

Note that an object A of  $\mathbb{A}_W$  is monogenic iff for any T <u>c</u> S(A) which generates A there exists t<sub>E</sub>T such that {t} generates A.

2.4.1 <u>LEMMA</u>: An object A of  $\mathbb{A}_W$  is monogenic iff for any family {A<sub>j</sub>} of subsystems of A indexed by a set J, U{S(A<sub>j</sub>) : j $\in$ J} = S(A) implies S(A<sub>j</sub>) = S(A) for some j $\in$  J. <u>Proof:</u> Assume that A is monogenic and generated by  $\{s_0\}$ . If US(A<sub>i</sub>) = S(A) then s<sub>0</sub>  $\in$  S(A<sub>i</sub>) for some j $\in$  J, and S(A<sub>i</sub>) = S(A).

Assume that for any family  $\{A_j\}$  indexed by a set J,  $US(A_j) = S(A)$  implies  $S(A_j) = S(A)$  for some jcJ. Define the family  $\{A_s\}$  for all scS(A), where  $A_s$  is the subsystem of A generated by  $\{s\}$ . Obviously  $US(A_s) = S(A)$  and therefore there exists  $s_0 \in S(A)$  for which  $S(A_s) = S(A)$ . Hence A is generated by  $\{s_0\}$ .

2.4.2 <u>COROLLARY</u>: For any monogenic object A of  $A_W$  and any automorphism F:  $A_W \rightarrow A_W$  of  $A_W$ , F(A) is also a monogenic object of  $A_W$ .

<u>Proof</u>: We recall that an automorphism F of  $\mathbb{A}_W$  is a functor F:  $\mathbb{A}_W \longrightarrow \mathbb{A}_W$  for which there exists a functor G :  $\mathbb{A}_W \longrightarrow \mathbb{A}_W$  such that both F.G and G.F are equal to the identity functor of  $\mathbb{A}_W$ .

The families of subsystems of A are represented faithfully by the families of monic morphisms of  $A_W$  with range A. Given a set J of monic morphisms  $A_j \xrightarrow{j} A$  we define a category T (which is a subcategory of  $A_W$ ) whose objects are all the monic morphisms

-11-

 $B \xrightarrow{b} A$  such that for any jeJ there exists a monic  $A_j \xrightarrow{D} B$  with

 $bb_i = j$ . The morphisms of **J** are of the form

$$(B_1 \xrightarrow{b_1} A) \xrightarrow{f} (B_2 \xrightarrow{b_2} A)$$

where  $B_1 \xrightarrow{f} B_2$  is a morphism of  $A_W$  with  $B_2 f = b_1$ .

For any set J of monic morphisms of  $A_W$  with range A, the categrory J has an initial object U(J), which is unique up to an isomorphism of J (which is an equivalence of monic morphisms in  $A_W$ (cf. Freyd 1964, MacLane 1965)). U(J) is a monic morphism of  $A_W$ with range A and whose image is precisely the union of the images of the morphisms in J.

We can rephrase now Lemma 2.4 1 : An object A of  $\mathbb{A}_{W}$ is monogenic iff for any set J of monic morphisms of  $\mathbb{A}_{W}$  with range A, if U(J) is an invertible morphism of  $\mathbb{A}_{W}$  (i.e., an isomorphism) then there is a jeJ which is invertible.

Since this characterization of the monogenic objects in  $\mathbf{A}_{\mathbf{w}}$  is preserved under the automorphisms of  $\mathbf{A}_{\mathbf{v}}$  the proof follows.

-12-

2.5 In the proof of Cor. 2.4.2 we have shown that the property of being a monogenic object of  $\mathbf{A}_{w}$ , which was defined originally by "looking inside A," is in fact definable by means of general properties of morphisms in categories. Knowing the way morphisms behave around an object A is sufficient in order to determine whether A contains a state from which all the rest of the states of A are accessible. In other words, the property of being a monogenic object in  $\mathbb{A}_w$  is <u>categorical</u>. In Chapter 4 we shall present a rigorous explication of this notion. The properties of  $M_{\mu\nu}$ , that we shall derive in the next chapter, will yield the result that all properties of objects of  ${\bf A}_{{\bf W}}$  (which are invariant under isomorphisms in  $\mathbb{A}_{\mu}$ ) are categorical (provided that W belong to a very broad class of monoids). That is, if W satisfies some weak conditions, then all the properties of the transition systems with input W can be derived from the categorical-algebra study of  $\mathbb{A}_{w^*}$ .

-13-

3.1 <u>LEMMA</u>: An object A of  $\mathbb{A}_W$  is monogenic iff there exists an epic morphism  $M_W \xrightarrow{e} A$ .

<u>Proof</u>: Assume that A is monogenic and generated by  $\{s_0\}$ .

Π

Define a morphism  $M_W \xrightarrow{f_{s_o}} A$  by  $f_{s_o}(1_W) = s_o$  (recall that  $M_W$  is free

on  $l_W$ ). Obviously,  $f_{s_0} : W \longrightarrow S(A)$  is surjective and therefore  $M_{w} \xrightarrow{f_{s_0}} A$  is epic.

On the other hand, if  $M_{W} \xrightarrow{e} A$  is epic then A is generated

by 
$$e(1_W)$$
 since  $e(\omega) = e(1_W)^* \omega$ .

3.2 We define a functor  $H_{M_W} : \mathbf{A}_{W} \longrightarrow \mathbf{S}$  by:

$$H_{\mathcal{H}_{W}}(A) \cong Hom_{\mathcal{H}_{W}}(M_{W},A)$$

$$H_{M_{W}}(A \xrightarrow{\mathbf{I}} B) = (I_{\underline{f}} : Hom_{M_{W}}(M_{W}A) \longrightarrow Hom_{M_{W}}(M_{W},B))$$

where 
$$l_{f}(M_{W} \xrightarrow{B} A) = (M_{W} \xrightarrow{f_{g}} B)$$

-14-

We define a transformation of functors  $g: S \longrightarrow H_{M_W}$  as

follows:

For any object A of  $\mathbb{A}_{W}$ ,  $g(A) : S(A) \longrightarrow H_{M}(A)$  is given by

 $[\mathfrak{g}(A)](\mathfrak{s}) : \mathbb{W} \longrightarrow \mathbb{S}(A) : \omega \longrightarrow \mathfrak{s} \cdot \omega \quad .$ 

In other words,  $[\varphi(A)](s)$  is the morphism  $M_W \xrightarrow{f_s} A$  which is determined by  $f_s(1_W) = s$ .

The function  $g(A) : S(A) \longrightarrow H_{M_{ful}}(A)$  is bijective. It

is injective since  $f_{s_1} = f_{s_2}$  implies  $f_{s_1}(1_W) = f_{s_2}(1_W)$ . It is sur-

jective since for any morphism  $M_{W} \xrightarrow{g} A$  we have  $f_{g(1_{W})} = g$ .

Furthermore, for any morphism  $A \xrightarrow{g} B$  of  $A_w$ , and for any  $s \in S(A)$ 

we have

$$g f_s = f_{g(s)}$$
,

where  $f_{g(s)} = [\rho(B)](g(s))$ .

For we clearly have

$$(gf_s)(\omega) = g(s \cdot \omega) = g(s) \cdot \omega = f_{g(s)}(\omega).$$

.

-15-

## From this follows directly, that for any morphism $A \xrightarrow{g} B$

Π

Π

{

of  $A_{\mu\nu}$ , the following diagram is commutative.

$$S(A) \xrightarrow{g} S(B)$$

$$P(A) \xrightarrow{\downarrow} P(B)$$

$$H_{M_{W}}(A) \xrightarrow{\longrightarrow} H_{M_{W}}(B)$$

$$H_{M_{W}}(g)$$

Thus we have proved:

3.2.1 <u>PROPOSITION</u>: The transformation  $\rho: S \longrightarrow H_M$  is a natural

equivalence of functors.

3.2.2 The pair  $(M_{W^{0}\rho}^{-1})$  is a <u>representation</u> of the forgetful functor S :  $A_{W} \longrightarrow S$  (cf. MacLane 1965).

3.2.3. Since S :  $A_W \longrightarrow S$  is an embedding functor (i.e., one-one on the morphisms) it follows that  $H_M$  is also an embedding and therefore  $M_W$ 

 $M_{W}$  is a generator of  $A_{W}$  (cf. Freyd 1964).

3.2.4 <u>COROLLARY</u>:  $M_W$  is a projective object of  $A_W$ . (An object P of

a category C is projective iff for any morphism  $P \xrightarrow{C} B$  and any epic morphism  $A \xrightarrow{e} B$  of C there exists a morphism  $P \xrightarrow{f} A$  of C for which the following diagram is commutative:



<u>Proof</u>: It is sufficient (and necessary) to show that if  $A \xrightarrow{e} B$  is an epic morphism of  $A_W$  then  $H_{M_W}(e) : H_{M_W}(A) \longrightarrow H_{M_W}(B)$  is surjective. From the commutative diagram for  $\boldsymbol{g} : S \longrightarrow H_{M_W}$  we derive

$$H_{M_W}(e) = \rho(B)e(\rho(A))^{-1}$$
. Hence  $H_{M_W}(e)$  is surjective.

3.3.3 <u>PROPOSITION</u>: The bijection  $p(M_W) : W \longrightarrow H_{M_W}(M_W)$  determines

an isomorphism of monoids

$$W \xrightarrow{R} End_{\mathbf{A}_{W}}(M_{W})$$

where  $\operatorname{End}_{A_{W}}(M_{W})$  is the monoid of the morphisms  $M_{W} \longrightarrow M_{W}$  of  $A_{W}$  with

-17-

respect to the composition of morphisms in  $\mathbf{A}_{w}$ .

Proof: Since 
$$f_{\omega}(\omega') = \omega \omega'$$
, it follows that  $f_{\omega} = f_{\omega} = f_{\omega} = \frac{1}{2}$ 

3.4 From Prop. 3.3 it follows that for any object A of  $\mathbf{A}_{W}$ , the set  $H_{M_{W}}$  (A) enjoys a structure of a transition system with input W by

combining  $H_{M_W}(A)$  with  $W \xrightarrow{R} End_{M_W}(M_W)$ .

Formally, we define a functor Mor :  $\mathbb{A}_{W} \longrightarrow \mathbb{A}_{W}$ , where for

 $\Box$ 

2

Π

[

any object A of  ${\bf A}_W$  we define Mor(A) by:

 $S(Mor(A)) = H_{M_{W}}(A),$   $f_{s} \cdot \omega = f_{s}f_{\omega} = f_{s} \cdot \omega \text{ for any } M_{W} \xrightarrow{f} A \text{ and } \omega \in W.$ 

For any morphism  $A \xrightarrow{g} B$  we define Mor(A)  $\xrightarrow{Mor(g)}$  Mor(B)

by  $Mor(g) = H_{M_W}(g)$ .

An immediate verfication shows that Mor(A) is an object of  $A_W$ , and that  $H_{M_W}(g)$  determines in fact a morphism of  $A_W$ . Furthermore,

it follows directly from the fact that  $H_{M_{12}}$  is a functor that Nor :  $A_{12} - A_{13}$ 

-18-

is also a functor. Likewise, the transformation  $\rho: S \longrightarrow H_M$ 

determines directly a transformation  $\rho_{A_{W}}$ : I  $\longrightarrow$  Mor from the identity

functor of  $\boldsymbol{\mathbb{A}}_{W}$  to Mor, and we have:

3.4.1 <u>THEOREM</u>: The transformation  $\rho_{\mathbf{A}_{W}}$ : I  $\longrightarrow$  Mor is a natural

equivalence of functors.

Ð

3.4.2 Intuitively speaking, the functor Mor constructs the "internal structure" of any object A of  $\mathbf{A}_W$  from a part of the category  $\mathbf{A}_W$  which lies around  $\mathbf{M}_W$  and between  $\mathbf{M}_W$  and A. Hence it is intuitively clear, that if  $\mathbf{N}_W$  can be recognized in  $\mathbf{A}_W$  (up to an isomorphism) by means of some categorical predicate, then the "internal structure" of any object can be reconstructed "categorically," and therefore any property of the transition systems with input W can be determined "categorically" as well.

-19-

3.5 <u>LENMA:</u> If W is a <u>unit-commutative</u> monoid (i.e., if  $uv = 1_W$  in W then  $vu = 1_W$ ) or a finite monoid then every epic morphism  $A \xrightarrow{e} M_W$ of  $A_W$ , where A is monogenic, is an isomorphism. a recent a

Π

2

 $\prod$ 

<u>Proof:</u> If W is a finite monoid then the cardinality of the set of states of any monogenic transition system with input W cannot exceed the cardinality of W. Hence  $e : S(A) \longrightarrow W$  must be bijective.

If W is a unit-commutative monoid and A is generated by  $\{s_0\}$ , then  $\{e(s_0)\}$  must generate  $\mathbb{M}_W$ , that  $is_i e(s_0)v = 1_W$  for some  $v_{\mathcal{E}}$  W, and therefore  $ve(s_0) = 1_W$ .

Assume that  $e(s_0 \cdot \omega_1) = e(s_0 \cdot \omega_2)$  for some  $\omega_1, \omega_2 \in W$ , then we have  $\omega_1 = ve(s_0)\omega_1 = ve(s_0 \cdot \omega_1) = ve(s_0 \cdot \omega_2) = ve(s_0)\omega_2 = \omega_2$ ,

and therefore  $s \cdot \omega_1 = s \cdot \omega_2$ , which shows that e is also injective.

3.5.1 <u>COROLLARY</u>: If W is a unit-commutative monoid or a finite monoid, then for any automorphism F of  $\mathbb{A}_{W}$ ,  $F(\mathbb{N}_{W})$  is isomorphic to  $\mathbb{N}_{W}$ .

-20-

<u>Proof:</u> From lemma 3.5 it follows that an object M of

 $\boldsymbol{A}_{\boldsymbol{W}}$  is isomorphic to  $\boldsymbol{M}_{\boldsymbol{W}}$  iff

(i) M is monogenic, and

(ii) for any monogenic object A of  $\mathbb{A}_W$  there exists an epic morphism  $M \xrightarrow{e} A$  of  $\mathbb{A}_W$ .

Since these properties of morphisms and objects of  $\mathbb{A}_{W}$  are preserved under the automorphisms of  $\mathbb{A}_{W}$ , the corollary follows.

3.5.2 Note that the class of unit-commutative monoids is broad enough to cover all the classes of monoids which are employed in automata theory. For example, the left-cancellative and the rightcancellative monoids are all unit-commutative. Hence the free monoids and the groups are unit-commutative. Note also that the cartesian products of unit-commutative monoids are unit-commutative, and therefore we can apply our results to multi-input transition systems as well.

-21-

4. CATEGORICAL PREDICATES AND TRANSPARENT CATEGORIES

4.1 A subcategory D of C is said to be <u>very full</u> iff for any morphism h of D and for any morphisms f and g of C such that fg = h, fh = g or hf = g holds in C it follows that f and g belong to D. Sec.

Sufficient States

10.000

and the second

Sec. 1

 $\left[ \right]$ 

0

A functor  $T : \mathbb{D} \longrightarrow C$  is said to be a <u>very full embedding</u> iff T is an embedding and the image of T is a very full subcategory of C.

Let **D** be a category and **D** a class of morphisms of **D**, we denote by (**D**,**D**,**C**) the class of all the images of the morphisms in **D** under any very full embedding  $T: \mathbb{D} \longrightarrow \mathbb{C}$ . That is,  $f \in (\mathbb{D}, D, \mathbb{C})$  iff there are a morphism  $d_{\mathbb{C}}D$  and a very full embedding  $T: \mathbb{D} \longrightarrow \mathbb{C}$  such that f = T(d),

A class K of morphisms of C is said to be <u>categorical</u> (in C) iff there is a category D such that K = (D, D, C) for some class D of morphisms of D.

4.2 <u>PROPOSITION:</u> A class K is categorical in C iff it is closed under all the automorphisms of C.

-22-

<u>Proof:</u> Since for any very full embedding functor  $T : D \longrightarrow C$ and for any automomphism F of C, F<sup>o</sup>T :  $D \longrightarrow C$  is also a very full embedding, it follows that every categorical class in C is closed under all the automomphisms of C.

On the other hand, let K be a class of morphisms of C which is closed under all automorphisms of C. Denote by D(K) the minimal very full subcategory of C which includes K, then K = (D(K), K, C).

In order to see this, let  $T : D(K) \longrightarrow C$  be any very full embedding and define  $F_T : C \longrightarrow C$  by

$$F_{T}(f) = \begin{cases} T(f) & \text{if fab}(K), \\ \\ f & \text{otherwise.} \end{cases}$$

Since D(K) is a very full subcategory of C,  $F_T$  maps D(K) into itself, and because T is a very full embedding,  $F_T$  maps D(K) onto itself in an injective manner. Furthermore,  $F_T$  must be a functor and it has an inverse, hence it is an automorphism of C.

Now, since K is closed under automorphisms it follows that  $(D(K), K, C) \subseteq K$ , and since clearly  $K \subseteq (D(K), K, C)$  we have the

-23-

the desired equality.

4.2.1 <u>COROLLARY</u>: A class K is categorical in C iff K = (D, D, C)for some very full subcategory D of C.

1 A 40

Π

Π

Π

C

4.2.2 <u>COROLLARY</u>: Let **D** be any category and D a class of some morphisms of **D**, then the class of all values of the morphisms in D under all embedding functors  $D \rightarrow C$  is categorical in C.

4.2.3 <u>COROLLARY</u>: The class of all values of the morphisms in D under all functors  $\mathbb{D} \longrightarrow \mathbb{C}$  is categorical in  $\mathbb{C}$ .

4.2.4 Note that we cannot dispense with the requirement of employing very full embeddings in the definition of the categorical classes in any arbitrary category. For example in the category N of natural numbers where the morphisms represent the natural partial order of natural numbers, every set of morphisms is categorical. However, the categorical classes achieved by means of 4.2.2 or 4.2.3 are always infinite or empty.

U

N

4.3.1 A class of morphisms of **C** is said to be <u>natural</u> iff it is closed under all those automorphisms of **C** which are naturally equivalent to  $I_{\mathbf{r}}$ , the identity functor of **C**.

Obviously, by Prop. 4.2 we have that every categorical class is natural. Note that a class of identity morphisms of **C** which is closed under the isomorphisms within **C** is always natural.

4.3.2 A category C is said to be transparent if all the natural classes in C are categorical.

4.4 Obviously, if all the automorphisms of C are naturally equivalent (i.e., to  $I_{c}$ ) then C is transparent.

Let us call a category C <u>autotrivial</u> iff all the automorphisms of C are naturally equivalent.

-25-

It is not known whether all transparent categories are autotrivial. All "natural" categories of mathematical systems that are known to be transparent are in fact autotrivial as well. 1000

s he x

Π

Π

The equivalence between the notion of transparent categories and that of autotrivial categories, in a special case, takes the form of the following problem in group theory:

Do all groups whose automorphisms are all (conjugate) class preserving have only inner automorphisms?

Any example of a group all of whose automorphisms are class preserving and which has an outer automorphism, yields a transparent category (with a single object and all its morphisms are invertible and in one-one correspondence with the elements of the group) which is not autotrivial. 5. THE DISTINGUISHABILITY OF M. AND THE TRANSPARENCE OF A.

5.1 The features exhibited by  $M_W$  in  $A_W$  are quite common in "natural" categories. As we shall see presently, they provide a reduction of the autotriviality of categories to the categoricity of certain classes of identity morphisms.

An object M of a category C is said to be a <u>generator</u> of C iff  $H_M : \mathbb{C} \longrightarrow \mathbb{S}$  (where  $H_M(A) = \operatorname{Hom}_{\mathbb{C}}(M,A)$ ) is an embedding. In this case, the values of  $H_M$  form a subcategory  $H_M(C)$  of S. In particular, M is said to be a <u>faithful</u> generator of C iff there exists a functor  $R_M : H_M(C) \longrightarrow C$  such that  $R_M \circ H_M : C \longrightarrow C$  is naturally equivalent to  $I_C$ .

5.1.1 <u>EXAMPLES</u>: The additive group Z of integers is a faithful generator . of both the category of obelian groups and the category of all groups (Freyd 1964). The single-element set U is a faithful generator of S the category of sets; in fact  $H_{11}$  is already naturally equivalent to the

-27-

identity functor of S.

From our results in Chapter 3 we know that  $M_W$  is a faithful generator of  $A_W$ .

ł

I

[

ſ

5.2 <u>LEMMA</u>: For any object A of C and any automorphism F of C with inverse G,  $H_A \circ F$  is naturally equivalent to  $H_{G(A)}$ 

<u>Proof:</u> We shall prove a stronger result; namely, for any two objects A and B of C and any automorphism F of C with inverse G, there exists a bijection

$$\P(A,B)$$
 : Hom<sub>r</sub> (G(A), B)  $\longrightarrow$  Hom<sub>r</sub> (A, F(B))

which is natural in both A and B.

Put differently, F and its inverse are adjoint. By Kan's characterization of adjoint functors (Kan 1958, MacLane 1965) the following is a proof that F and G are adjoint (to each other!):

Let us denote by  $s_{r_1}$  the identity morphism of an arbitrary object C of C.

(i) Every morphism A  $\xrightarrow{f}$  F(B) can be factored as  $f = F(h) \cdot e_A$ 

-28-

for some  $G(A) \xrightarrow{h} B$  (i.e., h = G(f)).

(ii) If  $F(h_1) \cdot e_A = F(h_2) \cdot e_A$  then obviously  $h_1 = h_2$ .

5.3 Let M be a faithful generator of **C**. From Lemma 5.2 we know that  $H_M \circ F$  is naturally equivalent to  $H_{G(M)}$  for any automorphism F of **C** with inverse G. Hence F, which is naturally equivalent to  $R_M \circ H_M \circ F$ , is naturally equivalent to  $R_M \circ H_{G(M)}$ . If G(M) is isomorphic to M, then  $H_M$  is naturally equivalent to  $H_{G(M)}$ , and therefore F, which is naturally equivalent to  $R_M \circ H_{G(M)}$ , is naturally equivalent to  $I_C$ .

5.3.1 An object A of C is said to be <u>distinguishable</u> (in C) iff for any automorphism F of C, F(A) is isomorphic to A. Put differently, A is distinguishable in C iff the natural class of all identity morphisms of the objects of C, which are isomorphic to A in C, is categorical.

Thus we have proved:

-29-

5.3.2 <u>THEOREM</u>: A category C with a faithful generator M is autotrivial iff M is distinguishable in C.

5.3.3 <u>COROLLARY</u>: A category C with a faithful generator is autotrivial iff it is transparent.

[]

5.4 Since  $M_W$  is a faithful generator of  $A_W$ ,  $A_W$  is transparent iff it is autotrivial. Furthermore  $A_W$  is autotrivial iff  $M_W$  is distinguishable in  $A_W$ . By 3.5.1 we know that if W is finite or unitcommutative there  $M_W$  is distinguishable in  $A_W$ . Thus we have:

5.4.1 <u>THEOREM</u>: If W is a finite monoid or a unit commutative monoid then  $\mathbb{A}_{W}$  is autotrivial (and therefore transparent).

6. DISCUSSION AND OPEN PROBLEMS

6.! Our result as expressed by Theorem 5.4.1 implies that, for a very broad class of monoids, the categorical study of a domain of all transition systems with input monoid of this class, is equivalent <u>in</u> <u>principle</u> to the "complete" study (or the "inside" study) of these systems. However, only experience may show us that in fact there is a psychological advantage to the categorical approach in the study of these systems 6.2 If G is a group then  $\mathbb{A}_{G}$  is the category of all <u>representations of</u> <u>G as operating on sets.</u> Since every group is in particular a unit-commutative monoid, we have that the categorical study of the representations of a fixed arbitrary group G is sufficient <u>in principle</u> for producing all the algebraic properties of the representations of G.

-31-

6.3 Our results so far, give rise to some general problems that deserve attention. For example, is it true that for any monoid W,  $M_W$ is distinguishable in  $A_W$ ?

More generally, what additional properties on faithful generators of categories, if any at all, are necessary in order to insure that they are distinguishable? In particular, is it true that every projective faithful generator is distinguishable?

6

6.4 Important and much more interesting categories of transition systems are those of <u>finite-state</u> transition systems (i.e., transition systems whose sets of states are finite). If the input monoid W is also finite then our results remain valid since  $M_W$  is also finite. If however W is infinite, as it is the case in the ordinary theory of finite automata (where the input monoids are finitely generated free monoids) then  $M_W$  is no longer applicable.

6.5 Another interesting restriction of  $A_w$  is to <u>abelian</u> transition

-32-

systems. A transition system A is said to be <u>abelian</u> iff  $s \cdot \omega_1 \omega_2 = s \cdot \omega_2 \omega_1$  holds for all  $s \in S(A)$  and  $\omega_1, \omega_2 \in W$ .

For any arbitrary monoid W there exists a homomorphism of monoids W  $\xrightarrow{ab}$  W<sup>ab</sup> where W<sup>ab</sup> is an abelian monoid with the following universality property : any homomorphism of W into an abelian monoid factors uniquely through ab. The direct construction of W<sup>ab</sup> must be evident.

> Denote by  $M_W^{ab}$  the following object of  $A_W$ :  $S(M_W^{ab}) = W^{ab}$ ,  $ab(\omega_1) \cdot \omega_2 = ab(\omega_1 \omega_2)$ .

Obviously,  $M_W^{ab}$  is abelian. If we denote by  $A_W^{ab}$  the full subcategory of  $A_W$  of abelian objects, one can easily follow the example of  $M_W$  in  $A_W$  and show that  $M_W^{ab}$  is a fait..fu' generator of  $A_W^{ab}$ . Furthermore, for any arbitrary monoid W,  $M_W^{ab}$  is distinguishable by the same properties which distinguish  $M_W$  in  $A_W$  in the case of unit-commutative input monoid W (cf. 3.5.1), hence for any arbitrary monoid W,  $A_W^{ab}$  is transparent and autotrivial. An equivalent proof of these properties of A follows

1.111 (1.111 (1.111))

[

directly from the fact that  $A_{W}^{ab}$  is a category isomorphic to  $A_{W}^{ab}$ .

- 34 -

#### BIBLIOGRAPHY

Į,

Freyd, P., "Abelian Categories: An Introduction to the Theory of Functors," Harper's Series in Modern Mathematics, New York (1964).

Give'on, Y., "Toward a Homological Algebra of Automata I: The Representation and Completeness Theorem for Categories of Abstract Automata," Technical Report, Department of Communication Sciences, ORA, The University of Michigan (1964).

Give'on, Y., "Toward a Homological Algebra of Automata," Technical Report, Department of Communication Sciences, ORA, The University of Michigan (1965).

Hartmanis, J. and Stearns, R. E., "Pair Algebra and Automata Theory," <u>Inform. Control</u>, 7, 485 - 507 (1964).

Kan, D. M., "Adjoint Functors," <u>Trans. Amer. Math. Soc.</u>, <u>87</u>, 295 - 329 (1958).

MacLane, S., "Categorical Algebra," Bull, Amer. Math. Soc., 71, 40 - 106 (1965).

Zeiger, H. P., "Loop-Free Synthesis of Finite State Machines," thesis, MIT (1964).

	and the second secon	Section in the section of the sectio		
DOCUMEN Complete elevatilization al title brids of shatest	T CONTROL DATA - RAD			
IGINATIN & ACTIVITY (Corporate author)	80. REP			
Logic of Computers Group		Unclassified		
The University of Michigan	25. 000			
Ann Arbor, Michigan 48104				
PURT TITLE				
TRANSPARENT CATEGORIES AND C	ATEGORIES OF TRANSITION S	YSTEMS		
SCRIPTIVE NOTES (Type of most and makelyn de	<b>m</b> )			
Give'on, Yehoshafat				
PORT DATE	Te TOTAL NO. OF PASES	74. NO. OF REFS		
May, 1905	25	<u>1 7</u>		
Nonr 1224(21)	ATTAE TO M			
PROJECT NO.	0210 <b>2-20-</b> 1			
	SA OTHER REPORT HOLD (A.	y other numbers that may be sealer		
UPPLEMENTARY NOTES	12. SPONSORING MILITARY ACTIVITY Office of Naval Research			
	Washington 25 D	C Navy		
	washington 2), D.	· · ·		

**....** 

E

0

Unclassified Security Classification

**3**7

Unclassification				-		n an airtean an a	an ann
Socurity Cleastingening		LINKA		LINK B			
KEY WORDS		ROLE	-	ROLE	TW	NOLE	. #¥.
algebraic automata theory categorical algebra							
	CTIONS						
of the contractor, subcontractor, grantee, Department of De- fense activity or other organization (corporate suthor) issuing the report. 22. REPORT SECURTY CLASSIFICATION: Enter the over- all security classification of the report. Indicate whether "Restricted Data" is included. Marking is to be in accord- ance with appropriate security regulations. 20. GROUP: Automatic downgrading is specified in DoD Di- rective 5200, 10 and Armed Forces Industrial Manual. Enter the group number. Also, when applicable, show that optional markings have been used for Group 3 and Group 4 as author- ised. 3. REPORT TITLE: Enter the complete report title in all capital letters. Titles in all cases should be unclassifica- tion, show title classification in all capitals in parenthesis immediately following the title. 4. DESCRIPTIVE NOTES: If appropriate, enter the type of report, e.g., interim, progress, summary, annual, or final. Give the inclusive dates when a specific reporting period is covered. 5. AUTHOR(S): Enter the name(s) of author(s) as shown on or in the report. Enter the strane, first name, middle initial. 11 military, show rank and branch of service. The name of the principal author is an absolute minimum requirement. 6. REPORT DATE: Enter the date of the report as day, month, year; or month, year. If more than one date appears on the report, use date of publication. 7a. TOTAL NUMBER OF PAGES: The total page count should follow normal paginstion procedures, L.s., enter the number of pages containing information. 75. NUMBER OF REFERENCER Enter the total number of references cited in the report. 8a. CONTRACT OR GRANT NUMBER: If appropriate, enter the applicable number of the contract or grant under which the report was written. 8b. c; h 8d. PROJECT NUMBER: Enter the stal number, subproject number, system numbers, task number, etc. 9a. ORIGINATOR'S REPORT NUMBER(S): Enter the offi- cial report sumber by which the document will be identified or controlled by the origining activity. This number must	<ul> <li>imposed by security classification, using standard statements such as: <ul> <li>(i) "Qualified requesters may obtain copies of this report from DDC."</li> <li>(2) "Foreign announcement and dissemination of this report by DDC is not authorized."</li> <li>(3) "U. S. Government agencies may obtain copies of this report directly from DDC. Other qualified DDC users shall request through</li></ul></li></ul>						his of DDC 

Unclassified Security Classification

\$

\$ • •

8

0

0

ſ

l

ŀ

l

L