

UNCLASSIFIED

AD NUMBER

AD466801

LIMITATION CHANGES

TO:

Approved for public release; distribution is unlimited.

FROM:

Distribution authorized to U.S. Gov't. agencies and their contractors;
Administrative/Operational Use; 16 JUL 1965.
Other requests shall be referred to Office of Naval Research, 875 North Randolph Street, Arlington, VA 22203-1995.

AUTHORITY

ONR memo dtd 28 Dec 1965

THIS PAGE IS UNCLASSIFIED

CATALOGED BY: DDC

AS AD NO.

466801

MASS SPECTROMETRIC STUDIES OF REACTIONS IN FLAMES -
QUANTITATIVE SAMPLING OF FREE RADICALS
FROM ONE-ATMOSPHERE FLAMES

TECHNICAL REPORT

16 July 1965

Contract No. Nonr-3599(00)


ARPA Order No. 23-63, Amendment No. 38

Program Code No. 4910

MRI Project No. 2551-P

For

Director
Advanced Research Projects Agency
Washington, D. C.



MRI

MIDWEST RESEARCH INSTITUTE

425 VOLKER BOULEVARD/KANSAS CITY, MISSOURI 64110/AC 816 LO 1-0202

NOTICE: When government or other drawings, specifications or other data are used for any purpose other than in connection with a definitely related government procurement operation, the U. S. Government thereby incurs no responsibility, nor any obligation whatsoever; and the fact that the Government may have formulated, furnished, or in any way supplied the said drawings, specifications, or other data is not to be regarded by implication or otherwise as in any manner licensing the holder or any other person or corporation, or conveying any rights or permission to manufacture, use or sell any patented invention that may in any way be related thereto.

MASS SPECTROMETRIC STUDIES OF REACTIONS IN FLAMES -
QUANTITATIVE SAMPLING OF FREE RADICALS
FROM ONE-ATMOSPHERE FLAMES

by

Thomas A. Milne
Frank T. Greene

TECHNICAL REPORT
16 July 1965

Contract No. Nonr-3599(00)
ARPA Order No. 23-63, Amendment No. 38
Program Code No. 4910

MRI Project No. 2551-P

For

Director
Advanced Research Projects Agency
Washington, D. C.



MIDWEST RESEARCH INSTITUTE

425 VOLKER BOULEVARD/KANSAS CITY, MISSOURI 64110/AC 816 LO 1-C202

This research was sponsored by the Chemistry Office
of the Advanced Research Projects Agency under
ARPA Order No. 23-63, Amendment No. 38, and moni-
tored by ONR under Contract No. Nonr-3599(00)

PREFACE

This report was prepared for the Advanced Research Projects Agency under ONR Contract No. Nonr-3599(00), monitored by Mr. Roland Jackel and Dr. Ralph Roberts of the Power Branch, ONR.

This report describes a single model of high pressure sampling and some results obtained with one-atmosphere flames.

The authors wish to thank Mr. George Vowels and Mr. Jacob Beachey for their help in gathering much of the data, Mr. Gordon Gross for stimulating discussions and Professors John Fenn and F. S. Sherman for information about the aerodynamics of free jet expansion.

Approved for:

MIDWEST RESEARCH INSTITUTE



Sheldon L. Levy, Director
Mathematics and Physics Division

16 July 1965

TABLE OF CONTENTS

	<u>Page No.</u>
Abstract	1
I. Introduction	2
II. Beam Formation From High-Pressure Sources	2
III. Experimental	3
IV. Sampling From One-Atmosphere Flames	5
A. Stable Products and Temperature-Dependent Fragmentation Patterns	5
B. Reactive Species	8
V. Conclusion	15
References	18

List of Figures

<u>Figure No.</u>	<u>Title</u>	<u>Page No.</u>
1	Line Reversal Flame Temperatures and Chlorine-Containing Species Profiles in Two Rich, $H_2-O_2-N_2$ Flames Directly Sampled at One Atmosphere	7
2	Species Profiles in a Lean $CO-O_2$ Flame Directly Sampled at 2770°K and One Atmosphere	10
3	Oxygen Atom and Hydroxyl Radical Profiles in a Lean H_2-O_2 Flame Directly Sampled at 2540°K and One Atmosphere	12
4	Sulfur-Containing Species Profiles in a Rich One-Atmosphere, H_2-O_2 Flame to Which a Few Per Cent of Either H_2S or SO_2 Had Been Added	16

TABLE OF CONTENTS (Concluded)

List of Tables

<u>Table No.</u>	<u>Title</u>	<u>Page No.</u>
I	Equilibrium Constants Determined From Stable Product Analyses in Several Flames Sampled Directly at One Atmosphere	5
II	Oxygen Dissociation Equilibria Measured by Direct Sampling of Several CO-O ₂ Flames at One Atmosphere	9
III	Oxygen Atoms Sampled Directly From Several One-Atmosphere H ₂ -O ₂ -N ₂ Flames	11
IV	Hydroxyl Radicals Sampled From Several One-Atmosphere H ₂ -O ₂ -N ₂ and CH ₄ -O ₂ Flames	13
V	Experimental Determination of the Equilibrium Constant for the Reaction $H_2O + O \rightleftharpoons 2OH$ in Several H ₂ -O ₂ -N ₂ Flames at One Atmosphere	13
VI	Experimental Determination of the Dissociation of H ₂ in Several H ₂ -O ₂ -N ₂ Flames at One Atmosphere	14
VII	Relative Ion Intensities Observed in a One-Atmosphere H ₂ -O ₂ -N ₂ , 2-1-1 Flame (T=2889°K) at 20 ev. Ionizing Energy	15

ABSTRACT

Direct mass spectrometric sampling studies of one-atmosphere flames are described. A simple model of the sampling process is summarized which accounts for the effects and results obtained. Stable product analysis and evidence for appreciable temperature dependence of fragmentation patterns are shown. Quantitative sampling of the free-radicals H, O, OH and Cl from the burnt gas region of one-atmosphere flames is demonstrated, with direct observation of excess free radical concentrations in the reaction zones of these flames.

I. INTRODUCTION

The first paper^{1/} in this series described the relationship between aerodynamic molecular beam formation and the problem of the direct mass spectrometric sampling of systems at atmospheric pressure and above. The experimental dependence of the beam intensity was given as a function of various system parameters, and the mass separation effect was quantitatively evaluated.

Since the publication of that paper, considerable theoretical and experimental work on the free jet expansion and aerodynamic molecular beams has been reported. Now a rather simple model of the sampling process can be formulated which accounts for the observed effects and gives a quantitative history of the sampling process. In this paper, we summarize this model and present data on beam effects and on the sampling of reactive species from 1-atmosphere flames.

II. BEAM FORMATION FROM HIGH-PRESSURE SOURCES

Starting from a stagnant condition, the gas to be sampled is accelerated as it approaches the sampling orifice, and reaches the speed of sound near the throat of the orifice. At this point, the pressure has dropped by as much as 15-20 per cent. From the sonic condition, the gas expands very rapidly as a "free" or supersonic jet. Along the center line of the jet, the expansion is essentially isentropic, reaching a nearly collisionless flow regime in a distance of 10-20 orifice diameters downstream. If the second slit (skimmer) is placed well downstream of the transition to "collisionless" flow, then shock phenomena are unimportant, and the attenuation experienced by the beam can be accounted for by self-scattering in the beam and by scattering from background gas between the orifice and the detector.

The justification for this model has been discussed in several recent publications. The ideal expansion history downstream of the sonic orifice can be calculated by means of the Mach number vs. distance calculations of Sherman^{2/} for various ratios of heat capacity, γ . Fenn et al.,^{3/} have verified these Mach number predictions over a fairly wide range of conditions and discussed various aspects of beam behavior and attenuation. The authors have discussed sampling history, scattering behavior, partial condensation effects and some preliminary sampling results in two recent symposia.^{4,5/} Scott^{6/} has measured velocity distributions in aerodynamic beams to provide a more detailed picture of the expansion; Deckers and Valleau^{7/} have discussed effects associated with collisions in the beam, and the Karlsruhe group

continues to work on various aspects of beam formation.^{8/} This work and the work of many others, of pertinence to the sampling problem, is extensively reviewed by Knuth.^{9/}

The effects upstream of the orifice are less well understood. No analytical solution exists for the subsonic flow upstream of the orifice. A point sink model has been used with linear interpolation near the orifice to represent the subsonic flow. The results of these calculations are in satisfactory agreement with the more recent experimental data of Sherman.^{2/} The subsonic and low Mach number supersonic flow, which is sensitive to the orifice geometry, may be the most critical for the quenching of very rapid equilibria.

A quantitative temperature pressure history of the expansion can be obtained from the combined subsonic and supersonic calculations. This history is given in previous publications.^{4,5/} For a monatomic gas at 2000°K and 1 atm., expanding through a 0.125-mm. orifice, the translational temperature drops to 30°K and the pressure to 10^{-5} atm. within about 1.5 μ sec. from the first appreciable change. Gases with larger heat capacities will expand somewhat more slowly.

This simple, ideal picture of the sampling process, although it cannot be completely correct, should not be substantially wrong. It appears from the calculated history of expansion that a reaction with a positive activation energy will be satisfactorily quenched.^{4/} Reactions with negative temperature coefficients will be quenched only by the drop in collision frequency, as shown by the partial condensation effects observed.^{10,11/} Real sampling situations may involve other factors. When sampling flames, one must consider the effects of flame-probe interactions, of reactions during sampling, and of γ changing with temperature. The combined effects can, therefore, best be tested by sampling flames of various kinds for species present in known equilibrium concentration. Some experiments of this type are described below.

III. EXPERIMENTAL

The beam-sampling system, which uses a Bendix mass spectrometer^{12/} as a detector, is identical to that described in Part I^{1/} and in Refs. 4 and 5, except that a beam "chopper" with a phase-sensitive detection system has been substituted for the mechanical shutter. A magnetically driven vibrating reed periodically interrupts the beam. Princeton Applied Research lock-in amplifiers provide phase-sensitive detection of the modulated beam. Details will be given in a separate article. Modulation at either 10 or 50 cps gave

identical results, indicating that within experimental error, no appreciable scattered or random gas was being modulated. Although molecules from the modulated beam scattered directly into the electron beam would not be distinguished by the frequency change, we believe that for our geometry this is not a significant source of signal. In a comparison of ratios of reactive to stable species in flames, such as O/O_2 , results taken with the manual shutter gave a smaller ratio of O/O_2 than with the modulated beam. This observation confirms earlier evidence that the shutter-dependent signal obtained with the manual shutter contains an appreciable component due to gas scattered from the beam in the ion source.

With the present system the background-to-beam discrimination is about 1,000 for moderately strong peaks. That is, one should be able to detect a beam signal which is 0.1 per cent of an interfering background peak. This figure was estimated on the basis of the peak-to-peak noise from various strength background peaks, assuming a S/N of 1.0. As one check on the discrimination ratio, a weak beam of argon was created by adding about 4 per cent argon to a stoichiometric $CO-O_2$ flame being sampled. The argon beam was then monitored while argon was leaked directly into the ion source. By the time the total argon signal had reached 1,000 times the original beam intensity, the argon beam intensity had dropped about 30 per cent, presumably because of the rather high argon pressures in the ion source.

Simple sheathed burners and torch flames have been used in the work described here. Some survey experiments, for which adiabatic flame temperatures were sufficiently accurate, employed either laboratory torches or simple tubes fed with premixed gases. For more careful work, sheathed burners were constructed from bundles of 20 gauge stainless steel tubes, capillary glass tubes, or cylindrical rolls of corrugated shim stock. Direct temperature measurements were made by standard line reversal techniques. The measured temperatures were generally from 50 - 100°K below the adiabatic temperatures. These sheathed flames gave convenient, flat flames for work in the burnt gas region. Because of the poor spatial resolution resulting from the large sampling orifices and very thin reaction zones, probing into the reaction zone of any of these 1-atm. flames led to only qualitative profiles.

The sampling orifices used were 90° cones spun from 0.12 and 0.024-mm. thick Pt-20 Rh sheet with 0.24 mm. diameter holes drilled in the apex. The relatively large orifices were used in anticipation of work with condensable species, where plugging is a problem. Although the orifices were soldered to a water-cooled plate, the tips of the orifices were white hot and not far from their softening points with the hotter H_2-O_2 and CH_4-O_2 flames.

IV. SAMPLING FROM ONE-ATMOSPHERE FLAMES

To test empirically the ability of the direct sampling technique to quench reactions and to assess the major difficulties involved, a number of convenient flame systems were surveyed without making a detailed study of any of them. The criterion for quantitative sampling adopted was that the measured equilibrium constant for the reactions being studied agree within a factor of two with the literature values, which were calculated from data taken from the JANAF Tables.^{13/} Experimental errors of this magnitude are usually to be expected for third-law mass spectrometric studies due to lack of knowledge of (a) mass spectrometer transmissions, (b) collection efficiencies and (c) ionization cross sections. Results are presented for stable products, free radicals and temperature-dependent fragmentation patterns.

A. Stable Products and Temperature-Dependent Fragmentation Patterns

A number of flames were sampled for the stable products H_2 , O_2 , H_2O , CO and CO_2 . Several CH_4-O_2 -AR and $CO-O_2$ sheathed flames were probed as a function of distance from the reaction zone. Temperatures were measured by line reversal for these flames. A few of the results reported in Ref. 4 are repeated in Table I for comparison. Stable species were measured in many other flames but have not been reduced to equilibrium constants, since the above data seemed to be sufficiently satisfactory to warrant going on to the more critical test of sampling very reactive species.

TABLE I

EQUILIBRIUM CONSTANTS DETERMINED FROM STABLE PRODUCT ANALYSES
IN SEVERAL FLAMES SAMPLED DIRECTLY AT ONE ATMOSPHERE

Flame Composition (Mole Ratios)				Exptl. Temp. (°K)	K _{eq}	
<u>CH₄</u>	<u>CO</u>	<u>O₂</u>	<u>Ar</u>		<u>Exptl.</u>	<u>Calc.</u>
1		2	8	2454	234	210
1		2.4	8	2457	151	210
					<u>CO + 1/2 O₂ → CO₂</u>	
1		2	8	2454	36.4	36.0
1		2.4	8	2457	28.9	36.0
	1	1		2769	4.9	7.6
	4	3		2824	4.5	6.0
	2	1		2834	4.8	5.7
	3	1		2807	4.4	6.4

12

Two points of interest emerged from these stable product data. First, agreement was obtained for the H_2 concentrations only after applying the first power of the molecular weight correction established in Part I of this series.^{1/} There is nothing in the simple model of the mass separation effect to indicate a temperature dependence, and none has been observed.^{1/} For the H_2 data in CH_4-O_2-Ar flames, the correction relative to O_2 is a factor of 16, the observed H_2^+ peak thus being rather weak.

One complication which arose in making the stable flame product measurements was the temperature-dependence of the fragmentation patterns of even these relatively small molecules. It was first observed that, with an ionizing electron energy of 50 ev., the fragmentation of CO_2 to CO^+ was appreciably temperature-dependent. This temperature-dependence became apparent when the $44^+/28^+$ ratio for a series of progressively leaner flames reached a limiting value. Two experiments were performed to determine the magnitude of this change. CO_2 was added to a lean H_2-O_2 flame which burned at about 1950°K. Since there is a negligible quantity of CO in this lean flame at equilibrium, the 28^+ observed at 50 ev. can be assigned to the fragmentation of hot CO_2 . Entrainment of N_2 was shown to be negligible. An approximately constant value of $44^+/28^+$ was obtained over the first three centimeters beyond the visible reaction zone, indicating equilibration of the reaction $CO_2 + 1/20 \rightleftharpoons CO_2$. The results showed that the fragmentation of CO_2 to CO^+ varies from about 12 per cent at room temperature to 17 per cent at 2000°K in our instrument.

A similar magnitude of temperature effect was deduced from the $44^+/28^+$ ratios observed at 25 and 50 ev. in the four $CO-O_2$ flames shown in Table I. Fragmentation was negligible at 25 ev. and yielded a CO_2/CO ratio in good agreement with equilibrium calculations. The 50 ev. data could then be corrected for the CO contribution to 28^+ and the hot CO_2 cracking patterns determined.

Another test of the effect of temperatures on fragmentation patterns was provided by sampling HCl from a series of rich $H_2-O_2-N_2$ flames which have been extensively studied by Sugden.^{14/} Profiles of $36^+/35^+$ (at an ionizing energy of 50 ev.) in the burnt gas regions of two such flames are shown in Fig. 1, together with measured flame temperatures and the room temperature value of $36^+/35^+$ for pure HCl. A third flame, F_5 in Sugden's nomenclature, exhibited behavior intermediate between U_5 and F_3 .

Two things should be noted. First, the limiting value of the $36^+/35^+$ ratio, well removed from the reaction zone, is smaller than for cold HCl and decreases when flame temperature increases. The HCl/Cl equilibrium ratio is very large for even the hottest of these flames, and hence the limiting $36^+/35^+$ should represent the fragmentation patterns of HCl at these temperatures. This apparent effect of temperature on HCl fragmentation is




Fig. 1 - Line Reversal Flame Temperatures and Chlorine-Containing Species Profiles in Two Rich, H₂-O₂-N₂ Flames Directly Sampled at One Atmosphere

comparable to the effect on CO₂. Temperature effects can be significant, even for relatively small molecules, and could seriously affect results with unknown systems. Flames could conceivably be convenient media in which to study such effects.

There is not contradiction between the observations of high-temperature fragmentation effects and the claim that the flame gases are cooled extremely during beam formation,^{4/} since, in the latter case, reference is to translational temperatures. Translational, rotational and vibrational degrees of freedom do not equilibrate to the same extent during the expansion. As a first approximation, it is assumed that there is no vibrational relaxation in such rapid expansion, but that extensive rotational relaxation and nearly perfect translational relaxation^{3,6/} occur. Thus, the translationally and rotationally cold CO₂ and HCl are still vibrationally hot, and it is probable that the populations of the vibrational levels control the fragmentation behavior. The temperature-dependence of fragmentation could conceivably provide a rough way of detecting large changes in vibrational temperatures in partly equilibrated expansions. One would want to choose larger molecules with larger temperature effects, however.

B. Reactive Species

Another point of interest in Fig. 1 is the marked drop in the 36⁺/35⁺ ratio as the reaction zone is approached in these flames. Since the temperature is decreasing toward the reaction zone (the apparent increase in Flame U₅ of Fig. 1 is due to chemi-excitation effects), this drop can only be interpreted as being due to a significant Cl atom concentration. The abrupt rise in the 36⁺/35⁺ ratio at small distances occurs when the flame suddenly visibly attaches to the cool sampling probe, and the unburned gas mixture is being sampled. Sugden has argued that in this type of flame the reaction $H + HCl \rightleftharpoons H_2 + Cl$ will be much more quickly equilibrated than the recombination of the excess of H atoms known to be present in the reaction zone.^{15/} We can use the observed 36⁺/35⁺ ratios, after correction for fragmentation, to deduce the HCl/Cl ratio at each point in the flame. Knowing H₂ in these rich flames by measurement or calculation, one can calculate the H atom concentration. As previously mentioned, Fig. 1 should only be interpreted qualitatively because of the large cooled orifice which is used and the very narrow reaction zone at 1 atm. Nevertheless, it is of interest to compare the calculated ratio of $[H]/[H]_{eq}$ for several of these flames. Maximum ratios of 17, 320, 446 and 4,000 were calculated for the progressively cooler flames, F₃, F₅, U₅, and U₈, respectively. These values are comparable to determinations made by spectroscopic means,^{16/} and give qualitative support to the validity of the direct sampling scheme.

In addition to the Cl atom measurements shown above, the equilibrium $\text{Cl}_2 \rightleftharpoons 2\text{Cl}$ was studied in several cool CO-O₂-Ar flames. The results, reported in Ref. 4, agreed with literature values to within a factor of two.

The most demanding test of quenching was felt to be the quantitative sampling of the species O, OH, and H. Oxygen atoms were studied first in CO-O₂ and H₂-O₂ flames identical to those in Table I. At 50 ev. ionizing electron energy, fragmentation of CO₂ and O₂ to O completely swamped the 16^+ from O atoms. At a nominal 20 ev. and at room temperature the fragmentation of CO₂, CO, and O₂ to O⁺ was negligible. In view of the simultaneous uncertainties in ability to sample O atoms and the suspected temperature-dependent fragmentation of O₂, CO₂, H₂O, or CO, the correlations of the O⁺ data from the several flames with the equilibrium calculations provide the best evidence that fragmentation was negligible at 20 ev. Table II shows the results for the four CO-O₂ flames studied.

TABLE II

OXYGEN DISSOCIATION EQUILIBRIA MEASURED BY DIRECT SAMPLING
OF SEVERAL CO-O₂ FLAMES AT ONE ATMOSPHERE

Flame Composition (Mole Ratios)		Exptl. Temp. (°K)	O → 1/2 O ₂	
CO	O ₂		Exptl.	Calc.
			K _{eq}	
1	1	2769	24.5	20.4
4	3	2824	22.0	15.9
2	1	2834	22.5	15.1
3	1	2807	20.9	17.0

Figure 2 gives typical profiles in one of the flames. The effects of entrainment are noted at large distances while the excess O atoms near the reaction zone are a prominent feature. The partial pressures of oxygen atoms were determined to vary from 0.01 to 0.025 atm., assuming equal atom and molecule cross sections at 20 ev.^{17/} and making a mass separation correction. The equilibrium constants are in quite satisfactory agreement at the measured flame temperatures.

Oxygen atoms were also sampled from several H₂-O₂-N₂ flames burned on simple torch tips. The results are shown in Table III together with the adiabatic flame temperatures. The agreement between measured and calculated atom partial pressures is quite good over large differences in the O₂/O equilibrium ratios in the lean and stoichiometric flames. The excess O atom




Fig. 2 - Species Profiles in a Lean CO-O₂ Flame Directly Sampled at 2770°K and One Atmosphere

concentrations observed could well be due to incomplete equilibration since profiles were not taken in the torch experiments. A more careful study, including profile and temperature measurements, was made for the 2-4 flame using a sheathed burner. The O/O₂ ratio was very nearly the same as that shown in Table II. The O atom profile through the 2-4 flame is shown in Fig. 3.

TABLE III

OXYGEN ATOMS SAMPLED DIRECTLY FROM SEVERAL
ONE-ATMOSPHERE H₂-O₂-N₂ FLAMES

Flame Composition (Mole Ratios)			Adiabatic Flame Temp. (°K)	Partial Pressures in %			O \longrightarrow 1/2 O ₂	
H ₂	O ₂	N ₂		O		O ₂	K _{eq}	
				Exptl.	Lit.	Lit.	Exptl.	Calc.
2	1	1	2889	1.6	1.3	2.7	10.1	13.2
2	3		2720	3.3	2.8	ca 50.0	21.3	25.6
2	1	2	2709	0.8	0.5	1.5	16.1	27.0
2	4		2570	2.4	1.6	ca 60.0	32.3	50.0

Hydroxyl radicals were sampled from an even greater variety of flames, with varying temperatures and mixture ratios, than were oxygen atoms. The same problems exist for OH determinations as for O due to lack of knowledge of the high temperature fragmentation of H₂O to 17⁺. The results from a number of flames burned on a simple torch tip are shown in Table IV. The 18⁺/17⁺ ratios at 50 ev. show virtually no temperature or composition dependence and are in agreement with the 18⁺/17⁺ ratio from pure water at room temperature. If the temperature dependence of fragmentation is the same at 20 ev. as at 50 ev., then H₂O makes a negligible 17⁺ contribution at 20 ev. in these flames. The 20 ev. ratios of 18⁺/17⁺ correlate rather well with the equilibrium values as shown in the last two columns and indicate quantitative sampling of OH radicals.

In the four flames of Table IV for which O atom data were reported, the equilibrium $H_2O + O \rightleftharpoons 2OH$ could be tested. The results are given in Table V.




Fig. 3 - Oxygen Atom and Hydroxyl Radical Profiles in a Lean H_2-O_2 Flame Directly Sampled at 2540°K and One Atmosphere

TABLE IV

HYDROXYL RADICALS SAMPLED FROM SEVERAL ONE-ATMOSPHERE
H₂-O₂-N₂ AND CH₄-O₂ FLAMES

Flame Composition (Mole Ratios)				Adiabatic Flame Temp. (°K)	H ₂ O ⁺ /OH ⁺		Calc. H ₂ O/OH
H ₂	CH ₄	O ₂	N ₂		50 ev.	20 ev.	
	1	4		2845	2.5	8.3	4.0
4		1		2830	2.9	35.8	24.9
2		3		2720	2.6	11.9	6.5
2		1	2	2709	3.1	19.7	14.2
5		1		2627	3.3	52.3	66.7
	1	1		2611	3.2	37.3	64.0
2		4		2570	2.8	11.4	8.3
2		1	3	2524	2.8	21.7	26.6
2		1	4	2334	3.0	32.5	55.0
2		5		2330	2.8	14.6	15.7
	1	8		2330	2.5	9.8	13.1

TABLE V

EXPERIMENTAL DETERMINATION OF THE EQUILIBRIUM CONSTANT
FOR THE REACTION H₂O + O ⇌ 2OH IN SEVERAL
H₂-O₂-N₂ FLAMES AT ONE ATMOSPHERE

Flame Composition (Mole Ratios)			Adiabatic Flame Temp. (°K)	K _{eq} for H ₂ O + O ⇌ 2OH	
H ₂	O ₂	N ₂		Exptl.	Calculated
2	1	1	2889	0.50	0.52
2	3		2720	0.33	0.44
2	1	2	2709	0.50	0.43
2	4		2570	0.22	0.37

The agreement is good, showing that O and OH are equilibrated. The data of Table III show that neither O nor OH concentrations alone greatly depart from equilibrium. As with the O atom data, OH radicals were sampled more carefully from two flames on sheated burners: a 2-4 and a 5-1 H₂-O₂ flame. For both flames, results agreed with the torch results of Table III at the measured flame temperatures which were within 50°K of the adiabatic temperature. The OH profile appears in Fig. 3 for the 2-4 flame. The concentrations of both radicals peak sharply when the sampling orifice appears to just touch the reaction zone.

Some direct measurements of H atoms were made at 20 ev. in four flames varying from rich to stoichiometric. The H⁺ results are considerably less reliable than the O and OH values because of the low sensitivity caused by the mass separation effect. Still, as shown in Table VI, in three out of four flames the equilibrium constant agrees with literature values to within a factor of two. Excess H atoms were observed, as expected, near the reaction zone. As yet no study of the H atom in sheathed burners has been made because of the extreme heating of the present thin orifices in these hot, H₂-containing flames.

TABLE VI

EXPERIMENTAL DETERMINATION OF THE DISSOCIATION OF H₂ IN
SEVERAL H₂-O₂-N₂ FLAMES AT ONE ATMOSPHERE

Flame Composition (Mole Ratios)			Adiabatic Flame Temp. (°K)	1/2 H ₂ → H	
H ₂	O ₂	N ₂		Exptl. K _{eq}	Calc.
3	1		3004	0.068	0.155
2	1	1	2889	0.10	0.112
4	1		2835	0.018	0.100
5	1		2627	0.025	0.043

In one very hot torch flame, an attempt was made to measure every species of even minor importance. Table VII shows the ion intensities observed at 20 ev., after correction for mass separation and normalization to the theoretical H₂O value. Even without corrections for relative cross sections, true flame temperatures, mass spectrometer transmission, or detector efficiency, the results reflect the expected composition.

TABLE VII

RELATIVE ION INTENSITIES OBSERVED IN A ONE-ATMOSPHERE
H₂-O₂-N₂, 2-1-1 FLAME (T-2889°K) AT 20 EV. IONIZING ENERGY

<u>Ion Observed</u>	<u>Per Cent</u>	
	<u>Exptl.</u>	<u>Calc.</u>
H ⁺	0.75	3.2
H ₂ ⁺	2.25	8.8
O ⁺	0.42	1.3
OH ⁺	3.2	5.6
H ₂ O ⁺	(47.5)	47.5
N ₂ ⁺	14.7	29.9
NO ⁺	0.12	1.0
O ₂ ⁺	0.70	2.7

Results obtained from one other flame system are rather interesting. When a few per cent of either H₂S or SO₂ was added to a rich 5-1 H₂-O₂ flame, species profiles virtually identical to those shown in Fig. 4 were obtained. The initially very high concentrations of SO, S, and SO₂ and the late appearance of SH and SH₂ are noteworthy. Further study of this system should lead to new insight into the mechanisms of combustion of sulfur compounds. Similarly, in a lean 2-4 H₂-O₂ flame with SO₂ or H₂S added, the SO₂⁺/SO⁺ ratio was nearly constant through the flame and only slightly lower than for cold, pure SO₂. These observations agree with equilibrium calculations which show that SO₂ is the major species. It may be that, with more careful study of appearance potentials and fragmentation patterns, one can shed light on the postulated existence of an isomer of SO₂, perhaps with a peroxide bonding, believed to be formed in the early stages of the combustion of sulfur compounds.^{18/}

V. CONCLUSION

The sampling results presented in this paper show that at least some highly reactive species can be quantitatively sampled from 1-atm. flames. The departures from the literature of the measured values of equilibrium constants are virtually all within the estimated experimental error. Such success is to be expected on the basis of the calculated rate of the free jet expansion. For more precise work, more study is needed on flame-probe




Fig. 4 - Sulfur-Containing Species Profiles in a Rich One-Atmosphere, H_2-O_2 Flame to Which a Few Per Cent of Either H_2S or SO_2 Had Been Added

actions, orifice size effects, expansion dynamics, and a better understanding of the effect of temperature on the fragmentation processes.

Two major problems must be resolved before this flame sampling technique can be extended to high-temperature, condensable species. One is the mundane but troublesome problem of physical plugging of the cooled sampling orifice by condensed phases of the material being sampled. This problem can be reduced by the use of sufficiently large orifices and minimum metal concentrations. This difficulty will not be encountered in situations in which the sample and sampling orifice are in thermal equilibrium. The second problem is that of the nucleation or partial condensation of species during the expansion, particularly if the species are initially near saturation. This area is being studied in a continuation of work reported earlier.^{10/}

Direct mass spectrometric sampling of high pressure sources should be useful in many scientific and practical problems. Sampling of the reaction zone of 0.01 to 0.1-atm. flames is now under way in order to study the kinetics of the high temperature reactions of both combustion intermediates and various other species. The extension of this direct sampling technique to other relatively high pressure kinetic and equilibrium studies should also be fruitful.

REFERENCES

1. Greene, F. T., J. Brewer, and T. A. Milne, J. Chem. Phys., 40, 1488 (1964).
2. Ashkenas, H., and F. S. Sherman, "Free Jet Testing in Low Density Wind Tunnels," presented at Fourth International Symposium on Rarefied Gas Dynamics (proceedings to be published); Toronto, University of Toronto, July 14-17, 1964.
3. Anderson, J. B., R. P. Andres, J. B. Ferry, and G. Maise, "Studies of Low Density Supersonic Jets," presented at Fourth International Symposium on Rarefied Gas Dynamics (proceedings to be published); Toronto, University of Toronto, July 14-17, 1964.
4. Milne, T. A., and F. T. Greene, "Mass Spectrometric Sampling of One-Atmosphere Flames," Tenth Symposium (International) on Combustion, The Combustion Institute, 153 (1965).
5. Greene, F. T., and T. A. Milne, "Mass Spectrometric Sampling of High Pressure - High Temperature Sources," presented at the Mass Spectrometry Conference, Paris, September 14-18, 1964; to be published in Advances in Mass Spectrometry, Vol. III, Institute of Petroleum.
6. Scott, J. E., Jr., H. S. Morton, Jr., H. A. Phipps, and J. F. Moonan, "Distribution Function Measurements on Rarefied Gas Flow Through an Orifice," presented at Fourth International Symposium on Rarefied Gas Dynamics (proceedings to be published); Toronto, University of Toronto, July 14-17, 1964.
7. Valleau, J. P., and J. M. Deckers, Can. J. Chem., 42, 225 (1964); ibid. 43, 6 (1965).
8. Bier, K., and O. Hagen, "Optimum Conditions for Generating Molecular Beams by Nozzles," presented at Fourth International Symposium on Rarefied Gas Dynamics (proceedings to be published); Toronto, University of Toronto, July 14-17, 1964.
9. Knuth, E. L., Appl. Mech. Review, 17, 751 (1964).
10. Greene, F. T., and T. A. Milne, J. Chem. Phys., 39, 3150 (1963).

11. Leckenby, R. E., E. J. Robbins, and P. A. Trevalion, Proc. Roy. Soc., 280A, 409 (1964).
12. Bendix Time-of-Flight Mass Spectrometer, Model No. 12-101 with S-14-107 Ion Source, Bendix Corporation, Cincinnati, Ohio.
13. JANAF Interim Thermochemical Tables, The Dow Chemical Company, Midland, Michigan, December 31, 1960. Revision through June 30, 1964.
14. Padley, P. J., and T. M. Sugden, Proc. Roy. Soc., A246 (1958).
15. Phillips, L. F., and T. M. Sugden, Can. J. Chem., 38, 1804 (1960).
16. Bulewicz, E. M., C. G. James, and T. M. Sugden, Proc. Roy. Soc., A235, 89 (1956).
17. Fite, W., and T. Brackmann, Phys. Rev., 113, 815 (1959).
18. McGarvey, J., and W. McGrath, Proc. Roy. Soc., 280A, 490 (1964).

DOCUMENT CONTROL DATA - R&D

(Security classification of title, body of abstract and indexing annotation must be entered when the overall report is classified)

1 ORIGINATING ACTIVITY (Corporate author) Midwest Research Institute, Kansas City, Missouri		2a. REPORT SECURITY CLASSIFICATION Unclassified	
		2b. GROUP	
3 REPORT TITLE MASS SPECTROMETRIC STUDIES OF REACTIONS IN FLAMES - QUANTITATIVE SAMPLING OF FREE RADICALS FROM ONE-ATMOSPHERE FLAMES			
4 DESCRIPTIVE NOTES (Type of report and Inclusive dates) Technical Report - 16 July 1965			
5 AUTHOR(S) (Last name, first name, initial) Milne, T. A., and Greene, F. T.			
6. REPORT DATE 16 July 1965		7a. TOTAL NO. OF PAGES 19	7b. NO. OF REFS 18
8a. CONTRACT OR GRANT NO. Nonr-3599(CO)		9a. ORIGINATOR'S REPORT NUMBER(S)	
b. PROJECT NO. 092-513			
c.		9b. OTHER REPORT NO(S) (Any other numbers that may be assigned this report)	
d.			
10. AVAILABILITY/LIMITATION NOTICES Qualified requesters may obtain copies of this report from DDC.			
11. SUPPLEMENTARY NOTES		12. SPONSORING MILITARY ACTIVITY Advanced Research Projects Agency Washington, D. C.	
13. ABSTRACT Direct mass spectrometric sampling studies of one-atmosphere flames are described. A simple model of the sampling process is summarized which accounts for the effects and results obtained. Stable product analysis and evidence for appreciable temperature dependence of fragmentation patterns are shown. Quantitative sampling of the free-radicals H, O, OH and Cl from the burnt gas region of one-atmosphere flames is demonstrated, with direct observation of excess free radical concentrations in the reaction zones of these flames.			

14. KEY WORDS	LINK A		LINK B		LINK C	
	ROLE	WT	ROLE	WT	ROLE	WT
Mass Spectrometry Thermochemistry High Temperature Combustion Metal Containing Flames Direct Sampling Aerodynamic Molecular Beams						

INSTRUCTIONS

1. ORIGINATING ACTIVITY: Enter the name and address of the contractor, subcontractor, grantee, Department of Defense activity or other organization (*corporate author*) issuing the report.

2a. REPORT SECURITY CLASSIFICATION: Enter the overall security classification of the report. Indicate whether "Restricted Data" is included. Marking is to be in accordance with appropriate security regulations.

2b. GROUP: Automatic downgrading is specified in DoD Directive 5200.10 and Armed Forces Industrial Manual. Enter the group number. Also, when applicable, show that optional markings have been used for Group 3 and Group 4 as authorized.

3. REPORT TITLE: Enter the complete report title in all capital letters. Titles in all cases should be unclassified. If a meaningful title cannot be selected without classification, show title classification in all capitals in parentheses immediately following the title.

4. DESCRIPTIVE NOTES: If appropriate, enter the type of report, e.g., interim, progress, summary, annual, or final. Give the inclusive dates when a specific reporting period is covered.

5. AUTHOR(S): Enter the name(s) of author(s) as shown on or in the report. Enter last name, first name, middle initial. If military, show rank and branch of service. The name of the principal author is an absolute minimum requirement.

6. REPORT DATE: Enter the date of the report as day, month, year; or month, year. If more than one date appears on the report, use date of publication.

7a. TOTAL NUMBER OF PAGES: The total page count should follow normal pagination procedures, i.e., enter the number of pages containing information.

7b. NUMBER OF REFERENCES: Enter the total number of references cited in the report.

8a. CONTRACT OR GRANT NUMBER: If appropriate, enter the applicable number of the contract or grant under which the report was written.

8b, 8c, & 8d. PROJECT NUMBER: Enter the appropriate military department identification, such as project number, subproject number, system numbers, task number, etc.

9a. ORIGINATOR'S REPORT NUMBER(S): Enter the official report number by which the document will be identified and controlled by the originating activity. This number must be unique to this report.

9b. OTHER REPORT NUMBER(S): If the report has been assigned any other report numbers (*either by the originator or by the sponsor*), also enter this number(s).

10. AVAILABILITY/LIMITATION NOTICES: Enter any limitations on further dissemination of the report, other than those

imposed by security classification, using standard statements such as:

- (1) "Qualified requesters may obtain copies of this report from DDC."
- (2) "Foreign announcement and dissemination of this report by DDC is not authorized."
- (3) "U. S. Government agencies may obtain copies of this report directly from DDC. Other qualified DDC users shall request through _____."
- (4) "U. S. military agencies may obtain copies of this report directly from DDC. Other qualified users shall request through _____."
- (5) "All distribution of this report is controlled. Qualified DDC users shall request through _____."

If the report has been furnished to the Office of Technical Services, Department of Commerce, for sale to the public, indicate this fact and enter the price, if known.

11. SUPPLEMENTARY NOTES: Use for additional explanatory notes.

12. SPONSORING MILITARY ACTIVITY: Enter the name of the departmental project office or laboratory sponsoring (*paying for*) the research and development. Include address.

13. ABSTRACT: Enter an abstract giving a brief and factual summary of the document indicative of the report, even though it may also appear elsewhere in the body of the technical report. If additional space is required, a continuation sheet shall be attached.

It is highly desirable that the abstract of classified reports be unclassified. Each paragraph of the abstract shall end with an indication of the military security classification of the information in the paragraph, represented as (TS), (S), (C), or (U).

There is no limitation on the length of the abstract. However, the suggested length is from 150 to 225 words.

14. KEY WORDS: Key words are technically meaningful terms or short phrases that characterize a report and may be used as index entries for cataloging the report. Key words must be selected so that no security classification is required. Identifiers, such as equipment model designation, trade name, military project code name, geographic location, may be used as key words but will be followed by an indication of technical context. The assignment of links, roles, and weights is optional.