UNCLASSIFIED

AD NUMBER

AD471208

LIMITATION CHANGES

TO:

Approved for public release; distribution is unlimited.

FROM:

Distribution authorized to U.S. Gov't. agencies and their contractors;

Administrative/Operational Use; 10 AUG 1965. Other requests shall be referred to Office of Naval Research, One Liberty Center, 875 North Randolph Street, Arlington, VA 22203-1995.

AUTHORITY

ONR ltr dtd 13 Sep 1977

THIS PAGE IS UNCLASSIFIED

THIS REPORT HAS BEEN DELIMITED AND CLEARED FOR FUBLIC RELEASE UNDER DOD DIRECTIVE 5200.20 AND NO RESTRICTIONS ARE IMPOSED UPON ITS USE AND DISCLOSURE.

DISTRIBUTION STATEMENT A

APPROVED FOR PUBLIC RELEASE; DISTRIBUTION UNLIMITED,

BEST AVAILABLE COPY

•

SECURITY MARKING

The classified or limited status of this report applies to each page, unless otherwise marked. Separate page printouts MUST be marked accordingly.

THIS DOCUMENT CONTAINS INFORMATION AFFECTING THE NATIONAL DEFENSE OF THE UNITED STATES WITHIN THE MEANING OF THE ESPIONAGE LAWS, TITLE 18, U.S.C., SECTIONS 793 AND 794. THE TRANSMISSION OR THE REVELATION OF ITS CONTENTS IN ANY MANNER TO AN UNAUTHORIZED PERSON IS PROHIBITED BY LAW.

NOTICE: When government or other drawings, specifications or other data are used for any purpose other than in connection with a definitely related government procurement operation, the U. S. Government thereby incurs no responsibility, nor any obligation whatsoever; and the fact that the Government may have formulated, furnished, or in any way supplied the said drawings, specifications, or other data is not to be regarded by implication or otherwise as in any manner licensing the holder or any other person or corporation, or conveying any rights or permission to manufacture, use or sell any patented invention that may in any way be related thereto.

ARPA Order No. 529



by

Michael J. Haggerty

Research Report No. PIBMRI-1289-65 Contract No. NONR 839(38)

Sponsored by

Advanced Research Projects Agency

Monitored by

Office of Naval Research Washington, D.C. August 10, 1965



POLYTECHNIC INSTITUTE OF BROOKLYN

MICROWAVE RESEARCH INSTITUTE DEPARTMENT OF PHYSICS ARPA Order No. 529

Research Report No. PIBMR1-1289-65 Contract No. NONR-839(38)

ON THE KINETIC THEORY OF DIFFUSION OF A PLASMA

COLUMN ACROSS A MAGNETIC FIELD

by

Michael J. Haggerty

Polytechnic Institute of Brooklyn Department of Physics 333 Jay Street Brooklyn 1, New York

Research Report No. PIBMRI-1289-65 Contract No. NONR-839(38)

August 10, 1965

Title Page Acknowledgement Summary Table of Contents 32 Pages of Text 1 Page of References

Michael J. Hagerty

Research Associate, Physics

mets dhe

H. J. Juretschke Acting Head, Dept. of Physics

> Reproduction in whole or in part is permitted for any purpose of the United States Government.

> > Sponsored by

Advanced Research Projects Agency

Monitored by

Office of Naval Research Washington, D.C.

Acknowledgments

The formal solution given in Section 3 was first derived (in a slightly different form) in 1962, as part of a doctoral program at the University of British Columbia, under the active supervision of Dr. L. G. de Sobrino. At that time, I was financially supported by a National Research Council of Canada Studentship.

More secently, I am indebted to Dr. T.-Y. Wu and Dr. D. D. H. Yee for many stimulating conversations. The sesarch was supported by the Office of Naval Research, Washington, D. C. under Contract No. NONR 839(38); ARPA Order No. 529.

Summary

The kinetic theory of Prigogine and Balescu, which was previously developed for a homogeneous plasma in a magnetic field, is now applied to the problem of diffusion of a plasma column across a magnetic field. The problem of deriving a kinetic equation is reduced to that of solving a two dimensional Fredholm-type integral equation with a complicated kernel. No restriction is made concerning the amplitude or length scale of the inhomogeneity across the magnetic field. The results are preliminary in nature, and suggestions are made for further refinements. Some comparisons with previously existing results are given.

TABLE OF CONTENTS

		Page	
Ackno	owledgments	ii	
Summa	ary	iii	
Table	of Contents	iv	
Section	n		
1.	Introduction	1	
2.	Physical Mcdel	2	
3.	Formal Solution of the Liouville Equation	4	
4.	Initial Conditions	8	
5.	Selection of the Dominant Terms	11	
6.	The Singular Integral Equation	16	
7.	The Fredholm-type Integral Equations	20	
8.	Comparison of Results with Existing Kinetic Equations	27	
9.	Conclusion	30	
Refere	References		

iv

SECTION 1

INTRODUCTION

The phenomenon of diffusion of a plasma across a magnetic field is of great importance in the problem of plasma confinement. There have been many theoretical papers on the subject; see for example the review articles by Golant⁽¹⁾ and Boeschoten⁽²⁾. So far, however, there has been very little work done on the kinetic theory of inhomogeneous nonequilibrium plasmas in magnetic fields.

We report here some pieliminary results on the diffusion of a plasma column across a uniform magnetic field. It is believed that no equivalent results have yet been published. The problem of deriving an equation for the one particle distribution function, including a proper treatment of screening effects, has been reduced to that of solving an integral equation of the F. edholm type, with a complicated kernel. The method used is an extension of that of Prigogine, Balescu, and coworkers.^{(3),(4)} It was applied to the homogeneous problem in preceding works ^{(5), (6), (7)}, to be referred to as papers I, II, and III respectively, We rely heavily on the notation summarized in the Appendix of paper I (although some superscripts \bot and z are suppressed, and $x_{j} \stackrel{g}{=}$ is replaced by \underline{O}). Some familiarity with paper I or with Balescu's book⁽⁴⁾ is necessary for a detailed examination of this report; the emphasis is on the differences between the present calculation and that of paper I.

Although the plasma is assumed to be homogeneous along the magnetic field, an arbitrary inhomogeneity is allowed in directions perpendicular to the field. The infinite order perturbation theory is tractable for this case because the guiding centers are constants of the unperturbed motion. We have, however, restricted ou selves here to the calculation of local effects in a single species plasma. It is an important problem to refine this model, and to properly take into account internal macroscopic electric fields.

More details concerning the physical model are given in Section 2. The diagram perturbation theory is summarized in Sections 3, 4, and 5. The main results are given in Sections 6 and 7. They are compared with existing kinetic equations in Section 8.

SECTION 2

PHYSICAL MODEL

The theory presented here is incomplete in the sense that a very primitive physical model is used.

We consider a single-species plasma in a region of volume Ω containing a uniform magnetic field <u>B</u>. The length of the region in the direction of <u>B</u> is Λ , and the cross section is $\Sigma = \Omega/\Lambda$. We will assume that the plasma is homogeneous in the direction of <u>B</u>. In addition, the limit

$$N \rightarrow \infty, \Lambda \rightarrow \infty, N/\Lambda \text{ constant},$$

 $\Sigma \text{ arbitrary}$ (2.1)

(cf. eq. (I.21)) will be taken. Thus end effects (for example, "short circuiting") are not considered. However, an arbitrary inhomogeneity in directions perpendicular to <u>B</u> will be allowed. There will be no restrictions on the amplitude or length scale of the inhomogeneities, although these scales may effect the plausibility of some of the other assumptions.

A neutralizing background charge of unspecified density variation across <u>B</u> is assumed to be present. This is an ill-defined artifice introduced solely for the purpose of restricting ourselves to the calculation of effects due to relatively <u>local</u> interactions among the plasma particles. The long range interactions and boundary effects which the perturbation expansion of paper I cannot take into account are assumed to effectively cancel each other: we again make assumption (I. 3. 26). We use the Hamiltonian (I. 2. 2) with W set equal to zero.

In the homogeneous case discussed in paper I, the generalization to a more realistic multi-component plasma would involve only the inclusion of additional indices and summation signs in appropriate places. In the present case, however, such a generalization may be a less trivial problem. It would involve a careful treatment of the transverse electric fields^{*} arising from the charge separation due to different diffusion rates of the

*Note that for the noninteracting case, the orbits in crossed <u>external</u> electric and magnetic fields are very simple; see for example Reference (8), Section 10.1.

various plasma components. This should lead to the development of a theory of ambipolar diffusion. While such a treatment is important, it is beyond the scope of the work reported here.

We will also assume that the one particle distribution function is initially "gyrotropic"; that is, isotropic in directions perpendicular to <u>B</u>. This may restrict the types of diffusion that the theory can predict. However, the theorem that the initial condition is preserved in time is much stronger than the corresponding result for the homogeneous plasma. In that case, once it became apparent that the higher order initial correlations would not appear in the final results, then there was no way in which a preferred direction perpendicular to <u>B</u> could appear. In the present case, we always have the preferred directions corresponding to the spatial density gradients.

SECTION 3

FORMAL SOLUTION OF THE

LIOUVILLE EQUATION

Our starting point is the same as in Section I. 3, except that we do not introduce periodic boundary conditions with respect to the $\{\underline{x}_j \ g^{\perp}\}\$ (hereafter called $\{\underline{Q}, \}\$). The fundamental reason why the treatment of arbitrary inhomogeneities perpendicular to <u>B</u> is tractable has already been noted in the paragraph following eq. (I. 3.13). The $\{\underline{Q}_j\}\$ are constants of the unperturbed motion, and hence can be created in the same way as the velocity cylindrical components $\{\underline{a}_j, \underline{p}_j^{\perp}\}\$. (From now on, the superscript z will be suppressed.) We had no reason to assume any sort of homogeneity in velocity space (except through the condition of stability); hence none is needed with respect to $\{\underline{Q}_i\}\$.

Thus eqs. (I. 3.14) and (I. 3.18) are replaced by

$$f_{N}(t) = f_{N}(\{z_{j}\}, \{\underline{\alpha}_{j}\}, \{\underline{\alpha}_{j}\}, \{\underline{\alpha}_{j}\}, \{\underline{\alpha}_{j}\}, \{\underline{\alpha}_{j}\}, t\})$$

$$= \sum_{k \in [n]} \sum_{j \in [n]} \rho' (t) \exp[i\sum_{j} (k_{j}z_{j} - n_{j}\theta_{j})], (3.1)$$

$$\{k \notin [n]\} \quad \{k \notin [n]\} \quad j \in [n]$$

$$= \rho'_{\{k}(0)\{\{n^{(0)}\}\} \quad [(\underline{\alpha}_{j}\}, \{\underline{\alpha}_{j}\}, \{\underline{\alpha}_{j}\}, \{\underline{\alpha}_{j}\}, t)$$

$$= -\frac{1}{2\pi i} \oint_{C} d\zeta \exp[-i\zeta t] \sum_{q=0}^{\infty} \{(-e^{2})^{q}$$

$$[(\sum_{j} k_{j}^{(0)} p_{j}/m) + u(\sum_{i} n_{j}^{(0)}) - \zeta]^{-1}$$

$$\sum_{\{k \in [1]\}\{n^{(1)}\}} \sum_{h < u} <\{k_{j}^{(0)}\} \{n_{j}^{(0)}\} |\delta L_{hu}| \ \{k_{j}^{(1)}\} \{n_{j}^{(1)}\} >$$

$$\begin{bmatrix} i \sum_{j} k_{j}^{(1)} p_{j} / m \} + w (\sum_{j} n_{j}^{(1)}) - \zeta \end{bmatrix}^{-1} \\ \cdots \sum_{i \neq k} \sum_{q \neq j} \sum_{n(q)} \sum_{i < m} \\ < \{ k_{j}^{(q-1)} \} \{ n_{j}^{(q-1)} \} | \delta L_{im} | \{ k_{j}^{(q)} \} \{ n_{j}^{(q)} \} > \\ \begin{bmatrix} (\sum_{j} k_{j}^{(q)} p_{j} / m) + w (\sum_{j} n_{j}^{(q)}) - \zeta \end{bmatrix}^{-1} \rho'_{\{k}(q) \} \{ n(q) \} (0) .$$
(3.2)

Here, the k's are restricted to values of the form $(2\pi/\hbar)$ (integer), and the n's are integers.

The matrix elements are defined as follows:

<
$$k_{j}$$
 { n_{j} } δL_{hu} { k'_{j} } n'_{j} >
= $\Lambda^{-N} (2\pi)^{-N} \int d^{N}z d^{N}\theta$

$$\exp\left[-i\sum_{j} (k_{j}z_{j} - n_{j}\theta_{j})\right] \delta L_{hu} \exp\left[i\sum_{j} (k_{j}'z_{j} - n_{j}'\theta_{j})\right].$$

They are operators with respect to $\{Q_j\}, \{a_j\}, and \{p_j\}\}$. An explicit expression analogous to eq. (I. 3. 25) can be calculated in the same way:

<
$$\{k_{j}\}$$
 $\{n_{j}\}$ $|\delta L_{hu}|$ $\{k'_{j}\}$ $\{n'_{j}\}$ >
= $[\prod_{j(\ddagger h, u)} \delta^{Kr}(k_{j} - k'_{j}) \delta^{Kr}(n_{j} - n'_{j})]$

 $\delta^{\mathbf{k} \mathbf{r}} (\mathbf{k}_{\mathbf{h}} + \mathbf{k}_{\mathbf{u}} - \mathbf{k}_{\mathbf{h}}' - \mathbf{k}_{\mathbf{u}}') \int_{\mathbf{0}}^{\infty} d\mathbf{l} \int_{\mathbf{0}}^{2\pi} d\beta$

6

$$[-\pi m \omega \Lambda (\ell^{2} + [k_{h} - k_{h}']^{2} + \chi^{2})]^{-1}$$

$$\exp\left[i(n_{h} + n_{u} - n_{h}' - n_{u}')\beta\right] \left\{J_{n_{h}} - n_{h}'(l_{a_{h}})\right\}$$

$$J_{-(n_{u} - n_{u}')}(k_{u}) [mw(k_{h} - k_{h}')(\frac{\partial}{\partial p_{h}} - \frac{\partial}{\partial p_{u}})$$

$$\left(\underline{\ell} \times \frac{\partial}{\partial \underline{Q}_{hu}}\right)^{z} + \frac{n_{h}}{a_{h}} \frac{\partial}{\partial a_{h}} + \frac{n_{u}}{a_{u}} \frac{\partial}{\partial a_{u}}\right]$$

$$J_{-(n_u - n'_u)}(la_u) \frac{n'_h}{a_h} \frac{\partial}{\partial a_k} J_{n_h - n'_h}(la_h)$$

$$-J_{n_{h}} - n_{h}' (la_{h}) \frac{n'_{u}}{a_{u}} \frac{\partial}{\partial a_{u}} J_{-(n_{u}} - n'_{u}) (la_{u}) \}$$

 $\exp\left[i\,\underline{l}\cdot\,\underline{Q}_{hu}\right],\tag{3.3}$

where $\underline{Q}_{hu} = \underline{Q}_{h} - \underline{Q}_{u}$, and $\underline{\ell}$ is the vector defined by the polar coordinates ℓ , B ($\underline{\ell} \cdot \underline{B} = 0$).

It is possible to perform the integration over β . Although the result will not be used, we write it for completeness:

$$< \{k_{j}\} \{n_{j}\} |\delta L_{hu}| \{k_{j}\} \{n_{j}'\} >$$

$$= \left[\prod_{j(\ddagger h, u)} \delta^{\mathbf{K}\mathbf{r}} \left(\mathbf{k}_{j} - \mathbf{k}_{j}' \right) \delta^{\mathbf{K}\mathbf{r}} \left(\mathbf{n}_{j} - \mathbf{n}_{j}' \right) \right]$$

PIBMRI _1289-65

 $\delta^{\mathbf{k}_{\mathrm{I}}}(\mathbf{k}_{\mathrm{h}}^{\dagger}+\mathbf{k}_{\mathrm{u}}^{\dagger}-\mathbf{k}_{\mathrm{h}}^{\prime}-\mathbf{k}_{\mathrm{u}}^{\prime})\ 2\pi\int_{0}^{\infty} \mathbf{l}d\mathbf{l}$

$$[-\pi m \omega \Lambda \mathscr{U}^{2} + [k_{h} - k_{h}']^{2} + {\binom{2}{n}}^{-1}$$

$$\int [m_{u}(k_{h} - k_{h}')(\frac{\partial}{\partial p_{h}} - \frac{\partial}{\partial p_{u}}) - \frac{n_{h}'}{a_{h}} \frac{\partial}{\partial a_{h}}$$

$$\frac{n'_{u}}{a_{u}} \frac{\lambda}{\lambda a_{u}} + \frac{n_{h} + n_{u} - n'_{h} - n'_{u}}{Q_{hu}} \frac{\lambda}{\lambda Q_{hu}}$$

$$+ \frac{1}{Q_{hu}} \frac{J'_{n_{h}} + n_{u} - n'_{h} - n'_{u}}{J_{n_{h}} + n_{u} - n'_{h} - n'_{u}} \frac{(\mathcal{Q}_{hu})}{(\mathcal{Q}_{hu})} \frac{\lambda}{\lambda \Phi_{hu}}]$$

$$(J_{n_{h}} - n'_{h}) - (n_{u} - n'_{u}) - (n_{u}$$

$$J_{-(n_{u} - n_{u}')}(n_{u}) \left[\frac{n_{h}}{a_{h}} \frac{\lambda}{\partial a_{h}} + \frac{n_{u}}{a_{u}} \frac{\lambda}{\partial a_{u}}\right]$$

$$J_{n_{h}} + n_{u} - n'_{h} - n'_{u} (\log_{hu}) \exp \left[i(n_{h} + n_{u} - n'_{h} - n'_{u})(\phi_{hu} + \frac{\pi}{2})\right], \quad (3.4)$$

where $\phi_{hu} = \arctan (Q_{hu}^y / Q_{hu}^x)$ (cf. eq. (I. 3.8) ff).

We also note that, according to Barabanenkov⁽⁹⁾, eqs. (6) and (7), the integration over \mathcal{L} is simple if $Q_{hu} > a_h + a_u$. One should however check the range of validity of those formulae before applying them.

SECTION 4

INITIAL CONDITIONS

For the physical model described in Section 2, the arguments in Sections I. 4 and I. 5 go through almost without change. All that is necessary is to replace Ω or $\Omega/8\pi^3$ by Λ or $\Lambda/2\pi$ in various places, and to include integrations over $\{Q_j\}$ with the velocity integrals.

The reduced distribution functions can be defined as before:

$$\mathbf{f}_{r,s}(\mathbf{z}_1,\ldots,\mathbf{z}_r; \{\mathbf{Q}_\sigma\}, \{\mathbf{a}_\sigma\}, \{\mathbf{\theta}_\sigma\}, \{\mathbf{p}_\sigma\}, \mathbf{f}_\sigma\})$$

$$= \frac{N!}{(N-r)!} \int_{(1,\dots,r)}^{\prime} d^{N-r} z \int_{\{\sigma\}}^{\prime} d^{N-s} Q d^{N-s} a d^{N-s} \theta d^{N-s} p$$

$$\begin{bmatrix} N \\ \Pi \\ h=s+1 \end{bmatrix} m^{2} \omega^{2} a_{h} f_{N}(\{z_{j}, \{Q_{j}\}, \{Q_{j}\},$$

$$\varphi_{\mathbf{s}} = \mathbf{f}_{\mathbf{o}, \mathbf{s}}, \varphi = \varphi_{\mathbf{1}} = \mathbf{f}_{\mathbf{o}, \mathbf{1}}; \ 0 \le \mathbf{r} \le \mathbf{s} \le \mathbf{N};$$

with the normalization condition

$$\int d^{s} Q d^{s} a d^{s} \theta d^{s} p (\prod m^{2} \omega^{2} a_{\sigma}) \varphi_{s}(\{\sigma\}, t) = 1.$$
(4.2)

They can be expressed in terms of the ρ' 's:

$$f_{r,s} = \Lambda^{N} (N/\Lambda)^{r} \sum_{k_{1} \cdots k_{r}} \sum_{\substack{n_{\sigma} \\ n_{\sigma} \\ k_{1} \cdots k_{r} \\ n}} \mathcal{Y}_{N-s}$$

$$\rho' = \lim_{k_{1} \cdots k_{r}} \{\sigma\} = \exp(-i\sum_{\sigma} n_{\sigma} \theta_{\sigma})$$

$$\exp(i[k_{1}z_{1} + \cdots + k_{r}z_{r}]),$$

(4.3)

which is analogous to eq. (I. 4. 7). Here,

$$\mathcal{Y}_{N-s} f = \int_{\{\sigma\}}^{\prime} d^{N-s} Q d^{N-s} a d^{N-s} p \left[\prod_{h=s+1}^{N} 2 \pi m^2 \omega^2 a_h \right] f .$$
(4.4)

The Ursell correlation functions $U_{s}(\{\sigma\})$ are defined by eqs. (I.4.11), with Ω replaced by Λ .

Our object is to obtain equations for the one particle distribution function φ . Initial conditions analogous to those of Section I.5 are imposed:

$$f_{1,s}(0) = f_{r,r}(0) \prod_{\tau=r+1}^{s} \varphi(Q_{\tau}, a_{\tau}, \theta_{\tau}, p_{\tau}; 0), \qquad (4.5)$$

$$\varphi(\underline{Q}_{\tau}, \underline{a}_{\tau}, \underline{\theta}_{\tau}, \underline{p}_{\tau}; 0) = \varphi(\underline{Q}_{\tau}, \underline{a}_{\tau}, \underline{p}_{\tau}; 0), \qquad (4.6)$$

 $U_s({\sigma}, 0)$ is finite in the limit (2.1), and

$$\lim_{\Lambda \to \infty} U_{s}(\ldots,\alpha,\ldots,\beta,\ldots;0) = 0 \text{ for } |z_{\alpha} - z_{\beta}| >> \pi_{corr}, \qquad (4.7)$$

$$f_N(\{z_j\},\ldots,0) = f_N(\{z_j+Z\},\ldots,0).$$
 (4.8)

It is assumed that there are no initial correlations among the guiding centers or velocities which are independent of the distances $\{ |z_i - z_j| \}$, in the limit $\Lambda \to \infty$.

The condition (4.6) is stronger than for the homogeneous case, because of the preferred directions corresponding to density gradients across <u>B</u>. However, a strong theorem can be established; namely, that the condition is preserved in time in the modified weak coupling approximation.

We can then easily establish the analogue of the result (I. 5.22): If

$$\rho \frac{1...r}{k...q} \begin{cases} \sigma \\ n \end{cases} \equiv \Lambda^{N} (\Lambda/2\pi)^{r-w} \rho' \frac{1...r}{k...q} \begin{cases} \sigma \\ n \end{cases},$$

$$k+...+q = 0, \ k \neq 0, \dots, q \neq 0, \qquad (4.9)$$

$$k+...+q = 0, k \neq 0, ..., q \neq 0,$$

then

 $y_{N-s}^{1...r} {\sigma}$ $(2\pi/\Lambda)^{W}$ (0) remains finite in the limit (2.1), (4.10)where w and w' have the same meaning as before. This follows from expressions such as that analogous to (I. 5. 13):

$$U_{2}(1, 2; 0) = (2\pi/\Lambda) \sum_{k(\ddagger 0)} \sum_{n} \mathcal{Y}_{N-2}$$

$$\rho_{k}^{1} - k \frac{2}{n_{1}} \frac{2}{n_{2}} = (0) \exp[-i \sum_{\sigma} n_{\sigma} \theta_{\sigma}] \exp[ik(z_{1} - z_{2})]. \qquad (4.11)$$

SECTION 5

SELECTION OF THE DOMINANT TERMS

The diagram representation of the formal solution (3.1), (3.2), (3.3) is almost identical to that for the homogeneous case. The "semi-connection" property (I. 77) remains valid, because although there is an extra term in (3.3) containing $\chi \partial/\partial Q_{hu}$, there is no dependence on Q_h or Q_u to the left of this factor.

By calculating the Λ dependences of the various diagram contributions to $\begin{cases} \sigma \\ N-s^{\rho} \\ k \\ n \end{cases}$ (t) (using condition (4.10)), we can establish that condition (4.10) remains valid for t > 0.

As for the e² dependence of the initial distributions, there is no essential change. In equations such as (I. 7. 20), one replaces $\int d^{N} \underline{x}^{g} / \Omega^{N}$ by $\int d^{N} z / \Lambda^{N}$, and $(4\pi^{2}/\Sigma) \sum_{n} by \int d \underline{a}^{n}$. (It is then possible to integrate over 8 to obtain a simpler

result; cf. eq. (3.4)) We therefore assume that

$$\rho'_{k} \{n\}^{(0)} \propto (e^2)^{r-w-w'} \rho'_{0} \{0\} \{0\}^{(0)}, \qquad (5.1)$$

which is similar to eq. (I. 7. 24).

Let us write the equation for the contribution to $\varphi(\alpha, t)$ corresponding to (I. 8. 3):

$$\varphi(\alpha, t) = \ldots - \mathcal{Y}_{N-1} (2\pi i)^{-1} \oint_C d\zeta \exp(-i\zeta t)$$

$$(-\zeta)^{-1} \sum_{k} \sum_{\substack{n_{\alpha}^{(1)} \\ \alpha}} \sum_{n_{h}^{(1)}} (-Ne^{2})$$

$$< \{0\} \{0\} | \delta L_{h\alpha} | k_h = -k, k_\alpha = k, n_\alpha^{(1)}, n_h^{(1)}, \{0\} ' >$$

$$(k[p_{\alpha} - p_{h}]/m + w[n_{\alpha}^{(1)} + n_{h}^{(1)}] - \zeta)^{-1} \sum_{\substack{n_{u}^{(2)}}} (-Ne^{2})$$

$$< k_{h} = -k, k_{\alpha} = k, n_{\alpha}^{(1)}, n_{h}^{(1)}, \{0\}' | \delta L_{uh} | k_{u} = -k, k_{\alpha} = k,$$

$$n_{\alpha}^{(1)}, n_{u}^{(2)}, \{0\}' > (k[p_{\alpha} - p_{u}]/m + w[n_{\alpha}^{(1)} + n_{u}^{(2)}] - \zeta)^{-1}$$

$$(-e^2) < k_u = k, k_a = k, n_a^{(1)}, n_u^{(2)}, \{0\}' | \delta L_{au} | \{0\} \{0\} > 0$$

$$(-\mathfrak{Y}^{-1})^{\circ} \{0\} \{0\}^{(0)},$$
 (5.2)

where $\{0\}'$ means that all suppressed variables are zero. We replace $n_{\alpha}^{(1)}$, $n_{u}^{(2)}$ by $n_{\alpha}^{}$, $-n_{h}^{}$, $-n_{u}^{}$, and make the following abbreviations in notation:

< 0
$$|\alpha h| > \equiv - Ne^{2}i \sum_{\substack{n_{\alpha} \\ n_{h}}} \sum_{\substack{n_{\alpha} \\ n_{h}}}$$

$$< \{0\} \{0\} \{\delta L_{h\alpha} | k_{h} = -k, k_{\alpha} = k, n_{\alpha}^{(1)} = n_{\alpha}, n_{h}^{(1)} = -n_{h}, \{0\} '>$$

(with certain terms that vanish under y_{N-1} omitted)

$$\equiv \sum_{\substack{n_{\alpha} \ n_{h} \ o}} \sum_{\substack{h \ d}} \int_{h}^{2\pi} \frac{2\pi}{d\beta_{h}} \exp(i[n_{n} - n_{\alpha}]\beta_{h})$$

$$\frac{e^{2}N \exp(i \underline{\mu}_{h} + \frac{Q}{h})}{i\pi m w \Lambda (\mu_{h}^{2} + \mu_{h}^{2} + \mu_{h}^{2})} [mwk \frac{\partial}{\partial p_{0}}]$$

$$-\left(\underline{\mathscr{L}}_{h} \times \frac{\partial}{\partial \underline{Q}_{\alpha}}\right)^{z} + \frac{n_{\alpha}}{a_{\alpha}} \frac{\partial}{\partial a_{\alpha}} \int_{n_{\alpha}} \left(\mathcal{L}_{h} a_{\alpha} \right) \int_{n_{h}} \left(\mathcal{L}_{h} a_{h} \right), \qquad (5.3)$$

<
$$|hu| > = -Ne^2 i \sum_{\substack{n \\ u}} < k_h = -k, k_a = k,$$

$$n_{\alpha}^{(1)} = n_{\alpha}, n_{h}^{(1)} = -n_{h}, \{0\}' | \delta L_{uh} | k_{u} = -k, k_{\alpha} = k, n_{\alpha}^{(2)} = n_{\alpha}, n_{u}^{(2)} = -n_{u}, \{0\}' > 0$$

(with similar terms omitted)

 $\equiv \sum_{\substack{n_{u} \ o}} \int_{u}^{\infty} \int_{u}^{u} d \int_{u}^{2\pi} d\beta_{u} \exp(i[n_{u} - n_{h}]\beta_{u})$

$$\frac{e^{2}N\exp(i \underbrace{\pounds_{u}} \underbrace{Q_{uh}})}{i\pi m \omega \Lambda (\underbrace{\mu_{u}}^{2} + k^{2} + \kappa^{2})} J_{n}(\underbrace{\mu_{u}}_{h} \underbrace{J_{n}}_{u}(\underbrace{\mu_{u}}_{u}))$$

$$[mwk\frac{\partial}{\partial p_{h}} - (\underline{\mathscr{L}}_{u} \times \frac{\partial}{\partial \underline{Q}_{h}})^{z} + \frac{n_{h}}{a_{h}} \frac{\partial}{\partial a_{h}}], \qquad (5.4)$$

$$< |u\alpha| 0 > \equiv -\frac{\Lambda e^2 i}{2\pi} < k_u = -k, k_a = k,$$

$$n_{\alpha}^{(2)} = n_{\alpha}, n_{u}^{(2)} = -n_{u}, \{0\}' | \delta L_{\alpha u} | \{0\} \{0\} >$$

$$\equiv \int_{0}^{\infty} \int d\mathcal{L} d\mathcal{L} \int_{0}^{2\pi} d\beta \exp \left(i\left[n_{\alpha} - n_{u}\right]^{\beta}\right)$$

$$\frac{e^{2} \exp(i \pounds Q_{\alpha u})}{2i\pi^{2} m \omega (\ell^{2} + k^{2} + \chi^{2})} J_{n_{u}} (\ell a_{u}) J_{n_{\alpha}} (\ell a_{\alpha})$$

$$\cdot \left[m \omega k \left(\frac{\partial}{\partial p_{u}} - \frac{\partial}{\partial p_{\alpha}} \right) - \left(\underbrace{\mathscr{L}}_{\alpha} \times \frac{\partial}{\partial Q_{u\alpha}} \right)^{z} + \frac{n_{u}}{a_{u}} \frac{\partial}{\partial a_{u}} - \frac{n_{\alpha}}{a_{\alpha}} \frac{\partial}{\partial a_{\alpha}} \right].$$
(5.5)

Then eq. (5.2) becomes:

$$\omega(\alpha, t) = \ldots + \mathcal{Y}_{N-1} \frac{i}{2\pi} \oint_{\mathbf{C}} d\zeta \exp(-i\zeta t)$$

$$(i\zeta^2)^{-1} \frac{2\pi}{\Lambda} \sum_{k} < 0 |\alpha h| > (i[G_{\alpha h} - \zeta])^{-1}$$

<
$$|hu| > (i[G_{\alpha u} - \zeta])^{-1} < |u\alpha| = 0 > \rho_{0} |0| |0| (0),$$
 (5.6)

where $\rho_{\{0\}}(0) \equiv \Lambda^{N} \rho'_{\{0\}}(0) = \alpha^{N} \rho'_{\{0\}}(0)$ can be replaced by $\eta \phi(j, 0)$ (cf. eq. (I. 10. 3)).

The arguments from eq. (I. 8. 8) to eq. (I. 8. 18) can then be repeated. In the asymptotic limit $t > > t_c$, eq. (5. 6) becomes

$$\varpi(\alpha, t) = \ldots + t \int_{-\infty}^{\infty} dk \quad \mathcal{Y}_{N-1} < 0 |\alpha h| > \delta_{\underline{G}}(G_{\alpha h}) < |hu| > \delta_{\underline{G}}(G_{\alpha u}) < |u\alpha| > \Pi \varphi(j, 0)$$

In selecting the dominant terms, we again make the modified weak coupling approximation. Out of all the diagrams proportional to particular powers of N/ Λ and t, we select the ones proportional to the lowest power of e^2 . Then the arguments of Section I.9 can be repeated (see, however, the footnote at the end of Section II.2).

The dominant diagrams are semiconnected chains of ring diagrams.

In particular, we have the result analogous to eq. (I. 9. 9):

If
$$k_j^{(\lambda)} = 0$$
, then $n_j^{(\lambda)} = 0$, for any j and λ . (5.8)

This implies that the isotropy condition (4.6) is preserved in time.

SECTION 6

16

THE SINGULAR INTEGRAL EQUATION

The major mathematical complication arising from the presence of the inhomogeneity across <u>B</u> is already apparent from eq. (5.7). The matrix elements contain two labels instead of one, corresponding to the labels to the right and left of the diagram vertex. This feature cannot be rectified by shifting the position of the Bessel functions, as was done for the homogeneous case, because of the presence of different dummy variables \mathcal{L}_h , \mathcal{L}_u , and \mathcal{L}_h instead of a single variable \underline{k}_h^+ .

This complication seems to prevent the derivation of a kinetic equation of the same simple form as for the homogeneous case. However, some progress has been made, as will be seen below.

It can be shown that the analogue of eq. (I. 10. 1) is

$$\varphi(\alpha, t) = \ldots + \mathcal{Y}_{N-1} t \int_{-\infty}^{\infty} dk < 0 |\alpha| > -\infty$$

$$\delta_{\alpha_1}(G_{\alpha_1}) < |12| > \delta_{\alpha_2}(G_{\alpha_2}) < |\alpha_3| > \delta_{\alpha_2}(G_{\alpha_2})$$

$$\delta_{G_{56}} < |57| > \delta_{G_{76}} < |68| > \delta_{G_{78}}$$

$$|\langle |79| \stackrel{*}{>} \delta_{(G_{98})} \langle |8 | 10| \rangle \delta_{(G_{910})} \langle |10 | 11| \rangle$$

$$\left| \left(\frac{\delta_{0}}{6_{11}} \right) < \left| 9 \right| 2 \right| \stackrel{*}{>} \frac{\delta_{0}}{6} \left(\frac{G_{12}}{12} \right) < \left| 11 \right| 13 \right| >$$

 $\cdot \delta_{(G_{12 \ 13})} < |13 \ 12| \ 0 > \mathbf{n} \ \varphi(j, 0),$

(6.1)

where the asterisks denote the com_r lex conjugate. The arguments leading to eq. (I. .15) are not affected by the more complicated expressions for the vertices; the cha of ring diagrams can be factorized as before. In particular the condition (4.5) is preserved in time, at least for r = 0.

However, we cannot express the results in a form exactly analogous to eqs. (I. 10.18), (I. 10.25). Instead, we define a new quantity $F(\alpha, \gamma)$, which is related to the binary correlation function by a Fourier transformation:

$$U_{2}(\alpha, \gamma) = \int_{-\infty}^{\infty} dk \sum_{n_{\alpha} = n_{\gamma}}^{\infty} F(\alpha, \gamma) \exp(ik[z_{\alpha} - z_{\gamma}]) \exp(-i[n_{\alpha}\theta_{\alpha} - n_{\gamma}\theta_{\gamma}]).$$

(6.2)

(We suppress the parameters k and t from the arguments from now on, since the equations no longer involve the initial time t = 0.) Then it can be shown that

$$\frac{\partial \varphi(\alpha)}{\partial t} = \int_{-\infty}^{\infty} dk \int_{\mathbf{h}}^{\infty} < 0 |\alpha \mathbf{h}| > \mathbf{F}(\alpha, \mathbf{h}), \qquad (6.3)$$

$$F(\alpha, h) = q(\alpha, h) + \delta_{\alpha}(G_{\alpha h}) \int_{u} < |hu| > m(h) F(\alpha, u)$$

$$+\delta_{\underline{(G_{\alpha h})}} \int_{\underline{u}} < |\alpha u| \stackrel{*}{>} \varphi(\alpha) F^{\stackrel{*}{}}(u, h), \qquad (6.4)$$

where

$$q(\alpha, h) \equiv \delta_{(G_{\alpha h})} < |h\alpha| \quad 0 > \varphi(h) \ \varphi(\alpha), \tag{6.5}$$

$$\int_{j} \mathbf{f} = \int_{\Sigma} d\underline{Q}_{j} \int_{0}^{\infty} da_{j} \int_{-\infty}^{\infty} dp_{j} 2\pi m^{2} \omega^{2} a_{j} \mathbf{f}.$$
 (6.6)

18

Note that if the matrix elements did not depend on the second labels, then one could express $\partial \varphi(\alpha)/\partial t$ in terms of $\int_{h} F(\alpha, h)$, and derive an integral equation for $\int_{h} F(\alpha, h)$.

The singular integral equation (6.4) is very complicated, and does not seem readily amenable to the treatment described in Section II. 4. By making transformations similar to those in Section 7 below, we could eliminate some of the variables by a sort of "barring" operation, and derive a very lengthy equation for a quantity $\Phi(\alpha, \underline{q}_{\gamma})$ which is independent of $\alpha_{\gamma}, p_{\gamma}$, and which determines $F(\alpha, \gamma)$ through the following relations:

 $\mathbf{F}(\alpha, \gamma) = \int d\underline{q}_{\alpha} \int d\underline{q}_{\gamma} \exp[i(\underline{q}_{\alpha} \cdot \underline{Q}_{\alpha} - \underline{q}_{\gamma} \cdot \underline{Q}_{\gamma})] \hat{\mathbf{F}}(\alpha, \gamma),$

(6.7)

and similarly for $q(\alpha, \gamma)$ and $\hat{q}(\alpha, \gamma)$;

$$\begin{split} &\hat{\mathbf{F}}(\alpha,\gamma) = \hat{\mathbf{q}}(\alpha,\gamma) + \left\{ \frac{4\pi e^2 N}{\mathrm{im}\psi\Lambda} \quad \delta_{-}(\mathbf{G}_{\alpha\gamma}) \int_{\mathbf{o}}^{\infty} \frac{\mathbf{q}_{\varepsilon} d\mathbf{q}_{\varepsilon}}{\mathbf{q}_{\varepsilon}^2 + \mathbf{k}^2 + \mathbf{x}^2} \right. \\ &\int_{\mathbf{o}}^{2\pi} d\Psi_{\varepsilon} J_{\mathbf{n}_{\gamma}} \left(\mathbf{q}_{\varepsilon} \alpha_{\gamma} \right) \exp(-\mathrm{in}_{\gamma} \Psi_{\varepsilon}) \left[\mathrm{m}\psi\mathbf{k} \frac{\partial}{\partial \mathbf{p}_{\gamma}} \right] \\ &+ \mathrm{i} \left[\mathbf{q}_{\varepsilon} \mathbf{x} \mathbf{q}_{\gamma} \right]^{\mathbf{z}} + \frac{\mathbf{n}_{\gamma}}{\mathbf{a}_{\gamma}} \frac{\partial}{\partial \mathbf{a}_{\gamma}} \right] \hat{\mathbf{\phi}}(\gamma; \mathbf{k}_{\gamma} = \mathbf{q}_{\varepsilon} - \mathbf{q}_{\gamma})) \\ &\sum_{\alpha} J_{\mathbf{n}_{\alpha}} - \nabla_{\alpha} \left(\mathbf{q}_{\alpha} \mathbf{a}_{\alpha} \right) J_{\nabla_{\alpha}} \left(\mathbf{q}_{\alpha} \mathbf{a}_{\alpha} \right) \exp\left(\mathrm{in}_{\alpha} \Psi_{\alpha} \right) \Phi(\alpha, \mathbf{q}_{\varepsilon}) \\ &+ \left\{ \alpha \longleftrightarrow \gamma \right\}^{*}, \end{split}$$
(6.8)

where $\hat{\phi}$ is the Fourier transform of ϕ :

$$\varphi(\gamma) = \int d\underline{k}_{\gamma} \exp\left(i\underline{k}_{\gamma} \cdot \underline{Q}_{\gamma}\right) \hat{\varphi}(\gamma), \qquad (6.9)$$

and $\underline{q}_{\varepsilon}$ is defined by the polar coordinates $q_{\varepsilon}, \psi_{\varepsilon}$. Such transformations have been used extensively by Rostoker and coworkers, and others; see for example Reference (10).

We do not bother to write out the equation for $\Phi(\alpha, \underline{q}_{\gamma})$, because a simpler set of equations can be found in another way. In the next section, we use the summation procedure developed by Résibois ⁽¹¹⁾ in order to derive equations equivalent to (6.3), (6.4), but having a more explicit form.

SECTION 7

THE FREDHOLM-TYPE INTEGRAL EQUATIONS

The Résibois summation procedure⁽¹¹⁾ has been described very clearly and concisely by Balescu (Reference (4), Appendix 10). Singular integral equations are avoided, and kinetic equations are obtained directly. The method is largely independent of the nature of the matrix elements (vertices). It generalizes immediately to the present physical model, provided that we can consider the integrations term by term in the n's. The n's are held fixed until we actually sum the "primitive diagrams".

The complication mentioned in the last section also appears here, but we are able to proceed much farther. We do not get simple geometric series, and therefore no simple kinetic equation is obtained. However, we can reduce the problem to that of solving a two dimensional integral equation of the Fredholm type but with a complicated kernel. Further progress has not been seriously attempted at the time of writing; it would seem that some approximation may be necessary for the derivation of a kinetic equation in a closed form.

We replace $\delta_{\alpha\gamma}(G_{\alpha\gamma})$ by the equivalent form $[i(G_{\alpha\gamma} - i\varepsilon)]^{-1}$, and write, according to the Résibois theorem,

$$F(\alpha, h) = \mathcal{Y}_{N-2} \frac{1}{2\pi} \oint_{C} d\zeta' \sum_{I,S} \mathcal{V}_{\alpha I}(\zeta') \mathcal{Y}^{hS}(i\varepsilon - \zeta')$$

$$(7.1)$$

where

$$\mathcal{V}_{\alpha\tau}(\zeta') = [i(G_{\alpha} - \zeta')]^{-1} < |\alpha 2| > *$$

$$\cdot [i(G_2 - \zeta')]^{-1} < |23| >^* \dots [i(G_1 - \zeta')]^{-1},$$

$$\gamma^{hS}(i\varepsilon - \zeta') = [i(-G_h - i\varepsilon + \zeta')]^{-1} < |h I + 1| >$$

$$[i(-G_{T+1}^{-} - i\epsilon + \zeta')]^{-1} < |I+1|I+2| >$$

$$\dots \left[i(-G_{a} - i\varepsilon + \zeta')\right]^{-1}.$$

Here, C' is a path of integration which consists of a straight line from $\zeta' = -\infty + i c/2$ to $\zeta' = +\infty + i c/2$, together with the usual semicircle in the lower half plane. (The orientation is opposite to that of path C in eq. (5.2).)

From eq. (7.1), it is a straightforward matter to obtain recurrence relations, and the resulting integral equations:

$$\mathbf{F}(\alpha, \mathbf{h}) = \frac{1}{2\pi} \oint_{\mathbf{C}'} d\zeta' \Gamma(\alpha, \mathbf{h}), \qquad (7.2)$$

$$\Pi(c \ h) = \left[i(\mathbf{G}_{\alpha} - c')\right]^{-1} \left[\Delta(\alpha, \mathbf{h}) + \int_{\mathbf{u}} \langle \alpha | \alpha | \rangle^{*} \varphi(\alpha) \Gamma(\mathbf{u}, \mathbf{h})\right],$$

$$\wedge (\alpha, \mathbf{h}) = \left[\mathbf{i} (-\mathbf{G}_{\mathbf{h}}^{-} \mathbf{i}_{\theta} + \zeta') \right]^{-1} \quad \left[\mathfrak{S}(\alpha, \mathbf{h}) + \int_{\mathbf{u}}^{\infty} \langle |\mathbf{h}\mathbf{u}| \rangle \varphi(\mathbf{h}) \wedge (\alpha, \mathbf{u}) \right] ,$$

$$\Theta(\alpha, h) = \langle |h\alpha| | 0 \rangle \varphi(h) \varphi(\alpha).$$
(7.5)

It is claimed that the set of equations (6.3), (7.2), (7.3), (7.4), (7.5) is simpler than the set (6.3), (6.4), (6.5). Note that if one can solve eq. (7.4), then there is no trouble in solving eq. (7.3), because it has essentially the same kernel.

Let us now investigate briefly the ζ' integration in eq. (7.2). It is relatively easy to perform the integration for the $[i(G_{\alpha} - \zeta')]^{-1} \Delta(\alpha, h)$ contribution (eq. (7.3)).

The factor $[i(-G_h - i\varepsilon + \zeta')]^{-1}$ in eq. (7.4) has no poles in the area enclosed by C! Hence it is very plausible to assume that $\Delta(\alpha, h)$ has no singularities there either; at worst there is some sort of stability condition involved. We can write

$$\frac{1}{2\pi} \oint_{\mathbf{C}'} d\zeta' [i(\mathbf{G}_{\alpha} - \zeta')]^{-1} \Delta(\alpha, \mathbf{h}) = \Delta(\alpha, \mathbf{h}) |\zeta' = \mathbf{G}_{\alpha}$$

$$\equiv \widetilde{\Delta}(\alpha, \mathbf{h}), \qquad (7.6)$$

where

$$\tilde{\Delta}(\alpha, h) = \delta_{\alpha}(G_{\alpha})[\Theta(\alpha, h) + \int_{u} \langle |hu| \rangle \phi(h) \tilde{\Delta}(\alpha, u)].$$
(7.7)

The other contribution to $\Gamma(\alpha, h)$ is more difficult to calculate. We merely note that the contribution from the semicircular portion of C'may be neglected, and then change notation. Replacing ζ' by $G_{\chi} + i\epsilon/2$, we get:

$$\mathbf{F}(\alpha, h) = \widetilde{\Delta}(\alpha, h) + \int_{-\infty}^{\infty} d\mathbf{G}_{\mathbf{x}} \left[\delta_{-}(\mathbf{G}_{\alpha\mathbf{x}}) \int_{\mathbf{u}} \langle \alpha \mathbf{u} \rangle \right] + \phi(\alpha) \Gamma(\mathbf{u}, h; \mathbf{x}) ,$$

(7.8)

$$\Gamma(\alpha, \mathbf{h}; \mathbf{x}) = \delta_{\mathbf{u}}(\mathbf{G}_{\alpha, \mathbf{x}}) \left[\Delta(\alpha, \mathbf{h}; \mathbf{x}) + \int_{\mathbf{u}} \langle \alpha \mathbf{u} \rangle^{*} \varpi(\alpha) \Gamma(\mathbf{u}, \mathbf{h}; \mathbf{x}) \right],$$

$$\Delta(\alpha, h; \mathbf{x}) = \delta_{\mathbf{G}} G_{\mathbf{x}h} \left[\odot(\alpha, h) + \int_{\mathbf{u}} \langle |h\mathbf{u}| \rangle \phi(h) \Delta(\alpha, h; \mathbf{x}) \right],$$

(7.10)

$$\widetilde{\Delta}(\alpha, h) = \Delta(\alpha, h; \alpha) = \int_{-\infty}^{\infty} dG_{\mathbf{x}} \delta(C_{\alpha \mathbf{x}}) \Delta(\alpha, h; \mathbf{x}).$$
(7.11)

We now examine the problem of simplifying the integral equations (7.3) and (7.4), or (7.9) and (7.10). All these equations have essentially the same kernel, and can in principle be solved in the same way. We consider eq. (7.10) as an example, and abbreviate $\Delta(\alpha, h; x)$ by $\Delta(h)$, since α and G_x appear only as parameters. Also, $\delta_{-}(G_{xh}) = (\alpha, h)$ will be written as $\Xi(h)$, so that

$$\Delta(\mathbf{h}) = \Xi(\mathbf{h}) + \delta_{\mathbf{h}}(\mathbf{G}_{\mathbf{h}}) \int_{\mathbf{u}} \langle |\mathbf{h}\mathbf{u}| \rangle \varphi(\mathbf{h}) \Delta(\mathbf{u}).$$
(7.12)

The following expansions are carried out for $\Xi(h)$ as well as for $\Delta(h)$:

 $\Delta(\mathbf{h}) = \int d\mathbf{g}_{\mathbf{h}} \exp(-i\mathbf{g}_{\mathbf{h}} \cdot \mathbf{Q}_{\mathbf{h}}) \hat{\boldsymbol{\Lambda}} (\mathbf{h}),$

(so that $\hat{\Delta}(h) = (2\pi)^{-2} \int dQ_h \exp(iq_h \cdot Q_h) \Delta(h)),$

$$\varphi(\mathbf{h}) = \int d\underline{\mathbf{k}}_{\mathbf{h}} \exp ((\mathbf{i}\underline{\mathbf{k}}_{\mathbf{h}} \cdot \underline{\mathbf{Q}}_{\mathbf{h}}) \hat{\varphi}(\mathbf{h});$$

$$\hat{\Delta}(h) = \sum_{\substack{n_h + \nu_h \\ \nu_h}} J_{n_h + \nu_h} (q_h a_h) J_{\nu_h} (q_h a_h) \exp(-in_h \psi_h) \hat{\Delta}'(h) ,$$

where $\psi_h = \arctan(q_h^y/q_h^x)$, and $\Delta'(h)$ depends on v_h instead of n_h . Then

$$\hat{\Delta}(\mathbf{h}) = \hat{\Xi}(\mathbf{h}) + \delta_{-}(\mathbf{G}_{\mathbf{x}\mathbf{h}}) \frac{4 \pi e^{2} N}{i m \omega \Lambda} \int_{0}^{\infty} \frac{q_{u} dq_{u}}{q_{u}^{2} + k^{2} + \kappa^{2}} \int_{0}^{2\pi} d\psi_{u} J_{n_{h}}(q_{u}a_{h}) \exp(-i n_{h}\psi_{u}) \int d\underline{k}_{h}$$
$$[m \omega k \frac{\partial}{\partial P_{h}} - i(\underline{q}_{u} \underline{k}_{h})^{2} + \frac{n_{h}}{a_{h}} \frac{\partial}{\partial a_{h}}] \delta(\underline{q}_{h} + \underline{k}_{h} - \underline{q}_{u}) \hat{\varphi}(\mathbf{h}) \overline{\Delta}(\underline{q}_{u}) ,$$

where

$$\overline{\Delta} (\underline{q}_{u}) \equiv \int_{0}^{\infty} da_{u} \int_{-\infty}^{\infty} dp_{u} 2\pi m^{2} \omega^{2} a_{u} J_{0}(\underline{q}_{u} a_{u}) \Delta'(\underline{u}) \Big|_{v_{u}} = 0 ,$$

(7.15)

and we have used the relation

$$\sum_{n} J_{n+\nu} J_{n} = \delta^{\mathbf{K}r}(\nu).$$

Furthermore, eq. (7.13) yields

$$\overline{\Delta} (\underline{q}_{h}) = \sum_{n_{h}} \int_{0}^{\infty} da_{h} \int_{-\infty}^{\infty} dp_{h} 2\pi m^{2} \omega^{2} a_{h}$$

$$J_{n_{h}}(q_{h}a_{h}) \exp(in_{h}\Psi_{h}) \hat{\Delta}(h). \qquad (7.16)$$

Therefore eq. (7.14) may be reduced by means of this barring operation to

 $\overline{\Delta}(\underline{\mathbf{q}}_{h}) = \overline{\Xi} (\underline{\mathbf{q}}_{h}) + \frac{4\pi e^{2}N}{i\pi m \omega \Lambda} \sum_{h=1}^{n}$

$$\int_{0}^{\infty} da \int_{-\infty}^{\infty} dp_{h}^{2\pi m^{2} \omega^{2} a} \int_{h}^{J} \int_{h}^{(q_{h} a_{h})} dp_{h}^{2\pi m^{2} \omega^{2} a} \int_{h}^{J} \int_{h}^{(q_{h} a_{h})} dp_{h}^{2\pi m^{2} \omega^{2} a} dp_{h}^{2\pi m^{2} \omega^{2} a} \int_{h}^{J} \int_{h}^{(q_{h} a_{h})} dp_{h}^{2\pi m^{2} \omega^{2} a} dp_{h}^{2\pi m^{2} \omega^{2} dp_{h}^{2\pi m^{2} \omega^{2} a} dp_{h}^{2\pi m^{2} \omega^{2} dp} dp_{h}^{2\pi m^{2} \omega^{2} dp} dp_{h}^{2\pi m^{2} \omega^{2} dp} dp_{h}^{2\pi m^{2} \omega^{2} dp} dp_{h}^{2\pi m^{2} \omega^{2} dp$$

$$(in_h \psi_h) \delta_{(G_x^{-[(kp_h/m) + n_h^{\omega}])}$$

$$\int_{0}^{\infty} \frac{q_u^{dq}_u}{q_u^2 + k^2 + k^2} \int_{0}^{2\pi} d\psi_u J_n(q_u^a h)$$

$$\cdot \exp(-in_{h} \psi_{u}) \left[mwk \frac{\partial}{\partial p_{h}} + i(\underline{q}_{u} \times \underline{q}_{h})^{z}\right]$$

$$+\frac{\mathbf{n}_{h}}{\mathbf{a}_{h}}\frac{\partial}{\partial \mathbf{a}_{h}}] \widehat{\boldsymbol{\omega}} (\mathbf{a}_{h},\mathbf{p}_{h},\underline{\mathbf{k}}_{h}=\underline{\mathbf{q}}_{u}-\underline{\mathbf{q}}_{h}) \overline{\boldsymbol{\Delta}} (\underline{\mathbf{q}}_{u}).$$
(7.17)

This equation can be written in a notation that may be more familiar:

$$\overline{\Delta} (\underline{q}) = \overline{\Xi} (\underline{q}) + \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} d\underline{q}' K(\underline{q}, \underline{q}') \overline{\Delta} (\underline{q}'), \qquad (7.18)$$

where

$$K(\underline{q},\underline{q}') = \frac{4\pi e^2 N}{i\Lambda(q'^2 + k^2 + \chi^2)} \int_{0}^{\infty} 2\pi p_{\mu} dp_{\mu} \int_{-\infty}^{\infty} dp_{\mu}$$

$$\sum_{n = -\infty}^{\infty} \delta_{G} - [kp_{\parallel}/m + nw])$$

$$J_n(\frac{qp_*}{mw}) J_n(\frac{q'p_*}{mw}) \exp(in[\psi - \psi'])$$

$$\cdot \left[k \frac{\partial}{\partial p} \right] + \frac{nmw}{p} \frac{\partial}{\partial p} + \frac{i}{mw} \left(g' \times g \right) \right]$$

$$\cdot \hat{\varphi}(\underline{p}, \underline{k} = g' - \underline{q}), \qquad (7.19)$$

and p is the kinetic momentum.

Equation (7.17) or (7.18) is of the Fiedholm type, but the kernel K is rather complicated. We give it some study in the next section, but no detailed examination has yet been carried out.

SECTION 8

COMPARISON OF RESUL IS WITH EXISTING KINETIC EQUATIONS

Our results reduce to the homogeneous kinetic equation(I. 11. 17), or (I. 11. 1), (II. 4. 9), when $\varphi(j)$ is independent of Q_j . The easiest way to show this is to return to eq. (7.1), and perform the integrations over the $\{Q_j\}$. We obtain factors $\delta(\pounds_h - \pounds_u), \ \delta(\pounds_u - \pounds), \ \text{etc.}, \ \text{so that the integrations over all but one of the}$ $\{\pounds_j\}$ are trivial; we may set $\pounds_h = \pounds_u = \ldots = \pounds$ and $\beta_h = \beta_u = \ldots = \beta$. Then the Résibois summation procedure yields geometric series, and the final result is precisely eq. (II. 4. 9).

In order to see that this is plausible without going through the details, one should note that eq. (7.17) reduces to

 $\overline{\Delta} (\underline{q}_{h}) = \overline{\underline{z}} (\underline{q}_{h})/\eta (x; \underline{k}_{\bullet} = \underline{q}_{h})$

when $\widehat{\phi}(h) = \delta(\underline{k}_{h})\phi(h)$, if the normalization of ψ is changed by a factor Σ . Roughly speaking, the term $q(\alpha)/\eta(\alpha)$ of eq. (II. 4. 9) comes from the term $\widetilde{\Delta}(\alpha, h)$ of eq. (7.8), and the integrand $\delta_{\Omega}(G_{\alpha u})[q_{(1)}(u)/\eta(u)]/\eta^{*}(u)$ in the other term comes from the factor $\Gamma(u, h; x)$ in eq. (7.8).

Let us now return to the diffusion problem. So far as we are aware, no equivalent results have yet been published. The only comparable results are those of Eleonskii, Zyryanov, and Silin⁽¹²⁾, especially their equation (17). They made a quantum mechanical calculation of the collision integral, using the first Born approximation and the Landau representation (Landau levels), and later took the classical limit. The collective effects were taken into account through the <u>ad hoc</u> introduction of a dielectric tensor. They simplified the results by assuming that $\varphi(j)$ is independent of Q_j^x , depending only on Q_j^y (or y_j^g , in our previous notation), a_j , p_j , and t. If we set the collective factor equal to 1 (so that the results are analogous to the Landau equation), their collision integral for the single species plasma may be written as follows:

$$\int dk \int dQ_{h}^{y} \int_{0}^{\infty} da_{h} \int_{-\infty}^{\infty} dp_{h} 2\pi m^{2} w^{2} a_{h} \sum_{n_{\alpha} = n_{h}}^{\infty} \sum_{n_{\alpha} = n$$

 $\left(\frac{e^4}{\pi^2}\right) \int_{-\infty}^{\infty} dh^x \left(k \frac{\partial}{\partial p_{\alpha}} - \frac{h^x}{mw} \frac{\partial}{\partial Q_{\alpha}^y} + \frac{h_{\alpha}}{mwa_{\alpha}} \frac{\lambda}{\partial a_{\alpha}}\right)$ $\left[Re \int_{-\infty}^{\infty} \left(\frac{kp_{\alpha}}{m} + n_{\alpha}w - \frac{kp_{h}}{m} - n_{h}w\right)\right]$ $\left(\int_{-\infty}^{\infty} dh^x \int_{-\infty}^{\infty} dh^y \left(k^2 + h^{x2} + h^y \int_{-\infty}^{y} h^y\right)^{-1}$ $exp(i[n_{\alpha}\beta_{\alpha} - n_{h}\beta_{h}])exp(i[h^y \alpha_{\alpha}^y - h^y \partial_{h}^y]^{-1}$ $\int_{-\infty}^{\infty} (h_{\alpha}a_{\alpha}) \int_{-n_{h}}^{\infty} (h_{h}a_{h}) \left(\frac{h_{\alpha}a_{\alpha}}{h} - \frac{h_{\alpha}}{mw} \frac{\lambda}{h}\right) - \frac{h_{\alpha}}{mw} \frac{\lambda}{h}$ $\left(\int_{-\infty}^{\infty} \frac{h_{\alpha}}{mw} - \frac{h_{h}}{mwa_{h}} \frac{\lambda}{h}\right) \left(\frac{h_{\alpha}}{h}w(h)\right) = 0$ $\left(\int_{-\infty}^{\infty} \frac{h_{\alpha}}{mw} - \frac{h_{\alpha}}{mw} \frac{\lambda}{h}\right) \left(\int_{-\infty}^{\infty} \frac{h_{\alpha}}{h}w(h)\right) = 0$

This is to be compared with our result (for $x \rightarrow 0$);

 $\operatorname{Re}\int dk \int dQ_{h}^{y} \int_{\Omega}^{\infty} da_{h} \int_{-\infty}^{\infty} dp_{h}^{2\pi m^{2}w^{2}a_{h}} \sum_{\substack{n_{\alpha} \\ n_{\alpha} }} \sum_{\substack{n_{\alpha} \\ n_{h}}} \sum_{\substack{n_{\alpha} \\ n_{\alpha} }} \sum_{\substack{n_{\alpha} \\ n$

(8.1)

$$\cdot \left(\frac{e^{4}}{\pi}\right) \int_{-\infty}^{\infty} d\mathbf{n} \left(k \frac{\partial}{\partial p_{\alpha}} - \frac{m}{mw} \frac{\partial}{\partial Q_{\alpha}^{y}} + \frac{n}{mwa_{\alpha}} \frac{\partial}{\partial a_{\alpha}}\right)$$
$$\cdot \int_{-\infty}^{\infty} \left(\frac{kp_{\alpha}}{m} + n_{\alpha}w - \frac{kp_{h}}{m} - n_{h}w\right) \left[\int d\mathbf{n}_{\alpha}^{y} \int d\mathbf{n}_{h}^{y}\right]$$
$$\cdot \left(k^{2} + \mathbf{n}_{\alpha}^{x2} + \mathbf{n}_{\alpha}^{y2}\right)^{-1} \left(k^{2} + \mathbf{n}_{\alpha}^{x2} + \mathbf{n}_{h}^{y2}\right)^{-1}$$

•
$$\exp(i[n_{\alpha} - n_{h}][\beta_{\alpha} - \beta_{h}])\exp(i[\beta_{\alpha} - \beta_{h}]Q_{\alpha h})$$

$$\cdot J_{n_{\alpha}} (\mathcal{L}_{\alpha} a_{\alpha}) J_{n_{\alpha}} (\mathcal{L}_{h} a_{\alpha}) J_{n_{h}} (\mathcal{L}_{\alpha} a_{h}) J_{n_{h}} (\mathcal{L}_{h} a_{h})]$$

$$(k\frac{\partial}{\partial p_{a}} - k\frac{\partial}{\partial p_{h}} - \frac{w}{mw}\frac{\partial}{\partial Q_{ah}^{y}} + \frac{n_{a}}{mwa}\frac{\partial}{\partial a_{a}}$$

$$-\frac{n_{h}}{m\omega a_{h}}\frac{\partial}{\partial a_{h}})\left[\frac{N}{\Lambda}\phi(h)\right]\phi(\alpha).$$
(8.2)

Expressions (8.1) and (8.2) are very similar, but seem to be slightly different. We have not yet traced the origin of the differences.

In (8.1) or (8.2), the coefficient of $\partial^2/\partial Q_{\alpha}^{y^2}$ is proportional to ω^{-2} , or B^{-2} . Hence the results may be in qualitative agreement with Fick's law with a collisional diffusion coefficient. (See References (1), (2), and (12) for more complete discussions.) Our hope is that by properly taking into account the collective effects, as we have done, we may eventually be able to understand better the phenomenon of anomalous diffusion, where the diffusion coefficient decreases less tapidly than B^{-2} for increasing magnetic field strength.

SECTION 9

30

CONCLUSION

We have provided here a basis for kinetic theoretical studies of the diffusion of a plasma across a magnetic field. The main result contained in this report is a demonstration that the problem of deriving a kinetic equation for the one particle distribution function can be reduced to the problem of solving an equation of the Fredholm type (7.18). The demonstration, which is believed to be original, contains no explicit restriction on the amplitude or length scale of the inhomogeneity across the magnetic field.

Much further work, however, remains to be done on the refinement of the physical model, and on the qualitative and quantitative examination of the properties of the resulting equations.

REFERENCES

- Golant, V. E. Usp. fiz. nauk <u>74</u>, 377(1963); English transl, : Soviet Phys. Uspekhi <u>6</u>, 191(1963).
- 2. Boeschoten, F. J. Nucl. Energy <u>C6</u>, 339(1964).
- 3. Prigogine, I. Non-equilibrium statistical mechanics (Interscience, New York, 1962).
- 4. Ealescu, R. Statistical mechanics of charged particles (Interscience, New York, 1963).
- 5. Haggerty, M.J. and de Sobrino, L. G. Can.J. Phys. <u>42</u>, 1969(1964). The symbol "I" refers to this paper.
- 6. Haggerty, M.J. Can, J. Phys. <u>43</u>, 122 (1965) [†]. The symbol "II" refers to this paper.
- 7. Haggerty, M.J. Can. J. Phys. (to be published, September or October 1965, and referred to as "III").
- 8. Rose, D.J. and Clark, M. Plasmas and controlled fusion (M.I.T. Press, Cambridge, Mass. and J. Wiley and Sons, New York, 1961),
- 9. Barabanenkov, Yu. N. Zh. Eksperim. i Teor. Fiz. <u>40</u>, 1476 (1961); English transl.: Soviet Phys. JETP 13, 1034 (1961).
- 10. Rostoker, N., Aamodt, R., and Eldridge, O. Ann. Phys. (N.Y.) <u>31</u>, 243 (1965).

11. Résibois, P. Phys. Fluids <u>6</u>, 817 (1963).

Eleonskii, V. M., Zyryanov, P.S., and Silin, V.P. Zh. Eksperim. i Teor.
 Fiz. 42,876 (1962); English transl.: Soviet Phys. JETP <u>15</u>, 619 (1962).

*Errata:

Page 1977, line 26: the last two lines of the expression for δL_{hu} should read as follows:

 $\{ [m_{\perp} \ \boldsymbol{\ell}^{z} \partial/\partial p_{h}^{z} - (\boldsymbol{\ell}^{\perp} \times \partial/\partial \underline{x}_{h}^{\boldsymbol{g} \perp})^{z} + \boldsymbol{\ell}^{\perp} (\cos [\theta_{h} - \beta] \partial/\partial a_{h} \}$

 $-[1/a_h]\sin[\theta_h-\beta_b]\partial/\partial\theta_h]-[h \leftrightarrow u] \}.$

Page 2000, line 6 and line 9: "(8.15)" should be "(8.14)".

Page 2008, line 12: "q = 1, 2, 2, 2, 1" should be "q = 1, 2, 2, 1, 1". See also the footnote on page 130 of paper II.

†Errata:

Page 123, line 5: "m²" should be "m⁶". line 10: "(2π)⁻¹" should be "(2π)⁻²". Page 125, Fig. 2: "Π" should be "s⁻¹". σ σ =1 Page 128, line 6: "m²" should be "m⁶". Page 128, last paragraph: "first order e²" should read "first order in e²". Page 129, line 21: " $\vartheta_h \delta(G_{\alpha h} \varphi^{eq}(\alpha) \varphi^{eq}(h)$ " should be " $\vartheta_h \delta(G_{\alpha h}) G_{\alpha h} \varphi^{eq}(\alpha) \varphi^{eq}(h)$ ". Page 129, line 26: " $\left(\frac{\varkappa_d^2}{\kappa^2 + \varkappa^2}\right)^{s-1}$ " should be " $\left(\frac{-\varkappa_d^2}{\kappa^2 + \varkappa^2}\right)^{s-1}$ ".

Security Classification					
DOCUMENT CO	NTROL DATA - RE	D			
(Security classification of title, body of abstract and index	ing annotation must be a	ntered vhen	the overall report to classified)		
Brooklyn I, New York		20. REPORT SECURITY CLASSIFICATION Unclassified 20. group			
REPORT TITLE					
On The Kinetic Theory of Diffusion o Fiel	f a Plasma Co d.	lumn A	cross a Magnetic		
DESCRIPTIVE NOTES (Type of report and inclusive dates)		<u></u>			
Research Report AUTHOR(3) (Leet name, first name, initief)					
Michael J. Haggerty					
REPORT DATE	74- TOTAL NO. OF	PAGES	75. NO. OF REFS		
August 10, 1965	32		12		
. CONTRACT OR GRANT NO.	S. ORIGINATOR'S R	EPORT NUM	IBER(S)		
NONR 839(38)		200 / 5			
ARPA Order No. 529	PIBMRI-I	289-05			
c.	\$5. OTHER REPORT NO(\$) (Any other numbers that may be seeigh this report)				
d.	J				
	Sponsored by Advanced Research Proje Agency and Monitored by Office of Naval Research, Washington, D.C.				
3. ABSTRACT The kinetic theory of Prigogine and I for a homogeneous plasma in a magn of diffusion of a plasma column acros deriving a kinetic equation is reduced Fredholm-type integral equation with made concerning the amplitude or let magnetic field. The results are pre- made for further refinements. Some results are given.	Balescu, which etic field, is r ss a magnetic d to that of sol a complicated ngth scale of the liminary in nations comparisons	was pr low appl field. ving a t d kernel he inhor cure, ar with pr	reviously developed lied to the problem The problem of wo dimensional l. No restriction is nogeneity across the ad suggestions are eviously existing		
D FORM 1472					

Unclassified

14 KEY WORDS		LINK A ROLE WT		LINK B		LINKC		
				HOLE WT		ROLE WT		
Plasma-diffusion dynamics kinetic theory Magnetoplasma.								
INSTR	UCTIONS							
 OKIGINATING ACTIVITY: Enter the function and address of the contractor, subcontractor, grantee, Depstment of De- fense activity or other organization (corporate author) issuing the report. REPORT SECURITY CLASSIFICATION: Enter the over- all security classification of the report. Indicate whether "Reatricted Data" is included. Marking Ia to be in accord- ance with appropriate security regulations. GROUP: Automatic downgrading is specified in DoD Di- rective 5200. IO and Armed Forces Industrial Manual. Enter the group number. Also, when applicable, show that optional markings have been used for Group 3 and Group 4 as author- ized. REPORT TITLE: Enter the complete report title in all capital lettera. Titles in all cases ahould be unclassified. If a meaningful title cannot be aelected without classifica- tion, ahow title classification in all capitals in parenthesis immediately following the title. DESCRIPTIVE NOTES: If appropriate, enter the type of report, e.g., interim, progress, summary, annual, or final. Give the inclusive dates when a specific reporting period is covered. AUTHOR(S): Enter the name(s) of author(s) as shown on or in the report. Enter tast name, first name, middle initial. If military, show rank end branch of service. The name of the principal author is an absolute minimum requirement. REPORT DATE: Enter the date of the report as day, month, year; or month, year. If more than one date appears on the report, use date of publication. TOTAL NUMBER OF PAGES: The total page count should follow normal pagination procedures, i.e., enter the number of pages containing information. NUMBER OF REFERENCES Enter the total number of references cited in the report. CONTRACT OR GRANT NUMBER: If appropriate, enter the applicable number of the contract or grant under which the report was written. & d. ONTRACT OR GRANT NUMBER: If appropriate, enter the applicable number of	 Imposed by security classification, using standard statements such as: "Qualified requesters may obtain copies of this report from DDC." "Foreign announcement and dissemination of thia report by DDC is not suthorized." "U. S. Government agencies may obtain copies of this report directly from DDC. Other qualified DDC users shall request through "" "U. S. military agencies may obtain copies of this report directly from DDC. Other qualified users shall request through "" "All diatribution of thia report is controlled. Qualified DDC users shall request through "" "All diatribution of thia report is controlled. Qualified DDC users shall request through "" If the report has been furnished to the Office of Technical borvices, Department of Commerce, for sale to the public, Indicate this fact and enter the price, if known. SUPPLEMENTARY NOTES: Use for additional explanatory notes. SPONSORING MILITARY ACTIVITY: Enter the name of the departmental project office or laboratory sponsoring (paying for) the research and development. Include address. ABSTRACT: Enter an abstract giving a brief and factual summary of the document indicative of the report, even though it may also appear cleawhere in the body of the technical report. If additional space is required, a continuation sheet shall be attached. It is highly desirable that the abstract of classified reports be unclassified. Each paragraph of the abstract shall end with an indication of the military security classification of the in-formation in the paragraph, represented as (TS). (S). (C), or (U). There is no limitation on the length of the abstract. How- 							
 military department identification, such as project number, subproject number, system numbers, task number, etc. 9a. ORIGINATOR'S REPORT NUMBER(S): Enter the official report number by which the document will be identified and controlled by the originating activity. This number must be unique to this report. 9b. OTHER REPORT NUMBER(S): If the report has been assigned any other report numbers (either by the originator or by the sponsor), also enter this number(s). 10. AVAILABILITY/LIMITATION NOTICES: Enter any limitations on further dissemination of the report, other than those 	14. KEY WORDS: Key words are technically meaningful terms or short phrases that characterize a report and may be used as index entries for cataloging the report. Key words must be selected so that no security classification ia required. Identi- fiers, such as equipment model designation, trade name, military project code name, geographic location, may be used as key words but will be followed by an indication of technical con- text. The assignment of links, rules, and weights is optional							

Unclassified

SUPPLEMENTARY

INFORMATION

D-471208

ę,

Research Report No. PIBMRI-1289-65 by Michael J. Haggerty (Department of Physics, Polytechnic Institute of Brooklyn, Contract No. NONR 839(38), ARFA Order No. 529,

August 10, 1965.)

On the Kinetic Theory of Diffusion of a Plasma Column across a Magnetic Field.

ERRATA (as of 12 January 1966):

- Page 2, fourth line after eq. (2.1): replace "effect" by "affect".
- Fage 2, third paragraph: replace "assumption (I.3.26)" by "assumptions (I.3.26) and (I.8.12)".
- Page 2, end of third paragraph: replace "with W set equal to zero" by "with W chosen so as to cancel the contribution of V(k^Z=0) to the interaction matrix elements."
- Page 2, second last line before footnote: replace "may be a less trivial problem" by "is a less trivial problem".

Page 7, add at bottom of page:

"It should be observed that the matrix elements that are diagonal with respect to the k's are not necessarily zero. However, in accordance with the discussion in Section 2, we ignore these contributions by replacing $(\ell^2 + [k_h - k_h^2]^2 + \kappa^2)^{-1}$ by 0 when $k_h = k_h^2$." Page 8, third line of eq. (4.1): " $\{z_i\}$ should read " $\{z_i\}$ ".

Page 9, left side of eq. (4.5) should read "fr.8(0)".

Page 11, third line from bottom: replace $\sum_{k} by \sum_{k \in k}$, and include a marginal note referring to the paragraph added at the bottom of page 7.

Page 12, line 3: " $k_n = k$ " should read " $k_n = -k$ ".

Page 14, second line of eq. (5.6): replace $\sum_{k} k \neq 0$.

Page 17, add a dagger after "binary correlation function", and write the following footne... at the bottom of the page: "'The appearance of U_2 at this stage, instead of $f_{2,2}$, implies that the self-consistent electric field is being ignored (of. Sections 2 and 3)."

Page 21, line 2: "Gg" should read "Gg".

Page 22, eq. (7.8): replace $"dG_{x}"$ by $"(dG_{x}/2\pi)"$.

Page 22, eq. (7.10): replace " $\phi(h) \Delta(\alpha,h;x)$ " by " $\phi(h) \Delta(\alpha,u;x)$ ".

Page 25, eq. (7.18): " $\overline{E}(\mathbf{q})$ " should read " $\overline{E}(\mathbf{q})$ ".

Page 25, add ") " at end of last line.

Page 26, last line of eq. (7.19): replace "k" by "k ".

Passe 28, eq. (8.1): replace $\left| \int d\ell_{\alpha}^{y} \dots J_{n_{h}}(\ell_{h}a_{h}) \right|^{2}$ by $\left| \int d\ell^{y}(k^{2} + \ell^{2})^{-1} \exp(i\left[n_{\alpha} - n_{h}\right]\beta) \exp(i\ell^{y}Q_{\alpha h}^{y}) J_{n_{\alpha}}(\ell a_{\alpha}) J_{n_{b}}(\ell a_{h}) \right|^{2}$.

Page 29, line 3: "k $\partial/\partial p_a$ " should be "k $\partial/\partial p_a$ ".

Page 29, lines 5 and 6: replace "Expressions (8.1) and (8.2) ... differences." by "The results (8.1) and (8.2) are identical."

Page 31, add at bottom of page: "Page 1996, eq. (7.21): replace Z(e²) by Z(0), and add '+ 0(e⁴)'. Eq. (7.22): add '+ 0(e⁴)'. Lines 15 to 17: replace 'and, as it happens, is independent of e², by 'to first order'."

Page 32, last correction: replace "d" by "D" in two places.

The present address of the author is:

Institute for Fluid Dynamics and Applied Mathematics University of Maryland, College Park, Maryland 20742, U.S.A.

- 3 -