# **UNCLASSIFIED**



# **AD NUMBER**

## AD-477 286

# **NEW LIMITATION CHANGE**

TO

DISTRIBUTION STATEMENT - A Approved for public release; distribution is unlimited

LIMITATION CODE: 1

FROM No Prior DOD Distr Scty Cntrl St'mt Assigned

# **AUTHORITY**

Office of Naval Research, via Notice dtd Jul 27, 1971

199903070

THIS PAGE IS UNCLASSIFIED

### FINAL REPORT

CONTRIBUTIONS TO THE MONO LAKE EXPERIMENTS

**Research** conducted for:

Department of the Navy Office of Naval Research Washington 25, D.C.

#### **VOLUME II**

THE LINEAR THEORY OF WATER WAVES GENERATED BY EXPLOSIONS

Prepared by: Robert W. Whalin

October 1965

AN LONG

STATES AND

distribution of

-1949F

State of

Contract No. Nonr 5006(00)

NESCO Report S-256-2

NATIONAL ENGINEERING SCIENCE COMPANY 711 South Fair Oaks Avenue Pasadena, California 91106 9999030309

Reproduced From Best Available Copy

#### FOREWORD

R

ANA NO

Andra Part

14.1944655

- states

di fina ka

199.55

hitte

12.584

344.40

4......

This is Volume II of the final report under Contract No. Nonr 5006(00). This report is issued in order to present the linear theory (in its most general form) of water waves produced by explosions and to specify the assumptions which prevail in performing the Mono Lake deep water predictions. The most general form of the linear theory has never --appeared in the literature prior to this report and it is evident that it has not been employed to its fullest extent.

This report was prepared under the direction of Dr. B. Le Méhauté, Associate Director and Project Manager for this program. Miss Lois Webb contributed to the development of the computing techniques and the programming was accomplished by Mr. H. Lee Butler.

### ABSTRACT

Name and

(in the

10.42

20.2

The objective of this project was to predict the water waves generated by the Mono Lake field tests. These tests consisted of detonating ten TNT explosions of approximately 9,500 pounds each, and measuring the water waves and run-up produced by these detonations.

Volume I of the final report contains these predictions and a brief description of the theoretical and empirical methods employed in performing the predictions. This volume presents the general form of the linear theory of water waves produced by explosions in water of constant depth. The theory is developed in detail so that the full extent of the assumptions made in performing the predictions for Mono Lake can be realized. It is also evident that several natural extensions to the presently employed theory can be made which will result in more realistic initial conditions and which should present a more reliable theory. Examples of the various solutions that depict the differences between a symmetric surface deformation, a symmetric time-dependent surface deformation, and an asymmetric time-dependent surface deformation are given.

### CONTENTS

a

8

Seator .

Ĩ

E

1."

۰.

		Page
1.	INTRODUCTION	1
2.	THEORETICAL FORMULATION	3
3.	ASYMPTOTIC SOLUTIONS	7
	3.1 Axially Symmetric Initial Impulse	7
	3.2 Axially Symmetric Surface Deformation	9
	3.3 Axially Symmetric Time-Dependent Surface Deformation	12
	3.4 Asymmetric Time-Dependent Surface Deformation	16
4.	METHOD OF SOLUTION IN THE AREA NEAR THE SOURCE	31
5.	CONCLUSIONS	39
6.	REFERENCES	41

·• /

#### 1. INTRODUCTION

The linear theory of water waves produced by explosions has been utilized by many investigators (Lamb 1945; Van Dorn 1959, 1963; Kranzer and Keller 1955; Prins 1956; Whalin 1965; Kajiura 1963; Unoki and Nakano 1953; Stoker 1947; Penney 1950; Fuchs 1952; Kirkwood and Seeger 1950) during the last 20 years. The basic mathematical foundation for the generalized linear theory has been in existence for a longer period of time. It involves the use of a timedependent Green's Function which satisfies the Laplace equation with the appropriate linearized boundary conditions and the recognized assumptions that linearization involves. However, this theory in its general form has escaped application with one notable exception (Kajiura, 1963), who applied the theory to calculate the leading wave of a tsunami and gave various examples of the theory to show that certain special cases reduced to the well known solutions (Kranzer and Keller, 1955) for explosion waves.

This report formulates the general theory and applies it to water waves produced by explosions. The asymptotic solutions are developed, along with examples, for several special cases and the full impact of the use of the time-dependent Green's Function is thoroughly discussed. The advantage of using this theory in the case of one detonation and multiple detonations is pointed out.

The axially symmetric water craters presently being used to predict explosion waves many times bear little resemblance to the actual craters created (Van Dorn 1963, Whalin 1965); these craters are generally termed "pseudo" craters. The use of a time-dependent initial deformation will undoubtedly result in much more realistic crater parameters. These parameters should be related to weapon yield and location as has been accomplished for stationary symmetric water craters (Whalin, 1965). The theory in its most general form can

be applied to predict the wave spectrum generated by an exotic shaped source resulting from multiple detonations which may be employed to produce wave amplification at a certain point or in a specified area.

The conclusion of this report is that the prediction methods presently employed and used for the Mono Lake program should be modified to account for a time-dependent source of disturbance. Further, the empirical relationships between the cavity parameters and the weapon size and location should be established as they have been for a stationary initial deformation. Furthermore, one can probably establish a model, which will account for the evaporation of water as well as the subsequent return of water from the plume collapse. In addition, the general solution for an arbitrary shaped initial deformation can be used to predict the wave spectrum generated by multiple explosions.

#### 2. THEORETICAL FORMULATION

The mathematical model employed in the generalized linear theory of water waves generated by explosions is based on the linear theory of surface waves in an inviscid incompressible fluid of constant depth. This model allows the source to be asymmetrical and time-dependent in its most general form. Consequently, a mathematical model is available in which the actual wave generation mechanism can be simulated much more precisely than that generally used in present applications.

The water is assumed incompressible, of constant depth d; the origin of the Cartesian coordinate system (x', y', z') is at the undisturbed free surface, the vertical axis z' is positive upwards, and the variables are transformed to dimensionless form (i.e., x = x'/d, y = y'/d,  $t = t' \sqrt{g/d}$ ,  $\eta = \eta'/d$ ,  $V = V'/\sqrt{gd}$ ;  $\omega = \phi'/d \sqrt{gd}$ ;  $p = (p'/\rho)/gd$ ; etc.).

The kinematic and dynamic conditions at the free surface are

$$\varphi_{\mathbf{z}} = \eta_{\mathbf{t}}, \ \mathbf{z} = 0 \tag{1}$$

$$\omega_{\perp} = -\eta - p, z = 0 \tag{2}$$

and the bottom condition is

$$\varphi_{z} = w_{B}, z = -1 \tag{3}$$

where  $w_B$  is the assumed bottom velocity corresponding to a bottom deformation. The above conditions are subject to the restrictions of assuming a linear approximation. That is, the deformation at the surface and bottom are assumed small compared with the wave length,  $\lambda'$ , and water depth, d, and the condition  $z'\lambda'^2/d^3 \leq 1$ .

The time-dependent Green's Function, G, is a solution of the Laplace equation

$$\nabla^2 G = 0, \ 0 > z > -1, \ t \ge \tau$$
 (4)

satisfying the free surface condition

$$G_{\mu\nu} + G_{z} = 0, z = 0$$
 (5)

and the bottom condition

والمعاد والمعترين والمراجع والمراجع والمحال

$$G_{1} = 0, z = -1$$
 (6)

It is required that G,  $G_x$ ,  $G_y$ ,  $G_z$ ,  $G_t$ ,  $G_{tx}$ ,  $G_{ty}$ , and  $G_{tz}$  be uniformly bounded for every t at x,  $y \rightarrow \infty$ . In addition, (G - 1/R) must be bounded at  $(x_0, y_0, z_0)$  where

$$R^{2} = (x - x_{o})^{2} + (y - y_{o})^{2} + (z - z_{o})^{2}$$
(7)

At time  $t = \tau$  the following initial conditions are assumed

 $G = G_{+} = 0, z = 0$ 

The above equation and conditions are sufficient to determine G uniquely. From Stoker, the Green's Function for the case of three-dimensional motion in water of finite depth is

 $G(x_{o}, y_{o}, z_{o}; \tau | x, y, z, t) = \int_{0}^{\infty} \frac{J_{o}(m\overline{r})}{\cosh m} \left[ \sinh m \left\{ 1 - |z - z_{o}| \right\} - \sinh m \left\{ 1 + (z + z_{o}) \right\} \right]$ 

$$+\frac{2}{\gamma^2}\left\{1-\cosh\gamma(t-\tau)\right\}\frac{m}{\cosh m}\cosh m(1+z)\cosh m(1+z_0)\right] dm,$$
  

$$0 > z, \qquad z_0 > -1$$

where

$$\overline{r}^2 = (x - x_0)^2 + (y - y_0)^2$$

and

محصور والمحصور المحمد والمحمد و

٤.

$$\gamma^{2} = m \tanh m \qquad (11)$$

Note that G is symmetric with respect to  $(x_0, y_0, z_0)$  and (x, y, z) and also t and  $\tau$ . Therefore

$$G(x_{0}, y_{0}, z_{0}; \tau | x, y, z; t) = G(x, y, z; t | x_{0}, y_{0}, z_{0}; \tau) = G(x_{0}, y_{0}, z_{0}; t | x, y, z; \tau)$$
(12)

From Green's formula and the above Green's function

$$\varphi_{\tau} (\mathbf{x}, \mathbf{y}, \mathbf{z}; \tau) = \frac{1}{4\pi} \int_{S} \int \left( G \varphi_{\tau z_{o}} - \varphi_{\tau} G_{z_{o}} \right)_{z_{o} = 0} dS_{o}$$
$$- \frac{1}{4\pi} \int_{S} \int \left( G \varphi_{\tau z_{o}} - \varphi_{\tau} G_{z_{o}} \right)_{z_{o} = -1} dS_{o}$$
(13)

where  $dS_0 = dx_0 dy_0$  and the integral on the lateral boundary in water vanishes because of the condition imposed on G.

The integration of Eq. 13 with respect to  $\tau$  from 0 to t and substitution of conditions in Eqs. 1, 2, 5, 6 and 8 for  $\varphi$  and G yield

$$\varphi(\mathbf{x}, \mathbf{y}, \mathbf{z}, t) - \varphi(\mathbf{x}, \mathbf{y}, \mathbf{z}, 0) = -\frac{1}{4\pi} \int_{S} \int \left[ \left( G \varphi_{\mathbf{z}_{0}} - G_{\tau} \eta \right)_{\tau=0} + \int_{0}^{t} \mu G_{\tau} d\tau \right]_{\mathbf{z}_{0}=0} dS_{0}$$
$$-\frac{1}{4\pi} \int_{S} \int \int_{0}^{t} \left( G \varphi_{\mathbf{z}_{0}\tau} \right)_{\mathbf{z}_{0}=-1} d\tau dS_{0}$$
(14)

5

(10)

**.** 

The first integral in Eq. 14 represents the contribution of the surface condition and the latter integral, that of the bottom condition.

At the surface z = 0, Eq. 9 becomes

$$G(x_0, y_0, z_0; \tau | x, y, 0, t) = \int_0^\infty \frac{2}{\gamma^2} \{1 - \cos \gamma (t - \tau)\} m \cosh m (1 + z_0) \frac{\int_0^0 (mx)}{\cosh m} dm$$

(15)

and since the surface elevation,  $\eta$ , is given by Eq. 2, the substitution of Eq. 14 into Eq. 2 yields

$$(\eta + p) = \frac{1}{4\pi} \int_{S} \int (\mathbf{I}_1 + \mathbf{I}_2 + \mathbf{I}_3) \, dS$$
 (16)

where

$$I_{1} = (G_{t} \varphi_{z_{0}} - G_{\tau t} \eta)_{\tau=0}; z_{0} = 0, z = 0, \qquad (17)$$

$$I_2 = \int_0^t p G_{\tau\tau t} d\tau + \left( p G_{\tau\tau} \right)_{\tau=t}^{, \text{or}}$$
(18)

$$I_{2} = \int_{0}^{t} p_{\tau} G_{\tau t} d\tau - (pG_{\tau t})_{\tau=0}; \quad s_{0} = 0, \ s = 0, \ and$$
(19)

$$I_3 = \int_0^t G_t \varphi_{z_0} \tau \, d\tau, \text{ or}$$
 (20)

$$I_{3} = -\int_{0}^{t} G_{t\tau} \varphi_{z_{0}} d\tau - (G_{t} \varphi_{z_{0}})_{\tau=0}; z_{0} = -1, z = 0.$$
(21)

Note that Eq. 17 represents the contribution of the initial velocity and elevation of the water at the surface, Eq. 18 or Eq. 19 the contribution of the surface pressure, and Eq. 20 or Eq. 21 the contribution from the bottom deformation.

#### 3. ASYMPTOTIC SOLUTIONS

a service of

Contraction of

B

- Heard

ave Au

F

144 M

**NAME** 

-axeve:

2.9 april

Contract of

C. Mar

This section presents the asymptotic solutions obtained by an application of the method of stationary phase to evaluate the integral in Eq. 16. The special cases for which the asymptotic solution is developed are a symmetric initial impulse, a symmetric initial deformation, a symmetric time-dependent initial deformation and an asymmetric timedependent initial deformation. In each case the source is given as an -arbitrary function where special cases are easily investigated. Examples of a special case appear for each assumed initial condition. The condition of a symmetric initial impulse (Section 3.1) and a symmetric initial deformation (Section 3.2) were solved by Kranzer and Keller (1955) and are reproduced here for completeness. However, the developments of Sections 3.3 and 3.4 are original and bear further investigation as far as application to actually performing predictions are concerned.

#### 3.1 Axially Symmetric Initial Impulse

The special case of an axially symmetric initial impulse was solved by Kranzer and Keller by the method of integral transforms. When the proper assumptions are employed their solution is obtained from Eq. 16. Cylindrical coordinates are used and the initial conditions become:

a) The water surface is initially undisturbed

 $\eta_{0}(\mathbf{r}_{0},\theta_{0},\tau)=0$ 

**b**)

The initial velocity at the surface at time t = 0 is.

$$\varphi_{z_{o}} = 0$$

$$z = 0 = 0$$

$$\tau = t = 0$$

Consequently I<sub>1</sub> =  $(G_t \varphi_{\mathbf{s}_0} - G_{\tau t} \eta)_{\tau=0; \mathbf{s}_0 = 0} = 0$ 

·: c)

(CTTT)

**M** 

E

C

È

E

E

E

------

101

Cartin and the state of the second second

The applied pressure at the surface is impulsive at  $\tau = 0^3$ , i.e.,  $p = I_0^{(7)}$ ; therefore

$$I_{2} = \int_{0}^{t} p G_{\tau\tau t} d\tau + (p G_{\tau\tau}), \quad z_{o} = 0, \quad z = 0$$

=  $I_o(r_o) G_{\tau\tau t}$ ,  $\tau = 0, z = 0, z_o = 0$ 

 $-d = I_3 = 0$ 

Upon substitution in Eq. 16 and expressing the wave amplitude in cylindrical coordinates

$$\eta(\mathbf{r},\theta,\mathbf{t}) = \frac{-1}{4\pi} \int_0^\infty 2\gamma \, \mathbf{m} \, \sin \gamma \mathbf{t} \left\{ \int_0^{2\pi} \int_0^\infty \mathbf{I}_0(\mathbf{r}_0) \, \mathbf{J}_0(\mathbf{m} \mathbf{r}) \, \mathbf{r}_0 d\mathbf{r}_0 d\theta_0 \right\} d\mathbf{m} \quad (22)$$

or upon integration with respect to  $r_0$  and  $\theta_0$  the wave amplitude becomes

$$\eta(\mathbf{r},\theta,t) = -\int_0^{\infty} m \sqrt{m \tanh m} \, \mathbf{I}_0(m) J_0(mr) \sin\left(\sqrt{m \tanh m} t\right) dm \quad (23)$$

where

$$\overline{I_0}(m) = \int_0^\infty I_0(r_0) J_0(mr_0) r_0 dr_0 \qquad (2.3)$$

The above integral is evaluated by the method of stationary phase upon the assumption that the frequency of oscillation of  $J_0(mr)$  and the circular function is large compared to that of  $\overline{I_0}(m)$ . Then  $J_0(mr)$  is approximated by an asymptotic expansion and upon integration

$$\eta(\mathbf{r}, \mathbf{t}) = \frac{\Gamma_0(\mathbf{m})\mathbf{m}}{\mathbf{r}} \sqrt{\frac{\varphi(\mathbf{m}) \tanh \mathbf{m}}{-\varphi(\mathbf{m})}} \sin (\mathbf{mr} - \sqrt{\mathbf{m} \tanh \mathbf{m}} \mathbf{t}) \quad (25)$$

where

$$\varphi(m) = \frac{1}{2} \sqrt{\frac{\tanh m}{m}} + \frac{1}{2 \cosh^2 m} \sqrt{\frac{m}{\tanh m}} = \frac{r}{t} \qquad (26)$$

Table I in Section 3.2 gives  $\prod_{o}(m)$  for various initial impulses  $I_{o}(z_{o})$ . Figure 1 shows an example of the wave amplitude as a function of time for the initial impulse in example 2 of Table I at one observation station.

#### 3.2 Axially Symmetric Surface Deformation

The asymptotic solution for an axially symmetric surface deformation was solved by Kranzer and Keller for several special cases. However, the theory is presented in the following pages from the more general mathematical formulation given in Section 2. When cylindrical coordinates are used, the initial conditions become:

a) The water surface is initially undisturbed and is defined by the function  $\eta_{c}(r_{c})$ .

b) The initial velocity of the surface at time t = 0 is

$$\varphi_{z_0} \Big|_{\tau = 0}^{z_0 = 0} = 0$$

c) 
$$I_1 = -G_{\tau t} \eta_0(r_0)|_{\tau=0, z_0=0, z=0}$$
  
=  $\left[ -\int_0^{\infty} 2\cos(\gamma t) m J_0(mr) dm \right] \eta_0(r_0)$   
d)  $I_2 = I_3 = 0$ 

Upon substitution in Eq. 16 and expressing the wave amplitude in cylindrical coordinates

$$\eta(\mathbf{r},\mathbf{t}) = \frac{-1}{4\pi} \int_0^\infty 2\mathbf{m} \cos\gamma t \left\{ \int_0^{2\pi} \int_0^\infty \eta_0(\mathbf{r}_0) J_0(\mathbf{m} \mathbf{r}) \mathbf{r}_0 d\mathbf{r}_0 d\theta_0 \right\} d\mathbf{m}$$
(27)



$$=\frac{-1}{4\pi}\int_{0}^{\infty} 2m\cos\gamma t \left[\int_{0}^{2\pi}\int_{0}^{\infty}\eta_{o}(\mathbf{r}_{o})\left\{J_{o}(m\mathbf{r}_{o})+2\sum_{n=1}^{\infty}J_{n}(m\mathbf{r}_{o})J_{n}(m\mathbf{r}_{o})\cos n(\theta-\theta_{o})\right\}\right]$$

r<sub>o</sub>dr<sub>o</sub>d0<sub>0</sub>] dm

(28)

however

or

$$\int_{0}^{2\pi} \cos n \left(\theta - \theta_{0}\right) d\theta_{0} = 0$$
 (29)

therefore

$$\eta(\mathbf{r},\mathbf{t}) = -\int_{0}^{\infty} \mathbf{m} J_{0}(\mathbf{m}\mathbf{r}) \cos \gamma \mathbf{t} \left[ \int_{0}^{\infty} \eta_{0}(\mathbf{r}_{0}) J_{0}(\mathbf{m}\mathbf{r}_{0}) \mathbf{r}_{0} d\mathbf{r}_{0} \right] d\mathbf{m}$$
(30)

where

$$\int_0^{\infty} \eta_0(r_0) J_0(mr_0) r_0 dr_0 = \overline{\eta}_0(m)$$
(31)

which is the zero order Hankel Transform of  $\eta_0(r_0)$ . The wave amplitude becomes

$$\eta(\mathbf{r}, \mathbf{t}) = -\int_0^{\infty} \overline{\eta}_0(\mathbf{m}) \mathbf{m} \cos\left(\sqrt{\mathbf{m} \tanh \mathbf{m}} \mathbf{t}\right) J_0(\mathbf{m}\mathbf{r}) d\mathbf{m}$$
(32)

The above integral is evaluated by the method of stationary phase where it is assumed that  $\overline{\eta}_0(m)$  has a much lower frequency of oscillation than  $J_0(mr)$  and the circular function. This assumption is a valid one for the types of initial deformations which are representative of underwater explosions, as can be observed from the examples in Table I.  $J_0(mr)$  is approximated by an asymptotic expansion for large argument and the integral becomes

$$\eta(r,t) = \frac{1}{r} \sqrt{\frac{m(r)(m)}{-\varphi'(m)}} \quad \overline{\eta}_{0}(m) \cos(mr - \sqrt{m \tanh m' t}) \quad (33)$$

where

1

18 8 A

ANY AN

. 34, 15

1. N. N.

$$\varphi(m) = \frac{1}{2}\sqrt{\frac{\tanh m}{m}} + \frac{1}{2\cosh^2 m}\sqrt{\frac{\ln}{\tanh m}} = \frac{r}{t}$$
(34)

The following table gives  $\overline{\eta}_{o}(\mathbf{m})$  for various initial deformations  $\eta_{o}(\mathbf{r}_{o})$ .

The wave amplitude may be computed either as a function of time with r constant or as a function of r with time constant.

Figure 2 shows the wave amplitude as a function of time for example 3 of Table I at one observation station.

#### 3.3 Axially Symmetric Time-Dependent Surface Deformation

There has been little investigation into the effect of using a timedependent symmetric surface deformation. This section presents the theory with several specific examples given in detail. The examples can be compared with those where the source is independent of time. The initial conditions become:

a) The water surface is initially undisturbed and is defined by

$$\begin{split} \eta_{0}(\mathbf{r}_{0}) & \sin \frac{\pi}{2} \frac{\tau}{\tau^{*}}, \quad 0 \leq \tau \leq \tau^{*} \\ \eta_{0}(\mathbf{r}_{0}, \tau) = \begin{cases} 0, & \tau > \tau^{*} \end{cases} \end{split}$$

It is assumed that the variables  $r_0$  and  $\tau$  are separable and  $r^*$  should be taken as the dimensionless period of the first expansion of the bubble or the water cavity for the case of a surface explosion.

Ъ)

The initial velocity of the surface at time  $\tau = 0$  is

$$\varphi_{\mathbf{z}_{0}} \begin{vmatrix} \mathbf{z}_{0} & \mathbf{z}_{0} \\ \mathbf{z}_$$



3-7381



TABLE I HANKEL TRANSFORM OF VARIOUS INITIAL DEFORMATIONS

A STATE

and a

alan sa

AND AND

- Andrews

Į.

c) 
$$I_1 = (G_t \varphi_{z_0}) |_{\tau=0, z_0=0, z=0}$$

$$G_t = \int_0^{\infty} \frac{2}{\gamma} \sin \gamma t m J_0(mr) dm$$

d)  $I_2 = I_3 = 0$ 

ີບ

0

0

Sec.

Sandrey .

Sec. 1

Upon substitution in Eq. 16 and expressing the wave amplitude in cylindrical coordinates

$$\eta(\mathbf{r},\mathbf{t}) = \frac{1}{4\pi} \int_{0}^{\infty} \frac{2\mathrm{m}\sin\gamma t}{\gamma} \left[ \int_{0}^{2\pi} \int_{0}^{\infty} (\mathrm{m}\vec{r}) \eta_{0}(\mathbf{r}_{0}) \left(\frac{+\pi}{2\tau^{*}}\right) \mathbf{r}_{0} \, \mathrm{d}\mathbf{r}_{0} \, \mathrm{d}\theta_{0} \right] \mathrm{d}\mathbf{m}$$
$$= \frac{+1}{4\tau^{*}} \int_{0}^{\infty} \frac{\mathrm{m}\sin\gamma t}{\gamma} \left[ \int_{0}^{2\pi} \int_{0}^{\infty} \left\{ J_{0}(\mathrm{m}\mathbf{r}) J_{0}(\mathrm{m}\mathbf{r}_{0}) + 2 \sum_{n=1}^{\infty} J_{n}(\mathrm{m}\mathbf{r}) J_{n}(\mathrm{m}\mathbf{r}_{0}) \cos n(\theta - \theta_{0}) \right\}$$
$$\eta_{0}(\mathbf{r}_{0}) \mathbf{r}_{0} \mathrm{d}\mathbf{r}_{0} \mathrm{d}\theta_{0} \right] \mathrm{d}\mathbf{m}$$

ъnd

$$= + \frac{2\pi}{4\tau^{*}} \int_{0}^{\infty} \frac{\min \gamma t}{\gamma} J_{o}(mr) \left[ \int_{0}^{\infty} \eta_{o}(r_{o}) J_{o}(mr_{o}) r_{o} dr_{o} \right] dm \qquad (35)$$

However, the integral with respect to  $r_c$  is merely the zero order Hankel Transform of the initial deformation  $\eta_o(r_o)$  and will be denoted by  $\overline{\eta_o}(m)$ .

$$\eta(\mathbf{r},\mathbf{t}) = \frac{+\pi}{2\tau^*} \int_0^\infty \frac{\mathrm{mJ}_0(\mathrm{mr})}{\sqrt{\mathrm{m \ tanh \ n}}} \quad \overline{\eta}_0(\mathrm{m}) \sin\left(\sqrt{\mathrm{m \ tanh \ m \ t}}\right) \mathrm{dm}$$
(36)

Upon evaluation by the method of stationary phase

$$\eta(\mathbf{r},\mathbf{t}) = \frac{+\pi}{2\tau^*} \frac{\overline{\eta_0}(\mathbf{m})}{\mathbf{r}\sqrt{\tanh \mathbf{m}}} \sqrt{\frac{\varphi(\mathbf{m})}{-\varphi'(\mathbf{m})}} \sin\left(\mathbf{mr} - \sqrt{\mathbf{m}\tanh \mathbf{m}} \mathbf{t}\right) \quad (37)$$

where  $\varphi$  and  $\varphi'$  are defined as before. Examples of  $\overline{\eta_0}(m)$  are found in Table I.

Figure 3 shows the wave amplitude as a function of time at 623 observation station where  $\overline{\eta}_{c}(m)$  is given by example 3 of Table I.

#### 3.4 Asymmetric Time-Dependent Surface Deformation

In addition to assuming the source of disturbance to be time-dependent, it is relatively straightforward to avoid the assumption of axial symmetry. However, in the case of explosion waves, this assumption is not necessary unless an unsymmetric cavity is produced by an incomplete detonation or by multiple detonations. The initial condition becomes:

a) The water surface is initially undisturbed and is defined by

 $\eta_{o}(\mathbf{r}_{o}, \theta_{o}, t) = \begin{cases} \eta_{o}(\mathbf{r}_{o})\theta(\theta_{o}) T(\tau), & 0 \le \tau \le \tau^{*}, \mathbf{r}_{o} \le \mathbf{R} \\ 0 & 0 \le \theta < 2\pi \\ t > \tau, & \mathbf{r}_{o} > \mathbf{R} \end{cases}$ 

It is assumed that the variables are separable and that 7<sup>\*</sup> is a parameter which should be related to the time of the initial expansion of the bubble.

b) The initial velocity of the surface at time  $\tau = 0$  is

$$\varphi_{\mathbf{z}_{0}} \begin{vmatrix} \mathbf{z}_{0} = 0 \\ \mathbf{z} = 0 \\ \tau = 0 \end{vmatrix} = \frac{d \eta_{0} (\mathbf{r}_{0}, \theta_{0}, \tau)}{d\tau} \begin{vmatrix} \mathbf{z}_{0} \\ \mathbf{z}_{0} \\ \mathbf{z}_{0} \end{vmatrix} = \eta_{0} (\mathbf{r}_{0}) \, \boldsymbol{\ell}(\theta_{0}) \, \mathbf{T}_{\tau}(0)$$

c)  $I_1 = G_t \varphi_{z_0} \Big|_{\tau=0, z_0=0, z=0}$ 

d)

and

16

 $I_2 = I_3 = 0$ 

 $G_t = \int_0^{\infty} \frac{2}{\gamma} \sin \gamma t m J_0(m\bar{r}) dm$ 



PB-3-7382

\*\*\*\*

a

ł.

Upon substitution in Eq. 16 and expressing the wave amplitude in cylindrical coordinates

$$\eta(\mathbf{r},\boldsymbol{\theta},\mathbf{t}) = \frac{T_{\tau}(0)}{2\pi} \int_{0}^{\infty} \frac{\min \gamma t}{\gamma} \left[ \int_{0}^{2\pi} \int_{0}^{\infty} \left\{ J_{o}(mr) J_{o}(mr_{o}) + 2 \sum_{n=1}^{\infty} J_{n}(mr) J_{n}(mr_{o}) \cos n(\theta - \theta_{o}) \right\} \cdot \eta_{o}(\mathbf{r}_{o}) \theta(\theta_{o}) \mathbf{r}_{o} d\mathbf{r}_{o} d\theta_{o} d\theta_{o} dm \quad (33)$$

Upon performing the integrations with respect to  $r_0$  and  $\theta_0$ 

$$\eta(\mathbf{r},\theta,t) = \frac{T_{\tau}(0)}{2\pi} \int_{0}^{\infty} \frac{\min \gamma t}{\gamma} \left[ J_{o}(mr) \overline{\eta_{o}}(m) I_{o} + 2 \sum_{n=1}^{\infty} J_{n}(mr) \overline{\eta_{n}}(m) I_{n}(\theta) \right] dm \quad (39)$$

Γ

[

E

E

E

E

-

AN WAR

Survey.

$$\overline{\eta_{o}(m)} = \int_{0}^{-} \eta_{o}(r_{o}) J_{o}(mr_{o}) r_{o} dr_{o} = \text{zero order Hankel transform}$$
of  $\eta_{o}(r_{o})$ 

 $\overline{\eta_{n}}(m) = \int_{0}^{\infty} \eta_{o}(r_{o}) J_{n}(mr_{o}) r_{o} dr_{o} = n^{\text{th}} \text{ order Hankel transform of } \eta_{o}(r_{o})$ 

$$I_{o} = \int_{0}^{2\pi} \theta(\theta_{o}) d\theta_{o}$$
(40)  
$$I_{n}(\theta) = \begin{cases} A_{n} \pi \sin n\theta \\ 2\pi \sin n\theta \end{cases}$$
(41)

and the A<sub>n</sub> are coefficients of the Fourier sine series  
representing 
$$\theta(\theta_0)$$
, i.e., if  $\theta(\theta_0) = \sum_{p=0}^{\infty} A_p \sin p\theta_0$  or if  
 $\theta(\theta_0) = \sum_{p=0}^{\infty} B_p \cos p\theta_0$ , then the B<sub>n</sub> are the coefficients of  
the Fourier cosine series and  $p = n$ .

The above integral is evaluated by the method of stationary phase and

$$\eta(r,\theta,t) = \frac{T_{\tau}(0)}{2\pi r \sqrt{\tanh m}} \sqrt{\frac{\varphi(m)}{-\varphi'(m)}} \left\{ I_0 \overline{\eta_0}(m) + 2 \sum_{n=1}^{\infty} \overline{\eta_n}(m) I_n(\theta) \right\}$$

$$\sin (mr - \sqrt{m \tanh m} t)$$
(42)

The following table illustrates the various parameters in the above solution for specific initial deformations.



Martin Charles

1. 1 (Q) (1. 1

1.54.13.

. 84.

Figures 4, 5, 6, 7, 8, 9, 10, 11, and 12 show the wave amplitude as a function of time at six observation stations which are equidistant, r, from the origin and at various locations  $\theta$  on the surface where the initial conditions are taken from example 3 of Table II. A comparison with Fig. 3 of the previous section shows the effect of considering an asymmetric time dependent source where the only difference is the function  $\theta(\theta)$ . It should be noted that the consideration of an asymetric source is of little value when considering the effects of a single --detonation unless it happens to be an incomplete and asymmetrical detonation. However, the method is certainly applicable to multiple detonations.

The physical significance of considering the initial deformation of this form should be clearly understood. It appears, at a cursory glance, that we have obtained an asymptotic solution, valid at large distances from the source, which yields the solution for a deformation that forms in a prescribed manner from time 0 to  $\tau^*$ , and that the solution is valid for all  $t \ge \tau^*$ . However, this is not precisely the case. Actually, we are solving an initial value problem where we prescribe the shape of the water surface and the velocity field at the surface at time  $t = \tau = 0$ . The velocity field is obtained from the prescribed time dependent deformation from time 0 to  $\tau^*$ . The result is that the solution obtained will be a solution for a class of initial conditions, precisely that class which has the same surface shape and velocity field at the surface at time  $\tau = 0$ . This class is described by the set S consisting of elements  $T(\tau)$  such that

$$T(0) = 0$$
 and  $T_{\tau}(0) = \frac{\pi}{2\tau^{*}}$ 

Examples of this set include

E State

1 Singrafe

Service of the servic

12.50

the series

A66 7.11

$$T(\tau) = \sin \frac{\pi}{2} \frac{\tau}{\tau^{\pm}} + \tau^{n} + \tau^{n} \cos a\tau, n \ge 2$$
$$= \sin \frac{\pi}{2} \frac{\tau}{\tau^{\pm}} + \tau^{n} e^{b\tau^{p}}, n \ge 2$$
$$= \frac{\pi}{2} \frac{\tau}{\tau^{\pm}}$$







and the

22.47

1.25

Sector -

STRACT

Wave Amplitude From an Asymmetric Time Dependent Surface Deformation

PB-3-7385







•





Figure 11

Wave Amplitude From an Asymmetric Time Dependent Surface Deformation

PB-3-1296



That is, the solution does not depend on the precise manner in which the deformation builds up from time 0 to  $\tau^*$  but only on how it starts to build up. Furthermore, if the solution were evaluated at  $t = \tau^*$ , it would not yield the precise deformation assumed (or thought to be assumed). However, it can be expected that the solution probably would not deviate significantly from  $\eta_0(r_0, \theta_0, \tau^*)$  since the assumed initial conditions are physically reasonable conditions and the mathematical model also is physically reasonable.

North N

#### 4. METHOD OF SOLUTION IN THE AREA NEAR THE SOURCE

The asymptotic solutions (method of stationary phase) presented in the previous section are not valid in the area near the explosion. The integral with respect to m must be evaluated in some other approximate manner and the method proposed by Whalin (1965) for a symmetric water cavity is also applicable with only slight modification. The equation to be solved, in general, is

$$= \eta(\mathbf{r},\theta,\mathbf{t}) = \frac{1}{2\pi} \int_0^\infty \frac{\min \gamma \mathbf{t}}{\gamma} \left[ \mathbf{I}_0 \mathbf{J}_0(\mathbf{mr}) \,\overline{\eta}_0(\mathbf{m}) + 2 \sum_{n=1}^\infty \mathbf{I}_n(\theta) \mathbf{J}_n(\mathbf{mr}) \overline{\eta}_n(\mathbf{m}) \right] d\mathbf{m}$$

where the quantities  $I_0$ ,  $\overline{\eta_0}(m)$ ,  $I_n(\theta)$ ,  $\overline{\eta_n}(m)$ , and  $\gamma$  are the same as previously defined. The proposed method of evaluation is to fit a fourth degree polynomical to each half oscillation of the integrand and integrate directly. This method requires that all computations be performed in double precision and the upper limit of integration may be chosen to bound the error to almost any predetermined limit.

The above integral is to be approximated in the following manner

$$\eta(r, \theta, t) = \frac{1}{2\pi} \int_0^{a} g(m) dm + \frac{1}{2\pi} \int_{a}^{\beta} \frac{f(m)}{m^{1/2}} dm + \frac{1}{2\pi} \int_{\beta}^{a} h(m) dm$$

where

1

- a small number such that the oscillatory functions of the integrand may be approximated by asymptotic expansions near m = 0.
- $\beta$  = a value of m large enough so that the contribution of the latter integral is negligible and in addition is a zero of the integrand. This must be carefully chosen and depends upon  $\overline{\eta_{n}}(r_{n})$ .
- g(m) = the integrand when approximated by asymptotic expansions for small m

Having chosen the proper limits a and  $\beta$ , the first integral may be integrated directly and the latter integral is negligible. The second integral will be approximated by a polynomical fit,  $f_i(m)$ , over each half oscillation of the integrand.

$$n^{f(m)} = \frac{m^{3/2} \sin \gamma t}{\gamma} \left\{ I_0 J_0(mr) \overline{\eta_0}(m) + 2 \sum_{n=1}^{\infty} I_n(\theta) J_n(mr) \overline{\eta_n}(m) \right\}$$

where

$$\int_{0}^{\infty} \int_{0}^{\infty} \int_{0$$

 $m_{i=1} \leq m \leq m_i, i = 2, 3, \dots, n^{I}$ 

where the  $m_i$  and the ordered zeros of the integrand, f(m), and n is the number of zeros of the integrand from a up to and including  $\beta$ . Upon substitution and subsequent integration, the second integral and consequently the wave amplitude becomes

$$\eta(\mathbf{r},\theta,\theta) \Big|_{\mathbf{a}}^{\beta} = \left\{ \frac{n^{A_{1}'}}{9} \left( m_{1}^{9/2} - \mathbf{a}^{9/2} \right) + \frac{n^{B_{1}'}}{7} \left( n^{m_{1}^{7/2}} - \mathbf{a}^{7/2} \right) + \frac{n^{C_{1}'}}{5} \left( n^{m_{1}^{5/2}} - \mathbf{a}^{5/2} \right) \right. \\ \left. - \frac{4I_{0}'}{9} \left( n^{m_{1}^{3/2}} - \mathbf{a}^{3/2} \right) + \frac{n^{B_{1}'}}{3} \left( n^{m_{1}^{1/2}} - \mathbf{a}^{1/2} \right) \right. \\ \left. + \frac{n^{D_{1}'}}{3} \left( n^{m_{1}^{3/2}} - \mathbf{a}^{3/2} \right) + \frac{n^{B_{1}'}}{n^{m_{1}^{2}}} \left( n^{m_{1}^{1/2}} - \mathbf{a}^{1/2} \right) \right. \\ \left. + \frac{n^{D_{1}'}}{9} \left( n^{m_{1}^{9/2}} - n^{m_{1-1}^{9/2}} \right) + \frac{n^{B_{1}'}}{7} \left( n^{m_{1}^{7/2}} - n^{m_{1-1}^{7/2}} \right) + \frac{n^{C_{1}'}}{5} \left( n^{m_{1}^{5/2}} - n^{m_{1-1}^{5/2}} \right) \right. \\ \left. + \frac{n^{D_{1}'}}{3} \left( n^{m_{1}^{3/2}} - n^{m_{1-1}^{3/2}} \right) + \frac{n^{B_{1}'}}{n^{m_{1}^{2}}} \left( n^{m_{1}^{1/2}} - n^{m_{1-1}^{1/2}} \right) \right] \right]$$

where

.

distant.

1

ź

20 m

<u> a a seger</u>ado

3

$$n^{A'_{1} = n^{A_{1}}} n^{A'_{1} = n^{A_{1}}} n^{A'_{1} = n^{A'_{1}}} n^{A'_{1} + n^{B'_{1}}} n^{A'$$

.

33

 $n^{A_1'} = n^{A_1}$  ${}_{n}B'_{1} = {}_{n}B_{1} - {}_{n}m_{1n}A_{1}$  ${}_{n}C'_{1} = {}_{n}C_{1} - {}_{n}m_{1} {}_{n}B_{1}$  ${}_{n}D'_{1} = {}_{n}D_{1} - {}_{n}m_{1} {}_{n}C_{1}$  $\mathbf{E}_{1}^{\prime} = -\mathbf{m}_{1}^{\prime} \mathbf{m}_{1}^{\prime}$  ${}_{n}A_{1} = \frac{1}{(a - n^{b})} \left[ \frac{\binom{n^{f_{5}} - n^{f_{7}}}{(a - n^{c})} - \frac{\binom{n^{f_{6}} - n^{f_{7}}}{(n^{b} - n^{c})}}{\binom{n^{b} - n^{c}}{(n^{b} - n^{c})}} \right]$  ${}_{n}B_{1} = \frac{1}{(a - n^{b})} \left[ \frac{(a + n^{b} + n^{c})}{(n^{b} - n^{c})} \cdot (n^{f}_{6} - n^{f}_{7}) - \frac{(n^{b} + n^{c} + n^{d})}{(a - n^{c})} (n^{f}_{5} - n^{f}_{7}) \right]$  ${}_{n}C_{1} = \frac{(\frac{b}{n}c + \frac{b}{n}\frac{d}{n} + \frac{c}{n}\frac{d}{n})}{(a - \frac{b}{n})(a - \frac{c}{n})} {}_{n}f_{5} - \frac{(\frac{a}{n}c + \frac{a}{n}\frac{d}{n} + \frac{c}{n}\frac{d}{n})}{(a - \frac{b}{n})(\frac{b}{n} - \frac{c}{n})} {}_{n}f_{6} + \frac{(\frac{a}{n}b + \frac{a}{n}\frac{d}{n} + \frac{b}{n}\frac{d}{n})}{(a - \frac{c}{n})(\frac{b}{n} - \frac{c}{n})} {}_{n}f_{7}$  ${}_{n}D_{1} = a \left[ -\frac{\binom{n}{n} - \binom{\alpha}{2} n^{d}}{\frac{n}{n} \binom{n}{n} - \frac{\alpha}{n} d} + \frac{\binom{n}{n} - \frac{\alpha}{3} n^{d}}{\frac{n}{n} \binom{n}{n} - \frac{\alpha}{n} d} \right] + \frac{(\binom{n}{n} - \frac{\alpha}{3} n^{d})}{\frac{n}{n} \binom{n}{n} - \frac{\alpha}{n} \binom{n}{n} \binom{n}{n} - \frac{\alpha}{n} \binom{n}{n} \binom{n}{n} - \frac{\alpha}{n} \binom{n}{n} \binom{n}{n} \binom{n}{n} - \frac{\alpha}{n} \binom{n}{n} \binom{n}$  $+ {}_{n}c \left[ \frac{\binom{n}{b} - {}_{n}\alpha_{2} n^{d}}{\frac{n}{b} \binom{n}{b} - {}_{n}d} - \frac{\binom{a}{a} - {}_{n}\alpha_{1} n^{d}}{a(a - {}_{n}d)} \right]$  $n^{f_{2}} = \frac{-4n^{f(\frac{n^{1}+3a}{4})}}{3(n^{m_{1}}+a)}$  $n^{b} = n^{m_1} + 3a$  $n^{f_3} = \frac{-2n^{f_1}(\frac{n^{m_1}+e}{2})}{n^{m_1}-e}$  $n^{c} = n^{m_1} + a$  $n^{d} = \frac{3_n m_1 + a}{4}$  $n_{4}^{f} = \frac{-4_{n}f(\frac{s_{n}m_{1}+a}{4})}{(n_{1}m_{1}-a)}$  $n^{f_{1}} = \frac{n^{f(a)}}{n^{m-a}} a$ 

-2444

Aller

N. Kiter

$$n^{f_{5}} = \frac{n^{f_{3}} - n^{f_{4}}}{n^{c} - n^{d}} \qquad n^{\alpha_{1}} = \frac{n^{f_{1}} / n^{f_{4}}}{n^{f_{5}} - n^{d}}$$

$$n^{\alpha_{1}} = \frac{n^{f_{2}} - n^{f_{4}}}{n^{b} - n^{d}} \qquad n^{\alpha_{2}} = \frac{n^{f_{2}} / n^{f_{4}}}{n^{2} - n^{d}}$$

$$n^{\alpha_{3}} = \frac{n^{f_{3}} - n^{f_{4}}}{n^{3} - n^{4}}$$

Examples of this integration method are shown in Figs. 13 to 15 for the case of a parabolic impulse and further details can be found in Whalin (1965). The method is relatively easy to use and the accuracy of the integration procedure is well established.



and a little of the

Figure 13

A Comparison of the Asymptotic Solution for a Parabolic Impulse With an Alternate Integration Method Contraction of the second

PA-3-7398



PA-3-7393



a

#### 5. CONCLUSIONS

The preceeding theoretical developments present the limit to which one may employ a linear constant depth model to the problem of water waves generated by explosions. The present state of the art is at a point where much can be gained by the natural extensions of the theory presented in Sections 2, 3, and 4.

Data and methods are available which will enable one to relate the parameters of a time-dependent symmetrical cavity to the size and location of the weapon. In addition, there are sufficient data available from the Mono Lake tests, Hydra, and other WES tests to analyze the usefulness of considering cavity parameters which result from this model. Figures 2 and 3 indicate the difference obtained in the wave train by considering the time dependence. This order of magnitude difference in the envelope amplitude is a result of the fact that a symmetric stationary cavity contains only potential energy and the assumed time-dependent cavity from time 0 to 7\* will contain approximately the same potential energy at time r\*. However, in addition, the fluid contains kinetic energy at this time and hence the resulting vast difference in the envelope amplitudes. A thorough investigation of this form of initial conditions should prove beneficial. Also, it has been pointed out by Van Dorn that one could probably even consider the effect of the plume collapse by the proper choice of initial conditions and skillful use of land cratering data.

The utility of considering an asymmetric time-dependent form as an initial condition is obvious. If the effects of multiple explosions are to be studied, then this is the only valid method. One can not, in general, merely add the solutions of several symmetric surface deformations after obtaining the asymptotic solution for each. The total deformation of the source area must be considered initially, then transformed and integrated approximately by the method of stationary phase (m.s.p) to

obtain the correct asymptotic solution. It also should be noted that the asymmetric model is applicable to the predictions of the wave spectrum generated by an earthquake when applied to circumstances consistent with the assumptions of the mathematical model.

Predictions at Mono Lake were made from example 3 of Table I which is a symmetric surface deformation surrounded by a lip such that the volume contained in the lip is equivalent to the volume of the cavity, i.e., no net addition or removal of fluid from the system.

40

ŧ.

#### 6. **REFERENCES**

- Fuchs, R. A. (1952), "Theory of Surface Waves Produced by Underwater Explosions," Univ. of Calif., Institute of Engineering Research, Berkeley, Calif., Technical Report Series No. 3, Issue No. 335, May 3.
- Kajiura, K. (1963), "The Leading Wave of a Tsunami," Bulletin of the Eat hquake Research Institute, Vol. 41.
- Kirkwood, J. G. and Seager, R. J. (1950), "Surface Waves from an Underwater Explosion," Underwater Explosion Research - The Gas Globe, Vol. II, Office of Naval Research, Department of the Navy, pp. 707-760.
- Kranzer, H. C. and Keller, J. B. (1955), "Water Waves Produced by Explosions," IMM-NYU 222, September.
- Lamb. H. (1945), <u>Hydrodynamics</u>, Sixth Edition, Dover Publications, New York, Articles 238, 241, and 255.
- Penngy, W. G. (1950), "Gravity Waves Produced by Surface and Underwater Explosions," Underwater Explosion Research, Vol. II -The Gas Globe, Office of Naval Research, Department of the Navy, pp. 679-700.
- Pinkston, J. M. Jr. (1964), "Surface Waves Resulting from Explosions in Deep Water," U.S. Army Engineer Waterways Experiment Station, Corps of Engineering, Vicksburg, Mississippi, DASA 1482-1, Technical Report No. 1-647, Report 1, July, Confidential.
- Prins, J. E. (1956), "Characteristics of Waves Generated by a Local Surface Disturbance," Wave Research Laboratory of the Institute of Engineering Research, Univ. of Calif., Berkeley, Series 99, Issue 1, August.
- Stoker, J. J. (1947), <u>Water Waves</u>, Interscience Publishers, Inc., New York.
- Unoki, S. and Nakano, M. (1953), "On the Cauchy-Poisson Waves Caused by the Eruption of a Submarine Volcano," Oceanographical Magazine, (1st paper) Vol. 4, No. 4, pp. 119-141, 1953a; (2nd paper) Vol. 5, 1 -, 1, pp. 1-13, 1953b; (3rd paper), Vol. 4, No. 3-4, pp. 139-150, 1953c.
- Van Dorn, W. G. (1963), "Water Waves from 10,000-1b. High-Explosive Charges," SIO Reference 63-20, June.

and the second second second

Van Dorn, W. G. (1959), "Impulsively Generated Waves," SIO Report No. 11.

- Van Dorn, W. G. (1964), "Explosion Generated Waves in Water of Variable Depth," Sears Foundation: Journal of Marine Research, Vol. 22, No. 2, May 15, pp. 123-141.
- Whalin, R. W. (1965b), "Water Waves Generated by Explosions: Propagation Theory for the Area Near the Explosion," Journal of Geophysical Research, Vol. 70, No. 22, November 15.

Whalin, R. W. (1965a), "Research on the Generation and Propagation of Water Waves Produced by Explosions, Part II: A Prediction Method," NMC-IEC, March, Confidential.

42

7.AV

£

.

2	an integration limit
n <sup>b</sup> i' n <sup>c</sup> i' n <sup>d</sup> i	functions of m <sub>i</sub>
$n^{f_{1'}n^{f_{2'}n^{f_{3,}}}}$	
n <sup>f</sup> 4' n <sup>f</sup> 5' n <sup>f</sup> 6'	functions of m
n <sup>£</sup> 7 )	
đ	water depth
g(m), f(m) h(m)	functions
g	acceleration of gravity
i, j, k, n	summation indices
m	dimensionless wave number
n <sup>m</sup> i	zeros of the integrand (i.e., the $m_i$ are the ordered zeros of $n^{f(m)}$
р	dimensionless pressure
p'	pressure
r	dimensionless radial distance from the origin
r'	radial distance from the origin
T	distance from a point $(r, \theta)$ to a source point $(r_0, \theta_0)$
r <sub>o</sub> , θ <sub>o</sub> , z <sub>o</sub>	space variables of a source point (cylindrical coordinate)
t	dimensionless time
t'	time
X, Y, 2	dimensionless space variables (Cartesian coordinate system)

x', y', z' space variables (Cartesian coordinate system) dimensionless bottom velocity w<sub>B</sub> A, <sub>n</sub>A<sub>1</sub>, <sub>n</sub>A<sub>i</sub>, constants  $n^{A'_{1}}$   $n^{A'_{1}}$ B, B<sub>1</sub>, B<sub>i</sub>, constants  $B'_{1'}, B'_{i}$  $C_{n}C_{1}, C_{i}$ constants  $\tilde{c}_{i}, c_{i}'$  $D_{i_{n}}D_{i_{1},n}D_{i_{1},n}$ constants  ${}_{n}D_{i'n}D_{i}'$  $\mathbf{E}_{i_{n}}\mathbf{E}_{i_{n}}\mathbf{E}_{i_{n}}$ constants  $_{n}E_{1}^{\prime}, _{n}E_{1}^{\prime}$ G time-dependent Green's Function

a function describing an initial impulse 10 I<sub>1</sub>, I<sub>2</sub>, I<sub>3</sub> integrals the number of zeros of the integran I<sub>n</sub> Γ, sero order Hankel transform of I n<sup>th</sup> order Bessel function J<sub>n</sub>(mr) distance between points (x, y, z) and  $x_0, y_0, z_0$ ; also a parameter of the initial disturbance R S a set; also a surface T(T) a function describing the time dependence of the initial deformation

44

v	dimensionless velocity
v'	velocity
$n^{\alpha_1}, n^{\alpha_2}, n^{\alpha_3}$	functions of m
β	an integration limit
Ŷ	"√m tanh m
η	dimensionless wave amplitude
η'	wave amplitude
$\overline{\eta_n}$	n <sup>th</sup> order Hankel transform of $\eta_0$
η	function describing the initial deformation
λ	dimensionless wave length
λ'	wave length
Ħ	constant
ρ	water density
7	a dimensionless time variable of the source
T*	a constant
τ'	a time variable of the source
¢	velocity potential (subscripts denote partial differentiation)

.

:

Ł

1

N B RUYA

[

[.

\_\_\_\_.

er nemer

a,

and the set of the set

and the second secon

### ACKNOWLEDGEMENTS

1

11.44910

E

Ľ

Ľ

I wish to acknowledge the assistance of Dr. W. G. Van Dorn of Scripps Institution of Oceanography for calling my attention to the paper of Kajiura (1963) and for several helpful discussions related to the contents of this report.

Security Classifiaction					
	COMPREL DATA · RED	and a second second theorem is a second s			
Growniy clocelfication of Hills, body of cho 1 Opinian and a children framework	terestine ed teurs materiana (mission) Ens tostia				
	20. 11	PORT SECURITY CLASSIFICATION			
National Engineering Scien	ce Company				
rasadena, California 9110	)5 N	/A			
3. REPORT TITLE					
Contributions to the Mono I	Lake Experiments				
4. DESCRIPTIVE NOTES (Type of report and mek	uelve detec)				
Final Report					
J. AUTHOR(3) (Lost name, Mast name, Initial)					
Whalin, R. W					
Webb, L. M.					
. REPORT DATE	74. 70746 NO. OF PAGES	78. NO. OF REFS			
October, 1965	193	51			
SO. CONTRACT OR GRANT NO.	94. OMENATOR'S REPORT N	umeer(s)			
Nonr 5006(00)	S 256-2				
A PROJECT NO.					
	] 같은, 주 특히 중간 중간 무료 문 부 원건(3) /4	AV AGINE BUILDER GAN MAN AA ARAI MAA			
<del>.</del>	De other report ners) (A	Ay effice multiple fiet may be easigned			
	D. OTNER REPORT NELL (A	ny office numbers that may be assigned			
G. 16. A V A IL ABILITY/LIMITATION NOTICES	D. OTHER REPORT NO(2) (A	hy office sumbore the may be designed			
d. 18. AVAILABILITY/LIMITATION NOTICES Qualified requesters may of	btain copies of this report fr	ny office numbers that may be easigned			
d. 18. AVAILASILITY/LIMITATION NOTICES Qualified requesters may of	btain copies of this report from	by other numbers ther may be easigned			
d. 18. A VA IL ABILITY/LIMITATION NOTICES Qualified requesters may of 11. SUPPLEMENTARY NOTES	btain copies of this report fr	by other numbers the may be decigned			
d. 18. AVAILABILITY/LIMITATION NOTICES Qualified requestors may of 11. SUPPLEMENTARY NOTES	btain copies of this report fro 12. SPG223CR:HS MILITARY AG Department of the	ny office numbers that may be easifyted			
d. 18. AVAILASILITY/LIMITATION NOTICES Qualified requesters may d 11. SUPPLEMENTARY NOTES	btain copies of this report fro 18. SPC23CR:MS MILITARY AC Department of the Office of Naval 1 Washington, D.	hy other numbers that may be easigned form DDC. STRUTY he Navy Research C. 20360			
d. 18. A VA IL ABILITY/LIMITATION NOTICES Qualified requesters may d 11. SUPPLEMENTARY NOTES 13. ABSTRACT	btain copies of this report fro lis spony military and Department of the Office of Naval 1 Washington, D.	hy other numbers ther may be easigned form DDC. Truity he Navy Research C. 20360			
d. 18. A VA IL ABILITY/LIMITATION NOTICES Qualified requesters may of 11. SUPPLEMENTARY NOTES 13. ADSTRACT GRAVITY WARDON OF COMPANY	btain copies of this report fro 12. SPG223CR:KS MILITARY AC Department of the Office of Naval 1 Washington, D. (	hy other numbers that may be easigned om DDC, truity te Navy Research C, 20360			
d. 16. A VA IL ABILITY/LIMITATION NOTICES Qualified requesters may of 11. SUPPLEMENTARY NOTES 13. ABSTRACT Gravity waves genera	btain copies of this report fro 12. SPOZESCRIMS MILITARY AG Department of the Office of Naval 1 Washington, D. ( Ated by underwater explosions	by other numbers that may be seel (net for DDC, trivity he Navy Research C, 20360 and the wave run-up			
d. 18. AVAILASILITY/LIMITATION NOTICES Qualified requesters may d 11. SUPPLEMENTARY NOTES 13. ABSTRACT Gravity waves genera are predicted for a series d	btain copies of this report fro lis specific ment of the Office of Naval 1 Washington, D. Atted by underwater explosions of tests to be performed in M	br DDC. Twity he Navy Research C. 20360 and the wave run-up ono Lake (California).			
d. 18. AVAILABILITY/LIMITATION NOTICES Qualified requestors may of 11. SUPPLEMENTARY HOTES 13. ABSTRACT Gravity waves genera are predicted for a series of The time histories of the wa	btain copies of this report fro lis specific and marked with the sport lis specific and the sport fro Department of the Office of Naval 1 Washington, D. ( ated by underwater explosions of tests to be performed in M ave profiles are given at vari	by other numbers that may be seeifned born DDC. Trwity he Navy Research C. 20360 and the wave run-up ono Lake (California). lous locations in the			
d. 18. AVAILABILITY/LIMITATION HOTICES Qualified requesters may of 11. SUPPLEMENTARY HOTES 13. ABSTRACT Gravity waves genera are predicted for a series of The time histories of the wallake in deep and shallow wa	btain copies of this report from is spony is spony is spony btain copies of this report from Department of the Office of Naval Washington, D. ited by underwater explosions of tests to be performed in M ave profiles are given at vari- iter where wave recorders with	bm DDC. The Navy Research C. 20360 and the wave run-up ono Lake (California). lous locations in the 11 be installed. The			
d. 16. AVAILABILITY/LIMITATION NOTICES Qualified requesters may of 11. SUPPLEMENTARY NOTES 13. ABSTRACT Gravity waves genera are predicted for a series of The time histories of the wallake in deep and shallow walling of a series of the series of the series of a series of a series of the series of a	btain copies of this report from 12. SPGESCRIMS MILITARY AC Department of the Office of Naval in Washington, D. Ated by underwater explosions of tests to be performed in M ave profiles are given at variations ater where wave recorders with	by other numbers that may be seel() ter proving the Navy Research C. 20360 and the wave run-up ono Lake (California). Hous locations in the 11 be installed. The			
d. 16. AVAILABILITY/LIMITATION NOTICES Qualified requestors may of 11. SUPPLEMENTARY NOTES 13. ABSTRACT Gravity waves genera are predicted for a series of The time histories of the we lake in deep and shallow wa times of arrival and heights	btain copies of this report from lis special special states and the special s	orn DDC. Trwity he Navy Research C. 20360 and the wave run-up ono Lake (California). lous locations in the 11 be installed. The ned at three locations			
d. 16. AVAILABILITY/LIMITATION NOTICES Qualified requesters may of 11. SUPPLEMENTARY HOTES 13. ABSTRACT Gravity waves genera are predicted for a series of The time histories of the wa lake in deep and shallow wa times of arrival and heights on the shoreline of Mono La	btain copies of this report from is spony btain copies of this report from Department of the Office of Naval 1 Washington, D. ( ated by underwater explosions of tests to be performed in M ave profiles are given at vari- ater where wave recorders with s of wave run-up are determinate.	by other numbers that may be seel(free from DDC. Troury Research C. 20360 and the wave run-up ono Lake (California). lous locations in the ill be installed. The med at three locations			
d. 18. AVAILABILITY/LIMITATION HOTICES Qualified requesters may of 11. SUPPLEMENTARY HOTES 13. ABSTRACT Gravity waves genera are predicted for a series of The time histories of the we lake in deep and shallow wa times of arrival and heights on the shoreline of Mono La This prediction is dor	btain copies of this report from the appoint of the second secon	by other numbers that may be seeifned orn DDC. Trwity he Navy Research C. 20360 and the wave run-up ono Lake (California). Hous locations in the ill be installed. The ned at three locations advanced theories and			
d. 16. AVAILABILITY/LIMITATION NOTICES Qualified requesters may of 11. SUPPLEMENTARY NOTES 13. ABSTRACT Gravity waves genera are predicted for a series of The time histories of the will lake in deep and shallow wa times of arrival and heights on the shoreline of Mono La This prediction is dor all available information in	btain copies of this report from the spony btain copies of this report from the spony btain copies of this report from the spony btain copies of this report from the spony the spony btain copies of this report from the spony the sp	om DDC. Troirv he Navy Research C. 20360 and the wave run-up ono Lake (California). lous locations in the 11 be installed. The ned at three locations advanced theories and pretical developments			
d. 16. AVAILABILITY/LIMITATION HOTICES Qualified requesters may of 11. SUPPLEMENTARY HOTES 13. ABSTRACT Gravity waves genera are predicted for a series of The time histories of the will lake in deep and shallow wa times of arrival and heights on the shoreline of Mono La This prediction is dor all available information in are presented in Volumes II	btain copies of this report from the spony btain copies of this report from the spony btain copies of this report from the spony be spony to files of Naval Washington, D. the spony to files are given at vari- ther where wave recorders will a of wave run-up are determinate. The by making use of the most that field, and some new the I and III, namely:	by other numbers that may be easigned orn DDC. STRUTY the Navy Research C. 20360 and the wave run-up ono Lake (California). Hous locations in the III be installed. The med at three locations advanced theories and oretical developments			
d. 16. AVAILABILITY/LIMITATION NOTICES Qualified requesters may of 11. SUPPLEMENTARY NOTES 13. ABSTRACT Gravity waves genera are predicted for a series of The time histories of the will lake in deep and shallow wa times of arrival and heights on the shoreline of Mono La This prediction is dor all available information in are presented in Volumes II 1) A linear theo	btain copies of this report from the approximation of the second of tests to be performed in M ave profiles are given at variants of the second of the sec	om DDC. Trivity he Navy Research C. 20360 and the wave run-up ono Lake (California). lous locations in the 11 be installed. The ned at three locations advanced theories and oretical developments plosions, making			
d 18 AVAILABILITY/LIMITATION NOTICES Qualified requesters may of 11 SUPPLEMENTARY NOTES 13 ABSTRACT Gravity waves genera are predicted for a series of The time histories of the will lake in deep and shallow wa times of arrival and heights on the shoreline of Mono La This prediction is dor all available information in are presented in Volumes II I) A linear theo use of a symmetric as	btain copies of this report from is specification with a specific the specific of the second is specific of the second of the specific of the second of th	by other numbers that may be easifyed by DDC. Trouvy Research C. 20360 and the wave run-up ono Lake (California). lous locations in the 11 be installed. The ned at three locations advanced theories and oretical developments plosions, making at surface disturbance			

2) A linear theory for the propagation of periodic waves over the continental slope.

DD .: 1473

( With

304 m.C.

I

I

UNCLASSIFIED Security Classification

HE LACED ST

UNCLASSIFIED

...

	LINK A		LINA B		LINXC				
KEF WORDS		ROLE	w1	ROLE	WT	ROLK	<b>#</b> T		
Water waves				· ·					
Underwater explosions									
Wave run-up									
Waves on a slope	•								
	×								
	· Ilevitorst								
ORIGINATING ACTIVITY: Enter the name and address	110. AVA	ILABILIT	Y/LN31	ATION N	OTICES	t Enter a	ny lim-		
he contractor, subcontractor, grantee, Department of De- se activity or othir organization (corporate author) issuing	itations (	on further by securi	dissemin ty classi	nation of I fication, 1	the report using sti	t, other th Inderd stat	an those Ioments		
REPORT SECURITY CLASSIFICATION: Enter the over-	such as	"Qualifie	d recuses	ters may (	istain co	pies of th	ie		
security classification of the report. Indicate whether estricted Data" is included. Marking is to be in accord- e with appropriate security regulations.	(2)	report fro "Foreign	m DBC."	ement and	dissom	instion of	this		
GROUP: Automatic downgrading is specified in DoD Di- tive \$200.10 and Armed Forces Industrial Manual. Enter	(3)	report by "U. S. G	DDC 18 1 DVeri.mett	t sgencier	mey ob	tain copie	s of		
group number. Also, when applicable, show that optional things have been used for Group 3 and Group 4 as author- d.	this report directly from DDC. Other qualified DDC users shall request through								
REPORT TITLE: Enter the complete report title in all	(4)	"U. S. m	litery eg	encies m	y obtain	copies of	this		
in meaningful tile cannot be selected without classifica- n, show tile classification in all capitals in parenthesis and state following the tile.	report directly from DDC. Other qualified users shall request through					"			
DESCRIPTIVE NOTES: If appropriate, enter the type of	(5)	"All dist	ribution ( C usors a	of this rep hall requi	ort is constitution	ontrolled gh	Qual-		
or the inclusive dates when a specific reporting period is						then of Te	m + <sup>84</sup>		
AUTHOR(S): Enter the name(s) of author(s) as shown on in the report. Enter last name, first name, middle initial.		If the report has been runished to the Office of Technical Services, Department of Commerce, for sale to the public, indi- cate this fact and enter the price, if known							
principal author is an absolute minimum requirement.	11. BUP tory note	Pleksn M	TARY N	OTER U	se for s	ditional e	spisate		
REPORT DATE: Enter the dele of the report as Gay, nth, year, or month, year. If more than one date epyears the report, use date of publication.	12. EPONODRINO HILITARY ACTIVITY: Enter the same of the departmental project office or laboratory sponsoring (per-								
TOTAL NUMBER OF PACES: The total page count ild follow normal pagination procedures, i.e., enter the 13. AISTRACT: Enter en abstract giving a brief and factual und follow normal pagination procedures, i.e., enter the									
NOW OF PAGES CONTAINIng INFORMATION NUMBER OF REFERENCES. Enter the total number of merces stud in the second.	it may a port. If abalt be	ndditions	r elsewho 1 space i	ere in the erequired	body ef i, a cont	the techni Invation of	cai te- heel		
CONTRACT OR GRANT NUMBER: If appropriate, enter applicable number of the contract or gram under which	li is porte he	highly de unclassi	atrable t	hat the ab	attact a ph of th	f classifie abatract	d rø- shall		
report was written. A. A. S. PROJECT NUMBER: Enter the appropriate	ond with of the in	i on Indici formation	tion of t in the p	ne militer pregruph,	7 80° Wi 700101	ry clausifi lied as (73	r ut (78 5), <b>(\$)</b> ,		
itary department identification, such as project number, project number, system numbers, task number, etc.	(C), or ( Ther	v). • Is an li	milation (	nn the len	gth <b>of</b> th	e sheiisci S words.	How-		
ChiGINATON'S REPORT NUMBER(8): Enter the offi- t report number by which the document will be identified	14. KEY	· #O#D#:	Key wo	rds are te	hnicell	y meaning!	ut terms		
s controlled by the originating activity. This number must unique to this report.	or shutt index or	physes 1 irise for	nat chard catalogin	a the repr	vepnet a vet. Key tration	ne mey bê · worda mu a reacteed	usen 88. al be ; liten:		
OTHER REPORT NUMBER(8). If the second has been signed any other report numbers (either by the originator	fiera, a	· usr 1741 1 µrh 66 Fili 14-1 : cula	nane es	sodet desi	gnatiss,	traile name	april 84		
by the approache), also enter this number(s).	tary wot	ta hut wil The asi	the full-	word by a of tinks.	n trits a rules, at	tion of the od whighth	hnii <b>8</b> 8 18		
				****	A 254 - E 4	***	<u>م التي نيم من المراجع الم</u>		
		<b></b>		UNCL	1221	時代。			

IJ

Ū

0

8

[

8

8

]]

1

9

]

]

J

[]

[]