

UNCLASSIFIED

AD NUMBER
AD491366
NEW LIMITATION CHANGE
TO Approved for public release, distribution unlimited
FROM Distribution: No foreign
AUTHORITY
USAPA ltr., 7 Oct 1971

THIS PAGE IS UNCLASSIFIED

UNCLASSIFIED

AD. 491 366

DEFENSE DOCUMENTATION CENTER  
FOR  
SCIENTIFIC AND TECHNICAL INFORMATION

CAMERON STATION ALEXANDRIA, VIRGINIA

Classification cancelled in accordance with  
Executive Order 10501 issued 5 November 1953

Document Service Center  
Armed Services Tech. Info Agency



UNCLASSIFIED

APR 30 1969  
LIBRARY

RETURN TO:  
ASIA REFERENCE CENTER  
LIBRARY OF CONGRESS  
WASHINGTON, D.C. 20540

This document is the property of the United States Government. It is furnished for the duration of the contract and shall be returned when no longer required, or upon recall by ASTIA to the following address:  
Armed Services Technical Information Agency, Arlington Hall Station,  
Arlington 12, Virginia

**NOTICE: THIS DOCUMENT CONTAINS INFORMATION AFFECTING THE NATIONAL DEFENSE OF THE UNITED STATES WITHIN THE MEANING OF THE ESPIONAGE LAWS, TITLE 18, U.S.C., SECTIONS 793 and 794. THE TRANSMISSION OR THE REVELATION OF ITS CONTENTS IN ANY MANNER TO AN UNAUTHORIZED PERSON IS PROHIBITED BY LAW.**

**AWA**

**BEST**

**AVAILABLE**

**COPY**

and

Station,

**AFFECTING THE**

**THE MEANING**

**793 and 794.**

**TENTS IN**

**IBITED BY LAW.**

307/22

NOV 19 1954

**FCG  
LTC**

**FILE COPY  
MAY BE LOANED TO  
LIBRARY OF CONGRESS  
TO BE RETURNED**

**NOTICE**  
This document is the property of the United States Government. It is loaned on a non-exclusive basis and shall be returned as specified, or upon request by ASTIA to the following address: ASTIA Reference Center, Technical Information Division, Library of Congress, Washington 25, D.C.

**RETURN TO:  
ASTIA REFERENCE CENTER  
LIBRARY OF CONGRESS  
WASHINGTON 25, D.C.**

**TECHNICAL INFORMATION CENTER  
SUBSIDIARY  
(MAY BE LOANED TO OTHERS)**

**ASTIA**

CONFIDENTIAL  
SECURITY INFORMATION

Copy No. 15

REPORT

ON

STUDY OF PURE EXPLOSIVE COMPOUNDS

Part IV

Calculation of Heat of Combustion

of

Organic Compounds from Structural Features

and Calculation of Power of High Explosives

to

OFFICE OF THE CHIEF OF ORDNANCE

Contract No. DA-19-021-ORD-47

C-53247

May 1, 1952

"This document contains information affecting the national defense of the United States within the meaning of the Espionage Act, U.S.C. 50:31 and 32. Its transmission or the revelation of its contents in any manner to an unauthorized person is prohibited by law."

CONFIDENTIAL

Arthur D. Little, Inc.

CONFIDENTIAL  
SECURITY INFORMATION

DEPARTMENT OF THE ARMY  
Boston Ordnance District  
Army Base  
Boston 19, Mass.

SUBJECT: "Fundamental Research on High Explosives. Part IV."  
Prepared by Arthur D. Little, Inc., under Contract  
DA-19-020-ORD-47

TO:

Attention:

1. You are furnished herewith Part IV of "Fundamental Research on High Explosives". This is the fourth of a series of reports prepared by Arthur D. Little, Inc., under a contract with the Ordnance Department comprising research and development in the field of military explosives.

2. This Report is furnished for your use under the limitations applicable to its security classification. If you do not desire to retain this Report, it is requested that it be returned to the Office, Chief of Ordnance, Ammunition Development Division, Research and Development Division, Washington 25, D. C.

3. Your comments, criticism and remarks relative to any and all aspects of the subject matter of this Report are invited, and should be directed to the Contracting Officer, Boston Ordnance District.

4. It is requested that the original of the inclosed receipt be signed and returned as indicated. The carbon copy may be retained for your files.

FOR THE DISTRICT CHIEF:

*A. C. YOKESAS*  
A. C. YOKESAS  
Lt Col, Ord Corps  
Assistant

1 Incl:  
Receipt

CONFIDENTIAL

Arthur D. Little, Inc.



TABLE OF CONTENTS

	<u>Page</u> <u>Number</u>
Letter of Transmittal	461
Summary	463
Recommendations	465
Section XVI. Calculation of Heats of Combustion of Organic Compounds	467
A. Abstract	467
B. Introduction	467
C. Discussion	467
1. Comparison with Other Methods	467
2. Congruence of Observation and Calculation	468
3. Relationship among Functional Groups	469
4. Multiple Groups on a Single Carbon Atom	475
D. Conclusions	476
E. Rules and Procedure for Calculating Heat of Combustion with Examples	478
F. Tables of Data	488
1. Scope	488
2. Method of Obtaining Least Squares Equations	488
3. Arrangement of Heat of Combustion Data	482
G. Historical	490
1. Methods of Summation of Bond Energies	490
2. Methods of Summation using a Base Value plus Group Corrections	492
3. Methods Based on Linear Equations	492
4. Methods Based on Group Substitutions	494
H. References	495
I. Glossary of Terms and Symbols	513
J. Heat of Combustion Data	514

TABLE OF CONTENTS (cont'd)

	<u>Page Number</u>
Section XVII. Calculation of Power (NRT) of Explosives Compounds and Mixtures	575
A. Abstract	575
B. Introduction	575
C. Discussion	577
1. Organic Compounds	590
2. Explosives Mixtures	591
3. Metallized Mixtures	591
4. Correlation of Spherical Lead Block Measurements and NRT	593
5. Application to New Compounds and Mixtures	596
D. Method of Calculation	596
E. Conclusions and Recommendations	613
F. References	615
Distribution List	617

LIST OF TABLES

<u>Table Number</u>	<u>Title</u>	<u>Page Number</u>
<u>Section XVI</u>		
58	Pisemotive Series. Contribution of Functional Groups to Heat of Combustion.	472
59	Illustration of Heat of Combustion Calculations.	481
60	Coefficients of Equations for Heat of Combustion.	484
61-138	Heat of Combustion Data.	510-555
139	List of Compounds	560
<u>Section XVII</u>		
140	Ratio of Specific Heats, $\frac{C_p}{C_v}$ , for Product Gases for Selected High Explosives.	578
141	Thermodynamic Data for Selected Reactions.	579
142	Thermodynamic Data for Selected Metals and their Corresponding Oxides.	579
143	Properties of Selected Pure Organic Compounds.	585
144	Properties of Selected Explosives Mixtures.	588
145	Properties of Selected Metallized Explosives Mixtures.	589
146	Pyre Compounds Showing Large Differences Between Observed Ballistic Mortar and Calculated nRT Values.	590
147	Correlation Between Calculated nRT and Spherical Lead Block Measurements.	595
148	Properties of Selected High Energy Compounds.	596
149	Average $C_v$ Over Range from 298°K to T (calories/mole/°K).	604

CONFIDENTIAL

Arthur D. Little, Inc.

CONFIDENTIAL  
SECURITY INFORMATION

LIST OF FIGURES

<u>Figure Number</u>	<u>Title</u>	<u>Page Number</u>
<u>Section XVI</u>		
79	Reduced Heat of Combustion (y') vs. Oxygen Needed (x')	470
<u>Section XVII</u>		
80	Comparison of Ballistic Mortar and Calculated nRT for Pure Explosive Compounds	581
81	Comparison of Ballistic Mortar and Calculated nRT for Explosive Organic Mixtures	582
82	Comparison of Ballistic Mortar and Calculated nRT for Explosive Metallic Mixtures	583
83	Statistical Distribution of Deviation from the Mean Comparing Ballistic Mortar and nRT for Explosive Substances	584
84	Comparison of Ballistic Mortar and Calculated nRT for Cyclotols (RDX/TNT)	592
85	Comparison of Calculated nRT, Ballistic Mortar and Spherical Lead Block for Tritonals (TNT/Al)	594
86	Calculation Sheet, RDX	596
87	Calculation Sheet (Reverse Side), RDX	599
88	Calculation Sheet, Ednatol	600
89	Calculation Sheet (Reverse Side), Ednatol	601
90	Calculation Sheet, Torpex	602
91	Calculation Sheet (Reverse Side), Torpex II	603

CONFIDENTIAL

Arthur D. Little, Inc.

# Arthur D. Little, Inc.

RESEARCH · ENGINEERING · INVESTIGATION



CAMBRIDGE 42, MASSACHUSETTS

Office of the Chief of Ordnance  
Pentagon Building  
Washington 25, D.C.

Contract No. DA-19-020-ORD-47  
C-58247

Gentlemen:

We are transmitting herewith Part IV, "Calculation of Heat of Combustion of Organic Compounds from Structural Features and Calculation of Power of High Explosives", continuing our fundamental study of explosives under the subject contract. In the earlier parts of this report we developed approximate methods for predicting performance in certain explosive tests from the concepts of oxygen balance and phosphoric groups, and from heat of explosion. These approximations were shown to be useful for both pure explosives and mixed explosives.

In the first section of this report a method for calculating heats of combustion of organic compounds, and hence heats of explosion, from structural features alone has been developed. The method is based entirely on observed heats of combustion for both explosive and non-explosive organic compounds. It represents a distinct improvement over the calculation method presented in Part II, and possesses several features which make it more adaptable and suited for our purposes than other methods proposed in the literature.

The second section of this report demonstrates the application of a system for predicting the power of an explosive, relative to TNT (or to any other substance). A comparison is made between the calculated power based on calculated heat of combustion, and the power observed in the ballistic mortar for pure compounds and for mixtures, both metallized and non-metallized. This method of prediction results in an accuracy much improved over the use of the previous concepts given in Parts I and III.

Recommendations are made for experimental work to confirm and expand the conclusions reached by this treatise.

Respectfully submitted,

*Arthur D. Little, Inc.*

CONFIDENTIAL  
SECURITY INFORMATION

SUMMARY

This report describes a new method for calculating heats of combustion of explosives and non-explosive organic compounds and a method for calculating relative powers of explosive compounds from heats of combustion.

It is shown that a straight-line relationship exists between molar heat of combustion ( $Q_c$ ) and molar oxygen balance for a large number of homologous series of organic compounds. From the slope and intercept coefficients of the straight lines obtained from observed data a method has been devised for calculating the heat of combustion at constant pressure (with liquid) of an organic compound, considering only the structural features of the molecule. Simple rules and examples of use are explained. The agreement between calculated and observed heat of combustion is generally better than 1%, although the variability is more pronounced in members of homologous series whose determinations are known with less assurance of accuracy.

It is shown that the coefficients of the equations can be used as a measure of the relative desirability of any functional group in an explosive compound. The coefficients have been arranged to give a quantitative idea of the energy contribution of any functional group in a molecule, based on the paraffin hydrocarbon as the zero energy potential. This means that the potentialities of a functional type can be assessed for explosive usefulness without necessity of preparing oxygen balanced compounds. Instead, an approximation can be made from a small number of organic substances containing the functional group of interest, but not necessarily any primary phosphoric group, thus eliminating the synthetic difficulties and attendant danger usually encountered

CONFIDENTIAL

CONFIDENTIAL  
SECURITY INFORMATION

with oxygen balanced molecules. After the potentialities of the group have been determined, it can then be decided whether the effort to prepare balanced explosives is worthwhile.

This system of calculation is compared with methods proposed by others. A historical section dealing with published heats of combustion completes the presentation of this subject matter.

The relative power of an explosive compound can be calculated from its heat of combustion by an iteration method employing the estimated temperature and the moles of gas produced in the detonation. The resulting figure (= nRT) compared to the value for TNI is equal to the power of the explosive as measured in the ballistic mortar or in the spherical lead block on an equivalent weight basis. A comparison of results with more than 100 pure compounds and more than 50 organic mixtures is in agreement with this conclusion. Greater harmony is achieved if the calculated rather than observed heat of combustion is used.

The results with a large number of metallized mixtures, while not giving as close agreement as the non-metallized substances do, show that the use of nRT to predict power for this type of mixture is definitely superior to the use of the concepts of oxygen balance or heat of explosion alone, as described in Part III of this report.

The nRT method, combined with the method for calculating heat of combustion can be used to predict the expected power, as it might be measured in the ballistic mortar compared to a standard explosive, of any organic compound or mixture whose structural features are known or postulated. A quantitative arrangement of the many types of functional groups from the viewpoint of power

CONFIDENTIAL

CONFIDENTIAL  
SECURITY INFORMATION

contribution in an explosive compound or mixture can also be obtained from use of this method for calculating power.

Thus, prior to synthesis, it can be determined with some degree of accuracy whether the effort to prepare oxygen balanced explosives containing any given functional type is worthwhile, and what functional types or mixtures might be expected to produce better power performance than the existing most powerful organic explosives.

RECOMMENDATIONS

1. Determination of the contribution to heat of combustion and explosion has been recommended for the nitroso group and for the following functional types of interest to the explosives program but not treated in this report:

azide

1,2,3-triazole and 1,2,4-triazole

furoxan and furazan

triazene, tetrazene, and other linear nitrogen chain

azine

hydrazone

imide

s-triazine

imine or anil

hydroxylamine

nitrite

perchlorate ester

CONFIDENTIAL



CONFIDENTIAL  
SECURITY INFORMATION

2. For general interest the contribution of the following groups should be estimated:

thiol

sulfide

disulfide

sulfonic acid

chloro

bromo

iodo

isocyanate

peroxide

3. Heat of combustion should be redetermined for 2,4,6-trinitroresorcinol (styphnic acid) and for  $\alpha, \alpha, \alpha$ -trifluoro-3,5-dinitrotoluene. 5-Nitraminotetrazole should be measured for heat of combustion.

None of the specific compounds and mixtures of negative oxygen balance whose observed powers are widely different from calculation are recommended for redetermination. Their low theoretical powers make them basically of no interest to a project endeavoring to improve on the present service explosives. Recommendations are made for a program of testing metallized mixtures containing high nitrogen compounds expected to have exceptionally high power, for testing certain high energy compounds, and for retesting certain substances of positive oxygen balance, such as hexanitroethane, which appear to disagree with the general theory of power calculated by NRI.

CONFIDENTIAL

SECTION XVI  
Calculation of Heats of Combustion of  
Organic Compounds

A. Abstract

A method of calculating heats of combustion of organic compounds from structural features is described. This method depends on the straight line relationship between molar oxygen balance and the molar heat of combustion of a homologous series. Intercept and slope values have been determined from published data for thirty-eight structural features. On the average, an accuracy of 1% is found by comparison of observed with calculated heats of combustion. The usefulness of the method for high explosives is illustrated and certain rules of procedure have been established.

B. Introduction

In an earlier report (70) a method of additive group energies was proposed for calculating heats of combustion of explosive compounds. This method was converted from the Springall and Roberts system of calculating heats of formation (110). Calculation gave good agreement with observation in most cases for organic compounds containing primary phosphoric groups (69) but it was later discovered that very poor agreement was frequently being obtained with non-explosive organic compounds whose observed values were undoubtedly correct. This poor agreement, as well as the lack of values for certain common structural features, led to a reexamination of the problem from a more empirical and less theoretical viewpoint than that of Springall and Roberts.

C. Discussion

1. Comparison with Other Methods

Holcomb et al described a relation between oxygen balance and heat of combustion on a weight basis for nitroalkanes (14) and later for nitroalcohols (31a). We have found that a similar relationship is applicable to a great many more functional types. The relationship is found to give a better straight line, however, between molar oxygen balance and molar heat of combustion, than between these parameters on a weight basis. Indeed such a weight relation does not exist for molecules with the empirical formula  $C_nH_{2n}$ . Olefins and cyclic paraffins all have the same oxygen balance on a weight basis (-342.2) and the weight heat of combustion becomes independent of this parameter. It was discovered subsequently that relationships between the molar heat of combustion and the number of carbon atoms in the molecule had been published by several investigators (11, 65, 100, 103, 113, 144, 146, and 148). None of them, though, had been or could be applied to any molecules containing more than a single type of functional group, and they were straight lines in any homologous series only above five carbon atoms.

CONFIDENTIAL

CONFIDENTIAL  
SECURITY INFORMATION

Relating the heat of combustion to the number of carbon atoms is neither as effective nor as flexible as relating it to the molar oxygen balance which is done here for the first time. When one considers that numerous polyfunctional compounds can have the same number of carbon atoms but different heats of combustion, necessitating an equation for each homologous series, it is apparent that the task is simplified by use of oxygen balance which leads to a single equation for any homofunctional type regardless of the number of like functional groups present.

The method of Kharasch (55) is as close as any to the system described in this report. Yet the difficulties in calculating the number of electrons in a molecule are definitely greater than in computing the oxygen balance, a figure which can be obtained from the molecular formula alone, independent of electronic structure. Furthermore, no corrections to be found for functional types not given by Kharasch are difficult to assign and assess with any assurance of accuracy (1). His method of derivation does not involve the straight-line relationship which smooths out errors as it does in the system described below.

The system presented here has one distinct advantage over all previously used methods. The linearity of  $Q_c$  for any homologous series with molar oxygen balance allows a reasonable estimation of the accuracy of any reported heat of combustion. Isolated points which do not fall on the best line or within a reasonable range of it arouse suspicion of inaccurate determination for one reason or another. The truthfulness of this statement has been upheld and justified by a number of instances in the pertinent field of explosive chemistry, in which a determination, discarded by us as being incorrect, has been revised by later determination or by other authors, bringing the observed heat of combustion in line with the calculated value. An example of similar reasoning applied to structure proof can be found in the recent literature (72). Thus many of the data in the Tables 61 through 130 show the effects of some selection, and the points chosen for estimating the best line by the least-squares method have been biased, but only for this reason.

The list of functional groups used may be considered incomplete. Additions will be made and reported at a future time. It may be that improved accuracy can be attained by reworking the accumulated data in a different fashion and employing punched card machine methods for this reevaluation. Such a possibility is under consideration.

## 2. Congruence of Observation and Calculation

The agreement between observation and calculation by the present system is generally within 1% or better, depending somewhat on the accuracy of the original data. Other factors affecting the agreement, and not taken into complete consideration in deriving the equations, are effect of positional isomerism, effect of neighboring groups and branching, and effect of not taking

CONFIDENTIAL

CONFIDENTIAL  
SECURITY INFORMATION

into account minor corrections necessary to convert observed data to a uniform basis of temperature, in vacuo measurements, Weidmann corrections, and the like. It has not seemed worthwhile to make all adjustments possible because the unknown accuracy of the original data, with a few exceptions, did not warrant the effort. The data used for this study and the resulting equations are given in Tables 61-130.

There is almost perfect agreement between calculated heats of combustion and observed values for normal paraffins and certain other hydrocarbons where the data are known with considerable accuracy (115). These data are taken as the basis for the whole system which follows. For other types of functional groups the agreement is less good but satisfactory enough for the purpose intended. A graphic illustration of some of the results is given in Figure 79. The accuracy and precision of results apparently depend only on the reliability and accuracy of combustion data actually measured. The greatest deviation from the straight line relation is generally exhibited by the lowest member, and this is particularly true for the polar type of compound without carbon-carbon bonds, e.g., methanol, formic acid, formaldehyde, methylamine, and tetranitroethane.

It may be argued that the calculation method devised and reported here does not give sufficient accuracy to the heat of combustion. It seems to be adequate, however, for the purpose intended, i.e., approximation of properties of new or unknown compounds, particularly for prediction of power. In the second section of this report it is shown how well such calculated values serve to relate power as computed by the nRT product directly to power as measured in the ballistic mortar. Indeed, in a number of instances the calculated heat of combustion leads to better agreement than does observed heat of combustion. At any rate the roughness of prediction is not so great but what satisfaction can be expected when used in the field of propellants and high explosives.

### 3. Relationship among Functional Groups

This system of estimating heats of combustion was devised specifically for use with high explosive materials, many of which contain groups not susceptible to estimation by any other simple system. The system as presented at this time, nevertheless, can be used for any compound containing the functional groups given in Table 60. Conceivably there is no limit to the number or types of groups or bonds which can eventually be included. The present method is not claimed to be indisputable, for there exist anomalies and difficulties still to be corrected. These may be accounted for if interactions could be completely estimated.

One use to which the equation coefficients can be put is establishing the relative contribution of any functional group to heat of combustion. The same contribution will be made to heat of explosion if the point of comparison is chosen to be zero oxygen balance to carbon dioxide and water, since at this point and at positive balance these two heat quantities are identical. Moreover, at the zero point the power of pure explosive compounds, a parameter which is of vital interest to the present investigation, is at the maximum.

CONFIDENTIAL




FIGURE 79  
REDUCED HEAT OF COMBUSTION (Y)  
VS  
OXYGEN NEEDED (x)

CONFIDENTIAL  
SECURITY INFORMATION

Thus there can be constructed what might be called a "plosometric" series to interrelate the relative desirability of a functional group in an explosive molecule. Paraffin is taken as the zero potential. The complete series is given in Table 58. Where possible or practical the values have been listed for the liquid state, and only the most reliable or highest value for types with practical isolation is given.

CONFIDENTIAL

TABLE 58

PLOSONOTIVE SERIES

Contribution of Functional Groups to Heat of Combustion










<u>Functional Type</u>		<u>Contribution kcal/mole</u>
(R=aliphatic residue or hydrogen)		
Nitrate	$\text{RONO}_2$	129
Substituted nitramide	$\text{RC(=O)NNO}_2$	120
Secondary nitramine	$\text{R}_2\text{NNO}_2$	113
Primary nitramine	$\text{RNNO}_2$	112
Nitramide	$\text{RC(=O)NNO}_2$	103*
Nitramine	$-\text{NNO}_2$	95*
Nitro	$\text{RNO}_2$	93
Primary nitrosamine	$\text{RNHNO}$	73
Azo compound	$\text{RN=NR}$	69
Azoxy compound	$\text{RN=NR}$ $\quad \quad \quad \downarrow$ $\quad \quad \quad \text{O}$	69
Hydrazine, disubstituted	$\text{RNHNHR}$	67
Nitrosamine	$-\text{NNO}$	56
Hydrazine, monosubstituted	$\text{RNHNH}_2$	50
Tetrazole		49
Hydrazide, monosubstituted	$\text{RC(=O)NHNHR}$	47
Oxime	$\text{R=NOH}$	45
Oxirane		41
Acetylene	$\text{RC}\equiv\text{CR}$	37
Furan		36
Azo bond-group	$-\text{N=N}-$	35*

TABLE 5B (cont'd)

PLC MOTIVE SERIES






<u>Functional Type</u>		<u>Contribution kcal/mole</u>
Azoxy bond-group		35*
Guandine, disubstituted	RNHC(=NH)NER	35
Hydrazine bond-group		33*
Hydrazide bond-group	RC(=O)N-N-	30
Carbylamine (isonitrile)	RNC	27
Thiophene		22
Nitroso	RNO	21
Guandine, monosubstituted	RNHC(=NH)NH <sub>2</sub>	18
Amine	RiNH <sub>2</sub>	18
Cycloheptane		17
Cyclopropane		16
Ester	RC(=O)OR	16
Ether	ROR	16
Fluorine	RF	14
Olefin	R <sub>2</sub> C=CR <sub>11</sub>	14
Aldehyde	RCHO	12
Amide, substituted	RC(=O)NER	11
Cyclobutane		10
Alcohol	ROH	9
Nitrile	RCN	9

CONFIDENTIAL



TABLE 5B (cont'd)

PLASMOGENIC SERIES

<u>Functional Type</u>		<u>Contribution kcal/mole</u>
Ketone	$RC(=O)R$	6
Acid anhydride	$RC(=O)OC(=O)R$	2
Guanidine	$-NC(=NH)N-$	1*
PARAFFIN	$R-R$	0
Cyclopentane		-2
Acid	$RCOOH$	-5
Amido	$RC(=O)N-$	-6*
Naphthalene		-6
Cyclohexane		-7
Benzene		-10
Salt		-16
Biphenyl		-31

\*For compounds with substituents on nitrogen, add appropriate group value.  
For example, primary nitramine,  $RN(NO_2)_2$ , contribution becomes 112.

CONFIDENTIAL  
SECURITY INFORMATION

The functional group which makes the best phosphorus is the nitro group ( $-\text{NO}_2$ ) and this attached to oxygen, nitrogen, or carbon appears highest on the list (93-129 kcal/mole). Nitrosamine ( $-\text{NNO}$ ) is also high (56 kcal/mole), consistent with its secondary phosphoric nature. Nitroso ( $-\text{NO}$ , 21 kcal/mole) appears considerably lower in the scale than would be expected. This may be due to the lack of sufficient data to give a completely reliable constant, and it is reflected in the unusually high slope for this functional type (see Tables 60 and 136). Aromatic rings, unsubstituted amides, and acids appear below paraffins (-11 to -5 kcal/mole) and are not desirable for contributing to heat of explosion or combustion. Energy is added to a molecule by replacing a single bond with a double bond (olefin, 14 kcal/mole) or with a triple bond (acetylene, 37 kcal/mole). Compounds with hydrazine, azo, and azoxy groups (67-69 kcal/mole) are rich in energy and, in addition, are high in nitrogen acting as a source of extra gas. Thus for power, all other factors being constant. The presence of such common groups as ether, alcohol, amine, fluorine, substituted guanidine, and ester (14-18 kcal/mole) is helpful rather than detrimental. Tetrazole (49 kcal/mole) is not as good as might be expected, probably because of loss of energy due to resonance, although it can still be classed as a secondary phosphorus. The same loss of energy is shown by benzene which, without resonance, ought to have been the equivalent of three olefinic bonds (42 kcal vs -10 kcal). Furan (36 kcal/mole) is not quite the sum of an ether and two double bonds (44 kcal/mole) perhaps for the same reason. Indeed, from such considerations a resonance energy value can be calculated. For the benzene ring it would amount to 52 kcal/mole and to 76 kcal/mole for naphthalene, in reasonable agreement with Wheland's assigned values of 41-51 kcal/mole and 77-86 kcal/mole respectively (151).

A table similar to Table 58 drawn up on a weight basis would be even more helpful in group evaluation from an explosive viewpoint, because of the dependence on weight rather than molar values for measuring explosive properties. Such a table can be constructed from the data on a molar basis and will be presented at a future date.

The contribution made by functional groups, as produced by the present report, converted to values which atom to atom bonds might contribute would be akin to the Pauling system of additive bond energies for heat of formation, and to the Arthur D. Little, Inc., method for heat of combustion (70), predecessor of this one. Some efforts in this direction have shown that this can be done. The conversion is easily made from the intercept and slope values shown in Table 60, but the results are not yet in a state for publication.

#### 4. Multiple Groups on a Single Carbon Atom

Attention is drawn to another point of interest. More than one of certain groups or atoms, such as nitro or halogen, can replace the hydrogen atoms on a single carbon atom. It has been found that compounds containing such a multiplicity of function make an energy contribution to heat of combustion greater than would be expected from simple addition using the values for one function per carbon as described by the system herein. Accordingly,

CONFIDENTIAL

CONFIDENTIAL  
SECURITY INFORMATION

special equations have been determined for the gem trinitro  $[R_2C(NO_2)_3]$  and gem trifluoro  $(RCF_3)$  groupings. The increase for the gem tri substitution over the sum of three single groups is small (0.7% and 1% for trinitro and trifluoro, respectively). It might be attributed to the lack of hydrogen atoms for bonding which normally consumes some energy, or to the energy of strain from increased bond length caused by repulsion of multiple highly electronegative groups on the same carbon atom.

The difference between the value for gem dinitro  $[R_2C(NO_2)_2]$  and the sum of two mononitro  $(R_2CNO_2)$  groups was so small that it could not be determined whether it was a real difference or due to errors in experimental measurement. A separate value for gem dinitro is not included, therefore, in Table 60. From the preliminary evaluation of the enhancement effect of multiple substitution during the early stage of study on the heat of combustion problem a much larger effect for both gem di- and tri- substitution was deduced. These original conclusions have been discarded now in favor of the present, more satisfactory equations.

#### D. Conclusions

From the evidence, given in this report it is concluded that the method of calculating heat of combustion from a straight line relationship with molar oxygen balance is valid and exceedingly useful. Of the many hundreds of compounds considered the only ones of interest which are in disagreement with calculation greater than 10% are the following:

	% difference from calcn.
Tetranitromethane	+44
$\alpha, \alpha, \alpha$ -trifluoro-3,5-dinitrotoluene	-31
2-nitroethanol nitrate	-20
<del>Nitroethanol nitrate</del>	-20
N-picryldiethanolamine dinitrate	+17
trinitroacetonitrile	+16
allylpentaerythritol trinitrate polymer	-12
pentanitroaniline	+11

Tetranitromethane has recently been determined with great accuracy by the National Bureau of Standards (83). It must be assumed that this compound is unique and does not fit the general scheme of the calculation method. Trifluoro-3,5-dinitrotoluene is sufficiently divergent to cast suspicion on the identity of the compound measured and should be reevaluated.

2,4,6-Trinitroresorcinol (styphnic acid) is only 3% less than calculated, and 5-nitraminotetrazole has never been measured, but current interest makes determination of both these desirable.

It thus becomes possible with this system to evaluate and screen functional groups for suitability in explosive compounds by using a limited number of non-explosive, even poorly balanced, compounds without the necessity and danger of preparing well-balanced explosives for actual testing. If any functional type is found desirable, then greater expenditure of effort can be made to prepare balanced compounds by introducing primary phosphoric groups.

CONFIDENTIAL

CONFIDENTIAL  
SECURITY INFORMATION


The present list of functional groups should be extended, partly for particular use with explosive compounds and partly for general use in other fields of science. The coefficients of many of them can be approximated from existing data. A more accurate estimate of nitroso (-NO) should be obtained, and the following should be considered:

1. Of interest for explosives

a) azide




b) 1,2,3-triazole and 1,2,4-triazole




c) furoxan and furazan



d) triazene, pentazene, and other linear nitrogen chains



e) 1,2,4,5-tetrazine and 1,2,3,4-tetrazine



f) azine




g) hydrazone



h) imide



i) s-triazine



j) imine or anil



k) hydroxylamine



l) nitrite



m) perchlorate ester



2. Of general interest

n) thiol



o) sulfide



p) disulfide



q) sulfonic acid



r) chloro



s) bromo



t) iodo



u) isocyanate



v) peroxide



CONFIDENTIAL

E. Rules and Procedure for Calculating Heat of Combustion with Examples

The present system described in this report is based on the assumption, adequately upheld by experimental evidence, that the molar heat of combustion of any organic homologous series bears a straight line relation, within an acceptable limit of error, to the number of atoms of oxygen lacking in the molecule which are required to burn the compounds to CO<sub>2</sub>, H<sub>2</sub>O, N<sub>2</sub>, HI, and SO<sub>2</sub>. A quick estimation of this number, called the molar oxygen balance, for the compound, C<sub>m</sub>H<sub>n</sub>N<sub>p</sub>O<sub>q</sub>X<sub>r</sub>S<sub>t</sub> can be made by

the equation: Oxygen balance =  $2(n+t) + \frac{(n-r)}{2} - q$ .

Linear equations for a large number of functional types have been obtained by the method of least squares. The slope and intercept values for these equations related to the paraffin equation as a basis are given in Table 60, and the heat of combustion data used for the estimation are given in Tables 61 through 138. At present calculations are limited to compounds containing the groupings given in Table 60. The data were taken from many sources, and the only correction applied to the original data, aside from correcting obvious errors, was to convert the results reported to a basis of constant pressure, where necessary.

Fundamental Rule

The heat of combustion of a compound is computed from the oxygen balance of the compound on a molar basis and the equation derived from the summation of the slope and intercept factors for each structural feature, including paraffin as one of the features in every case. Thus  $Q_c = \sum a_i + x \sum b_i$

in which  $a_i$  = intercept coefficient for each functional type.

$b_i$  = slope coefficient for each functional type.

$x$  = atoms of external oxygen required for combustion.

The system, in effect, calculates and applies the equation for the homologous series which has the same functional groups as the compound in question.

In Table 59 a few examples of the use of the equation illustrate the additional rules which have been followed in the derivation of the coefficients given in Table 60. Examples 5 and 6 illustrate larger and more complicated molecules which can be handled easily by the present system.

Rule 1.

The values for normal paraffins ( $a = 5.7$  and  $b = 52.08$ ) are always part of the equation, and it is considered that never more than one of this feature is ever present.

CONFIDENTIAL  
SECURITY INFORMATION

Rule 2.

If a chain is branched, or there is more than one alkyl substituent attached to a cyclic structure the values for branched paraffin are added to the equation, but never more than one of these features is ever considered to be present. Thus benzene and toluene are not branched, but xylene, i-propylbenzene, and hexamethylcyclohexane each has one branched paraffin feature. (See example 1, Table 59).

Rule 3.

If more than one functional group of the same type is present, the intercept term (a') is multiplied by the number of such groups, but the slope term is the same as for a single such group. The only exceptions to this rule are the normal paraffin and branched paraffin features. This rule is illustrated in example 1, Table 59, with six nitrate groups.

Rule 4.

When there is a choice between two possible groups, the values for the group which appears later in the table of coefficients (Table 50) are taken. In example 2, Table 59, although a formamide can be considered as both an aldehyde and an amide, only the amide values are taken in order to avoid duplicating groups. Incidentally, the amide value is about the same as the sum of an aldehyde and an amine, which is the alternate choice that could be made in the example shown. Exceptions to this rule of duplication are the urea type of group (-N-C(=O)-N-) and the biuret type of group

$(-N-\underset{\text{O}}{\underset{||}{C}}-N-\underset{\text{O}}{\underset{||}{C}}-N-)$  both of which are considered to be diamides. In addition the

biuret group has an amine group as one of the components of the equation. It is hoped that such questions of duplication, which arise only with nitrogen containing compounds, can be resolved or avoided by use of bond contributions rather than group contributions. This may be done at a later date.

Rule 5.

Where a choice of coefficients is possible, depending on physical state, the ones taken are those corresponding to the physical state of the compound under consideration. Some exceptions to this rule are made in Table 60 when one set of coefficients is less reliable than the other, such as acid anhydride (solid is preferred to liquid), ester, nitrile, and secondary amine for which liquid is preferred to solid. Benzene (solid) and primary amine (solid) are reserved for hydrocarbons and aromatic amines, respectively. Derivation of coefficients for the liquid and solid phases have been based on normal paraffin (liquid) in all cases and for the gas phase have been based on normal paraffin (gas) in all cases.

CONFIDENTIAL

CONFIDENTIAL  
SECURITY INFORMATION

Rule 6.

With salts of organic bases and inorganic acids, and with hydrates, an additional step is required. The heat of combustion of the component base is calculated independently, by the general scheme, then added to the heat of combustion of the inorganic acid along with the value for each salt link, to give the heat of combustion of the salt. This is illustrated in example 3, Table 59. For inorganic components, such as nitric acid, the heat of combustion is computed from the heat of formation obtained from other sources. A similar procedure is adapted for salts of organic acids and inorganic bases. For organic salts, with carbon-carbon bonds, the ordinary rules for calculation can be followed. (See example 4, Table 59).

Rule 7.

For certain nitrogen containing groups there must be added to the contribution given in Table 60 additional values for amine or hydrazine when a carbon-nitrogen or nitrogen-nitrogen bond has been formed by substitution. Such is the case for amide, hydrazine, hydrazide, azo, azoxy, guanidine, tetrazole, nitramine, nitramide, nitrosamine, and nitrosamide. Thus a primary nitramine is considered to have both a nitramine and a primary amine contribution, and an N-substituted nitramide also has a primary amine contribution. To calculate an organic azo compound of the type Ar-N=N-Ar' two primary amine contributions besides the azo, aromatic and paraffin features must be included. As in examples 2 and 3, Table 59, an N-substituted amide or guanidine should be considered to possess appropriate amine features in addition to the amide and guanidine contributions.

Rule 8.

For compounds which have functional groups not covered specifically in Table 60 approximations sometimes can be made by using values for a closely related group. This rule may be applied particularly to the physical state aspect. Little error will be introduced in most organic compounds if the value for the liquid state is used for solids, and vice versa. (See rule 5). Compounds containing other structural features cannot be estimated at this time.

The data for homologous series of compounds in the same physical state were used to estimate the best straight line. In several instances enough data were available to give equations for more than one physical state for compounds containing a given functional group. The differences between the equations, then, would be the heat of fusion or heat of vaporization for the homofunctional types. The differences due to heat of fusion for organic compounds are small, particularly for the less polar types, but it is felt that account should be taken of this factor where it is possible to do so, reducing at least one source of error. However, the error of using liquid data for solid compounds is small.


CONFIDENTIAL

TABLE 59

Illustration of Heat of Combustion Calculation

$$Q_C^D = \sum a' + x \sum b'$$

1. Hexamethylolbenzene hexanitrate  
(183)




$Q_C^D$  (obs) = 1396.1 kcal/mole Ref. 71




x = 12

	a'	b'	
	(intercept)	(slope)	
1 normal paraffin (liq)	+ 5.7	+52.08	(Rule 1)
1 branched paraffin (liq)	- 3.7	+ 0.09	(Rule 2)
1 benzene (liq)	-10.1	+ 0.07	(Rule 5)
6 $ONO_2$ (s) $\overline{6 \times 128.4}$	+770.4	+ 0.5 <sup>3</sup>	(Rule 3)
$Q_C^D = 762.3 + (12 \times 0) (52.77) = 1395.7$ kcal/mole			

2. Dimethylformamide




$Q_C^D$  (obs) not found



x = 8.5

	a'	b'	
	(intercept)	(slope)	
1 normal paraffin (liq)	+ 5.7	+52.08	(Rule 1)
1 amide (liq)	- 5.6	+ 0.47	(Rule 4)
1 secondary amine (liq)	+18.3	- 0.12	(Rule 7)
$Q_C^D$ (calc) = 18.4 + (52.43) (8.5) = 464.1 kcal/mole			

3. Diaminoguanidine nitrate



$Q_C^D$  (obs) = 329.8 kcal/mole (59)



x = 5.5 (for diaminoguanidine)

(Rule 6)



TABLE 59 (cont'd)

	a' (intercept)	b' (slope)	
1 normal paraffin (liq)	5.7	52.08	(Rule 1)
1 guanidine (s)	0.7	+ 0.46	(Rule 7)
2 hydraxine bonds (s)	65.0	- 0.10	(Rule 7)
$Q_C^D$ (calc) = $71.4 + (51.52)(5.5) = 354.8$ kcal/mole for diaminoguanidine			
1 salt link	-14.0		(Rule 6)
1 HNO <sub>3</sub>	- 7.2		
$Q_C^D$ (calc) = $354.8 - 14.0 - 7.2 = 333.6$ kcal/mole for diaminoguanidine nitrate			


4. Carbohydrazide hydrogen oxalate



x = 5.0 (for entire molecule)



(Rule 6)

$Q_C^D$  (obs) = 314.6 kcal/mole Ref. 96



	a' (intercept)	b' (slope)	
1 normal paraffin (liq)	+ 5.7	+52.08	(Rule 1)
2 hydrazide (s) (2 x 28.7) =	+57.4	- 0.09	(Rule 3,7)
2 acid (s) (2 x -3.8) =	- 7.6	- 0.01	(Rule 3,5)
1 salt	-14.0	-	(Rule 6)
$Q_C^D = 41.5 + (51.98)(5) = 304.0$ kcal/mole			

5. Vitamin A (Carotene)




x = 54

	a' (intercept)	b' (slope)	
1 normal paraffin (liq)	5.7	52.08	(Rule 1)
1 branched paraffin (liq)	- 3.7	+ 0.09	(Rule 2)
1 cyclohexane (liq)	- 7.4	0.00	(General)
5 normal olefin (liq) 5 x (14.2) =	+71.0	- 0.01	(Rule 3)
1 primary alcohol (liq)	+ 9.2	- 0.05	(Rule 5)
$Q_C^D = 71.8 + (52.11)(54) = 2868.7$ kcal/mole			

CONFIDENTIAL

TABLE 59 (cont'd)

6. Phenolphthalein



$x = 43$

	<u>a'</u> (intercept)	<u>b'</u> (slope)
1 normal paraffin (liq)	5.7	52.08 (Rule 1)
1 branched paraffin (liq)	- 3.7	+ 0.09 (Rule 2)
3 benzenes (liq)	-10.1	+ 0.07 (Rules 5 and 3)
2 aromatic hydroxyl (s)	+ 7.0	- 0.29 (Rule 3)
1 ester (liq)	+16.1	- 0.42 (Rule 5)

$$Q_C^D = 15.0 + (51.53) (43) = 2230.8 \text{ kcal/mole}$$






CONFIDENTIAL  
SECURITY INFORMATION

TABLE 60

COEFFICIENTS OF EQUATIONS FOR HEAT OF COMBUSTION

$$\frac{D}{C} = \sum a_i + x \sum b_i$$






Normal paraffin (l)      a = 5.7, b = 52.06  
Normal paraffin (g)      a = 5.5, b = 52.48

Functional Type	a <sup>i</sup> (intercept)	b <sup>i</sup> (slope)	
(R=aliphatic chain or hydrogen)			
Paraffin branched (liq)	- 3.7	+0.09	
Cyclopropane (liq)	+16.2	-0.13	
Cyclobutane (liq)	+10.3	+0.11	
Cyclopentane (liq)	- 1.7	0.00	
Cyclohexane (liq)	- 7.4	0.00	
Cycloheptane (liq)	+17.1	-0.99	
Olefin			
normal (liq)	+14.2	-0.01	$R_2C=CR_2$
normal (gas)	+14.2 <sup>a</sup>	0.00 <sup>a</sup>	
Acetylene			
normal (liq)	+37.3	0.00	$RC\equiv CR$

CONFIDENTIAL

**CONFIDENTIAL**  
**SECURITY INFORMATION**







TABLE 60 (cont'd)

Functional Type	a' (intercept)	b' (slope)	
Benzene (liq) (gas) (solid)	-10.1 - 7.0 <sup>a</sup> -16.5 <sup>b</sup>	+0.07 0.00 <sup>a</sup> +0.45 <sup>b</sup>	
Biphenyl (liq)	-34.3	+0.37	
Naphthalene (solid)	- 6.1	-0.59	
Alcohol			
primary (liq)	+ 9.2	-0.05	RCH <sub>2</sub> OH
primary (gas)	+18.3 <sup>a</sup>	-0.24 <sup>a</sup>	
secondary (liq)	+ 4.5	-0.44	(R <sub>2</sub> C) <sub>n</sub> COH
tertiary (liq)	+ 2.6	-0.25	(R <sub>3</sub> C) <sub>n</sub> COH
mixed (liq)	+10.3	-0.63	
mixed (solid)	+ 3.8	+1.27	
Aromatic hydroxyl (solid)	+ 7.0	-0.29	
Ether (liq) (gas)	+15.5 +28.1	+0.02 -0.05 <sup>a</sup>	R <sub>2</sub> COCH <sub>2</sub>
Oxirane (ethylene oxide) (liq)	+41.2	-1.05	
Furan (liq)	+35.5	-1.17	
Aldehyde (liq) (gas)	+11.5 +20.9	-0.09 -0.68	RC(=O)H
Ketone (liq)	+ 5.5	-0.19	RC(=O)R

**CONFIDENTIAL**

CONFIDENTIAL  
SECURITY INFORMATION

TABLE 60 (cont'd)

Functional Type	a' (intercept)	b' (slope)	
Acid (liq)	- 4.7	+0.07	RCOOH
(solid) <sup>c</sup>	- 3.8	-0.01	
Acid anhydride (liq) <sup>d</sup>	+ 9.2	-0.03	RC=O
(solid)	+ 2.4	-0.01	
Ester (liq)	+16.1	-0.4 <sup>e</sup>	RC(=O)OR
(solid) <sup>e</sup>	+16.1	-0.62	
Nitrile (liq)	+ 9.3	-0.01	RC≡N
(solid) <sup>e</sup>	+ 8.9	+0.29	
Carbylamine (Isonitrile)	+26.5	+0.57	RNC
<b>Amine</b>			
primary (liq)	+17.7	-0.31	R <sub>3</sub> CN <sub>2</sub>
primary (gas)	+18.0 <sup>a</sup>	-0.49 <sup>a</sup>	
primary (solid)	+ 4.3 <sup>f</sup>	-0.08 <sup>f</sup>	
secondary (liq)	+18.3	-0.12	(R <sub>3</sub> C) <sub>2</sub> NH
secondary (solid)	-44.9 <sup>g</sup>	+1.82 <sup>g</sup>	
tertiary (liq)	+20.5	+0.08	(R <sub>3</sub> C) <sub>3</sub> N
Amide (liq) <sup>h</sup>	- 6.5	+0.57	RC(=O)N-
(solid) <sup>h</sup>	- 6.0	+0.16	
Hydrazine (solid) <sup>h</sup>	+32.5	-0.10	
Hydrazide (solid) <sup>h</sup>	+30.2	-0.15	RC(=O)N-N-
Azo (solid) <sup>h</sup>	-35.4	+0.11	
Azoxy (solid) <sup>h</sup>	+32.5	+1.44	
Guanidine (solid) <sup>h</sup>	+ 0.7	-0.46	
Tetrazole (solid) <sup>h</sup>	+48.7	+0.32	
Oxime (solid)	+45.3	-0.12	R <sub>2</sub> C=N-OH
<b>Nitro</b>			
aliphatic (liq)	+38.4	-0.38	R <sub>3</sub> CNO <sub>2</sub>
aliphatic (solid)	+92.8	-0.65	
aromatic (liq)	+97.9	-0.39	
aromatic (solid)	+92.2	-0.40	

CONFIDENTIAL

CONFIDENTIAL  
SECURITY INFORMATION

TABLE 60 (cont'd)

Functional Type	a' (intercept)	b' (slope)	
Gas trinitro (liq) (solid)	+292.5 +288.2	+0.29 -0.32	$\text{RC}(\text{NO}_2)_3$
Nitrate (liq) (solid)	+129.4 +128.4	+0.23 +0.53	$\text{R}_2\text{CONO}_2$
Nitramine (solid) <sup>h</sup>	+ 95.3	+0.81	$\begin{array}{c} \text{---NO}_2 \\   \\ \text{R---N} \end{array}$
Nitramide (solid) <sup>h</sup>	+103.6	+0.10	$\begin{array}{c} \text{R---C---NO}_2 \\    \\ \text{O} \end{array}$
Fluorine (liq) (solid)	+ 10.9 + 14.2	+0.61 -0.19	$\text{R}_2\text{CF}$
Gas trifluoro (liq)	+ 36.1	-0.44	$\text{RCF}_3$
Nitrosamine (solid) (liq)	+ 56.0	+1.05	$\begin{array}{c} \text{---NO} \\   \\ \text{R---N} \end{array}$
Thiophene (solid) (liq)	+ 21.5	+0.98	$\begin{array}{c} \text{R---C} \quad \text{---C---R} \\ // \quad \quad // \\ \text{R} \quad \quad \quad \text{C---R} \\ \backslash \quad \quad / \\ \text{S} \end{array}$
Nitroso (solid)	+ 20.9	+1.93	$\text{R}_2\text{CHO}$
Salt formation	- 16.1	-	
Hydrate formation	+ 4.5	-	

- a) Based on normal paraffin (gas) value.  
 b) For hydrocarbons only.  
 c) From dibasic acids.  
 d) Based on small amount of data and not as reliable as value for solid.  
 e) Based on limited amount of data and less reliable than value for liquid.  
 f) From aromatic amines only.  
 g) Of questionable accuracy. Based on limited amount of data.  
 h) For substituents on the nitrogen atom(s), not appropriate bond-group values.  
 Add primary amine values for one substituent on nitrogen, secondary amine values for two substituents on the same nitrogen.

CONFIDENTIAL

CONFIDENTIAL  
SECURITY INFORMATION

F. Tables of Data  
1. Scope

An attempt has been made to obtain from the literature, open and classified, all heats of combustion which have been published between 1929, when the bibliography by Kharasch (55) appeared, and 1951. Much of the material found appears in Tables 61 through 130, the omissions being compounds which do not fall into any homologous series reported there. With some exceptions, notably in the nitro and nitrate functions, heats of combustion before 1929 which have been used are the ones as given by Kharasch (55), but all of the heats reported by Kharasch have not been used or reproduced in the present treatise. However, all known heats of combustion of organic compounds containing the nitro or the nitroso group attached to carbon, oxygen or nitrogen have been included in this report even though there is duplication of compounds.

The primary source of the open literature data was Chemical Abstracts, for the years 1929-1951. In many cases the original publication cited by Chem. Abstr. was consulted for evaluation of the author's work, but this was not a universal rule. Classified literature covered included OSRD reports of Division 8, SPIA/MJ reports, Picatinny Arsenal Technical Reports, various current reports of Navy Installations and contractors, and original work done for Arthur D. Little, Inc. by its subcontractors.

2. Method of Obtaining Least Squares Equations

The equation,  $Q_p^D = a + bx$ , was obtained for the paraffin hydrocarbons from the formulas:

$$b = \frac{n \sum xy - \sum x \sum y}{n \sum x^2 - (\sum x)^2}$$

$$a = \frac{\sum y - b \sum x}{n}$$

in which  $x$  = molar oxygen balance

$y$  = heat of combustion at constant pressure

$n$  = number of items.

The equations for the remaining homologous series were obtained by the same formulas from what might be called "reduced data", i.e. the values of  $Q_p^D$  remaining after subtracting 5.7 and the contribution of functional groups in the molecule other than the one being processed. To obtain  $x'$  and  $y'$  for least squares computation the reduced value for  $Q_p^D$  and the molar oxygen balance of the compound were then each divided by the number of like functional groups which were being calculated. Thus the equations were obtained literally for functional groups of a single type regardless of the presence of other groups in the compounds. This approach allowed the use of a larger body of available data than possible if only the data for pure, homofunctional types had to be used.

CONFIDENTIAL

CONFIDENTIAL  
SECURITY INFORMATION

3. Arrangement of Heat of Combustion Data

Tables 61-138 contain the data of interest to this project. They have been arranged in general order of increasing complexity of functional type, although the equations were not all determined in the order in which they appear in the tables. The columns are self-explanatory. The ADL numbers are those assigned in Part I of this report (69) and as extended in Table 139. Many of the compounds have been renamed here to conform to accepted usage which avoids ambiguity. The numbers originally assigned in Part I (69), however, have not been changed.

Each of the Tables 61-138 shows the equation for the homologous series represented and the summation figures for its derivation. All combustion data above it were used for its calculation. The combustion data below the equation were not used but have been included for completeness, or because the data are pertinent to the field of explosives. An occasional entry made below the equation was found too late to include in the calculations.

The heat of hydration was determined as an average from pairs of data for hydrates and the anhydrous compound (Table 134). The heat of salt formation was obtained as the difference between the heat of formation (or combustion) of the salt and the sum of the heats of formation (or combustion) of the basic and acidic components of the salt. (Table 135).

CONFIDENTIAL



### G. Historical

Methods of calculating heats of formation and of combustion for organic compounds have been used for many years. They can be divided into four general types:

1. Methods of summation, using bond, resonance, and group energies. These generally have a sound basis in theory and have been well-developed for fairly wide application. The energy values are obtained from considerations of observed thermochemical data.
2. Methods of summation, using a base value to which are added corrections for substituent functional groups. These are generally empirical and based on observed data.
3. Methods based on linear equations obtained from relationships between heat value and some intrinsic feature of a series of homologous compounds, such as number of carbon atoms in the molecule or paracher. These are empirical methods using observed data but of limited applicability.
4. Methods involving effects observed upon substituting one group for another. These have been widely used but are extremely limited or completely unsuitable for prediction of thermochemical properties of new or unknown structures.

#### 1. Methods of Summation of Bond Energies

The method which has probably received the greatest popular attention is that based on the additivity of bond energies as outlined by Pauling (89) for heats of formation, and expanded and extended by Wheland (20) and Klages (59) to heats of combustion. The bond values given by Pauling (89) were determined from thermochemical considerations of small molecules as given by Bichowsky and Rossini (16)

Wheland (20) in explaining the theory of resonance and comparing values for different resonant structures, converted Pauling's bond energies of formation to values which could be used to estimate heat of combustion. Klages (59) later improved the accuracy and added to the values of Wheland, enlarging the scope and applicability of this system.

Springall and Roberts (130) obtained a set of bond, resonance, and group energies for heats of formation especially suitable for explosive compounds containing carbon, hydrogen, nitrogen, and oxygen. Pauling's bond energies were taken as a basis, and group energies were determined from known heats of formation for a great many explosive compounds.

In the present continuing investigation it was felt that heat of combustion was a more valuable and accessible function than heat of formation. Part II of this report by Arthur D. Little, Inc., (70) presented bond and group energies for calculating heats of combustion which were determined for the special field of explosives along the lines of Springall and Roberts (130).

CONFIDENTIAL  
SECURITY INFORMATION

Anderson, Beyer, and Watson (3) reported a system of group contributions for calculating heats of formation, entropies, and heat capacities of organic compounds. The development of their system was based on theoretical consideration and "heats of atomization".

All of these methods, except that of Arthur B. Little, Inc., as described in Part II (70), produce results for compounds in the gaseous state and appropriate corrections must be assumed to convert these to values for the standard state of the compounds of which the great majority are solid or liquid. All the preceding systems of bond and group energies become exceedingly difficult to apply to complex groups and heterocyclic structures.

A good discussion of the early methods proposed for calculating heats of formation in the gaseous state from bond energies is given by Rossini (114). Major contributions to these methods, besides those described above (3, 69, 110), have been made by Anderson and Gilbert (5), Coates and Sutton (25), and Cole and Gilbert (26). Resonance energies of ring structures were introduced into the system of bond energies by Wheland (20), and additional values have been obtained by Willis (2, 151) and by workers at the Naval Ordnance Test Station (77).

Efforts to improve the accuracy of calculated heats of formation have been made by Laidler (62, 63) who used a system similar to Pauling's and Wheland's but based it on bond strengths and "heats of atomization." Laidler's method suffers from being too complicated for use with polyfunctional compounds and the accuracy claimed for functional types other than hydrocarbons is misleading, since values given are based on a limited amount of unconfirmed data in the early literature. Others have attempted to apply group contribution methods to heat of formation taking into account neighboring effects (37, 97, 127). These are all rather theoretical treatments, Platt (97) limiting his calculations to paraffin hydrocarbons, and Souder (127) to hydrocarbon vapors. Franklin (37) attempted to include many of the more common functional groups but used very limited data to obtain his figures. His suggestions are also very difficult to apply to complicated configurations.

Stern and Klebs (131, 132), from a study of heats of combustion of pyrroles, determined thermochemical characteristics for a large number of common functional groups attached to the heterocyclic nucleus.

Of more theoretical interest are the investigations of bond strengths by Cotrell and Sutton (30), Walsh (150), and Skinner and Springall (124). Cotrell (28) also calculated binding energies of hydrocarbons and resonance energies of simpler aromatic hydrocarbons. Roberts and Skinner (112) derived heats of formation of a number of alkyl radicals and resonance energies from thermal and other data.

CONFIDENTIAL

CONFIDENTIAL  
SECURITY INFORMATION

Methods for calculating heats of formation from structural groups and molecular formulas, employing standard equations, have been suggested (19, 22, 105, 106), and heats of combustion have been calculated by similar methods (20, 21). All of these results were limited to gaseous hydrocarbons. Another author (42) adopted 8, the lowest even calculated value for bond energies published, as the unit of the chemical bond and calculated heats of formation from energies of all other bonds grouped as multiples of this unit.

2. Methods of Summation using a Base Value plus Group Corrections

One of the best and most thoroughly described methods for calculating heats of combustion from the structure of an organic compound is that described by Kharasch (55). Kharasch and Jner (46) have given the background and basis for his method of calculation which depends on adding the number of electrons in a molecule multiplied by the combustion value of each electron, to the corrections for structural and functional features. Aerojet Corp. (1) has extended the system to include primary phosphoric types (69) of compounds. The method is more empirical and less theoretical than those described previously in this report, and in general gives good agreement with observed values. The physical state is liquid which involves fewer corrections and assumptions to compute values for most compounds. It is not adapted to heterocyclic compounds, however, and does not always work well with the more complicated functional groups.

3. Methods Based on Linear Equations

The Kharasch theory of molar summation has been converted to a weight basis (52). Results are expressed in a unique series of graphs for a single functional type, with heat of combustion plotted against percent carbon in the compounds. Heats of combustion and formation are tabulated in the form of general equations for the different classes of compounds. The method has not been used, however, with poly- and heterofunctional molecules.

One author (68) has attempted to establish a relationship between atomic number, effective nuclear charge, and heat of combustion for compounds containing C, H, and O, and another (43) has proposed a relationship between heats of combustion and composition of organic compounds by use of a single simple formula. It is not easy to see how these systems have general applicability and validity.

The values of the paracher and the heats of combustion or formation of organic compounds have been correlated by a straight line relationship (95). The values of the slope and intercept are different for various homologous series and classes of compounds, but it is claimed they can be determined easily if the values of paracher and combustion are known for one of the representatives of a homologous series. Slope and intercept values for nine homologous series are given. Satisfactory agreement between experimental and calculated values is illustrated with saturated alcohols.

CONFIDENTIAL

Numerous articles have been published giving empirical methods for calculating heats of combustion from observed data. Heat of combustion of successive members of homologous series have been obtained by Verkade and co-workers from which was calculated the heat of combustion for the methylene (-CH<sub>2</sub>-) group. Thus the straight chain dicarboxylic acids (113) and their dimethyl esters (116), the homologous normal primary aliphatic alcohols (114), and monoalkyl-substituted malonic acids (115) have been studied. Ramee and Perks (11) did much the same thing for paraffin hydrocarbons. Schjanberg (123) calculated the energy of cleavage of the C-C1 link in a series of aliphatic monocarboxylic acids and their esters, relating it to the change in position of the halogen.

Rossini and co-workers applied similar techniques. Relative to the number of carbon atoms in a series of normal aliphatic primary alcohols to their heats of combustion, he found that a straight line would express the heat of combustion for alcohols longer than pentanol (113). The equation given is

$$Q_c = 21.60 + 157.00 n \quad (\text{gaseous state})$$
$$= 12.00 + 157.00 n \quad (\text{liquid state})$$

in which n, the number of carbon atoms, must be greater than 5. Similar equations applicable to the estimation of heats of combustion of paraffin hydrocarbons were calculated (103), giving

$$Q_c = 57.909 + 157.443 n \quad (\text{gaseous state})$$
$$= 57.430 + 156.263 n \quad (\text{liquid state})$$

in which n, again, is the number of carbon atoms, and is greater than 5. An equation obtained for alkylbenzenes (100) is:

$$Q_c = -158.990 + 157.443 n \quad (\text{gaseous state})$$

in which n is the number of carbon atoms, and is greater than 9.

A straight line for approximating the heat of combustion of gasolines from the percent of hydrogen has been proposed recently (53a).

Lentle (65) gave straight line equations for calculating heats of combustion of eleven homologous series of compounds with commonly encountered functional groups. Agreement with observation was good, but no attempt was made to expand the system to poly- or hetero-functional compounds. For calculating heats of formation he gave a treatment similar to Sugden's procedure.

Holcomb and co-workers found a straight line relationship between oxygen balance and heat of combustion (on a weight basis) for nitroalkanes (14) and later for nitroalcohols (31a). Earlier a straight line relationship was reported between the weight heat of combustion and oxygen balance of explosives, in which different lines were found for different types (23). The value of substituting an aromatic alkyl ether for an aromatic hydroxyl group in explosives was also determined.

CONFIDENTIAL  
SECURITY INFORMATION

4. Methods Based on Group Substitutions

One of the most common methods of predicting heats of formation and combustion is based on the effects observed upon substituting one group for another. Although frequent publications have been made comparing classes of compounds differing only in a single functional group, prediction of heat values is limited to such comparisons, and has not generally been extended to prediction from additive structural features, or to unknown compounds. Thus Swietoslowski (Svyentoslavskii) (135) obtained thermochemical characteristics of many groupings including  $-N=O$ ,  $-N-N-$ ,  $-N=N-$ , and  $-NO_2$ . By a similar technique the effect of a nitro group on heat of combustion has been found from a study of aromatic nitro derivatives (137) and there has been calculated the heats of combustion of certain groups such as carbonyl, hydroxyl, olefin, etc. (12). A series of furans and benzene has been compared (64); a study of furazan and furoxan compounds (60) showed a relation similar to the azo and azoxy derivatives according to McMurt (67). The contribution of the oxime group has been calculated (61) and energy changes in the formation of chelate rings have been measured (17). Badoche has correlated the effects on heat of combustion of introducing nitro groups into the benzene ring (8), of introducing hydroxyl and methyl groups into the benzene ring (9), and of introducing hydroxyl groups into the benzene ring (10) already containing nitro groups. Matignon investigated the difference between a nitroso substituted on carbon and on nitrogen (75) and applied the information gained from the effect of adding a methyl group to a nitrogen atom in cyclic ureas to deduce and establish formulas of constitution (74). The effect of introducing successive nitro groups into aromatic rings and the relationship of the heat of formation to the number of nitro groups present has been studied (110).

CONFIDENTIAL

H. References

1. Aerojet Engineering Corp., Report No. 417 A, Contract N7onr-462 by Gold and Carlston (23 December 1949), pp. 31-36.  
CONFIDENTIAL.
2. A. Albert and J. B. Willis, *Nature*, 157, 341 (1946); *C.A.*, 40, 3311 (1946). "Resonance energies of acridine and phenazine".
3. J. W. Anderson, G. H. Beyer, and K. M. Watson, *Nat. Petroleum News*, 36, 476 (1944). "Thermodynamic properties of organic compounds - Estimation from group concentrations". Also see O. A. Hougen and I. M. Watson, "Chemical process principles", Part II, John Wiley, New York (1947) pp. 758-765.
4. C. Anderson, I. G. Cole, and E. C. Gilbert, *J. Am. Chem. Soc.*, 72, 1263-1264 (1950). "The heats of combustion of some isomeric nitrostilbenes and their isomerization energies".
5. C. H. Anderson and E. C. Gilbert, *J. Am. Chem. Soc.*, 64, 2369 (1942). "The apparent energy of the N-H bond as calculated from heats of combustion".
6. Army-Navy Solid Propellant Group, Bulletin of 4th Meeting. (19-20 April 1948) and Addendum. CONFIDENTIAL
7. M. Badoche, *Bull. soc. chim.* [5], 4, 549-558 (1937); *C.A.*, 31, 4196 (1937). "Determination of heats of combustion of several nitro derivatives of naphthalene".
8. M. Badoche, *Bull. soc. chim.*, 6, 570-579 (1939); *C.A.*, 33, 5736 (1939). "Determination of heats of combustion of nitro derivatives of the benzene series".
9. M. Badoche, *Bull. soc. chim.*, 8, 212-220 (1941). *C.A.*, 36, 1840 (1942). "Heats of combustion of phenol, m-cresol, and of their ethers".
10. M. Badoche, *Bull. soc. chim.*, 9, 86-95 (1942); *Chem. Zentr.*, 1942, II, 2013; *C.A.*, 38, 2558 (1944). "The heat of combustion of nitrated phenol derivatives".
11. H. Fance and G. S. Parks, *J. Am. Chem. Soc.*, 55, 3223 (1933). "Thermal data on organic compounds. XII. The heats of combustion of nine hydrocarbons".
12. H. F. Barker, *J. Phys. Chem.*, 29, 1345-1363 (1925); *C.A.*, 20, 326 (1926). "Calorific value and constitution".

CONFIDENTIAL

CONFIDENTIAL  
SECURITY INFORMATION

13. M. Beckers, Bull. soc. chim. Belg., 40, 518-610 (1931); C.A., 26, 2626 (1932). "Methods and equipment used at the Bureau of Physico-chemical Standards. V. VI. Heats of combustion".
14. Berthelot and Matignon, Compt. rend., 113, 246 (1891). "Heats of combustion and formation of nitrobenzenes".
15. Berthelot and Matignon, Ann. chim. et phys. (6), 39, 565 (1895). "Nitromethane and its homologues".
16. F. K. Bichowsky and F. D. Rossini, "Thermochemistry of the chemical substances", Reinhold, N. Y. (1936)
17. G. B. Mondino, R. Manzoni-Angioli, and M. Rolla, Ricerca sci., 8, No. 5-6, 4 pps. (reprint) (1937); C.A., 22, 4846 (1938). "Experimental contributions to the problem of chelation. VI. Thermochemical investigations of acetylphenols and acetylnesoreindols".
18. L. Brull, Gazz. chim. ital., 65, 19-28 (1935); C.A., 29, 3587 (1935). "Heats of combustion of some derivatives of biphenyl".
19. M. Brutzkus, Compt. rend., 204, 490-3 (1937); C.A., 31, 2579 (1937). "Intrinsic value of linkages C-C and C-H in hydrocarbons".
20. M. Brutzkus, Compt. rend., 204, 574-5 (1937); C.A., 31, 2916 (1937). "Thermochemistry of hydrocarbons".
21. M. Brutzkus, Compt. rend., 204, 657-61 (1937); C.A., 31, 3373 (1937). "Yields of combustion of hydrocarbons and energy values of the linkages of their atoms".
22. M. Brutzkus, Compt. rend., 204, 1802-4 (1937); C.A., 31, 5666 (1937). "Heat of formation of hydrocarbons".
23. E. Burlet, M. Thomas, and M. Badocq, Mem. poudres, 25, 226-282 (1939); C.A., 34, 1849 (1940). "Calorimetry of explosives".
24. Chemical Rubber Publishing Co., Cleveland, Ohio. "Handbook of Chemistry and Physics".
25. G. E. Coates and L. E. Sutton, J. Chem. Soc., 1918, 1187. "The heats of formation of the -N-N- and -O-N- bonds".
26. I. G. Cole and E. C. Gilbert, J. Am. Chem. Soc., 72, 5423-5427 (1951). "The heats of combustion of some nitrogen compounds and the apparent energy of the N-N bond".

CONFIDENTIAL

CONFIDENTIAL  
SECURITY INFORMATION

27. J. Coops, D. Mulder, J. W. Dienske, and J. Smittenberg, Rec. trav. chim., 65, 128 (1946); C.A., 40, 3972 (1946). "The heats of combustion of a number of hydrocarbons (preliminary communication)". See also Rec. trav. chim., 66, 153-160 (1947); C.A., 41, 6460 (1947). "Heat of combustion. IV. Technique for the determination of heats of combustion of volatile liquids".
28. T. L. Cottrell, J. Chem. Soc., 1948, 1148-1149; C.A., 43, 2352 (1949). "Binding energies in hydrocarbons".
29. T. L. Cottrell and J. E. Gill, J. Chem. Soc., 1951, 1798-1800. "The preparation and heats of combustion of some azine nitrates".
30. T. L. Cottrell and L. E. Sutton, Quart. Rev., 2, 260-276 (1948); C.A., 43, 1230 (1949). "The interpretation of bond properties".
31. R. S. Crog and H. Hunt, J. Phys. Chem., 46, 1162-1163 (1942); C.A., 37, 1324 (1944). "Heats of combustion. II. Heats of combustion of ethyl methyl ketone and ethylamine oxide".
- 31a. R. M. Currie, C. O. Bennett, and D. E. Holcomb, Ind. Eng. Chem., 44, 329-331 (1952). "Heats of combustion of some nitro alcohols".
32. H. S. Davis and O. F. Wiedeman, Ind. Eng. Chem., 37, 483 (1945). "Physical properties of acrylonitrile".
33. M. Delapine, Bull. soc. chim., 15, 1199 (1896). "Hexamethylenetetramine and its nitroso derivatives: thermochemistry".
34. M. Delapine and M. Badoche, Compt. rend., 211, 777-81 (1942); C.A., 38, 5138 (1944). "The thermochemistry of  $\text{C}_6\text{H}_{12}\text{O}$ ,  $(\text{CH}_2)_6\text{N}_4$ , and their derivatives".
35. F. dePauw, Z. ges. Schiess- u. Sprengstoffw., 32, 10-12, 36-37, 60-63 (1937); C.A., 31, 3696 (1937). "Heat of combustion of smokeless powder".
36. Fasalar, "Dissertation", Brauda (1930).
37. J. L. Franklin, Ind. Eng. Chem., 41, 1070-1076 (1949). "Prediction of heat and free energies of organic compounds".
- 37a. J. L. Franklin and H. E. Lumpkin, J. Am. Chem. Soc., 74, 1023-1026 (1952). "Some C-S, H-S and S-S bond strengths by the electron impact method".
38. Cl. Fromageot and E. Danni, Bull. soc. chim., 49, 929-936 (1931); C.A., 25, 5076 (1931). "The heat of combustion of methylglyoxal".

CONFIDENTIAL



CONFIDENTIAL  
SECURITY INFORMATION

39. W. E. Garner and C. L. Abernethy, Proc. Roy. Soc. (London), 99A, 213-235 (1921); C.A., 15, 3748 (1921). "Heats of combustion and formation of nitro-compounds. Part I. Benzene, toluene, phenol and methylaniline series".
40. O. Glensner and V. Kausner, Z. Naturforsch., 3b, 159 (1948); C.A., 43, 3682 (1949). "Carbon monoxide compounds of the nonmetals. II. Properties of carbonyl cyanide".
41. W. Gliwitsky, Feuerschutz, 19, 80-82 (1939); C.A., 33, 815 (1939). "Naphthalene dust and vapor explosions".
42. A. Gosselin, Bull. soc. chim., 12, 61-67 (1945); C.A., 40, 4267 (1946). "The unit of the chemical bond (energy)".
43. A. Harlin, IVA, 17, No. 1, 28 (1946); C.A., 40, 5525 (1946). "Elementary analysis and heat of combustion".
44. D. E. Holcomb and C. L. Dorsay, Jr., Ind. Eng. Chem., 41, 2788-2792 (1949). "Thermodynamic properties of nitroparaffins".
45. W. H. Hubbard, J. W. Krowlton, and H. M. Huffman, J. Am. Chem. Soc., 70, 3259 (1948). "The heat of combustion of 2,2'-bis-(4-tert-butyl-phenyl)-propane".
46. W. Huckel, A. Gercke and A. Gross, Ber., 66B, 563 (1933); C.A., 27, 4811 (1935). "Cyclodecane".
47. H. M. Huffman, J. Am. Chem. Soc., 60, 1171-1177 (1938). "Thermal data. II. The heats of combustion of hippuric and succinic acids and a proposal for the use of hippuric acid as a secondary standard in combustion calorimetry".
48. H. M. Huffman, J. Am. Chem. Soc., 62, 1009 (1940). "Thermal data. XII. The heats of combustion of urea and guanidine carbonate and their standard free energies of formation".
49. H. M. Huffman, E. L. Ellis, and S. W. Fox, J. Am. Chem. Soc., 58, 1728 (1936). "Thermal data. VI. The heats of combustion and free energies of seven organic compounds containing nitrogen".
50. A. H. Hughes, R. J. Corruccini, and E. C. Gilbert, J. Am. Chem. Soc., 61, 2639 (1939). "Studies on hydrazine: The heat of formation of hydrazine and of hydrazine hydrate".
51. H. Hunt and P. F. Stehlo, Purdue University, Private Communication.
52. E. Janscke, Z. Elektrochem., 40, 462-469 (1934); C.A., 23, 6364 (1934); cf. C.A., 25, 3226 (1931). "The heats of combustion and formation of organic compounds in relation to their structure".

CONFIDENTIAL

CONFIDENTIAL  
SECURITY INFORMATION

53. R. S. Jessup, J. Research Nat. Bur. Standards, 20, 559 (1936).  
"Heat of combustion of isoprene".
- 53a. W. H. Jones and C. E. Starr, Ind. Eng. Chem. Anal. Ed., 13, 267-290 (1941). "Determination of the heat of combustion of gasolines".
54. H. Kast. Chem. Zeit., 1925, 621 (as given by Barlot, Ref. 27).
55. M. S. Kharasch, Bur. Standards J. Research, 2, 359-430 (1929).  
"Heats of combustion of organic compounds".
- 55-9. E. Berner, Arch. Math. Naturwissenschaften, 26, No. 6, 5-134 (1926);  
C.A., 21, 3814 (1927). "A contribution to the thermochemistry of organic compounds".
- 55-24. Berthelot and G. Andre, Ann. chim. et phys. (7), 17, 433 (1899).  
"New studies on the heats of formation and combustion of various nitrogen compounds and others".
- 55-25. Berthelot and Delépine, Compt. rend., 122, 920 (1899). "Lactic acid".
- 55-26. Berthelot and Delépine, Ann. chim. et phys. (7), 21, 289 (1900).  
"Heat of combustion of very volatile liquids".
- 55-34. Berthelot and Matignon, Compt. rend., 114, 1145 (1892). "Heats of combustion and formation of alcohol and formic and acetic acids".
- 55-37. Berthelot and Matignon, Ann. chim. et phys. (6), 30, 547 (1893).
- 55-41. Berthelot and M. P. Petit, Ann. chim. et phys. (6), 16, 107 (1869).  
"Heats of combustion and formation of nitriles".
- 55-44. Berthelot and Rivals, Ann. chim. et phys. (7), 7, 29 (1896).  
"New studies on thermochemical relations between aldehydes, alcohols, and acids".
- 55-51. P. Bruylants and A. Christiaen, Bull. soc. chim. Belg., 34, 144 (1925). "Thermochemistry of the butene nitriles".
- 55-52. Bureau of Standards, Cir. 11, 3d. ed., (1917). "The standardization of bomb calorimeters".
- 55-54. J. Coops and P. E. Verkade, Rec. trav. chim., 44, 983 (1925).  
"Calorimetric researches. IX. The heat of combustion of d- and meso-tartaric acids, succinic acid and a number of their derivatives".

CONFIDENTIAL

CONFIDENTIAL  
SECURITY INFORMATION

- 55-58. M. Delapine, Compt. rend., 131, 684 (1900). "Acetals of monohydroxy alcohols".
- 55-60. M. Delapine, Compt. rend., 132, 777 (1901). "Heat of formation of acetals compared to that of isomeric compounds".
- 55-64. A. G. Emery and F. G. Benedict, Am. J. Physiol., 28, 301, (1911). "The heat of combustion of compounds of physiological importance".
- 55-65. P. A. Favre and J. T. Silbermann, Ann. chim. et phys. (3), 3h, 357 (1852). "Researches on the quantities of heat released in chemical and molecular reactions".
- 55-66. Fischer and Wrede, Sitzber. kgl. preuss. Akad. Wiss., 1904, 687.
- 55-77. H. Guillemand, Ann. chim. et phys. (8), 14, 311 (1908). "Studies on the tautomerism of cyanogen derivatives".
- 55-78. J. Quinchant, Compt. rend., 121, 354 (1895). "Heat of combustion of some B-keto esters". Ibid., 122, 943 (1896). "Heat of combustion of cyano derivatives." All data collected and evaluated in Ann. chim., 10, 51 (1918). "Study of the acid function in methenic and methinic derivatives".
- 55-80. F. Henning, Z. physik. Chem., 97, 467 (1921). "Heats of combustion of benzoic acid, naphthalene, and cane sugar. A critical review".
- 55-93. Ph. Landrieu, Compt. rend., 140, 867 (1905). "Heat of formation of oximes".
- 55-97. P. Lemoult, Compt. rend., 143, 772 (1906). "Heat of combustion and formation of some cyclic nitrogen compounds".
- 55-98. P. Lemoult, Compt. rend., 143, 902 (1906). "Carbysamines and nitriles".
- 55-99. P. Lemoult, Ann. chim. et phys. (8), 10, 395 (1907). "Theoretical and experimental studies on the heats of combustion and formation of organic compounds. I. Primary, secondard, and tertiary amines".
- 55-100. P. Lemoult, Ann. chim. et phys. (8), 13, 562 (1908). "2. Hydrazine compounds".
- 55-101. P. Lemoult, Ann. chim. et phys. (8), 14, 184 (1908). "3. Azoxy compounds". Ibid., 14, 289 (1908). "4. Azo compounds".
- 55-102. P. Lemoult, Compt. rend., 148, 1602 (1909). "Comparison of nitriles and carbysamines".

CONFIDENTIAL

CONFIDENTIAL  
SECURITY INFORMATION

- 55-110. W. Louguinine, *Ann. chim. et phys.* (5), 21, 139 (1880). "Heat released in the combustion of some isomeric alcohols of the fatty series as well as that of oenanthal".
- 55-113. W. Louguinine, *Ann. chim. et phys.* (5), 23, 384 (1881). "Heats of combustion of some alcohols of the allylic series and of aldehydes isomeric with them".
- 55-117. W. Louguinine, *Compt. rend.*, 100, 63 (1885). "Heat of combustion of some substances of the fatty series".
- 55-118. W. Louguinine, *Compt. rend.*, 101, 1061 (1885). "Heat of combustion of some substances of the fatty series". *Ibid.*, 101, 1154 (1885). "Heat of combustion of some esters of organic acids".
- 55-119. W. Louguinine, *Ann. chim. et phys.* (6), 8, 128 (1886). "Measure of the heat of combustion of esters of some organic acids as a means of determining the heat of formation of these acids".
- 55-120. W. Louguinine, *Ann. chim. et phys.* (6), 11, 220 (1887). "Heat of combustion of fatty acids and of fats in which they occur".
- 55-121. W. Louguinine, *Ann. chim. et phys.* (6), 23, 179 (1891). "Study of the heat of combustion of some organic acids and anhydrides".
- 55-130. C. Matignon, *Ann. chim. et phys.* (6), 28, 70 (1893). "Ureides derived from acids with one and two carbon atoms". *Ibid.*, 28, 289 (1893). "Polybasic ureides and uric acid". *Ibid.*, 28, 498 (1893). "Some general results of the thermochemical study of ureides." These articles contain also the data reported by Matignon, *Compt. rend.*, 110, 1267 (1890); 113, 108 (1891).
- 55-131. C. Matignon and Deligny, *Compt. rend.*, 121, 422 (1895). "Nitro substitution".
- 55-138. C. Mourou and E. Andre, *Ann. Chim.* (9), 1, 113 (1914). "Thermochemistry of acetylenic compounds".
- 55-139. C. Mourou and J. Ch. Fougrand, *Compt. rend.*, 170, 1025 (1920). "New studies on carbon subnitride. Action of halogens, halohydric acids, and alcohols".
- 55-141. C. Mourou, A. Boutaric, and C. Dufraisse, *J. chim. phys.*, 18, 333 (1920). "Some physicochemical constants of acrolein".
- 55-142. J. A. Muller, *Ann. chim. et phys.* (8), 20, 116 (1910). "Heats of combustion and specific weights of methylamines".
- 55-147. P. Petit, *Ann. chim. et phys.* (6), 18, 145 (1889). "Researches on some nitrogen compounds of the aromatic series".

CONFIDENTIAL  
SECURITY INFORMATION

- 55-151. Reyer, Diss., Braunschweig (1923); quoted in Landolt-Bornstein Tabellen (1923).
- 55-153. T. W. Richards and H. S. Davis, J. Am. Chem. Soc., 42, 1599 (1920). "The heats of combustion of benzene, toluene, aliphatic alcohols, cyclohexanol, and other carbon compounds".
- 55-154. T. W. Richards and R. H. Jesse, J. Am. Chem. Soc., 32, 268 (1910). "The heats of combustion of the octanes and xylenes".
- 55-160. W. A. Roth, Rec. trav. chim., 45, 246 (1926). (Value by Lemoult corr. by Roth.) "A note on the work of Swietoslowski and Popow: 'Corrections of the thermochemical results of Lemoult'".
- 55-163. W. A. Roth, Unpub. work. Authority for these values in Landolt-Bornstein Tabellen- (1923).
- 55-172. W. A. Roth and Muller, Landolt-Bornstein Tabellen (1927).
- 55-175. W. A. Roth and G. J. Ostling, Ber., 46, 309 (1913). "Thermochemical studies in the alicyclic series."
- 55-179. J. Schmidlin, Ann. chim. et phys. (8), 7, 195 (1906). "Chemical and thermochemical studies on the constitution of pyrazoline".
- 55-182. F. Stohmann, Z. physik. Chem., 6, 334-357 (1890). "Heats of combustion of organic compounds".
- 55-183. F. Stohmann, Z. physik. Chem., 10, 410-424 (1892). "Heats of combustion of organic compounds".
- 55-184. F. Stohmann and E. Hausmann, J. prakt. Chem. (2), 55, 263 (1897). "Calorimetric researches. 36. Calorific value of amide and anilide of the first member of the dibasic acid series". For details consult Ber. Verhandl. K. sachs. Ges. Wiss., 49, 1 (1897).
- 55-185. F. Stohmann and Kleber, J. prakt. Chem. (2), 43, 1 (1891).
- 55-187. F. Stohmann, Cl. Kleber and H. Langbein, J. prakt. Chem. (2), 40, 77 (1889). "Calorimetric researches. 17. Heat of combustion of benzene and other hydrocarbons of aromatic series".
- 55-189. F. Stohmann, Cl. Kleber and H. Langbein, J. prakt. Chem. (2), 40, 128 (1889). "Calorimetric researches. 18. Heat of combustion of carboxylic acids of the aromatic series".
- 55-190. F. Stohmann, Cl. Kleber and H. Langbein, J. prakt. Chem. (2), 40, 202 (1889). "Calorimetric researches. 19. Heat of combustion of acids of the oxalic acid series and of fumaric and maleic acids".

CONFIDENTIAL  
SECURITY INFORMATION

- 55-192. F. Stohmann, G. Kleber, H. Langhein, and P. Offenhammer, J. prakt. Chem. (2), 48, 99 (1894). "Calorimetric researches. 30. Heat of combustion of aliphatic acids". For details of the work consult Ber. Verhandl. K. sachs. Ges. Wiss., 45, 604 (1893).
- 55-193. F. Stohmann and H. Langbein, J. prakt. Chem. (2), 42, 361 (1890). "Calorimetric researches. 22. Fats and a fatty acid".
- 55-195. F. Stohmann and H. Langbein, J. prakt. Chem. (2), 45, 505 (1892). "Calorimetric researches. 26. Heat of combustion of carbohydrates, polyalcohols, and phenols".
- 55-201. F. Stohmann, P. Rodatz and H. Herzberg, J. prakt. Chem. (2), 33, 464-472 (1886). "Calorimetric researches. 7. Heat of combustion of phenol".
- 55-203. F. Stohmann, P. Rodatz and W. Herzberg, J. prakt. Chem. (2), 25, 22-42 (1867). "Calorimetric researches. 11. Heat of combustion of ethers of phenol".
- 55-204. F. Stohmann, J. prakt. Chem. (2), 35, 40, 136 (1887). "Calorimetric researches. 12. Heat of combustion of benzene homologs".
- 55-208. F. Stohmann and R. Schmidt, J. prakt. Chem. (2), 52, 59 (1895). "Calorimetric researches. 34. Heat of combustion of amides and anilides of monobasic acids". For details consult Ber. Verhandl. K. sachs. Ges. Wiss., 47, 1 (1895).
- 55-209. F. Stohmann and R. Schmidt, J. prakt. Chem. (2), 52, 345 (1895). "Calorimetric researches. 35. Heat of combustion of hippuric acid, its homologues, and of anisuric acid."
- 55-215. W. Swietoslowski, J. Am. Chem. Soc., 42, 1092 (1920). "The restatement and correction of the thermochemical data on organic compounds. 1. The data of P. W. Zubow".
- 55-216. W. Swietoslowski, J. chim. phys., 22, 583 (1925). "Remarks concerning the paper of W. Jaeger and H. V. Steinwehr titled: On the establishment of the heat of combustion of normal substances".
- 55-217. W. Swietoslowski, Private comm.
- 55-221. W. Swietoslowski and H. Starcowska, Bull. soc. chim. Belg. (4), 31, 654 (1922). "Heat of combustion of benzoic acid, cane sugar, and naphthalene".
- 55-223. J. Thomsen, Z. physik. Chem., 52, 343 (1905). "Numerical results of a systematic study of the heats of combustion and formation of liquid organic compounds".

CONFIDENTIAL

CONFIDENTIAL  
SECURITY INFORMATION

- 55-225. A. Vialer, Ann. chim. et phys. (7), 21, 470 (1900). "Contribution to the thermochemical study of quinones. Study of the constitution of quinhydrone".
- 55-226. P. E. Verkade, Amst. Akad. Verh., 35, 496 (1926). Values by Verkade, van de Lee and Coops.
- 55-227. P. E. Verkade, Proefschr. Delft (1915).
- 55-238. E. Voisenet, Bull. soc. chim. Belg. (4), 17, 34 (1915). "Heats of formation of acrolein and of methacrolein".
- 55-239. H. v. Wartenberg, A. Muchlinski and G. Riedler, Z. angew. Chem., 37, 457 (1924). "Investigations of formaldehyde". See also H. v. Wartenberg and Lerner-Steinberg, Z. angew. Chem., 38, 591-592 (1925). "Heat of formation of formaldehyde".
- 55-243. P. W. Zubov, J. Russ. Phys.-Chem. Soc. (7), 28, 687 (1896).
- 55-245. P. W. Zubov and W. Swietoslowski, Bull. soc. chim. Belg., 37, 271 (1925). "Heat of combustion of three oxides".
56. M. S. Kharasch and R. Sner, J. Phys. Chem., 29, 625 (1925). "The electronic conception of valence and heats combustion of organic compounds".
57. O. M. Kibler and H. Hunt, J. Phys. and Colloid Chem., 53, 955-956 (1949); C.A., 43, 7318 (1949). "Heats of combustion. V. Five nitrogen-containing compounds".
58. J. Kielland, Ber., 71E, 220-226 (1938); C. A., 32, 2865 (1938). "Rapid colorimetric determination of potassium with hexanitro-diphenylamine (dipicrylamine) as well as some thermodynamic data concerning the amine and its potassium salt".
59. F. Klages, Chem. Ber., 82, 358-375 (1949) C.A., 44, 1323 (1950). "Improvement in the additive calculation of heats of combustion and the calculation of mesomerism energy from heats of combustion".
60. J. W. Knowlton and F. D. Rossini, J. Research Nat. Bur. Standards, 43, 113 (1949). "Heats of combustion and formation of cyclopropane".
61. P. N. Kogerman, Sitzber. Naturforsch. Ges. Univ. Tartu, 41, No. 3-4, 62pp. (1934); C.A., 29, 3297 (1935). "Addition reactions and polymerization of 2,3-dimethylbutadiene".
62. Catholic University of America, Contract No. 10260 Progress Report July 1 - December 31, 1947, by K. J. Laidler. "The estimation of heats of formation of organic compounds. Part I". RESTRICTED

CONFIDENTIAL

CONFIDENTIAL  
SECURITY INFORMATION

63. Catholic University of America, Contract NOrd 10260, Final Report, 30 June 1950, by K. J. Laidler. "The estimation of heats of formation of organic compounds". RESTRICTED
64. P. Landrieu, P. Baylocc, and John R. Johnson, Bull. soc. chim., 45, 36-49 (1929); C.A., 23, 2967 (1929). "Thermochemical studies in the furan series".
65. R. Lantie, Bull. soc. chim. [5], 5, 1306-1321 (1938); C.A., 33, 1582 (1939). "Heats of formation and combustion".
66. A. Lesan and G. Lepoutre, Compt. rend., 226, 1976-1978 (1948); C.A., 42, 8061 (1948). "Heats of combustion of naphthols".
67. P. Ledoult, Ann. chim. et phys. [8], 1, 496 (1903).
68. Lanhard, Z. Naturforsch., 4a, 257-261 (1949); C.A., 43, 8844 (1949). "Relationships between atomic number, effective nuclear charge, and heat of combustion".
69. Arthur D. Little, Inc., Contract W-19-020-ORD-6436 (2 January, 1947). "Study of pure explosive compounds. Part I. Correlation of organic structure with explosive properties from existing data". CONFIDENTIAL See also W. C. Lothrop and G. R. Handrick, Chem. Rev., 44, 419-445 (1949). "The relationship between performance and constitution of pure organic explosive compounds".
70. Arthur D. Little, Inc., Contract W-19-020-ORD-6436 (2 April, 1947). "Study of pure explosive compounds. Part II. Correlation of thermal quantities with explosive properties." CONFIDENTIAL
71. Arthur D. Little, Inc., Contract W-19-020-ORD-6436 and DA-19-020-ORD-12 (1 March, 1951). "Second report on synthesis and testing of high explosives". pp. 346-347. CONFIDENTIAL
72. W. C. Lothrop, G. R. Handrick and R. M. Hainer, J. Am. Chem. Soc., 73, 3581 (1951). "The structure and infrared absorption spectra of polynitrophenylmethylnitramines and polynitroanilines".
73. Mathieson Alkali Works, Inc., Lit. Report No. 76. "Uses and reactions of hydrazine". Revised April 1949.
74. C. Matignon, Ann. chim. et phys. (6), 26, 510 (1893). "General results of the thermochemical study of ureas".
75. C. Matignon and Deligny, Compt. rend., 125, 1103 (1897). "Difference between the nitroso substituted on carbon and on nitrogen".
76. M. Matui and T. Abo, Bull. Tokyo Univ. Eng., 8, 339-357 (1939); Chem. Zentr., 1941, I, 804; C.A., 37, 3 (1943). "Method of operation of an adiabatic calorimeter".

CONFIDENTIAL



CONFIDENTIAL  
SECURITY INFORMATION

77. W. S. McEwan and M. W. Rigg, *J. Am. Chem. Soc.*, 73, 4725-4727 (1951). "The heats of combustion of compounds containing the tetrazole ring".
78. C. B. Miles and H. Hunt, *J. Phys. Chem.*, 45, 1346-1359 (1941); *C.A.*, 36, 2469 (1942). "Heats of combustion. I. The heat of combustion of acetone".
79. J. J. Miller and H. Hunt, *J. Phys. Chem.*, 49, 25-31 (1945); *C.A.*, 39, 1352. "Heats of combustion. III. Heats of combustion of some polynitroparaffins".
80. H. Milone, *Gazz. chim. ital.*, 61, 153-158 (1931); *C.A.*, 25, 3624-3625 (1931). "Dioxines. LXXVI."
81. H. Milone and G. Venturello, *Gazz. chim. ital.*, 66, 808-812 (1936); *C.A.*, 32, 4646 (1937). "Dioximes. CIVIII".
82. H. Moureu and M. Dods, *Bull. soc. chim. Belg.*, 53, 4, 637-647 (1937); *C.A.*, 31, 4884 (1937). "Heats of formation of ethylene oxide, of ethanedioic and some of their homologues".
- 82a. National Bureau of Standards, "Selected values of chemical thermodynamic properties. Series I." U. S. Dept. of Commerce, Washington, D. C. (30 June 1947).
83. Naval Ordnance Laboratory, Private Communication 4 October 1951 (Determined by National Bureau of Standards).
84. Naval Powder Factory, Quarterly Report 4 by F. C. Inames, 15 December 1948. CONFIDENTIAL.
85. OSRD 702. G. B. Kistiakowsky, "The heats of combustion of explosives". (15 July 1942) CONFIDENTIAL.
86. OSRD 5306. G. Stegeman, "Heat of combustion of explosive substances". (4 July 1945) CONFIDENTIAL.
87. G. S. Parks and G. E. Moore, *J. Chem. Phys.*, 7, 1066 (1939). "The heat of combustion of isopropanol".
88. G. S. Parks, T. J. West, R. F. Naylor, P. S. Fujii and L. A. McClaine, *J. Am. Chem. Soc.*, 68, 2524 (1946). "Thermal data on organic compounds. XVIII. Modern combustion data for 14 hydrocarbons and 5 polyhydroxy alcohols."
89. L. Pauling, "The nature of the chemical bond", Cornell University Press, Ithaca, New York (1940) pp. 47-56.
90. Picatinny Arsenal Laboratory Report 134460 (20 February 1951). CONFIDENTIAL.

CONFIDENTIAL

CONFIDENTIAL  
SECURITY INFORMATION

91. Picatinny Arsenal Tech. Report 1409 (3 April 1944). "Compounds of high nitrogen content in propellant powders". CONFIDENTIAL
92. Picatinny Arsenal Tech. Report 1694 by O. E. Sheffield (26 May 1948) "Long range basic research leading to the development of ideal propellants. Compounds of high nitrogen content propellant powders". CONFIDENTIAL.
93. Picatinny Arsenal Tech. Report 1740 by W. R. Tomlinson (20 June 1949). "Properties of explosives of military interest". CONFIDENTIAL
94. Picatinny Arsenal Tech. Report 1752 by S. Helf (7 Nov. 1949). "Compounds of high nitrogen content in propellant powders". CONF.
95. Picatinny Arsenal Tech. Report 1755 by T. C. Castorina (30 Dec. 1949). "Long range basic research leading to the development of ideal propellants. Explosive plasticizers for nitrocellulose". CONFIDENTIAL
96. Picatinny Arsenal Tech. Report 1841 by S. Helf, I. Minsky and B. D. Guida (1 October 1951). "Long range basic research leading to the development of superior propellants. Compounds of high nitrogen content in propellant compositions". CONFIDENTIAL
97. J. H. Platt, J. Chem. Phys., 15, 419-420 (1947). "Influence of neighbor bonds on additive bond properties in paraffins".
98. T. M. Polonskii, Nauch. Zapiski Khar'kov Poligraf. Inst., 1940 No. 5, 103-5; Khim. Referat. Zhur., 4, No. 4, 26-7 (1941); C.A., 37, 4618 (1943). "The connection between the parachor values and the heats of combustion".
99. A. Pongratz, S. Behmert-Suss, and K. Scholtis, Ber., 77, 651 (1944); C.A., 40, 6068 (1946). "Thermochemistry of the benzidine rearrangement".
100. E. J. Prosen, W. H. Johnson and F. D. Rossini, J. Research Nat. Bur. Standards, 36, 455-461 (1946); C.A., 40, 5630 (1946). "Heats of combustion and formation at 25° of the alkylbenzenes through C<sub>10</sub>H<sub>14</sub>, and of the high normal monoalkylbenzenes".
101. E. J. Prosen, F. W. Maron, and F. D. Rossini, J. Research Nat. Bur. Standards, 42, 259 (1949). "Heat of isomerization of the two butadienes".
102. E. J. Prosen and F. D. Rossini, J. Research Nat. Bur. Standards, 31, 59 (1945). "Heats of formation and combustion of 1,3-butadiene and styrene".

CONFIDENTIAL

CONFIDENTIAL  
SECURITY INFORMATION

103. E. J. Prosen and F. D. Rossini, J. Research Nat. Bur. Standards, 24, 263-269 (1945). "Heats of combustion and formation of the paraffin hydrocarbons at 25°C."
104. N. A. Faschin, Z. Physik. Chem., 124, 16 (1926). "The binary system naphthalene-*m*-dinitrobenzene".
105. M. Rebeš, Monatsh., 73, 57-81 (1940); Sitzber. Akad. Wiss. Wien, Math.-naturw. Klasse. Abt. IIb, 149, 1-25 (1940); C.A., 36, 702 (1942). "Distribution of the heats of formation of organic molecules into bond increments. I. Hydrocarbons".
106. M. Rebeš, Monatsh., 73, 257-68 (1941); C.A., 36, 5084 (1942). "Distribution of the heats of formation of organic molecules into bond increments. II. Normal chains and isomers".
107. J. W. Richardson and G. S. Parks, J. Am. Chem. Soc., 61, 3543-3546 (1939). "Thermal data on organic compounds. III. Modern combustion data for some non-volatile compounds containing carbon, hydrogen, and oxygen".
108. W. H. Rinkenbach, Ind. Eng. Chem., 18, 1195 (1926).
109. W. H. Rinkenbach, Ind. Eng. Chem., 19, 925-9 (1927). "Preparation and properties of diethylene glycol dinitrate".
110. W. H. Rinkenbach, J. Am. Chem. Soc., 52, 116 (1930). "The heats of combustion and formation of aromatic nitro compounds".
111. D. E. Roberts and R. S. Jessup, J. Research Nat. Bur. Standards, 40, 281 (1948); C.A., 42, 5327 (1948). "Heat of combustion of phenyl-beta-naphthylamine (N-phenyl-2-naphthylamine)".
112. J. S. Roberts and H. A. Skinner, Trans. Faraday Soc., 45, 339-357 (1949); C.A., 43, 6479 (1949). "Dissociation energies of carbon bonds, and resonance energies in hydrocarbon radicals".
113. F. D. Rossini, Bur. Standards J. Research, 13, 189 (1934). "Heats of combustion and of formation of the normal aliphatic alcohols in the gaseous and liquid states, and the energies of their atomic linkages".
114. F. D. Rossini, Ind. Eng. Chem., 29, 1424-1430 (1937). "Heats of formation of simple organic molecules".
115. F. D. Rossini, K. S. Pitzer, W. J. Taylor, J. P. Ebert, J. E. Kilpatrick, C. W. Beckett, M. G. Williams, and H. G. Werner, "Selected values of properties of hydrocarbons", National Bureau of Standards, U. S. Government Printing Office, Washington, D.C. (1947).
116. W. A. Feth and K. Isecke, Ber., 77B, 537-539 (1944); C.A., 40, 5328 (1945). "The heats of combustion and formation of tetramethane".

CONFIDENTIAL  
SECURITY INFORMATION

117. W. A. Roth and Machaleidt, "Dissertation", Braunschweig (1930).
118. W. A. Roth and E. Rist-Schumacher, Z. Elektrochem., 50, 7-9 (1946); C.A., 36, 6183 (1944). "Thermochemistry of the sulfonic acids and acid chlorides".
119. P. P. Emtsov and L. A. Server'yanov, J. Russ. Phys.-Chem. Soc., 50, 140-144 (1918); C.A., 18, 1401 (1924). "The heats of combustion and of formation of certain explosive nitro compounds from their elements".
120. L. Ruzicka and P. Schlapfer, Helv. Chim. Acta, 16, 262 (1932); C.A., 27, 3919 (1933). "Carbon rings. XIII. Heats of combustion of high-membered cyclic compounds".
121. D. J. Sallee and J. R. Gray, J. Am. Chem. Soc., 70, 2650 (1948). "Heats of combustion of some organic nitrogen compounds".  
See also: J. Am. Chem. Soc., 71, 5925 (1949)
122. Sarasa and Vieille, Compt. rend., 23, 269 (1881). "Heats of formation of explosives; numerical results".
123. E. Schjanberg, Z. physik. Chem., A 172, 197-233 (1935); C.A., 29, 3589 (1935). "The heat of combustion and refraction data for various chlorine-substituted fatty acids and esters. The relation between chemical constitution, heat of combustion, and molecular refraction".
124. A. Schmidt, Z. Ges. Schiess- u. Sprengstoffe., 29, 259-266, 296-301 (1934). "Thermochemical tables for explosive chemistry".
125. M. Simonetta, Chimica e Industria (Milan), 29, 37 (1947); C.A., 41, 7165 (1947). "The thermal problem in the chain of reactions between ethylene oxide and methanol".
126. H. A. Skinner and H. D. Springall, Nature, 162, 343-344 (1948); C.A., 43, 10 (1949). L. H. Long, Nature, 162, 344-345 (1948); C.A., 43, 10 (1949). "Definitions of bond energy in physical chemistry".
127. M. Souders, C. S. Matthews, and C. O. Hurd, Ind. Eng. Chem., 41, 1048-1055 (1949). "Entropy and heat of formation of hydrocarbon vapors".
128. Solid Propellant Information Agency, SPIA/M3 Propellant Powder Ingredients Manual (1951, unpublished).
  - 128-1. Determined at Naval Powder Factory.
  - 128-2. Determined at Aerojet Engineering Corp.
  - 128-3. Data given by British Advisory Council reports.
  - 128-4. Determined by Picatinny Arsenal.

CONFIDENTIAL

CONFIDENTIAL  
SECURITY INFORMATION

129. R. Spitzer and H. M. Huffman, *J. Am. Chem. Soc.*, 69, 211 (1947). "Heats of combustion of cyclopentane, cyclohexane, cycloheptane and cyclooctane".
130. British Ministry of Supply Report AC-7048 (July 1944) by H. D. Springall and E. Roberts. "On the calculation of heats of formation, and of the explosive powers of compounds from bond, resonance, and "group" energies". BRITISH SECRET
131. A. Stern and G. Klebs, *Ann.*, 500, 91-108 (1932); *C.A.*, 27, 1346 (1933). "Application of calorimetry to pyrroles and pyrrole dyes. II. Calorimetric determination of simple and multiple-nucleus pyrrole derivatives".
132. A. Stern and G. Klebs, *Ann.*, 504, 287 (1933); *C.A.*, 27, 4796. "Application of calorimetry to pyrroles and pyrrole dyes. III. Calorimetric determination of simple and polymucleated pyrrole derivatives".
133. M. V. Sullivan and H. Kunz, *J. Phys. and Colloid Chem.*, 53, 497 (1949). "Heats of combustion. IV. The heats of combustion of five aromatic amines".
134. S. Szwarc, *Svensk Kem. Tidn.*, 58, 71-81 (1946); *C.A.*, 40, 3972 (1946). "Determination of combustion heats of organosulfur compounds."
135. V. Svyatoslavskii, *Z. physik. Chem.*, 72, 49-83 (1910); *C.A.*, 4, 2101 (1910). "Thermochemical investigations of organic compounds. III. Nitrogen compounds". Also given by same author in *J. Russ. Phys.-Chem. Soc.*, 41, 387-429 (1909); *C.A.*, 6, 1907 (1911), and *J. Russ. Phys.-Chem. Soc.*, 41, 920-925 (1909); *C.A.*, 5, 1414 (1911).
136. F. Swarts, *Rec. trav. chim.*, 32, 59-89 (1913). "Heat of formation of some organic fluorine compounds".
137. F. Swarts, *Rec. trav. chim.*, 33, 281-98 (1914); *Bull. acad. roy. belg.*, 1914, 18-44; *C.A.*, 9, 445 (1915). "The heat of combustion of some aromatic nitro derivatives".
138. F. Swarts, *J. chim. phys.*, 17, 3-70 (1919). "Thermochemical studies on organic fluorine compounds".
139. W. Swietoslowski and J. Bobinska, *Koczniki Chem.*, 9, 723 (1929); *C.A.*, 24, 1790 (1930). "The heats of combustion of camphor, azobenzene, and hydrazobenzene".
140. J. Taylor and C. K. L. Hall, *J. Phys. Coll. Chem.*, 51, 593 (1947). "Determination of the heat of combustion of nitroglycerin and the thermochemical constants of nitrocellulose".
141. J. Taylor, C. K. L. Hall, and H. Thomas, *J. Phys. and Colloid Chem.*, 51, 580-592 (1947); *C.A.*, 41, 3967 (1947). "The thermochemistry of propellant explosives".

CONFIDENTIAL  
SECURITY INFORMATION

142. A. Testmann and N. Armandi, *Atti accad. sci. Torino, Classe sci. fis. nat. nat.*, 77, 278-281 (1942); *Chem. Zentr.*, 1943, I, 1848-1849; *E. A.*, 38, 3041 (1944). "Preparation of benzitrosorbital and its explosive properties".
143. K. Tocioka and Takahashi, *Rep. Res. Dep. Nav. Expl.* 1934, p. 122. *Tables An. Int. de Constantes (1931-1934) 1st Part, Vol. XI*, p. 12-51. (As cited by Burlot, Ref. 27).
144. P. E. Verkade and J. Coops, *Rec. trav. chim.*, 46, 903-917 (1927); *C.A.*, 22, 571 (1928). "Calorimetric researches XIV. Heats of combustion of successive members of homologous series: the normal primary aliphatic alcohols".
145. P. E. Verkade and J. Coops, Jr., *Proc. Acad. Sci. Amsterdam*, 36, 76-87 (1933); *C.A.*, 27, 3134 (1933). "Alterations in the properties of monoalkylmalonic acids. Malkin's views concerning the alternation phenomena".
146. P. E. Verkade, J. Coops, and H. Hartman, *Rec. trav. chim.*, 45, 585-606 (1926); *C.A.*, 21, 1216 (1927). "Calorimetric researches. XIII. Heats of combustion of successive terms of homologous series".
147. P. E. Verkade and H. Hartman, *Rec. trav. chim.*, 52, 945-68 (1933); *C.A.*, 28, 30 (1934). "Calorimetric investigations. XII. Heats of combustion of methyl-, ethyl- and phenyl-substituted succinic acids and of their anhydrides ... (a) experimental data".
148. P. E. Verkade, H. Hartman, and J. Coops, *Rec. trav. chim.*, 45, 373-393 (1926); *C.A.*, 20, 2937 (1926). "Calorimetric researches. X. Heats of combustion of successive terms of homologous series: dicarboxylic acids of the oxalic acid series".
149. G. Waddington, J. W. Knowlton, D. W. Scott, G. D. Oliver, S. S. Todd, W. N. Hubbard, J. C. Smith and M. H. Huffran, *J. Am. Chem. Soc.*, 71, 797 (1949). "Thermodynamic properties of thiophene".
150. A. D. Walsh, *Trans. Faraday Soc.*, 43, 60-77 (1947); *C.A.*, 41, 4978 (1947). "Remarks on the strengths of bonds".
151. O. W. Wasland, "The theory of resonance", John Wiley and Sons, New York (1944) pp. 52-87.
152. H. Wieland, K. Heymann, T. Tsatsas, D. Juchacz, G. Varvoglis, G. Labriola, O. Döbbelestein, and H. S. Boyd-Barrett, *Ann.*, 514, 145-181 (1934); *C.A.*, 29, 1800 (1935). "Occurrence of free radicals in chemical reactions. VIII"

CONFIDENTIAL

CONFIDENTIAL  
SECURITY INFORMATION

153. J. B. Willis, Trans. Faraday Soc., 43, 97-102 (1947); C.A., 11, 5008 (1947). "Heats of combustion of some organic bases and their salts. The resonance energy of acridine and phenazine".
154. P. H. Wise, K. T. Serijan, and I. A. Goodman, Technical Note 2081 for National Advisory Committee for Aeronautics (Copied for OCO 12 September, 1950). "Correlation of physical properties with molecular structure for dicyclic hydrocarbons. I. 2- Alkylbiphenyls. 1,1-diphenylalkane,  $\alpha,\omega$ -diphenylalkane, 1,1-dicyclohexylalkane, and  $\alpha,\omega$ -dicyclohexylalkane series". - Also appears as: K. T. Serijan and P. H. Wise, J. Am. Chem. Soc., 73, 4766-4769 (1951). "Dicyclic hydrocarbons. III. Biphenyl- and dicyclohexylalkanes through C<sub>18</sub>".

CONFIDENTIAL  
SECURITY INFORMATION

1. Glossary of Terms and Symbols

A functional group is any substituent for a hydrogen of a normal paraffin.

A monofunctional compound is one containing a single functional group.

A polyfunctional compound contains more than one functional group which may be alike or different.

A homofunctional compound contains only one functional type and may be either mono or polyfunctional.

A heterofunctional compound contains more than one functional type and is necessarily polyfunctional.

Molar oxygen balance is equal to the atoms of external oxygen required to complete the combustion of a compound to carbon dioxide, water, and nitrogen (and to hydrohalic acid and sulfur dioxide when X and S are present).

$Q_C$  = heat of combustion ( $H_2O$  liquid), heat evolved;  $Q_C$  of Part II (Ref. 70);  $-\Delta H$  of Lewis and Randall.

superscript p = constant pressure as in  $Q_C^p$

superscript v = constant volume as in  $Q_C^v$

x = oxygen balance on a molar basis

$x'$  = x divided by number of homofunctional groups

y =  $Q_C^p$

$y'$  = ( $Q_C^p - 5.7$ ) divided by number of homofunctional groups

a = intercept of a homologous series; the value of y when x = 0. For paraffins the intercept is 5.7.

b = slope of a homologous series, or  $\frac{y_2'}{x_2'} - \frac{y_1'}{x_1'}$ . For normal paraffins (liq) the slope is 52.08.

$a'$  = (a - 5.7) for liquid and solid state, (a - 5.5) for gas state; the value of  $y'$  when  $x' = 0$ .

$b'$  = (52.08 - b) for liquid and solid states, (52.48 - b) for gas state.

CONFIDENTIAL



CONFIDENTIAL  
SECURITY INFORMATION

J. Heat of Combustion Data

<u>Table</u>		<u>Page</u>
61	Normal Paraffins (liquid)	516
62	Normal Paraffins (gas)	516
63	Normal Paraffins Heat of Vaporisation at 25°C	517
64	Branched Paraffins (liquid)	517
65	Cyclopropanes (liquid and solid)	518
66	Cyclobutanes (liquid)	518
67	Normal Alkylcyclopentanes (liquid)	518
68	Normal Alkylcyclohexanes (liquid)	519
69	Cycloheptanes (liquid)	519
70	Other Saturated Alicyclic Hydrocarbons	519
71	Normal Olefins (liquid)	520
72	Normal Olefins (gas)	520
73	Branched Olefins (liquid)	521
74	Diolefins (liquid)	522
75	Normal Acetylenes (liquid)	522
76	Normal Alkylbenzenes (liquid)	523
77	Normal Alkylbenzenes (gas)	523
78	Normal Alkylbenzenes Heat of Vaporization	524
79	Branched Alkylbenzenes (liquid and solid)	524
80	Polycyclics (solid)	525
81	Biphenyls (liquid)	525
82	Naphthalenes (solid)	526
83	Primary Alcohols (liquid)	526
84	Primary Alcohols (gas)	527
85	Secondary Alcohols (liquid)	527
86	Tertiary Alcohols (liquid)	527
87	Mixed Polyalcohols (liquid)	528
88	Mixed Polyalcohols (solid)	528
89	Aromatic Hydroxyls (solid)	528
90	Ethers (liquid)	529
91	Aliphatic Ethers (gas)	529
92	Oxiranes (Ethylene Oxides) (liquid)	529
93	Furans (liquid)	530
94	Aldehydes (liquid)	530
95	Aldehydes (gas)	530
96	Ketones (liquid)	531
97	Acids (liquid)	531
98	Acids (solid)	532
99	Acid Anhydrides (liquid)	533
100	Acid Anhydrides (solid)	533
101	Esters (liquid)	534
102	Esters (solid)	534
103	Nitriles (liquid)	535
104	Nitriles (solid)	535
105	Carbonyl Amines (Isonitriles) (liquid)	535
106	Primary Amines (liquid)	536

CONFIDENTIAL

CONFIDENTIAL  
SECURITY INFORMATION

Heat of Combustion Data (cont'd)

<u>Table</u>		<u>Page</u>
107	Primary Amines (gas)	536
108	Primary Amines (solid)	536
109	Secondary Amines (liquid)	537
110	Secondary Amines (solid)	537
111	Tertiary Amines (liquid)	537
112	Azides (liquid)	538
113	Azides (solid)	538
114	Hydrazines (solid)	539
115	Hydrazides (solid)	539
116	Azo's (solid)	540
117	Azox's (solid)	540
118	Guanidines (solid)	541
119	Tetrazoles (solid)	542
120	Oximes (solid)	542
121	Aliphatic Nitros (liquid)	543
122	Aliphatic Nitros (solid)	544
123	Aromatic Nitros (liquid)	545
124	Aromatic Nitros (solid)	545
125	Aliphatic Gem Trinitros (liquid)	548
126	Aliphatic Gem Trinitros (solid)	548
127	Nitrates (liquid)	549
128	Nitrates (solid)	550
129	Nitramines (solid)	551
130	Nitrides (solid)	552
131	Fluorox (liquid)	553
132	Fluorox (solid)	553
133	Gen Trifluorox (liquid)	554
134	Nitrosamines (solid and liquid)	554
135	Thiophenes (solid and liquid)	555
136	Nitrosos (solid)	555
137	Heat of Hydration	556
138	Salt Link	557

CONFIDENTIAL

TABLE 61

NORMAL PARAFFINS (liquid)

ADL NO.	FORMULA	NAME	MOL. WT.	OXYGEN REQUIRED	OXYGEN BALANCE	G.CAL/MOLE		Y'	X'	REF.
						Observed	Calculated			
	C <sub>3</sub> H <sub>8</sub>	Propane	44.09	10	-262.0	553.8	553.8			115-1
	C <sub>4</sub> H <sub>10</sub>	n-Butane	58.12	13	-357.9	682.8	682.7			115-1
	C <sub>5</sub> H <sub>12</sub>	n-Pentane	72.15	16	-351.8	838.8	839.0			115-1
	C <sub>6</sub> H <sub>14</sub>	n-Hexane	86.17	19	-352.8	925.0	925.2			115-1
	C <sub>7</sub> H <sub>16</sub>	n-Heptane	100.20	22	-351.3	1151.3	1151.4			115-1
	C <sub>8</sub> H <sub>18</sub>	n-Octane	114.22	25	-350.2	1307.5	1307.7			115-1
	C <sub>9</sub> H <sub>20</sub>	n-Nonane	128.25	28	-348.3	1493.8	1483.9			115-1
	C <sub>10</sub> H <sub>22</sub>	n-Decane	142.28	31	-348.6	1620.1	1625.1			115-1
	C <sub>11</sub> H <sub>24</sub>	n-Undecane	156.30	34	-346.0	1746.3	1775.1			115-1
	C <sub>12</sub> H <sub>26</sub>	n-Dodecane	170.33	37	-347.6	1912.6	1931.6			115-1
	C <sub>13</sub> H <sub>28</sub>	n-Tridecane	184.35	40	-347.2	2080.9	2068.9			115-1
	C <sub>14</sub> H <sub>30</sub>	n-Tetradecane	198.38	43	-346.8	2215.1	2215.1			115-1
	C <sub>15</sub> H <sub>32</sub>	n-Pentadecane	212.41	46	-346.5	2401.1	2401.1			115-1
	C <sub>16</sub> H <sub>34</sub>	n-Hexadecane	226.43	49	-346.4	2577.6	2577.6			115-1
	C <sub>17</sub> H <sub>36</sub>	n-Heptadecane	240.46	52	-346.0	2713.9	2713.8			115-1
	C <sub>18</sub> H <sub>38</sub>	n-Octadecane	254.48	55	-345.8	2870.2	2870.1			115-1
	C <sub>19</sub> H <sub>40</sub>	n-Nonadecane	268.51	58	-345.6	3026.4	3026.3			115-1
	C <sub>20</sub> H <sub>42</sub>	n-Eicosane	282.54	61	-345.5	3182.7	3182.6			115-1
		$y = 5.67 + 52.00x$								
								$I_x = 657.0$		
								$I_y = 31,361.2$		
								$2xy = 1,419,340.00$		
								$I_x^2 = 27,081.00$		
								$n = 18$		
	CH <sub>4</sub>	Methane	16.04	4	-399.0	-	211.0			115-1
	C <sub>2</sub> H <sub>6</sub>	Ethane	30.07	7	-372.5	-	270.2			115-1

TABLE 62

NORMAL PARAFFINS (gas)

ADL NO.	FORMULA	NAME	MOL. WT.	OXYGEN REQUIRED	OXYGEN BALANCE	G.CAL/MOLE		Y'	X'	REF.
						Observed	Calculated			
	C <sub>3</sub> H <sub>8</sub>	Propane	44.09	10	-372.5	372.8	372.8			115-1
	C <sub>4</sub> H <sub>10</sub>	n-Butane	58.12	13	-342.9	530.61	530.6			115-1
	C <sub>5</sub> H <sub>12</sub>	n-Pentane	72.15	16	-347.9	687.99	687.7			115-1
	C <sub>6</sub> H <sub>14</sub>	n-Hexane	86.17	19	-351.8	816.14	816.2			115-1
	C <sub>7</sub> H <sub>16</sub>	n-Heptane	100.20	22	-351.3	1002.57	1002.6			115-1
	C <sub>8</sub> H <sub>18</sub>	n-Octane	114.22	25	-351.3	1160.01	1160.1			115-1
	C <sub>9</sub> H <sub>20</sub>	n-Nonane	128.25	28	-350.2	1317.45	1317.5			115-1
	C <sub>10</sub> H <sub>22</sub>	n-Decane	142.28	31	-348.3	1474.90	1474.9			115-1
	C <sub>11</sub> H <sub>24</sub>	n-Undecane	156.30	34	-348.6	1632.34	1627.1			115-1
	C <sub>12</sub> H <sub>26</sub>	n-Dodecane	170.33	37	-346.0	1789.78	1789.8			115-1
	C <sub>13</sub> H <sub>28</sub>	n-Tridecane	184.35	40	-347.6	1947.23	1947.3			115-1
	C <sub>14</sub> H <sub>30</sub>	n-Tetradecane	198.38	43	-347.2	2104.67	2104.7			115-1
	C <sub>15</sub> H <sub>32</sub>	n-Pentadecane	212.41	46	-346.8	2262.11	2262.1			115-1
	C <sub>16</sub> H <sub>34</sub>	n-Hexadecane	226.43	49	-346.8	2419.55	2419.6			115-1
	C <sub>17</sub> H <sub>36</sub>	n-Heptadecane	240.46	52	-346.2	2577.0	2577.0			115-1
	C <sub>18</sub> H <sub>38</sub>	n-Octadecane	254.48	55	-346.0	2734.4	2734.5			115-1
	C <sub>19</sub> H <sub>40</sub>	n-Nonadecane	268.51	58	-345.8	2891.88	2891.9			115-1
	C <sub>20</sub> H <sub>42</sub>	n-Eicosane	282.54	61	-345.6	3049.33	3049.3			115-1
		$y = 5.50 + 52.48x$								
								$I_x = 626.0$		
								$I_y = 31,006.6$		
								$2xy = 1,425,435.43$		
								$I_x^2 = 27,054.0$		
								$n = 19$		
	CH <sub>4</sub>	Methane	16.04	4		212.80	215.4			115-1

TABLE 63  
NORMAL PARAFFIN- $\alpha$ -OLEFINS OF CARBOHYDRATES, LIQUID

ACL NO.	FORMULA	NAME	MW. WT.	OXYGEN REQUIRED	OXYGEN BALANCE	$\Delta^{\circ}$ OBSERVED		Y'	Y''	REF.
						KCAL/MOLE	CAL/GRAM			
	C <sub>6</sub> H <sub>14</sub>	n-Hexane	86.17			3.3	66.7			115-1
	C <sub>7</sub> H <sub>16</sub>	n-Heptane	98.18			5.1	72.4			115-1
	C <sub>8</sub> H <sub>18</sub>	n-Octane	114.23			6.1	78.1			115-1
	C <sub>9</sub> H <sub>20</sub>	n-Nonane	128.26			7.0	83.7			115-1
	C <sub>10</sub> H <sub>22</sub>	n-Decane	142.28			8.7	89.2			115-1
	C <sub>11</sub> H <sub>24</sub>	n-Undecane	156.30			9.9	94.8			115-1
	C <sub>12</sub> H <sub>26</sub>	n-Dodecane	170.33			11.1	100.5			115-1
	C <sub>13</sub> H <sub>28</sub>	n-Tridecane	184.35			12.3	106.1			115-1
	C <sub>14</sub> H <sub>30</sub>	n-Tetradecane	198.38			13.5	111.7			115-1
	C <sub>15</sub> H <sub>32</sub>	n-Pentadecane	212.40			14.6	117.3			115-1
	C <sub>16</sub> H <sub>34</sub>	n-Hexadecane	226.43			15.8	122.9			115-1
	C <sub>17</sub> H <sub>36</sub>	n-Heptadecane	240.45			17.0	128.5			115-1
	C <sub>18</sub> H <sub>38</sub>	n-Octadecane	254.48			18.2	134.1			115-1
	C <sub>19</sub> H <sub>40</sub>	n-Nonadecane	268.50			19.4	139.7			115-1
	C <sub>20</sub> H <sub>42</sub>	n-Eicosane	282.53			20.5	145.3			115-1
	C <sub>21</sub> H <sub>44</sub>	n-Heneicosane	296.55			21.7	150.9			115-1
	C <sub>22</sub> H <sub>46</sub>	n-Biosane	310.58			22.9	156.5			115-1
						24.1	162.1			115-1
						AVERAGE		86.3		
	C <sub>2</sub> H <sub>6</sub>	Ethane	30.07			2.6	(Calc.)			115-1
	C <sub>1</sub> H <sub>4</sub>	Methane	16.04			1.1	(Calc.)			115-1
		a) $\Delta^{\circ} (g) - \Delta^{\circ} (l)$ These values were subtracted from $\Delta^{\circ} (gas)$ to give $\Delta^{\circ} (liq)$ for paraffins, olefins and acetylenes where necessary to obtain data given in Tables 67, 68, 72, 73, and 75.								

TABLE 64  
BRANCHED PARAFFINS (Liquid)

ACL NO.	FORMULA	NAME	MW. WT.	OXYGEN REQUIRED	OXYGEN BALANCE	$\Delta^{\circ}$ KCAL/MOLE		Y'	Y''	REF.
						OBSERVED	CALCULATED			
	C <sub>4</sub> H <sub>10</sub>	2-Methylpropane	58.12	13	-57.9	681.6	682.2	676.9	23	115-2
	C <sub>5</sub> H <sub>12</sub>	2-Methylbutane	72.15	15	-354.8	817.3	824.7	831.6	17	115-2
		2,2-Dimethylpropane				815.2		829.5	16	115-2
	C <sub>6</sub> H <sub>14</sub>	2-Methylpentane	86.17	19	-352.8	963.7		980.0	10	115-2
		3-Methylpentane				794.3		799.6	19	115-2
		2,3-Dimethylbutane				993.1		997.4	19	115-2
		2,2-Dimethylbutane				991.5		987.8	19	115-2
	C <sub>7</sub> H <sub>16</sub>	3-Methylhexane	100.20	22	-351.3	1150.0	1149.7	1144.3	22	115-2
		4-Methylhexane				1150.6		1144.9	22	115-2
		2,3-Dimethylpentane				1151.1		1145.1	22	115-2
		2,4-Dimethylpentane				1149.7		1143.4	22	115-2
		2,2-Dimethylpentane				1148.7		1143.0	22	115-2
		3,3-Dimethylpentane				1147.9		1141.7	22	115-2
		2,3,3-Trimethylpentane				1145.8		1141.1	22	115-2
		2,2,3-Trimethylpentane				1144.3		1140.4	22	115-2
	C <sub>8</sub> H <sub>18</sub>	2-Methylheptane	114.22	25	-350.2	1356.3	1356.2	1350.6	25	115-2
		3-Methylheptane				1356.9		1351.2	25	115-2
		4-Methylheptane				1357.1		1351.4	25	115-2
		1-Ethylhexane				1357.4		1351.7	25	115-2
		2,3-Dimethylhexane				1356.9		1351.2	25	115-2
		2,4-Dimethylhexane				1356.8		1350.1	25	115-2
		2,5-Dimethylhexane				1356.0		1299.3	25	115-2
		3,6-Dimethylhexane				1357.0		1351.3	25	115-2
		2-Methyl-3-ethylpentane				1357.6		1351.9	25	115-2
		2,3,6-Trimethylpentane				1356.1		1350.6	25	115-2
		3,5-Dimethylhexane				1356.6		1298.9	25	115-2
		3,3-Dimethylhexane				1356.7		1350.0	25	115-2
		3-Methyl-3-ethylpentane				1356.9		1351.1	25	115-2
								1x' = 622.0		
								2y' = 32,345.0		
								1x'y' = 734,118.8		
								1x'y' = 10,122.0		
								n = 25		

TABLE 65  
CYCLOPROPANES (liquid and solid)

ADL NO	FORMULA	NAME	MOL. WT.	OXYGEN REQUIRED	OXYGEN BALANCE	O <sub>c</sub> KCAL/MOLE		Y'	X'	REF.
						OBSERVED	CALCULATED			
	C <sub>3</sub> H <sub>6</sub>	Cyclopropane	42.08	9	-130.2	625.3	639.5	100.6	9	60
	C <sub>3</sub> H <sub>4</sub> O <sub>2</sub>	1,1-Cyclopropanedicarboxylic acid	120.10	"	-110.7	652.9	661.9	60.9	"	55-187
	C <sub>3</sub> H <sub>4</sub> O <sub>2</sub>	cis-1,2-Cyclopropanedicarboxylic acid (a)	"	"	"	"	"	65.9	"	55-187
	C <sub>3</sub> H <sub>4</sub> O <sub>2</sub>	1,1,2-Cyclopropanedicarboxylic acid (a)	216.12	"	-66.0	682.7	676.3	62.2	9	55-187
	C <sub>3</sub> H <sub>4</sub> O	Cyclopropyl methyl ketone	68.11	13	-157.1	677.3	685.4	67.7	9	55-174
	C <sub>3</sub> H <sub>4</sub> O <sub>2</sub>	Cyclopropanedimethanol	106.13	"	-217.3	691.6	700.3	66.7	13	55-215
	C <sub>3</sub> H <sub>4</sub> O <sub>2</sub>	Tetrahydro-1,1,2,2-cyclopropanetetracarboxylate (a)	276.22	21	-122.5	707.6	716.9	60.3	13	55-215
								112.5	21	55-187

$y' = 16.22 + 51.95x'$

$z' = 101.0$   
 $y' = 5,372.7$   
 $z' = 67,384.1$   
 $z'' = 1705.0$   
 $n = 9$

TABLE 66  
CYCLOBUTANES (liquids)

ADL NO	FORMULA	NAME	MOL. WT.	OXYGEN REQUIRED	OXYGEN BALANCE	O <sub>c</sub> KCAL/MOLE		Y'	X'	REF.
						OBSERVED	CALCULATED			
	C <sub>4</sub> H <sub>8</sub>	Cyclobutanecarboxylic acid	104.11	12	-191.8	611.0	616.4	632.2	12	55-175
	C <sub>4</sub> H <sub>8</sub> O	Cyclobutanemethanol	86.13	14	-262.1	639.2	637.6	637.6	12	55-215
	C <sub>4</sub> H <sub>8</sub> O <sub>2</sub>	Methyl cyclobutanecarboxylate	114.14	15	-210.3	747.8	755.2	731.6	13	55-215
	C <sub>4</sub> H <sub>8</sub> O	Cyclobutyl methyl ketone	98.34	16	-260.9	807.1	802.7	793.6	15	55-215
						856.8	851.5	848.6	16	55-215
								69.0		
								3657.4		
								51,071.2		
								155.0		
								n = 2		
	C <sub>4</sub> H <sub>8</sub> O <sub>2</sub>	1,1-Cyclobutanedicarboxylic acid (a)	114.12	12	-133.2	612.0	616.0			55-187
	C <sub>4</sub> H <sub>8</sub> O <sub>2</sub>	cis-1,2-Cyclobutanedicarboxylic acid (a)	"	"	"	"	"			55-187
	C <sub>4</sub> H <sub>8</sub> O <sub>2</sub>	1,1,2-Cyclobutanedicarboxylic acid (a)	"	"	"	"	"			55-215
	C <sub>4</sub> H <sub>10</sub>	Methylcyclobutane	70.13	15	-342.2	639.2	706.2	799.9		55-215

TABLE 67  
NORMAL ALKYL CYCLOPENTANES (liquids)

ADL NO	FORMULA	NAME	MOL. WT.	OXYGEN REQUIRED	OXYGEN BALANCE	O <sub>c</sub> KCAL/MOLE		Y'	X'	REF.
						OBSERVED	CALCULATED			
	C <sub>5</sub> H <sub>10</sub>	Cyclopentane	70.13	15	-312.2	765.5	765.7	780.8	15	115-3
	C <sub>5</sub> H <sub>12</sub>	Methylcyclopentane	86.16	18	"	911.1	941.5	915.4	18	115-3
	C <sub>5</sub> H <sub>14</sub>	Ethylcyclopentane	98.18	21	"	1097.5	1097.7	1091.8	21	115-3
	C <sub>5</sub> H <sub>16</sub>	n-Propylcyclopentane	112.21	24	"	1251.7	1254.0	1246.0	24	115-3
	C <sub>5</sub> H <sub>18</sub>	n-Butylcyclopentane	126.23	27	"	1410.1	1410.2	1410.1	27	115-3
	C <sub>5</sub> H <sub>20</sub>	n-Pentylcyclopentane	140.26	30	"	1566.5	1566.4	1560.7	30	115-3
	C <sub>5</sub> H <sub>22</sub>	n-Hexylcyclopentane	154.29	33	"	1722.7	1722.7	1717.0	33	115-3
	C <sub>5</sub> H <sub>24</sub>	n-Heptylcyclopentane	168.31	36	"	1878.9	1878.2	1871.2	36	115-3
	C <sub>5</sub> H <sub>26</sub>	n-Octylcyclopentane	182.34	39	"	2035.2	2035.2	2028.5	39	115-3
	C <sub>5</sub> H <sub>28</sub>	n-Nonylcyclopentane	196.37	42	"	2191.4	2191.4	2185.7	42	115-3
	C <sub>5</sub> H <sub>30</sub>	n-Decylcyclopentane	210.39	45	"	2347.6	2347.6	2341.9	45	115-3
	C <sub>5</sub> H <sub>32</sub>	n-Undecylcyclopentane	224.42	48	"	2503.8	2503.9	2498.1	48	115-3
	C <sub>5</sub> H <sub>34</sub>	n-Dodecylcyclopentane	238.44	51	"	2660.1	2660.1	2654.5	51	115-3
	C <sub>5</sub> H <sub>36</sub>	n-Tridecylcyclopentane	252.47	54	"	2816.3	2816.1	2810.6	54	115-3
	C <sub>5</sub> H <sub>38</sub>	n-Tetradecylcyclopentane	266.49	57	"	2972.5	2972.6	2966.8	57	115-3
	C <sub>5</sub> H <sub>40</sub>	n-Pentadecylcyclopentane	280.52	60	"	3128.6	3128.3	3123.1	60	115-3

$y' = -1.67 + 52.02x'$

\*Corrected to liquid from gas values given in reference 15 using heat of vaporization = 60.3 cal/g at 25°C in Table 65.

$z' = 660.0$   
 $y' = 31,221.5$   
 $z' = 1,330,154.7$   
 $n = 2,560.0$   
 $n = 16$

TABLE 66  
NORMAL ALKYL CYCLOHEXANES (11q-18)

ADL NO.	FORMULA	NAME	MOL. WT.	OXYGEN REQUIRED	OXYGEN BALANCE	O <sub>2</sub> KCAL/MOLE		Y'	X'	REF.
						OBSERVED	CALCULATED			
	C <sub>6</sub> H <sub>12</sub>	Cyclohexane	84.16	10	-142.2	944.9	945.7	944.2	15	115-4
	C <sub>8</sub> H <sub>16</sub>	Methylcyclohexane	98.18	21	"	1001.1	1002.9	1006.4	21	115-4
	C <sub>10</sub> H <sub>16</sub>	Ethylcyclohexane	112.19	24	"	1124.1	1124.9	1141.5	24	115-4
	C <sub>12</sub> H <sub>18</sub>	n-Propylcyclohexane	126.19	27	"	1204.3	1204.4	1198.6	27	115-4
	C <sub>14</sub> H <sub>20</sub>	n-Butylcyclohexane	140.26	30	"	1363.6	1363.6	1355.1	30	115-4
	C <sub>16</sub> H <sub>22</sub>	n-Pentylcyclohexane	154.29	33	"	1476.9	1476.9	1471.2	33	115-4
	C <sub>18</sub> H <sub>24</sub>	n-Hexylcyclohexane	168.31	36	"	1573.14	1573.1	1587.4	36	115-4
	C <sub>20</sub> H <sub>26</sub>	n-Heptylcyclohexane	182.34	39	"	2027.14	2027.4	2024.7	39	115-4
	C <sub>22</sub> H <sub>28</sub>	n-Octylcyclohexane	196.36	42	"	2185.6	2185.6	2177.9	42	115-4
	C <sub>24</sub> H <sub>30</sub>	n-Nonylcyclohexane	210.39	45	"	2181.8	2181.8	2136.1	45	115-4
	C <sub>26</sub> H <sub>32</sub>	n-Decylcyclohexane	224.42	48	"	2108.0	2108.1	2192.3	48	115-4
	C <sub>28</sub> H <sub>34</sub>	n-Undecylcyclohexane	238.44	51	"	2444.2	2444.7	2404.5	51	115-4
	C <sub>30</sub> H <sub>36</sub>	n-Dodecylcyclohexane	252.47	54	"	2910.5	2910.5	2844.8	54	115-4
	C <sub>32</sub> H <sub>38</sub>	n-Tridecylcyclohexane	266.49	57	"	2966.7	2966.8	2961.0	57	115-4
	C <sub>34</sub> H <sub>40</sub>	n-Tetradecylcyclohexane	280.52	60	"	3123.0	3123.0	3117.3	60	115-4

$y' = -7.43 + 52.0x'$   
 $\Delta x' = 505.0$   
 $\Delta y' = 30,335.0$   
 $\Delta x'y' = 1,315,068.70$   
 $\Delta x'^2 = 78,335.0$   
 $n = 15$

Corrected to liquid from gas values given in Reference 115 by subtracting 0.6 cal/g. at 16 mmHg. 61.

TABLE 69  
CYCLOHEPTANES (11q-16)

ADL NO.	FORMULA	NAME	MOL. WT.	OXYGEN REQUIRED	OXYGEN BALANCE	O <sub>2</sub> KCAL/MOLE		Y'	X'	REF.
						OBSERVED	CALCULATED			
	C <sub>7</sub> H <sub>14</sub> O	Cycloheptanone	112.17	19	-271.0	996.7	996.1	992.1	19	55-215
	C <sub>7</sub> H <sub>14</sub>	Cycloheptane	96.17	20	-332.7	1049.9	1049.6	1049.2	20	55-215
	C <sub>7</sub> H <sub>14</sub> O	Cycloheptanol	114.18	"	-280.3	1050.2	1049.3	1049.8	20	55-215
	C <sub>7</sub> H <sub>14</sub>	Cycloheptane	96.11	21	-312.5	1099.1	1095.7	1093.4	21	129
	C <sub>7</sub> H <sub>14</sub> O <sub>2</sub>	Cycloheptanecarboxylic acid (s)	142.19	"	-236.3	1047.9	1091.7	1246.2	21	55-215
	C <sub>7</sub> H <sub>14</sub> O	Methylcycloheptanone	112.21	24	-348.2	1214.5	1249.0	1248.8	24	55-215
	C <sub>7</sub> H <sub>14</sub> O	Cycloheptyl methyl carbinol	142.32	26.5	-292.3	1342.2	1344.2	1343.0	26	55-215
	C <sub>7</sub> H <sub>14</sub>	Ethylcycloheptane	126.23	27	-342.2	1436.8	1402.2	1401.1	27	55-215

$y' = 17.12 + 51.05x'$   
 $\Delta x' = 178.0$   
 $\Delta y' = 4911.0$   
 $\Delta x'y' = 208,613.0$   
 $\Delta x'^2 = 1002.0$   
 $n = 8$

C <sub>7</sub> H <sub>14</sub> O	Cycloheptyl methyl ketone	140.22	25	-255.3	1276.3	1342.8	-	-	-	55-215
----------------------------------	---------------------------	--------	----	--------	--------	--------	---	---	---	--------

TABLE 70  
OTHER SATURATED ALICYCLIC HYDROCARBONS

ADL NO.	FORMULA	NAME	MOL. WT.	OXYGEN REQUIRED	OXYGEN BALANCE	O <sub>2</sub> KCAL/MOLE		Y'	X'	REF.
						OBSERVED	CALCULATED			
	C <sub>7</sub> H <sub>14</sub> O <sub>2</sub>	Cycloheptanecarboxylic acid	142.29	21	-236.3	1069.7 (7)				55-215
	C <sub>7</sub> H <sub>14</sub>	Cycloheptane	(1)							
	C <sub>7</sub> H <sub>14</sub>	Cycloheptane	112.21	24	-342.7	1250.4				129
	C <sub>7</sub> H <sub>14</sub> O	Cycloheptanol	114.18	30	"	1570.8				46
	C <sub>7</sub> H <sub>14</sub> O	Cycloheptanone	(2)			2167				120
	C <sub>7</sub> H <sub>14</sub> O	Cycloheptanone	(3)			2678				120
	C <sub>7</sub> H <sub>14</sub> O	Cycloheptanone	(4)			4674				120

TABLE 71  
 NORMAL OLDFINS (liquid)

ADL NO.	FORMULA	NAME	MOL. WT.	OXYGEN REQ (lb/lb-H)	OXYGEN BALANCE	OF, KCAL/M-M		Y'	X'	HEF.
						OBTAINED*	CALCULATED			
	C <sub>3</sub> H <sub>6</sub>	Propene	42.08	9	-342.2	285.4	108.7	103.7	9	115-9
	C <sub>4</sub> H <sub>10</sub>	1-Butane	58.12	12	"	444.9	642.7	639.7	12	115-9
	C <sub>5</sub> H <sub>12</sub>	1-Pentane	70.13	15	"	604.8	894.9	794.1	15	115-9
	C <sub>6</sub> H <sub>14</sub>	1-Hexane	84.16	18	"	857.0	1251.1	951.3	18	115-9
	C <sub>7</sub> H <sub>16</sub>	1-Heptane	98.18	21	"	1113.2	1611.3	1197.5	21	115-9
	C <sub>8</sub> H <sub>18</sub>	1-Octane	112.21	24	"	1269.6	1767.6	1263.7	24	115-9
	C <sub>9</sub> H <sub>20</sub>	1-Nonane	126.23	27	"	1525.7	2125.8	1423.0	27	115-9
	C <sub>10</sub> H <sub>22</sub>	1-Decane	140.26	30	"	1781.8	2582.0	1579.2	30	115-9
	C <sub>11</sub> H <sub>24</sub>	1-Undecane	154.29	33	"	1731.1	2738.2	1734.4	33	115-9
	C <sub>12</sub> H <sub>26</sub>	1-Dodecane	168.31	36	"	1891.4	1891.4	1883.7	36	115-9
	C <sub>13</sub> H <sub>28</sub>	1-Tridecane	182.34	39	"	2050.6	2050.6	2043.9	39	115-9
	C <sub>14</sub> H <sub>30</sub>	1-Tetradecane	196.36	42	"	2209.8	2209.8	2201.1	42	115-9
	C <sub>15</sub> H <sub>32</sub>	1-Pentadecane	210.39	45	"	2369.1	2369.1	2357.4	45	115-9
	C <sub>16</sub> H <sub>34</sub>	1-Hexadecane	224.42	48	"	2529.3	2529.3	2513.6	48	115-9
	C <sub>17</sub> H <sub>36</sub>	1-Heptadecane	238.44	51	"	2679.6	2679.6	2669.9	51	115-9
	C <sub>18</sub> H <sub>38</sub>	1-Octadecane	252.47	54	"	2831.7	2831.7	2822.7	54	115-9
	C <sub>19</sub> H <sub>40</sub>	1-Nonadecane	266.49	57	"	2982.0	2982.0	2977.8	57	115-9
	C <sub>20</sub> H <sub>42</sub>	1-Eicosane	280.52	60	"	3134.2	3134.2	3138.5	60	115-9
		$y' = 14.20 + S. 01a'$								
								$1y' = 37,550.6$		
								$1x' = 1,351,477.60$		
								$1z' = -25,785.0$		
								$n = 18$		
	C <sub>2</sub> H <sub>6</sub>	Ethane	30.05	6	-342.2	-	332.3	-	-	115-9
		*Corrected to liquid state from gas values given in Reference 115 by subtracting 86.3 cal/g as in Table 6j.								

 TABLE 72  
 NORMAL OLDFINS (gas)

ADL NO.	FORMULA	NAME	MOL. WT.	OXYGEN REQ (lb/lb-H)	OXYGEN BALANCE	OF, KCAL/M-M		Y'	X'	HEF.
						OBTAINED*	CALCULATED			
	C <sub>3</sub> H <sub>6</sub>	Propene	42.08	9	-342.2	161.99	491.97	186.49		115-9
	C <sub>4</sub> H <sub>10</sub>	1-Butane	58.12	12	"	219.74	619.74	614.74		115-9
	C <sub>5</sub> H <sub>12</sub>	1-Pentane	70.13	15	"	396.85	876.85	801.95		115-9
	C <sub>6</sub> H <sub>14</sub>	1-Hexane	84.16	18	"	564.96	964.96	954.74		115-9
	C <sub>7</sub> H <sub>16</sub>	1-Heptane	98.18	21	"	721.69	1121.73	1116.19		115-9
	C <sub>8</sub> H <sub>18</sub>	1-Octane	112.21	24	"	879.13	1279.17	1273.63		115-9
	C <sub>9</sub> H <sub>20</sub>	1-Nonane	126.23	27	"	1036.58	1436.61	1431.08		115-9
	C <sub>10</sub> H <sub>22</sub>	1-Decane	140.26	30	"	1194.92	1594.95	1588.52		115-9
	C <sub>11</sub> H <sub>24</sub>	1-Undecane	154.29	33	"	1351.46	1751.49	1745.95		115-9
	C <sub>12</sub> H <sub>26</sub>	1-Dodecane	168.31	36	"	1509.91	1949.93	1943.40		115-9
	C <sub>13</sub> H <sub>28</sub>	1-Tridecane	182.34	39	"	1668.35	2148.37	2140.85		115-9
	C <sub>14</sub> H <sub>30</sub>	1-Tetradecane	196.36	42	"	1827.77	2347.31	2341.29		115-9
	C <sub>15</sub> H <sub>32</sub>	1-Pentadecane	210.39	45	"	1987.23	2547.25	2541.71		115-9
	C <sub>16</sub> H <sub>34</sub>	1-Hexadecane	224.42	48	"	2146.64	2747.69	2741.18		115-9
	C <sub>17</sub> H <sub>36</sub>	1-Heptadecane	238.44	51	"	2306.13	2947.63	2941.61		115-9
	C <sub>18</sub> H <sub>38</sub>	1-Octadecane	252.47	54	"	2465.57	3147.57	3141.97		115-9
	C <sub>19</sub> H <sub>40</sub>	1-Nonadecane	266.49	57	"	3011.91	3011.91	3005.51		115-9
	C <sub>20</sub> H <sub>42</sub>	1-Eicosane	280.52	60	"	3168.45	3168.45	3162.95		115-9
		$y' = 14.15 + S2. 01a'$								
								$1y' = 671.0$		
								$1x' = 1,301,977.60$		
								$1z' = -25,785.0$		
								$n = 18$		
	C <sub>2</sub> H <sub>6</sub>	Ethane	30.05	6	-342.2	337.2	334.5	-	-	115-9

TABLE 73  
ANALYSIS OF PMS (Liquid)

API NO.	FORMULA	NAME	MOLE WT	OXYGEN REQUIRED	OXYGEN BALANCE	O <sub>2</sub> KCAL/MOLE		Y	X'	REF.
						OBSERVED*	CALCULATED			
C <sub>4</sub> H <sub>8</sub>		cis-2-Butene	56.10	12	-162.2	641.2	641.2			115-5
		trans-2-Butene				642.2	642.2			115-5
C <sub>5</sub> H <sub>10</sub>		2-Methylpropane	70.13	15		611.2	611.2			115-5
		cis-2-Pentene				797.4	797.4			115-5
		trans-2-Pentene				795.4	795.4			115-5
		2-Methyl-1-butene				795.2	795.2			115-5
		1-Methyl-1-butene				795.8	795.8			115-5
C <sub>6</sub> H <sub>12</sub>		2-Methyl-2-butene	84.16	18		955.6	955.6			115-5
		cis-2-Hexene				954.6	954.6			115-5
		trans-2-Hexene				955.6	955.6			115-5
		cis-1-Hexene				954.6	954.6			115-5
		trans-1-Hexene				953.6	953.6			115-5
		2-Methyl-1-pentene				956.7	956.7			115-5
		3-Methyl-1-pentene				956.7	956.7			115-5
		4-Methyl-1-pentene				952.7	952.7			115-5
		2-Methyl-2-pentene				952.8	952.8			115-5
		cis-1-Methyl-2-pentene				952.8	952.8			115-5
		trans-3-Methyl-2-pentene				953.7	953.7			115-5
		cis-1-Pentyl-2-pentene				952.9	952.9			115-5
		trans-4-Methyl-2-pentene				954.2	954.2			115-5
		2-Ethyl-1-butene				954.3	954.3			115-5
2,3-Dimethyl-1-butene	954.9	954.9	115-5							
C <sub>7</sub> H <sub>14</sub>		1,1-Dimethyl-1-butene	96.18	21		1112.0	1111.6			115-5
		2,1-Dimethyl-2-butene				1111.0	1111.0			115-5
		cis-2-Heptene				1111.0	1111.0			115-5
		trans-2-Heptene				1112.0	1112.0			115-5
		cis-1-Heptene				1111.0	1111.0			115-5
		trans-1-Heptene				1110.0	1110.0			115-5
		2-Methyl-1-hexene				1110.0	1110.0			115-5
		3-Methyl-1-hexene				1112.5	1112.5			115-5
		4-Methyl-1-hexene				1112.5	1112.5			115-5
		5-Methyl-1-hexene				1109.6	1109.6			115-5
		2-Methyl-2-hexene				1109.7	1109.7			115-5
		cis-3-Methyl-2-hexene				1109.7	1109.7			115-5
		cis-1-Methyl-2-hexene				1110.9	1110.9			115-5
		trans-1-Methyl-2-hexene				1110.3	1110.3			115-5
		trans-2-Methyl-2-hexene				1109.3	1109.3			115-5
		cis-2-Methyl-3-hexene				1110.3	1110.3			115-5
		trans-2-Methyl-3-hexene				1109.2	1109.2			115-5
		cis-3-Methyl-3-hexene				1109.2	1109.2			115-5
		trans-3-Methyl-3-hexene				1109.2	1109.2			115-5
		2-Ethyl-1-pentene				1109.2	1109.2			115-5
		1-Ethyl-1-pentene				1107.6	1107.6			115-5
		2,3-Dimethyl-1-pentene				1111.1	1111.1			115-5
2,1-Dimethyl-1-pentene	1109.0	1109.0	115-5							
3,1-Dimethyl-1-pentene	1106.3	1106.3	115-5							
3,2-Dimethyl-1-pentene	1110.3	1110.3	115-5							
1,4-Dimethyl-1-pentene	1110.8	1110.8	115-5							
1,4-Dimethyl-1-pentene	1109.2	1109.2	115-5							
3-Ethyl-2-pentene	1109.2	1109.2	115-5							
2,3-Dimethyl-2-pentene	1107.4	1107.4	115-5							
2,4-Dimethyl-2-pentene	1106.9	1106.9	115-5							
cis-1,4-Dimethyl-2-pentene	1107.6	1107.6	115-5							
trans-3,4-Dimethyl-2-pentene	1107.6	1107.6	115-5							
cis-1,4-Dimethyl-2-pentene	1107.6	1107.6	115-5							
trans-1,4-Dimethyl-2-pentene	1107.6	1107.6	115-5							
2-Methyl-2-ethyl-1-butene	1106.6	1106.6	115-5							
2,3,1-Trimethyl-1-butene	1109.6	1109.6	115-5							
C <sub>7</sub> H <sub>16</sub>		Di-n-butylsulfide	112.21	24		1265.4	1265.0			11
C <sub>7</sub> H <sub>16</sub>		Diisobutylene	110.26	30		1502.2	1501.0			55-65

\*For branched olefins:  $y = 16.2 + 52.2x$

\*Corrected to liquid state from gas values given in Reference 115 by subtracting 56.3 cal/g as in Table 6).



TABLE 74  
DIOLIFINS (liquid)

ADL NO.	FORMULA	NAME	MOL WT	OXYGEN REQUIRED	OXYGEN BALANCE	O <sub>2</sub> KCAL/MOLE		Y'	X'	REF.
						OBSERVED	CALCULATED			
	C <sub>8</sub> H <sub>16</sub>	1,2-Butadiene	54.09	11	-125.4	615.6	606.9			101
	C <sub>8</sub> H <sub>16</sub>	1,3-butadiene	54.09	11	-125.4	604.0	606.9			106
	C <sub>8</sub> H <sub>16</sub>	Isoprene	68.11	15	-128.9	755.3	755.9			51
	C <sub>10</sub> H <sub>18</sub>	1,5-Hexadiene	68.14	17	-111.1	916.4	916.9			27
		2,3-Dimethyl-2,3-butadiene				926.0				61
						911.7	912.4			61
		for n-diolifins: $y' = 38.1 + 52.0x'$ for branched diolifins: $x' = 15.7 + 52.16x'$								

TABLE 75  
ACETYLENES (liquid)

ADL NO.	FORMULA	NAME	MOL WT	OXYGEN REQUIRED	OXYGEN BALANCE	O <sub>2</sub> KCAL/MOLE		Y'	X'	REF.
						OBSERVED	CALCULATED			
	C <sub>2</sub> H <sub>2</sub>	Propyne	40.06	6	-219.5	479.7	469.6	156.0	8	115-6
	C <sub>3</sub> H <sub>4</sub>	1-Butyne	54.09	11	-125.4	616.2	616.9	616.5	11	115-6
	C <sub>4</sub> H <sub>6</sub>	1-Pentyne	68.11	14	-123.0	772.7	772.1	766.5	14	115-6
	C <sub>5</sub> H <sub>8</sub>	1-Hexyne	82.14	17	-111.1	928.4	923.4	922.7	17	115-6
	C <sub>6</sub> H <sub>10</sub>	1-Heptyne	96.17	20	-112.7	1084.6	1064.6	1075.9	20	115-6
	C <sub>7</sub> H <sub>12</sub>	1-Octyne	110.19	23	-114.0	1240.4	1210.8	1235.1	23	115-6
	C <sub>8</sub> H <sub>14</sub>	1-Nonyne	124.22	26	-111.9	1397.0	1377.1	1391.3	26	115-6
	C <sub>9</sub> H <sub>16</sub>	1-Decyne	138.24	29	-115.5	1553.3	1532.3	1547.6	29	115-6
	C <sub>10</sub> H <sub>18</sub>	1-Undecyne	152.27	32	-116.1	1709.7	1689.7	1705.2	32	115-6
	C <sub>11</sub> H <sub>20</sub>	1-Dodecyne	166.30	35	-116.7	1866.1	1846.1	1861.1	35	115-6
	C <sub>12</sub> H <sub>22</sub>	1-Tridecyne	180.32	38	-117.2	2022.9	2002.0	2016.3	38	115-6
	C <sub>13</sub> H <sub>24</sub>	1-Tetradecyne	194.35	41	-117.5	2179.2	2158.2	2172.5	41	115-6
	C <sub>14</sub> H <sub>26</sub>	1-Pentadecyne	208.37	44	-117.7	2335.7	2314.7	2329.0	44	115-6
	C <sub>15</sub> H <sub>28</sub>	1-Hexadecyne	222.40	47	-113.1	2492.7	2470.3	2485.0	47	115-6
	C <sub>16</sub> H <sub>30</sub>	1-Heptadecyne	236.43	50	-118.4	2649.4	2627.0	2641.2	50	115-6
	C <sub>17</sub> H <sub>32</sub>	1-Octadecyne	250.45	53	-118.6	2806.3	2783.2	2797.5	53	115-6
	C <sub>18</sub> H <sub>34</sub>	1-Nonadecyne	264.48	56	-115.6	2963.4	2939.9	2954.7	56	115-6
	C <sub>19</sub> H <sub>36</sub>	1-Eicosenyne	278.50	59	-119.0	3120.6	3096.7	3111.7	59	115-6
		$y' = 37.32 + 52.02x'$								
	C <sub>2</sub> H <sub>2</sub>	Acetylene	26.04	5	-107.2	-	303.4	-	-	-
		*Calculated from gaseous data given in reference 115 by subtracting 86.3 cal/g in Table 63.								
								$2x' = 603.0$ $3y' = 32,075.4$ $12'y' = 1,361,661.9$ $12x'^2 = 24,362.0$ $n = 18$		

TABLE 76  
NORMAL ALKYL BENZENES (continued)

AQL NO.	FORMULA	NAME	MOL. WT.	OXYGEN REQUIRED	OXYGEN BALANCE	G., KCAL/MOLE		Y'	X'	REF.
						CONSERVED	CALCULATED			
	C <sub>6</sub> H <sub>6</sub>	Benzene	78.11	15	-307.3	781.0	777.6			115-7
	C <sub>8</sub> H <sub>10</sub>	Ethylbenzene	106.16	21	-316.5	1061.0	1056.5	18		115-7
	C <sub>9</sub> H <sub>12</sub>	n-Propylbenzene	126.19	24	-319.5	1261.7	1256.2	21		115-7
	C <sub>10</sub> H <sub>14</sub>	n-Butylbenzene	136.21	27	-321.9	1463.4	1457.9	24		115-7
	C <sub>11</sub> H <sub>16</sub>	n-Pentylbenzene	146.24	30	-324.8	1665.9	1660.4	27		115-7
	C <sub>12</sub> H <sub>18</sub>	n-Hexylbenzene	156.26	33	-325.8	1871.4	1865.9	30		115-7
	C <sub>13</sub> H <sub>20</sub>	n-Heptylbenzene	176.29	36	-325.7	2071.9	2066.4	33		115-7
	C <sub>14</sub> H <sub>22</sub>	n-Octylbenzene	196.32	39	-327.9	2272.4	2266.9	36		115-7
	C <sub>15</sub> H <sub>24</sub>	n-Nonylbenzene	206.34	42	-325.9	2483.9	2478.4	39		115-7
	C <sub>16</sub> H <sub>26</sub>	n-Decylbenzene	216.37	45	-329.7	2684.4	2678.9	42		115-7
	C <sub>17</sub> H <sub>28</sub>	n-Undecylbenzene	236.40	48	-330.5	2884.9	2879.4	45		115-7
	C <sub>18</sub> H <sub>30</sub>	n-Dodecylbenzene	256.42	51	-331.1	3085.4	3079.9	48		115-7
	C <sub>19</sub> H <sub>32</sub>	n-Tridecylbenzene	266.45	54	-331.7	3285.9	3280.4	51		115-7
	C <sub>20</sub> H <sub>34</sub>	n-Tetradecylbenzene	276.47	57	-332.2	3486.4	3480.9	54		115-7
	C <sub>21</sub> H <sub>36</sub>	n-Pentadecylbenzene	286.50	60	-332.8	3686.9	3681.4	57		115-7
	C <sub>22</sub> H <sub>38</sub>	n-Hexadecylbenzene	306.52	63	-333.2	3887.4	3881.9	60		115-7
								649.0		
								33,630.0		
								1,521,627.0		
								29,398.0		
								n = 16		
	C <sub>6</sub> H <sub>6</sub>	Benzene	78.11	15	-307.3	781.0	777.6			115-7
	C <sub>8</sub> H <sub>10</sub>	Diphenylmethane	146.22	12	-334.3	1451.2	1446.7			68
	C <sub>10</sub> H <sub>14</sub>	1,1-Diphenylethane	172.28	35	-307.3	1806.1	1810.7			27
	C <sub>10</sub> H <sub>14</sub>	1,2-Diphenylpropane	166.28	35	-309.8	1770.7	1807.1			154
	C <sub>10</sub> H <sub>14</sub>	1,3-Diphenylpropane	166.28	35	-309.8	1770.7	1807.1			154
	C <sub>12</sub> H <sub>18</sub>	2,2-Diphenylbutane	210.30	41	-311.9	2127.6	2123.0			154

\* Calculated by subtracting heat of vaporization from gas values given in reference 115 and in Table 75.

TABLE 77  
NORMAL ALKYL BENZENES (ms)

AQL NO.	FORMULA	NAME	MOL. WT.	OXYGEN REQUIRED	OXYGEN BALANCE	G., KCAL/MOLE		Y'	X'	REF.
						CONSERVED	CALCULATED			
	C <sub>6</sub> H <sub>6</sub>	Benzene	78.11	15	-307.3	789.00	755.7			115-7
	C <sub>8</sub> H <sub>10</sub>	Toluene	92.12	18	-312.0	945.85	913.1			115-7
	C <sub>9</sub> H <sub>12</sub>	Ethylbenzene	106.16	21	-316.5	1101.13	1104.5			115-7
	C <sub>10</sub> H <sub>14</sub>	n-Propylbenzene	126.19	24	-319.5	1258.24	1258.0			115-7
	C <sub>10</sub> H <sub>14</sub>	n-Butylbenzene	136.21	27	-321.9	1415.34	1415.5			115-7
								558.0		
								29,159.84		
								1,425,335.61		
								27,234.0		
								n = 12		

TABLE 75

## NORMAL ALKYL BENZENES - HEAT OF VAPORIZATION

ADL NO.	FORMULA	NAME	MOL WT.	OXYGEN REQUIRED	OXYGEN BALANCE	G.P. KCAL/MOLE		V <sup>1</sup>	K <sup>1</sup>	REF.
						OBSERVED	CALCULATED			
	C <sub>6</sub> H <sub>6</sub>	Benzene	78	15	-	8,162	8,118			11-7
	C <sub>8</sub> H <sub>10</sub>	Toluene	92	18	-	9,631	9,096			11-7
	C <sub>10</sub> H <sub>14</sub>	Ethylbenzene	106	21	-	10,287	10,562			11-7
	C <sub>12</sub> H <sub>18</sub>	n-Propylbenzene	120	24	-	11,033	11,016			11-7
	C <sub>14</sub> H <sub>20</sub>	n-Butylbenzene	134	27	-	11,779	12,006			11-7
		$y = 3.248 + 0.324x$				$1x = 10,510$				
						$1y = 50,310$				
						$1xy = 1465,6230$				
						$1x^2 = 225,100$				
						$n = 5$				
	C <sub>16</sub> H <sub>22</sub>	n-Hexylbenzene	148	30	-	-	12,973			11-7
	C <sub>18</sub> H <sub>26</sub>	n-Heptylbenzene	162	33	-	-	13,750			11-7
	C <sub>20</sub> H <sub>30</sub>	n-Octylbenzene	176	36	-	-	14,592			11-7
	C <sub>22</sub> H <sub>34</sub>	n-Nonylbenzene	190	39	-	-	15,491			11-7
	C <sub>24</sub> H <sub>38</sub>	n-Decylbenzene	204	42	-	-	16,446			11-7
	C <sub>26</sub> H <sub>42</sub>	n-Undecylbenzene	218	45	-	-	17,453			11-7
	C <sub>28</sub> H <sub>46</sub>	n-Dodecylbenzene	232	48	-	-	18,510			11-7
	C <sub>30</sub> H <sub>50</sub>	n-Tridecylbenzene	246	51	-	-	19,702			11-7
	C <sub>32</sub> H <sub>54</sub>	n-Tetradecylbenzene	260	54	-	-	20,954			11-7
	C <sub>34</sub> H <sub>58</sub>	n-Pentadecylbenzene	274	57	-	-	22,278			11-7
	C <sub>36</sub> H <sub>62</sub>	n-Hexadecylbenzene	288	60	-	-	23,683			11-7
	C <sub>38</sub> H <sub>66</sub>		302	63	-	-	25,170			11-7

TABLE 76

## BRANCHED ALKYL BENZENES (Liquid and Solid)

ADL NO.	FORMULA	NAME	MOL WT.	OXYGEN REQUIRED	OXYGEN BALANCE	G.P. KCAL/MOLE		V <sup>1</sup>	K <sup>1</sup>	REF.
						OBSERVED	CALCULATED			
	C <sub>8</sub> H <sub>10</sub>	o-Tolene (1)	92.15	21	-326.5	1055.2	1055.7			11-8
	"	m-Tolene (1)	"	"	"	1057.7	"			11-8
	"	p-Tolene (1)	"	"	"	1055.7	"			11-8
	C <sub>9</sub> H <sub>12</sub>	Mesitylene (3)	120.17	24	-319.1	1243.5	1243.6			11-8
	"	1-Propylbenzene (1)	"	"	"	1256.5	"			11-8
	"	1-Methyl-2-ethylbenzene (1)	"	"	"	1257.5	"			11-8
	"	1-Ethyl-2-ethylbenzene (1)	"	"	"	1255.7	"			11-8
	"	1-Methyl-3-ethylbenzene (1)	"	"	"	1246.5	"			11-8
	"	1,2,3-Trimethylbenzene (1)	"	"	"	1262.5	"			11-8
	"	1,2,4-Trimethylbenzene (1)	"	"	"	1251.5	"			11-8
	"	1,1,5-Trimethylbenzene (1)	"	"	"	1241.2	"			11-8
	C <sub>10</sub> H <sub>14</sub>	t-Butylbenzene (1)	134.21	27	-321.9	1329.5	1329.3			11-8
	"	1,2,3,4-Tetraethylbenzene (Isodurene) (1)	"	"	"	1357.8	"			11
	"	1,2,3,4-Tetraethylbenzene (Durene) (1)	"	"	"	1336.3	"			11
	"	1,2,3,4-Tetraethylbenzene (Tetraethyl) (1)	"	"	"	1372.8	"			11
	"	1,3-Diisopropylbenzene (1)	"	"	"	1375.2	"			11-8
	"	3,5-Diisopropylbenzene (1)	"	"	"	1379.5	"			11-8
	"	1,3-Diisopropylbenzene (1)	"	"	"	1367.5	"			11-8
	C <sub>11</sub> H <sub>16</sub>	1-Ethyl-2-propylbenzene (1)	152.25	30	-323.5	1428.7	1429.1			63
	C <sub>12</sub> H <sub>18</sub>	1-Ethyl-3-propylbenzene (1)	162.26	33	-324.9	1506.5	1511.8			63
		$y = -5.2 + 92.26x$ (From n-alkylbenzenes + branched paraffins)								

TABLE 90  
POLYAROMES (solid)

#DL NO.	FORMULA	NAME	MOLE WT	OXYGEN REQUIRED	OXYGEN BALANCE	G <sub>c</sub> - KCAL/MOLE		Y*	X*	REF.
						OBSERVED	CALCULATED			
<b>Diphenyls</b>										
C <sub>12</sub> H <sub>10</sub>	1,2-Diphenylmethane	182.25	38	-337.3	1627.5	1811.2	899.9	17.60		88
C <sub>12</sub> H <sub>12</sub>	1,2-Diphenylbutane	210.30	42	-311.9	2127.6	2120.4	1061.3	20.96		134
<b>Triphenyls</b>										
C <sub>18</sub> H <sub>14</sub>	Triphenylmethane	244.32	45	-401.2	2472.7	2372.5	76.8	11.31		86
C <sub>18</sub> H <sub>16</sub>	1,1,1-Triphenylethane	278.34	57	-353.5	2532.4	2530.1	82.2	15.33		27
	1,1,2-Triphenylethane				2520.5		81.1	16.11		27
<b>Tetra- and Penta-phenyls</b>										
C <sub>24</sub> H <sub>18</sub>	Tetraphenylmethane	308.36	57	-427.7	2921.5	2931.8	72.1	14.27		107
C <sub>24</sub> H <sub>20</sub>	Tetra-phenylmethane	320.41	60	-399.5	3021.3	3091.4	771.7	15.00		27
C <sub>24</sub> H <sub>22</sub>	1,1,1,1-Tetra-phenylethane	334.44	63	-361.4	3252.3	3267.0	811.7	15.75		27
	1,1,1,2-Tetra-phenylethane				3278.2		810.6	15.75		27
<b>Pentaphenyls</b>										
C <sub>30</sub> H <sub>22</sub>	1,1,1,1,2-Pentaphenylethane	410.51	77	-360.1	3987.4	3967.9	796.3	15.40		27
	$\gamma^* = -16.52 + 52.53x$						2x <sup>2</sup> = 162.14			
							2y <sup>2</sup> = 8,352.0			
							2xy <sup>2</sup> = 116,850.981			
							2x <sup>2</sup> y = 2,450,194.2			
							a = 16			
C <sub>24</sub> H <sub>22</sub>	1,1,1,2-Tetra-phenylethane	334.44	63	-361.4	3181	3267.0	-	-	-	152

TABLE 91  
BIPHENYLS (liquid)

#DL NO.	FORMULA	NAME	MOLE WT	OXYGEN REQUIRED	OXYGEN BALANCE	G <sub>c</sub> - KCAL/MOLE		Y*	X*	REF.
						OBSERVED	CALCULATED			
C <sub>12</sub> H <sub>10</sub>	2-Methylbiphenyl	164.23	32	-321.3	1657.5	1657.3	1754.2	12		16
	3-Methylbiphenyl	"	"	"	1654.4	"	1648.7	12		18
	4-Methylbiphenyl	"	"	"	1647.2	"	1641.5	12		18
C <sub>14</sub> H <sub>12</sub>	3,3'-Dimethylbiphenyl	182.25	35	-307.3	1801.2	1814.1	1795.7	15		16
	2,2'-Dimethylbiphenyl	"	"	"	1812.5	"	1806.9	15		15L
C <sub>14</sub> H <sub>14</sub>	2-n-Propylbiphenyl	194.8	38	-309.8	1974.7	1967.5	1755.0	13		15L
	$\gamma^* = 11.30 + 52.65x$						2x <sup>2</sup> = 204			
							2y <sup>2</sup> = 10,511.8			
							2xy <sup>2</sup> = 359,974.8			
							2x <sup>2</sup> y = 8966.0			
							a = 6			
C <sub>16</sub> H <sub>14</sub>	Biphenyl	(a) 154.20	29	-340.9	1675.0	1674.4				16
C <sub>16</sub> H <sub>16</sub>	4,4'-Dimethylbiphenyl	(a) 172.25	35	-327.3	1792.3	1810.1				18

TABLE 52  
NAPHTHALENES (solid)

ADL NO.	FORMULA	NAME	MOLE WT.	OXYGEN REQUIRED	OXYGEN BALANCE	Q <sub>v</sub> - KCAL/MOLE		-y'	x'	HEP
						OBSERVED	CALCULATED			
C <sub>10</sub> H <sub>8</sub> O	1-Naphthol	124.16	23.0	-275.3	1187.8	1164.7	1114.8	23.0	66	
		"	"	"	1185.4	"	1129.4	23.0	66	
C <sub>10</sub> H <sub>6</sub> O	2-Naphthol	"	"	"	1181.0	"	1124.0	23.0	66	
		"	"	"	1187.2	"	1161.2	23.0	66	
C <sub>10</sub> H <sub>6</sub>	Naphthalene	124.16	24.0	-299.6	1233.3	1235.4	1227.6	24.0	41	
		"	"	"	1234.5	"	1228.6	24.0	76	
		"	"	"	1232.1	"	1236.4	24.0	43	
		"	"	"	1212.5	"	1226.5	24.0	55-80	
		"	"	"	1231.3	"	1228.1	24.0	55-52	
		"	"	"	1227.9	"	1224.2	24.0	55-241	
		"	"	"	1221.6	"	1229.1	24.0	55-190	
		"	"	"	1224.9	"	1224.9	24.0	55-157	
C <sub>11</sub> H <sub>8</sub> O <sub>2</sub>	2-Naphthoic acid	172.17	"	-223.0	"	"	"	"	"	
C <sub>10</sub> H <sub>7</sub> N	-naphthylamine	143.18	24.5	-273.3	1253.5	1261.0	1258.9	24.5	55-99	
C <sub>10</sub> H <sub>7</sub> N	"naphthylamine	"	"	"	1261.0	"	1252.4	24.5	55-79	
C <sub>10</sub> H <sub>7</sub> N	"	"	"	"	1261.0	"	1255.3	24.5	153	
C <sub>10</sub> H <sub>12</sub> N <sub>2</sub> O <sub>2</sub>	2-Naphthylamine nitrate	206.20	"	-390.1	1410.7	1434.7	1257.4	24.5	153	
C <sub>11</sub> H <sub>8</sub> N	1-Naphthnitrile	153.17	25.5	-264.4	1274.2	1271.6	1211.5	25.5	55-105	
C <sub>11</sub> H <sub>8</sub> N	2-Naphthnitrile	"	"	"	1324.0	"	1291.1	25.5	55-152	
C <sub>11</sub> H <sub>12</sub>	2-Methylnaphthalene	142.19	27.0	-303.3	1318.9	1319.9	1273.2	27.0	107	
C <sub>11</sub> H <sub>10</sub> O <sub>2</sub>	1-Naphthyl acetate	185.20	"	-232.0	1362.1	1374.7	1371.6	27.0	66	
C <sub>11</sub> H <sub>10</sub> O <sub>2</sub>	2-Naphthyl acetate	"	"	"	1398.5	"	1368.0	27.0	66	
C <sub>16</sub> H <sub>12</sub> N	4-Imagyl-2-naphthylamine	219.27	38.5	-260.7	1367.1	1368.3	1275.1	38.5	111	
y' = -6.08 + 51.0x'								3x' = 552.5		
								2y' = 28,311.4		
								2x'y' = 722,311.5		
								7x'' = 14,658.75		
								n = 22		
C <sub>18</sub> H <sub>16</sub> O <sub>4</sub>	1,8-Naphthalenedicarboxylic acid	216.16	24.0	-177.6	1292.1	1277.7			55-127	
C <sub>18</sub> H <sub>12</sub> O <sub>4</sub>	2-Naphthyl benzoate	246.27	38.0	-244.9	1666.0	1623.9			7	
C <sub>18</sub> H <sub>16</sub> O <sub>2</sub>	2-Naphthol formal	190.14	28.0	-254.7	2593.5	2277.7			55-60	

TABLE 53  
PRIMARY ALCOHOLS (liquid)

ADL NO.	FORMULA	NAME	MOLE WT.	OXYGEN REQUIRED	OXYGEN BALANCE	Q <sub>v</sub> - KCAL/MOLE		-y'	x'	HEP
						OBSERVED	CALCULATED			
C <sub>2</sub> H <sub>6</sub> O	Methyl alcohol	32.06	3	-157.6	173.6	170.9	167.9	3	113	
C <sub>3</sub> H <sub>8</sub> O	Ethyl alcohol	60.07	6	-208.4	326.7	326.9	321.0	6	113	
C <sub>4</sub> H <sub>10</sub> O	n-Propyl alcohol	60.09	9	-239.6	382.2	383.0	376.5	9	113	
C <sub>5</sub> H <sub>12</sub> O	n-Butyl alcohol	74.12	12	-257.0	636.1	639.2	634.4	12	113	
C <sub>6</sub> H <sub>14</sub> O	n-Amyl alcohol	88.15	15	-272.3	794.1	797.1	788.6	15	113	
C <sub>7</sub> H <sub>16</sub> O	n-Hexyl alcohol	102.17	18	-273.9	950.6	951.3	944.9	18	113	
C <sub>8</sub> H <sub>18</sub> O	n-Heptyl alcohol	116.20	21	-299.2	1107.1	1107.4	1101.4	21	113	
C <sub>9</sub> H <sub>20</sub> O	n-Octyl alcohol	130.22	24	-294.5	1263.6	1263.5	1257.4	24	113	
C <sub>10</sub> H <sub>22</sub> O	n-Nonyl alcohol	144.25	27	-299.5	1419.6	1419.6	1414.0	27	113	
C <sub>10</sub> H <sub>22</sub> O	n-Decyl alcohol	158.28	30	-303.1	1576.9	1575.7	1571.2	0	113	
y' = 9.17 + 57.0x'								3x' = 165.0		
								2y' = 8676.3		
								3x'y' = 101,790.7		
								2x'' = 3455.0		
								n = 10		

TABLE 84  
PRIMARY ALCOHOLS (LIQ.)

ADL NO.	FORMULA	NAME	MOL. WT.	OXYGEN REQUIRED	OXYGEN BALANCE	Q <sub>v</sub> KCAL/MOLE		Y'	X'	REF.
						OBSERVED	CALCULATED			
	C <sub>1</sub> H <sub>3</sub> O	Methyl alcohol	32.06	3	-113.8	181.78	180.5	177.09	3	113
	C <sub>2</sub> H <sub>5</sub> O	Ethyl alcohol	46.07	6	-208.4	336.78	337.2	331.28	6	113
	C <sub>3</sub> H <sub>7</sub> O	n-Propyl alcohol	60.09	9	-239.6	471.29	469.7	467.70	9	113
	C <sub>4</sub> H <sub>9</sub> O	n-Butyl alcohol	74.12	12	-259.7	610.90	609.4	604.10	12	113
	C <sub>5</sub> H <sub>11</sub> O	n-Amyl alcohol	88.15	15	-272.1	806.75	807.4	801.25	15	113
	C <sub>6</sub> H <sub>13</sub> O	n-Hexyl alcohol	102.17	18	-281.9	961.60	961.4	956.10	18	113
	C <sub>7</sub> H <sub>15</sub> O	n-Heptyl alcohol	116.20	21	-289.7	1120.60	1120.2	1115.10	21	113
	C <sub>8</sub> H <sub>17</sub> O	n-Octyl alcohol	130.22	24	-294.5	1277.60	1277.5	1272.10	24	113
	C <sub>9</sub> H <sub>19</sub> O	n-Nonyl alcohol	144.25	27	-299.5	1434.60	1434.2	1429.10	27	113
	C <sub>10</sub> H <sub>21</sub> O	n-Decyl alcohol	158.28	30	-303.3	1591.60	1591.0	1586.10	30	113
								Y' = 165.0		
								Y' = 8002.71		
								X' = 184.021.77		
								Z' = 345.0		
								a = 10		

TABLE 85  
SECONDARY ALCOHOLS (LIQ.)

ADL NO.	FORMULA	NAME	MOL. WT.	OXYGEN REQUIRED	OXYGEN BALANCE	Q <sub>v</sub> KCAL/MOLE		Y'	X'	REF.
						OBSERVED	CALCULATED			
	C <sub>3</sub> H <sub>8</sub> O	1-Propyl alcohol	60.09	9	-239.6	471.8	471.0	469.1	9	87
	C <sub>4</sub> H <sub>10</sub> O	Cyclohexanol	100.15	17	-271.6	890.7	889.5	892.1	17	55-153
	C <sub>4</sub> H <sub>10</sub> O	2-Methylcyclopentanol	"	"	"	890.7	889.5	893.4	17	55-215
	C <sub>4</sub> H <sub>10</sub> O	2-Methylcyclohexanol	100.15	17	-271.6	890.7	889.5	892.9	17	55-215
	C <sub>5</sub> H <sub>12</sub> O	3-Methylcyclohexanol	114.18	20	-280.9	1030.4	1031.4	1026.1	20	55-215
	C <sub>5</sub> H <sub>12</sub> O	1-Ethyl-2-cyclopentanol	"	"	"	1030.4	"	1031.0	20	55-215
	C <sub>5</sub> H <sub>12</sub> O	1,3-Dimethyl-2-cyclopentanol	"	"	"	1030.4	"	1028.4	20	55-215
	C <sub>5</sub> H <sub>12</sub> O	1,3-Dimethyl-2-cyclohexanol	128.21	23	-287.0	1195.0	1190.5	1197.7	23	55-215
	C <sub>5</sub> H <sub>12</sub> O	1,3-Dimethyl-2-cyclohexanol	"	"	"	1183.4	"	1183.1	21	55-215
	C <sub>6</sub> H <sub>14</sub> O	1-Ethyl-1,2-cyclohexanol (Asymmetric propargyl alcohol)	202.28	36	-288.0	1901.1	1899.0	1865.7	36	55-118
								Z' = 202.0		
								Y' = 10,528.0		
								X' = 235,706.6		
								Z' = 4537.0		
								a = 10		

TABLE 86  
TERTIARY ALCOHOLS (LIQ.)

ADL NO.	FORMULA	NAME	MOL. WT.	OXYGEN REQUIRED	OXYGEN BALANCE	Q <sub>v</sub> KCAL/MOLE		Y'	X'	REF.
						OBSERVED	CALCULATED			
	C <sub>4</sub> H <sub>10</sub> O	t-Butyl alcohol	74.12	12	-259.0	629.3	627.7	621.6	12	55-215
	C <sub>5</sub> H <sub>12</sub> O	Dimethyl ethyl carbinol	88.15	15	-272.3	784.5	781.9	774.9	15	55-215
	C <sub>5</sub> H <sub>12</sub> O	Allyl dimethyl carbinol	100.16	17	-271.6	900.1	899.9	860.0	17	55-215
	C <sub>5</sub> H <sub>12</sub> O	Methyl diethyl carbinol	102.17	18	-281.9	927.0	918.6	921.3	18	55-215
	C <sub>5</sub> H <sub>12</sub> O	Triethyl carbinol	116.20	21	-289.2	1082.0	1080.0	1071.3	21	55-215
	C <sub>5</sub> H <sub>12</sub> O	Methyl dipropyl carbinol	130.22	24	-294.7	1237.7	1237.0	1227.0	24	55-215
	C <sub>5</sub> H <sub>12</sub> O	Diethyl isopropyl carbinol	116.20	21	-289.5	1198.5	1198.0	1182.8	21	55-215
	C <sub>5</sub> H <sub>12</sub> O	1,3-Dimethyl-3-cyclopentanol	114.18	20	-280.1	1031.0	1035.3	1029.9	20	55-215
	C <sub>5</sub> H <sub>12</sub> O	Allyl methyl ethyl carbinol	"	"	"	1050.1	1050.7	1030.4	20	55-215
	C <sub>5</sub> H <sub>12</sub> O	1,3-Dimethyl-2-cyclohexanol	128.21	23	-287.0	1176.5	1162.9	1193.2	23	55-163
						1172.5	"	1198.2	23	55-215
						1201.5	"	1182.2	23	55-215
						1207.1	"	1187.4	23	55-215
	C <sub>6</sub> H <sub>14</sub> O	1-Ethyl-1,3-cyclohexanol	144.25	26	-292.5	1322.4	1316.3	1324.1	26	55-215
						1324.7	"	1325.4	26	55-215
						1303.0	"	1311.4	26	55-215
	C <sub>7</sub> H <sub>16</sub> O	Allyl n-propyl carbinol	158.28	28	-290.5	1472.1	1468.3	1460.0	28	55-215
	C <sub>7</sub> H <sub>16</sub> O	Allyl ethyl heptyl carbinol	170.29	32	-290.7	1666.7	1664.4	1641.1	32	55-215
								Z' = 106.0		
								Y' = 20,817.1		
								X' = 425,706.00		
								Z' = 9160.0		
								a = 10		

TABLE 87  
MIL-D POLYALCOHOLS (liquid)

ADL NO.	FORMULA	NAME	MOL. WT.	OXYGEN REQUIRED	OXYGEN BALANCE	Q <sub>v</sub> KCAL/MOLE		Y'	X'	REF.
						OBSERVED	CALCULATED			
	C <sub>2</sub> H <sub>6</sub> O	Ethylene alcohol	62.07	5	-128.9	284.3	-283.5	139.2	2.50	88
		1,2-Propanediol	76.09	8	-168.2	436.7	434.5	224.5	4.00	82
		1,2-Butanediol	90.12	11	-195.3	593.8	572.2	294.1	5.50	82
		1,3-Butanediol	"	"	"	595.8	"	275.1	5.50	82
		2,3-Butanediol	"	"	"	592.5	"	291.9	5.50	82
		2-Methyl-1,3-propanediol	"	"	"	289.9	"	392.1	5.50	82
	C <sub>3</sub> H <sub>8</sub> O <sub>3</sub>	Glycerol	92.09	7	-181.5	395.5	396.7	130.0	2.33	88
		v' = 10.26 + 51.35a' for a single OH						Y <sub>1</sub> ' = 31.83 Y <sub>2</sub> ' = 1,658.0 Z <sub>1</sub> ' = 7,965.750 Z <sub>2</sub> ' = 149.6769 n = 7		

TABLE 88  
MIXED POLYALCOHOLS (solid)

ADL NO.	FORMULA	NAME	MOL. WT.	OXYGEN REQUIRED	OXYGEN BALANCE	Q <sub>v</sub> KCAL/MOLE		Y'	X'	REF.
						OBSERVED	CALCULATED			
	C <sub>6</sub> H <sub>13</sub> O <sub>6</sub>	Pentarythritol	136.15	12	-111.0	651.2	651.1	163.9	1.00	55-195
		Syccaritrol	122.12	9	-179.2	500.2	501.1	123.6	2.25	80
	C <sub>6</sub> H <sub>13</sub> O <sub>5</sub>	Arabitol	152.15	11	-115.7	611.8	611.6	121.2	2.20	55-195
	C <sub>6</sub> H <sub>11</sub> O <sub>5</sub>	Dulcitol	182.17	13	-114.2	720.5	722.1	119.2	2.17	88
		Mannitol	"	"	"	722.5	"	119.5	2.17	88
	C <sub>7</sub> H <sub>15</sub> O <sub>7</sub>	Glucosaccharol	214.20	15	-113.1	635.8	632.6	118.6	2.14	55-195
		v' = 3.8 + 53.35a' for a single OH						Z <sub>1</sub> ' = 13.9 Y <sub>1</sub> ' = 166.0 Z <sub>2</sub> ' = 1,504.723 Z <sub>3</sub> ' = 37.0999 n = 6		

TABLE 89  
AROMATIC HYDROXYS (solid)

ADL NO.	FORMULA	NAME	MOL. WT.	OXYGEN REQUIRED	OXYGEN BALANCE	Q <sub>v</sub> KCAL/MOLE		Y'	X'	REF.
						OBSERVED	CALCULATED			
	C <sub>2</sub> H <sub>6</sub> O	Ethanol	92.11	14	-219.0	730.6	721.7	731.0	14.0	9
	C <sub>9</sub> H <sub>9</sub> O <sub>2</sub>	n-Propylphenyl alcohol	124.13	16	-206.2	815.4	810.7	810.1	16.0	55-195
		m-Cresol	106.13	17	-251.5	313.5	314.2	656.7	17.0	9
	C <sub>8</sub> H <sub>10</sub> O	p-Cresol	"	"	"	295.0	"	686.2	17.0	12
		o-Cymenol	"	"	"	381.7	"	235.9	17.0	12
	C <sub>8</sub> H <sub>10</sub> O	p-Cymenol	122.16	20	-242.0	1018.1	1017.5	1040.0	20.0	55-201
		o-Cymenol	"	"	"	1037.4	"	1042.1	20.0	55-201
	C <sub>8</sub> H <sub>10</sub> O	p-Tolueneol	124.16	22	-270.2	1035.3	"	1040.2	20.0	55-201
		m-Tolueneol	114.16	"	-270.1	1185.4	1185.6	1198.0	21.0	55-201
	C <sub>8</sub> H <sub>10</sub> O	o-Tolueneol	"	"	"	1187.5	"	1200.4	23.0	66
		2-Methylphenol	"	"	"	1187.2	"	1197.8	23.0	55-225
	C <sub>8</sub> H <sub>10</sub> O	Thymol	150.21	26	-245.9	1340.7	1340.6	1153.4	24.0	55-191
		Pyrocatechol	110.11	13	-198.9	684.8	685.7	314.2	6.5	55-195
	C <sub>8</sub> H <sub>10</sub> O	Resorcinol	"	"	"	683.0	"	310.1	6.5	55-195
		Hydroquinone	"	"	"	683.2	"	314.4	6.5	55-225
	C <sub>10</sub> H <sub>10</sub> O <sub>2</sub>	2-Hydroxy-1-naphthol	124.13	16	-206.2	815.1	819.3	819.2	8.0	55-201
		Phenylhydroquinone	126.21	25	-240.7	1397.1	1334.7	657.6	12.5	55-225
	C <sub>14</sub> H <sub>10</sub> O <sub>2</sub>	2,2'-Methyl-4,4'-diphenol	231.29	36	-242.3	1844.1	1847.8	947.8	16.0	47
		1,1'-Methylenedi(2-naphthol)	202.31	28	-255.7	2195.2	2237.6	1254.1	24.0	55-20
	C <sub>8</sub> H <sub>8</sub> O <sub>2</sub>	Pyrogallol	126.11	12	-152.9	936.7	937.2	214.1	4.0	55-195
		v' = 51.79a'						Z <sub>1</sub> ' = 300.0 Y <sub>1</sub> ' = 19,212.5 Z <sub>2</sub> ' = 37,39.55 Z <sub>3</sub> ' = 11 n = 22		
	C <sub>12</sub> H <sub>16</sub> O	2-Phenylphenol	154.20	25	-253.2	1642.5	1113.2	-	-	14

TABLE 90  
ETHERS (liquid)

IDL NO.	FORMULA	NAME	MOL. WT.	OXYGEN REQUIRED	OXYGEN BALANCE	Q <sub>v</sub> - KCAL/MOLE		Y'	X'	REF.
						OBSERVED	CALCULATED			
	C <sub>2</sub> H <sub>6</sub> O	2-Methoxyethanol	76.09	8.0	-168.2	141.2	427.6	126.7	4.0	125
	C <sub>4</sub> H <sub>10</sub> O	Diethylene glycol	106.12	10.0	-159.8	554.7	549.0	58.4	10.0	82
	C <sub>4</sub> H <sub>10</sub> O	Diethyl ether	74.12	12.0	-257.0	641.7	647.6	64.0	12.0	55-20
	C <sub>5</sub> H <sub>12</sub> O	Anisole	106.13	12.0	-257.5	901.9	899.7	907.1	17.0	9
	C <sub>5</sub> H <sub>12</sub> O	Miscetole	122.15	20.0	-252.0	1054.4	1055.5	1059.4	20.0	9
	C <sub>5</sub> H <sub>12</sub> O	m-Cresol methyl ether	136.19	24.0	-270.2	1213.4	1211.7	1217.8	23.0	55-203
	C <sub>5</sub> H <sub>12</sub> O	p-Cresol methyl ether	136.19	24.0	-270.2	1213.4	1211.7	1217.8	23.0	55-203
	C <sub>5</sub> H <sub>12</sub> O	p-Cresol isopropyl ether	150.21	24.0	-276.9	1313.1	1315.9	1377.5	23.0	55-203
	C <sub>5</sub> H <sub>12</sub> O	Thymol methyl ether	164.24	24.0	-282.5	1524.3	1525.8	1578.8	29.0	55-203
	C <sub>5</sub> H <sub>12</sub> O	Thymol ethyl ether	178.26	32.0	-287.2	1679.9	1682.9	1682.9	32.0	55-203
	C <sub>6</sub> H <sub>14</sub> O	Diethylene glycol monomethyl ether	120.15	13.0	-173.1	720.2	721.8	751.0	6.5	125
	C <sub>6</sub> H <sub>14</sub> O	Triethylene glycol	151.19	15.5	-164.0	857.8	857.5	868.2	7.75	82
	C <sub>6</sub> H <sub>14</sub> O	Diglycol dimethyl ether	118.17	17.0	-220.2	775.0	784.1	837.5	8.5	55-172
	C <sub>8</sub> H <sub>18</sub> O	Tetraethylene glycol	196.24	21.0	-171.2	1114.9	1165.6	124.7	7.0	82
		$y' = 15.17 + 52.20x'$								
	C <sub>10</sub> H <sub>22</sub> O	Diampyl ether	158.28	10	-303.3	1503.3	1507.2	-	-	55-65

TABLE 91  
ALIPHATIC ETHERS (gas)

IDL NO.	FORMULA	NAME	MOL. WT.	OXYGEN REQUIRED	OXYGEN BALANCE	Q <sub>v</sub> - KCAL/MOLE		Y'	X'	REF.
						OBSERVED	CALCULATED			
	C <sub>2</sub> H <sub>6</sub> O	Dimethyl ether	60.09	6	-236.4	317.6	318.2	342.1	6	55-223
	C <sub>3</sub> H <sub>8</sub> O	Methyl ethyl ether	60.09	9	-239.6	501.4	509.5	497.9	9	55-223
	C <sub>3</sub> H <sub>8</sub> O	Methyl isopropyl ether	74.09	10	-248.3	600.8	595.2	558.0	10	55-223
	C <sub>4</sub> H <sub>10</sub> O	Methyl allyl ether	74.10	11	-254.1	821.8	824.5	604.3	11	55-223
	C <sub>4</sub> H <sub>10</sub> O	Diethyl ether	74.12	12	-259.0	660.3	662.8	654.4	12	55-204
	C <sub>4</sub> H <sub>10</sub> O	Diallyl ether	94.14	15	-260.9	906.6	906.8	967.1	15	55-223
		$y' = 28.1 + 52.13x'$								

TABLE 92  
GIARME (Ethylene Oxide) (liquid)

IDL NO.	FORMULA	NAME	MOL. WT.	OXYGEN REQUIRED	OXYGEN BALANCE	Q <sub>v</sub> - KCAL/MOLE		Y'	X'	REF.
						OBSERVED	CALCULATED			
	C <sub>2</sub> H <sub>4</sub> O	Ethylene oxide	44.05	5	-181.6	302.4	302.1	295.7	5	82
	C <sub>3</sub> H <sub>6</sub> O	1,2-Dioxolane oxide	58.06	8	-220.1	551.4	455.1	457.7	8	82
	C <sub>4</sub> H <sub>8</sub> O	1,2-Butene oxide	72.10	11	-216.1	610.2	600.2	501.5	11	82
	C <sub>4</sub> H <sub>8</sub> O	1,3-Butene oxide	100.12	17	-217.6	911.7	911.4	907.2	17	55-215
		$y' = 41.2 + 51.02x'$								
	C <sub>6</sub> H <sub>10</sub> O	2-Methyl-2,3-butene oxide	86.13	14	-260.1	749.8	761.3	-	-	55-245





TABLE 66  
 (Continued)

COUNTRY	CITY	NO. OF STATIONS	NO. OF SEWERS	NO. OF PUMPS	OF REAL ESTATE		V.	K.	RE.
					IMPROVED	UNIMPROVED			
U.S.A.	Albany	10	11	100.0	107.7	107.7	121.0	7	71
	Albany	12	11	100.0	107.7	107.7	121.0	11	31
	Albany	14	11	100.0	107.7	107.7	121.0	12	50-225
	Albany	15	11	100.0	107.7	107.7	121.0	13	50-215
	Albany	16	11	100.0	107.7	107.7	121.0	14	50-215
	Albany	17	11	100.0	107.7	107.7	121.0	15	50-215
	Albany	18	11	100.0	107.7	107.7	121.0	16	50-215
	Albany	19	11	100.0	107.7	107.7	121.0	17	50-215
	Albany	20	11	100.0	107.7	107.7	121.0	18	50-215
	Albany	21	11	100.0	107.7	107.7	121.0	19	50-215
							Exp. = 121.0 Imp. = 107.7 K.V. = 114.9 RE. = 121.0 n = 8		

TABLE 67  
 (Continued)

COUNTRY	CITY	NO. OF STATIONS	NO. OF SEWERS	NO. OF PUMPS	OF REAL ESTATE		V.	K.	RE.
					IMPROVED	UNIMPROVED			
U.S.A.	Albany	10	11	100.0	107.7	107.7	121.0	7	71
	Albany	12	11	100.0	107.7	107.7	121.0	11	31
	Albany	14	11	100.0	107.7	107.7	121.0	12	50-225
	Albany	15	11	100.0	107.7	107.7	121.0	13	50-215
	Albany	16	11	100.0	107.7	107.7	121.0	14	50-215
	Albany	17	11	100.0	107.7	107.7	121.0	15	50-215
	Albany	18	11	100.0	107.7	107.7	121.0	16	50-215
	Albany	19	11	100.0	107.7	107.7	121.0	17	50-215
	Albany	20	11	100.0	107.7	107.7	121.0	18	50-215
	Albany	21	11	100.0	107.7	107.7	121.0	19	50-215
							Exp. = 121.0 Imp. = 107.7 K.V. = 114.9 RE. = 121.0 n = 8		

CONFIDENTIAL

TABLE 9B  
ACIDS (solid)

MOL WT.	FORMULA	NAME	MOL WT.	OXYGEN % (CALC)	OXYGEN BALANCE	G.C. KCAL/MOLE		Y'	X'	REF.
						GENERATED	CALCULATED			
<b>Monocarboxylic</b>										
	$C_{10}H_{20}O_2$	Capric acid	172.26	28	-260.1	1158.1	1149.9	1152.1	28	65-192
	$C_{11}H_{22}O_2$	Undecanoic acid	186.29	31	-266.1	1215.9	1216.1	1210.7	31	65-192
	$C_{12}H_{24}O_2$	Dodecanoic acid	200.31	34	-271.6	1272.8	1272.3	1269.4	34	118
	$C_{13}H_{26}O_2$	Tridecanoic acid	214.34	36	-276.3	1329.5	1328.7	1326.1	36	65-192
	$C_{14}H_{28}O_2$	Tetradecanoic acid	228.36	40	-287.0	1391.5	1392.1	1385.8	40	65-192
	$C_{18}H_{36}O_2$	Stearic acid	284.4	52	-292.5	2704.8	2709.6	2699.1	52	65-64, 55-192, 55-64, (Av.)
	$C_{19}H_{38}O_2$	Arachidic acid	312.42	60	-295.7	3045.9	3042.0	3040.2	60	65-192
	$C_{22}H_{44}O_2$	Behenic acid	352.57	64	-300.7	3538.8	3538.8	3532.7	64	65-192
<b>Dicarboxylic-Straight Chain</b>										
	$C_4H_6O_4$	Malonic acid	106.05	8	-61.5	206.5	206.5	150.4	7.6	118
	$C_5H_8O_4$	Succinic acid	118.09	7	-94.8	156.2	162.7	151.3	1.5	17
	$C_6H_{10}O_4$	Glutaric acid	132.1	10	-121.1	512.9	518.9	214.6	5.0	148
	$C_7H_{12}O_4$	Adipic acid	146.13	13	-152.3	662.9	671.1	311.6	6.5	148
	$C_8H_{14}O_4$	Pimelic acid	160.17	15	-199.8	827.7	831.1	431.0	8.0	148
	$C_9H_{16}O_4$	Suberic acid	174.19	19	-174.5	981.2	981.5	523.8	9.5	148
	$C_{10}H_{18}O_4$	Azelaic acid	188.2	22	-187.0	1144.7	1143.7	608.9	11.0	148
	$C_{11}H_{20}O_4$	Sebacic acid	188.2	25	-197.8	1297.3	1297.3	625.8	12.5	148
	$C_{12}H_{22}O_4$	Dodecanedioic acid	214.27	28	-207.1	1459.5	1455.1	725.0	14.0	148
	$C_{14}H_{26}O_4$	Hexadecanedioic acid	254.31	31	-215.4	1610.7	1612.3	860.5	15.5	148
	$C_{16}H_{30}O_4$	Octadecanedioic acid	294.34	34	-244.7	1768.6	1768.5	881.5	17.0	148
<b>Monocarboxylic-Branches Chain</b>										
	$C_4H_8O_2$	Methylmalonic acid	118.09	7	-94.8	161.5	159.6	123.5	3.5	65-192
	$C_5H_{10}O_2$	Isobutyramic acid	132.11	10	-127.1	531.2	526.1	250.3	6.0	148
	$C_5H_{10}O_2$	Methylsuccinic acid	132.11	10	-127.1	531.2	526.1	250.3	6.0	148
	$C_5H_{10}O_2$	Methylglutaric acid	132.11	10	-127.1	531.2	526.1	250.3	6.0	65-192
	$C_6H_{12}O_2$	2-Methylglutaric acid	146.13	13	-152.3	672.4	671.1	311.6	7.5	65-192
	$C_6H_{12}O_2$	3-Methylglutaric acid	146.13	13	-152.3	672.4	671.1	311.6	7.5	65-192
	$C_6H_{12}O_2$	2-Methyladipic acid	146.13	13	-152.3	672.4	671.1	311.6	7.5	65-192
	$C_6H_{12}O_2$	3-Methyladipic acid	146.13	13	-152.3	672.4	671.1	311.6	7.5	65-192
	$C_7H_{14}O_2$	2-Methylpimelic acid	160.17	15	-199.8	827.7	831.1	431.0	8.0	65-192
	$C_7H_{14}O_2$	3-Methylpimelic acid	160.17	15	-199.8	827.7	831.1	431.0	8.0	65-192
	$C_8H_{16}O_2$	2-Methylsuberic acid	174.19	19	-174.5	981.2	981.5	523.8	9.5	65-192
	$C_8H_{16}O_2$	3-Methylsuberic acid	174.19	19	-174.5	981.2	981.5	523.8	9.5	65-192
	$C_9H_{18}O_2$	2-Methylazelaic acid	188.2	22	-187.0	1144.7	1143.7	608.9	11.0	65-192
	$C_9H_{18}O_2$	3-Methylazelaic acid	188.2	22	-187.0	1144.7	1143.7	608.9	11.0	65-192
	$C_{10}H_{20}O_2$	2-Methylsebacic acid	188.2	25	-197.8	1297.3	1297.3	625.8	12.5	65-192
	$C_{10}H_{20}O_2$	3-Methylsebacic acid	188.2	25	-197.8	1297.3	1297.3	625.8	12.5	65-192
	$C_{12}H_{24}O_2$	2-Methyl-dodecanedioic acid	214.27	28	-207.1	1459.5	1455.1	725.0	14.0	65-192
	$C_{12}H_{24}O_2$	3-Methyl-dodecanedioic acid	214.27	28	-207.1	1459.5	1455.1	725.0	14.0	65-192
	$C_{14}H_{28}O_2$	2-Methyl-octadecanedioic acid	254.31	31	-215.4	1610.7	1612.3	860.5	15.5	65-192
	$C_{14}H_{28}O_2$	3-Methyl-octadecanedioic acid	254.31	31	-215.4	1610.7	1612.3	860.5	15.5	65-192
	$C_{16}H_{32}O_2$	2-Methyl-hexadecanedioic acid	294.34	34	-244.7	1768.6	1768.5	881.5	17.0	65-192
	$C_{16}H_{32}O_2$	3-Methyl-hexadecanedioic acid	294.34	34	-244.7	1768.6	1768.5	881.5	17.0	65-192

TABLE 97  
 ACID ANHYDRIDES (liquid)

AID NO.	FORMULA	NAME	MOL WT.	OXYGEN OBSERVED	OXYGEN BALANCE	Q <sub>v</sub> - KCAL/MOLE		Y <sup>a</sup>	X <sup>a</sup>	REF.
						OBSERVED	CALCULATED			
	$C_4H_6O_3$	Acetic anhydride	102.09	8	-12.4	631.9	631.3	126.2	8	55-223
	$C_6H_8O_5$	Glutaric anhydride	128.12	12	-104.5	665.6	669.1	127.2	13	147
	$C_8H_{10}O_5$	Phthalic anhydride	178.14	16	-174.1	746.6	743.6	146.9	14	55-148
		$y^a = 7.18 + 52.02x^a$						$F_{21}^a = 35.0$ $F_{22}^a = 1849.3$ $\Sigma y^a = 27,650.60$ $\Sigma x^a = 129.0$ $n = 3$		

 TABLE 100  
 ACID ANHYDRIDES (solid)

AID NO.	FORMULA	NAME	MOL WT.	OXYGEN OBSERVED	OXYGEN BALANCE	Q <sub>v</sub> - KCAL/MOLE		Y <sup>a</sup>	X <sup>a</sup>	REF.
						OBSERVED	CALCULATED			
	$C_4H_4O_3$	Succinic anhydride	100.07	8	-111.9	369.9	372.6	164.2	7	147
	$C_6H_8O_5$	Glutaric anhydride	128.12	12	-104.2	523.0	528.6	522.3	10	55-183
	$C_8H_{10}O_5$	Methylsuccinic anhydride	144.14	14	-124.2	544.2	546.9	545.1	10	147
	$C_{10}H_{14}O_5$	asym-Dimethylsuccinic anhydride	174.16	16	-142.3	681.3	682.5	680.1	13	147
	$C_{10}H_{14}O_5$	cis-sym-Dimethylsuccinic anhydride	174.16	16	-142.3	681.7	682.5	679.5	13	147
	$C_{10}H_{14}O_5$	trans-sym-Dimethylsuccinic anhydride	174.16	16	-142.3	677.9	677.0	676.7	13	147
	$C_{10}H_{14}O_5$	Trimethylsuccinic anhydride	190.18	18	-154.1	817.0	818.0	813.6	16	147
	$C_{12}H_{18}O_5$	Tetraethylsuccinic anhydride	196.18	18	-164.6	993.7	995.4	993.0	14	147
	$C_{14}H_{22}O_5$	asym-Diethylsuccinic anhydride	212.20	20	-184.6	993.5	993.5	995.8	19	147
	$C_{14}H_{22}O_5$	cis-sym-Diethylsuccinic anhydride	212.20	20	-184.6	993.4	993.4	994.7	19	147
	$C_{14}H_{22}O_5$	trans-sym-Diethylsuccinic anhydride	212.20	20	-184.6	996.8	996.8	993.1	19	147
	$C_{16}H_{26}O_5$	Trimethylsuccinic anhydride	184.23	25	-177.1	1311.2	1307.4	1307.0	25	147
	$C_{18}H_{30}O_5$	Tetraethylsuccinic anhydride	212.20	31	-211.7	1621.5	1621.4	1619.7	31	147
	$C_{18}H_{30}O_5$	Dimethylsuccinic anhydride	144.14	12	-124.9	1673.9	1673.9	1664.2	32	55-227
	$C_{18}H_{30}O_5$	Butyric anhydride	202.25	18	-202.6	1985.3	1986.0	1979.2	35	55-227
		$y^a = 2.61 + 52.07x^a$						$F_{21}^a = 230.0$ $F_{22}^a = 14,226.0$ $\Sigma y^a = 319,681.8$ $\Sigma x^a = 6510.0$ $n = 15.0$		

TABLE 101  
ESTWAS (liqwd)

ADJ. NO.	FORMULA	NAME	MOL. WT.	OXYGEN REQUIRED	OXYGEN BALANCE	Q <sub>p</sub> KCAL/MOLE		Y'	X'	REF.
						OBSERVED	CALCULATED			
<u>Of Monocarboxylic Acids</u>										
	C <sub>2</sub> H <sub>4</sub> O <sub>2</sub>	Methyl formate	60.05	1	-108.5	2,31.1	227.4	227.4	4.0	55-26
	C <sub>3</sub> H <sub>6</sub> O <sub>2</sub>	Methyl acetate	74.08	7	-151.7	350.6	181.4	174.9	7.0	123
		Ethyl formate	"	"	"	391.7	"	"	7.0	55-26
	C <sub>4</sub> H <sub>8</sub> O <sub>2</sub>	Ethyl acetate	88.10	10	-181.6	515.1	338.4	329.4	16.0	123
		Propyl propionate	"	"	"	536.9	"	511.2	10.0	123
	C <sub>6</sub> H <sub>12</sub> O <sub>2</sub>	Ethyl propionate	102.13	13	-201.7	650.9	663.4	685.2	13.0	123
		Methyl n-butyrate	"	"	"	693.3	"	694.6	13.0	123
		i-Propyl acetate	"	"	"	693.0	"	687.4	13.0	123
		Methyl i-butyrate	"	"	"	693.2	"	685.0	13.0	123
	C <sub>8</sub> H <sub>16</sub> O <sub>2</sub>	Ethyl n-butyrate	116.16	16	-220.4	815.9	641.4	691.0	13.0	55-154
		n-Propyl propionate	"	"	"	815.3	"	810.2	15.0	123
		n-Butyl acetate	"	"	"	815.7	"	811.6	16.0	123
		i-Butyl acetate	"	"	"	815.7	326.1	811.3	16.3	123
		i-Propyl propionate	"	"	"	815.0	"	810.4	16.0	123
		Ethyl i-butyrate	"	"	"	815.7	"	812.1	16.0	55-118
	C <sub>10</sub> H <sub>20</sub> O <sub>2</sub>	n-Butyl propionate	130.18	19	-231.5	1003.2	1263.3	954.2	19.0	123
		n-Propyl n-butyrate	"	"	"	1003.9	"	996.2	19.0	123
		Ethyl valerate	"	"	"	1017.5	"	1011.8	19.0	55-65
		i-Butyl acetate	"	"	"	1020.4	1001.3	996.7	19.0	123
		i-Propyl propionate	"	"	"	1020.3	"	996.0	19.0	123
		i-Propyl n-butyrate	"	"	"	1020.1	"	996.4	19.0	123
	C <sub>8</sub> H <sub>16</sub> O <sub>2</sub>	n-Butyl n-butyrate	116.21	22	-244.1	1157.8	1159.3	1152.1	22.1	123
		i-Butyl propionate	"	"	"	1156.9	"	1152.9	22.0	123
		i-Butyl n-butyrate	"	"	"	1157.1	"	1152.1	22.0	123
	C <sub>10</sub> H <sub>20</sub> O <sub>2</sub>	i-Butyl n-butyrate	158.23	25	-252.8	1312.4	1311.0	1306.2	25.0	43
<u>Of Dicarboxylic Acids</u>										
	C <sub>4</sub> H <sub>6</sub> O <sub>4</sub>	Dimethyl malonate	118.11	13	-121.1	554.0	574.4	274.2	5.0	46
	C <sub>6</sub> H <sub>10</sub> O <sub>4</sub>	Dimethyl succinate	146.14	13	-121.3	707.1	739.5	350.7	6.5	126
		Dimethyl oxalate	"	"	"	715.0	"	715.2	7.5	55-119
	C <sub>8</sub> H <sub>14</sub> O <sub>4</sub>	Dimethyl glutarate	160.17	15	-159.7	863.2	864.5	428.8	8.0	126
		Dimethyl malonate	"	"	"	863.4	"	863.4	8.0	55-119
	C <sub>10</sub> H <sub>18</sub> O <sub>4</sub>	Dimethyl adipate	174.19	19	-174.5	1019.6	1019.5	483.6	9.5	126
		Dimethyl succinate	"	"	"	1007.3	"	900.5	9.5	55-119
	C <sub>12</sub> H <sub>22</sub> O <sub>4</sub>	Dimethyl pimelate	188.22	21	-187.0	1176.0	1176.5	585.2	11.0	126
	C <sub>14</sub> H <sub>26</sub> O <sub>4</sub>	Dimethyl suberate	202.24	23	-197.8	1332.3	1329.5	662.3	12.5	126
		Methyl dimethylsuccinate (syn)	"	"	"	1323.9	1327.5	662.4	12.5	55-123
		Methyl dimethylsuccinate (trans)	"	"	"	1327.9	"	557.4	12.5	55-2
	C <sub>12</sub> H <sub>22</sub> O <sub>4</sub>	Dimethyl sebacate	216.27	29	-207.1	1486.3	1484.3	743.3	14.0	126
	C <sub>14</sub> H <sub>26</sub> O <sub>4</sub>	Dimethyl dodecanoate	230.30	31	-215.4	1644.7	1649.2	824.5	15.5	126
		y' = 14.08 + 57.66x'								
								Σ x' = 535.0		
								Σ y' = 28,265.0		
								Σ x'y' = 441,621.95		
								Σ x'^2 = 8,342.0		
								n = 39.0		

TABLE 102  
ESTWAS (solid)

ADI. NO.	FORMULA	NAME	MOL. WT.	OXYGEN REQUIRED	OXYGEN BALANCE	Q <sub>p</sub> KCAL/MOLE		Y'	X'	REF.
						OBSERVED	CALCULATED			
	C <sub>4</sub> H <sub>6</sub> O <sub>4</sub>	Dimethyl oxalate	118.09	7	-94.8	420.8	398.1	197.6	3.5	126
	C <sub>10</sub> H <sub>18</sub> O <sub>4</sub>	Dimethyl dimethylsuccinate (trans)	202.24	25	-197.8	1324.1	1323.0	660.0	12.5	75-9
	C <sub>10</sub> H <sub>18</sub> O <sub>4</sub>	Tetramethyl ethanetetra-carboxylate	262.21	19	-115.9	1025.1	1027.7	459.9	4.75	55-123
	C <sub>11</sub> H <sub>18</sub> O <sub>4</sub>	Tetramethyl ethylene-dimalonate	276.24	22	-127.4	1201.4	1202.1	490.9	5.0	55-187
	C <sub>14</sub> H <sub>26</sub> O <sub>4</sub>	Tetramethyl ethanetetra-carboxylate	318.32	31	-155.8	1661.2	1665.3	711.9	7.75	55-7
		y' = 16.13 + 51.56x'								
								Σ x' = 44.0		
								Σ y' = 1830.3		
								Σ x'y' = 15,067.6		
								Σ x'^2 = 281.375		
								n = 5		

TABLE 103  
NITRILES (Liquid)

ADL NO.	FORMULA	NAME	MOL. WT.	OXYGEN REQUIRED	OXYGEN BALANCE	G <sup>o</sup> KCAL/MOLE		Y'	X'	REF.
						OBSERVED	CALCULATED			
	C <sub>2</sub> H <sub>3</sub> N	Acetonitrile	41.05	5.5	-214.4	309.4	301.1	296.7	5.5	55-102
	C <sub>3</sub> H <sub>3.5</sub> N	Chloroacetonitrile	57.05	4.0	-125.2	250.7	250.7	242.0	4.0	55-24
	C <sub>3</sub> H <sub>5</sub> N	Propyl nitrile	53.06	7.5	-206.2	263.4	119.9	100.5	7.5	52
	C <sub>3</sub> H <sub>7</sub> N	Propyl nitrile	69.04	8.5	-205.9	174.4	157.6	155.7	8.5	55-102
	C <sub>4</sub> H <sub>7</sub> N	Methyl cyanoacetate	67.09	9.5	-137.2	171.6	170.1	151.4	8.5	55-78
	C <sub>4</sub> H <sub>9</sub> N	Allyl cyanide	67.09	10.5	-250.4	174.9	174.8	155.1	10.5	55-11
	C <sub>4</sub> H <sub>9</sub> N	Crotonitrile	67.09	9.5	-137.2	171.9	171.9	152.1	10.5	55-51
	C <sub>4</sub> H <sub>9</sub> N	n-Butyronitrile	69.10	11.5	-246.3	613.3	613.8	601.6	11.5	55-102
	C <sub>4</sub> H <sub>9</sub> N	Isobutyronitrile	69.10	11.5	-246.3	613.0	625.1	613.0	11.5	55-78
	C <sub>4</sub> H <sub>9</sub> N	1-Toleronitrile	83.13	14.5	-278.1	772.1	757.6	768.8	14.5	55-102
	C <sub>4</sub> H <sub>9</sub> N	Amlylpropionitrile	121.38	21.5	-283.9	1154.2	1174.8	1121.2	21.5	55-138
	C <sub>6</sub> H <sub>5</sub> N	o-Cyanobiphenyl	179.21	30.5	-272.3	1524.3	1583.1	1600.6	30.5	18
	C <sub>6</sub> H <sub>5</sub> N	Succinonitrile	80.09	10.0	-197.8	345.7	345.1	270.0	5.0	55-42
		y' = 9.33 + 52.67x'								
								Σ x' = 150.0		
								Σ y' = 7931.8		
								Σ x'y' = 154,474.4		
								Σ x'^2 = 2346.0		
								n = 13		
	C <sub>2</sub> F <sub>3</sub> O	Carbonyl cyanide	80.05	5.0	-99.9	332.0	289.2	-	-	50
	C <sub>2</sub> H <sub>3</sub> NO	Acetonitrile oxynitrile	71.05	7.5	-179.4	421.1	425.7	-	-	55-24
	C <sub>2</sub> H <sub>5</sub> N <sub>2</sub>	Propyl cyanazotate	127.14	14.5	-182.5	109.0	780.0	-	-	55-78

TABLE 104

NITRILES (solid)

ADL NO.	FORMULA	NAME	MOL. WT.	OXYGEN REQUIRED	OXYGEN BALANCE	G <sup>o</sup> KCAL/MOLE		Y'	X'	REF.
						OBSERVED	CALCULATED			
	C <sub>4</sub> H <sub>9</sub> N <sub>2</sub>	Cyanazotate acid	86.06	5.5	-101.5	293.8	290.3	257.4	5.5	55-78
	C <sub>4</sub> H <sub>9</sub> N <sub>2</sub> O	Cyanazotate acid	102.06	7.0	-133.2	376.3	377.7	374.1	7.0	55-78
	C <sub>4</sub> H <sub>9</sub> N <sub>2</sub> O	Methyl acetylcyanazotate	141.12	1.5	-141.7	685.0	683.2	665.3	12.5	55-78
	C <sub>4</sub> H <sub>9</sub> N <sub>2</sub>	Nitroacetonitrile	66.06	9.0	-169.5	391.8	390.1	391.6	3.5	55-42
	C <sub>4</sub> H <sub>9</sub> N <sub>2</sub>	Nitroacetonitrile	72.12	14.0	-221.0	695.9	704.1	346.9	8.5	55-42
	C <sub>4</sub> H <sub>9</sub> N <sub>2</sub>	Phthalonitrile	128.13	18.5	-224.8	955.4	957.3	479.3	9.0	121
		y' = 6.90 + 52.37x'								
								Σ x' = 64.0		
								Σ y' = 2357.7		
								Σ x'y' = 17,200.77		
								Σ x'^2 = 371.0		
								n = 6		
	C <sub>2</sub> F <sub>4</sub>	Carbon suboxide	76.06	3.0	-148.3	511.8	483.0	-	-	55-139

TABLE 105

CARBYLAMINES (Liquid)

ADL NO.	FORMULA	NAME	MOL. WT.	OXYGEN REQUIRED	OXYGEN BALANCE	G <sup>o</sup> KCAL/MOLE		Y'	X'	REF.
						OBSERVED	CALCULATED			
	C <sub>2</sub> H <sub>5</sub> N	Methyl carbylamine	41.05	5.5	-214.4	317.4	321.9	311.7	5.5	55-93
	C <sub>2</sub> H <sub>5</sub> N	Methyl carbylamine	41.05	5.5	-214.4	320.1	320.1	314.4	5.5	55-77
	C <sub>2</sub> H <sub>5</sub> N	Ethyl carbylamine	55.08	8.5	-216.9	177.1	479.8	471.4	8.5	55-84
	C <sub>2</sub> H <sub>5</sub> N	Ethyl carbylamine	55.08	8.5	-216.9	180.5	474.8	474.8	8.5	55-77
	C <sub>2</sub> H <sub>5</sub> N	Allyl carbylamine	67.09	10.5	-250.4	605.1	597.2	589.3	10.5	55-77
	C <sub>2</sub> H <sub>5</sub> N	Propyl carbylamine	81.10	11.5	-266.1	619.6	617.8	611.9	11.5	55-77
	C <sub>2</sub> H <sub>5</sub> N	1-Butyl carbylamine	83.13	14.5	-278.1	796.0	791.3	782.7	14.5	55-77
	C <sub>2</sub> H <sub>5</sub> N	1-Propyl carbylamine	97.16	17.5	-298.2	949.5	911.6	914.9	17.5	55-77
	C <sub>2</sub> H <sub>5</sub> N	Butyl carbylamine	111.18	19.5	-294.3	1026.5	1050.1	1049.5	19.5	55-77
		y' = 26.63 + 52.65x'								
								Σ x' = 101.5		
								Σ y' = 5983.6		
								Σ x'y' = 73,178.4		
								Σ x'^2 = 1,344.25		
								n = 9		

TABLE 106  
PRIMARY AMINES (liquid)

ADL NO.	FORMULA	NAME	MOL. WT.	OXYGEN REQUIRED	OXYGEN BALANCE	Cp. KCAL/MOLE		Y'	X'	REF.
						OBSERVED	CALCULATED			
	C <sub>2</sub> H <sub>7</sub> N	Methylamine	31.06	4.5	-231.8	255.1	254.1	250.1	4.5	55-99
	C <sub>2</sub> H <sub>9</sub> N	Ethylamine	45.08	7.5	-266.2	408.5	407.0	402.5	7.5	55-99
	C <sub>3</sub> H <sub>9</sub> N	Propylamine	57.09	9.5	-266.2	524.8	524.5	507.0	9.5	55-99
	C <sub>4</sub> H <sub>11</sub> N	n-Butylamine	73.14	10.5	-266.2	583.3	581.7	557.6	10.5	55-99
	"	1-Butylamine	"	"	"	710.6	715.5	704.9	13.5	55-99
	"	sec-Butylamine	"	"	"	713.6	712.9	710.5	13.5	55-99
	"	tert-Butylamine	"	"	"	713.0	"	709.8	13.5	55-99
	"	n-Pentylamine	"	"	"	715.0	"	712.8	13.5	55-99
	C <sub>6</sub> H <sub>13</sub> N	Aniline	93.12	15.5	-266.3	810.6	809.1	813.9	15.5	55-99
	C <sub>8</sub> H <sub>17</sub> N	1-Octylamine	87.16	16.5	-266.2	852.9	851.1	853.3	16.5	55-99
	C <sub>9</sub> H <sub>19</sub> N	Dodecylamine	107.15	18.5	-276.2	867.6	863.1	920.7	18.5	55-101 55-117 (Av.)
	"	n-Toluidine	"	"	"	961.3	"	967.1	18.5	55-117
	"	m-Toluidine	"	"	"	955.3	"	958.1	18.5	55-117
	C <sub>10</sub> H <sub>17</sub> N	Dodecylamine	101.19	19.5	-306.3	1022.2	1023.1	1016.5	19.5	55-99
	C <sub>10</sub> H <sub>19</sub> N	1-Methyl-2-piperidylamine	113.20	21.5	-301.9	1116.7	1118.3	1120.0	21.5	55-215
	C <sub>10</sub> H <sub>19</sub> N	2,6-Dimethylpiperidine	121.18	"	-283.9	1108.0	1115.3	1119.7	21.5	55-99
	C <sub>10</sub> H <sub>19</sub> N	Mapylamine	115.21	22.5	-312.5	1176.9	1176.9	1171.2	22.5	55-99
	C <sub>10</sub> H <sub>16</sub> N <sub>2</sub> O	Ethylenediamine hydrate	78.12	8.0	-163.7	452.6	454.0	221.2	8.0	16
		y' = 17.67 + 5L27x'								
								I x' = 262.5		
								I y' = 13,776.5		
								I x'y' = 28,805.95		
								I x'^2 = 4372.25		
								n = 18		

TABLE 107  
PRIMARY AMINES (gas)

ADL NO.	FORMULA	NAME	MOL. WT.	OXYGEN REQUIRED	OXYGEN BALANCE	Cp. KCAL/MOLE		Y'	X'	REF.
						OBSERVED	CALCULATED			
	CH <sub>5</sub> N	Methylamine	31.06	4.5	-231.8	258.1	257.5	252.6	4.5	55-223 55-160 (Av.)
	C <sub>2</sub> H <sub>7</sub> N	Ethylamine	45.08	7.5	-256.2	433.1	433.4	407.5	7.5	55-223
	C <sub>3</sub> H <sub>9</sub> N	Propylamine	57.09	9.5	-261.2	528.1	531.5	509.5	9.5	55-223
	C <sub>4</sub> H <sub>11</sub> N	n-Butylamine	59.13	10.5	-261.2	572.3	599.4	566.8	10.5	55-223
		y' = 17.96 + 5L.99x'								
								I x' = 32.0		
								I y' = 1735.5		
								I x'y' = 14,975.35		
								I x'^2 = 277.0		
								n = 4		

TABLE 108  
PRIMARY AMINES (solid)

ADL NO.	FORMULA	NAME	MOL. WT.	OXYGEN REQUIRED	OXYGEN BALANCE	Cp. KCAL/MOLE		Y'	X'	REF.
						OBSERVED	CALCULATED			
	C <sub>9</sub> H <sub>11</sub> NO	1-Adrenergolol	109.12	14.5	-212.6	790.0	787.7	767.6	14.5	55-97
	C <sub>9</sub> H <sub>11</sub> NO	p-Andradoline	123.15	17.5	-227.4	914.0	911.0	911.0	17.5	55-97
	C <sub>9</sub> H <sub>13</sub> N	n-Toluidine	107.15	18.5	-276.2	954.1	953.2	951.5	18.5	55-117
	C <sub>10</sub> H <sub>15</sub> N	Pseudoecardine	135.20	24.5	-259.9	1265.9	1274.1	1270.1	24.5	55-99
	C <sub>10</sub> H <sub>15</sub> N	1-Naphthylamine	143.18	"	-275.8	1263.0	1263.4	1278.1	24.5	55-99
	"	2-Naphthylamine	"	"	"	1261.0	"	1275.2	24.5	55-99
	C <sub>10</sub> H <sub>13</sub> N	2-Amino-1-phenyl	109.22	29.5	-278.9	1532.4	1523.6	1547.1	29.5	18
	"	1-Amino-1-phenyl	"	"	"	1524.1	"	1530.8	29.5	18
	C <sub>10</sub> H <sub>15</sub> N	p-Phenylenediamine	106.14	16.0	-235.7	817.0	817.3	821.6	16.0	57
	C <sub>10</sub> H <sub>13</sub> N	Serentine	104.23	30.0	-260.5	1555.4	1554.1	1570.0	15.0	99
	"	"	"	"	"	1556.0	"	1575.1	15.0	100
	"	"	"	"	"	1560.9	"	1577.7	15.0	55-117
		y' = 4.26 + 5L.00x'								
								I x' = 236.0		
								I y' = 12,323.3		
								I x'y' = 258,538.85		
								I x'^2 = 5339.0		
								n = 12		

TABLE 109  
SECONDARY AMINES (liquid)

ADL NO.	FORMULA	NAME	MOL. WT.	OXYGEN REQUIRED	OXYGEN BALANCE	Q <sub>v</sub> - KCAL/MOLE		Y'	X'	REF.
						OBSERVED	CALCULATED			
	C <sub>2</sub> H <sub>7</sub> N	Methylamine	45.08	7.5	-256.2	115.7	113.7	111.0	7.5	55-99
	C <sub>3</sub> H <sub>9</sub> N	Ethylamine	73.14	11.5	-295.3	122.8	125.5	121.1	11.5	55-99
	C <sub>4</sub> H <sub>11</sub> N	Nethylamine	107.15	18.5	-276.4	976.9	976.5	980.0	18.5	55-99
	C <sub>5</sub> H <sub>13</sub> N	Propylamine	115.20	24.5	-299.9	1299.6	1299.6	1292.3	24.5	55-99
	C <sub>6</sub> H <sub>15</sub> N	N-1-butylamine	129.24	29.5	-315.7	1341.4	1347.5	1340.2	29.5	55-99
	C <sub>7</sub> H <sub>17</sub> N	N-1-pentylamine	157.29	31.5	-320.8	1660.6	1629.0	1650.6	31.5	55-99
		$y' = 13.34 + 51.96x'$								
								X' = 139.5		
								Y' = 2316.8		
								X'Y' = 167,050.1		
								X'² = 3168.75		
								n = 7		
	C <sub>8</sub> H <sub>19</sub> N	N-hexylamine	127.18	21.5	-283.9	1121.5	1132.6	-	-	55-99

TABLE 110  
SECONDARY AMINES (solid)

ADL NO.	FORMULA	NAME	MOL. WT.	OXYGEN REQUIRED	OXYGEN BALANCE	Q <sub>v</sub> - KCAL/MOLE		Y'	X'	REF.
						OBSERVED	CALCULATED			
	C <sub>10</sub> H <sub>23</sub> N	Dodecylamine	169.28	29.5	-278.9	1534.2	1532.3	1544.2	29.5	55-102
	C <sub>11</sub> H <sub>25</sub> N	Tridecylamine	197.37	35.5	-287.9	1530.2	1530.2	1512.7	29.5	55-99
	C <sub>12</sub> H <sub>27</sub> N	N-Tetradecylamine	225.47	38.5	-280.9	1813.0	1846.5	1865.1	35.5	55-99
		N-Pentadecylamine				2003.8	1999.5	2034.3	38.5	55-99
		N-Hexadecylamine				1998.0		2023.5	36.5	55-99
		$y' = 14.9 + 53.90x'$								
								X' = 171.5		
								Y' = 2019.3		
								X'Y' = 113,825.15		
								X'² = 5965.23		
								n = 5		

TABLE 111  
TERTIARY AMINES (Liquid)

ADL NO.	FORMULA	NAME	MOL. WT.	OXYGEN REQUIRED	OXYGEN BALANCE	Q <sub>v</sub> - KCAL/MOLE		Y'	X'	REF.
						OBSERVED	CALCULATED			
	C <sub>3</sub> H <sub>9</sub> N	Trimethylamine	59.11	10.5	-286.2	578.6	573.9	572.9	10.5	55-99
	C <sub>4</sub> H <sub>11</sub> N	Triethylamine	101.19	19.5	-308.3	1036.8	1031.1	1031.1	19.5	55-99
	C <sub>5</sub> H <sub>13</sub> N	Diethylamine	121.18	21.5	-281.9	1182.7	1189.0	1182.7	21.5	55-102
	C <sub>6</sub> H <sub>15</sub> N	Diethylamine	149.23	27.5	-254.8	1171.6	1152.1	1151.1	27.5	55-102
	C <sub>7</sub> H <sub>17</sub> N	Tri-n-butylamine	185.34	37.5	-327.7	1973.6	1981.2	1976.2	37.5	55-99
	C <sub>8</sub> H <sub>19</sub> N	Tri-1-amylamine	227.42	46.5	-327.1	2459.3	2452.1	2452.1	46.5	55-99
		$y' = 20.46 + 52.16x'$								
								X' = 163.0		
								Y' = 8675.0		
								X'Y' = 278,616.7		
								X'² = 5277.5		
								n = 6		
	C <sub>10</sub> H <sub>23</sub> N	Tripropylamine (a)	245.31	43.5	-283.7	2267.8	2257.8(116)	-	-	55-102
	C <sub>11</sub> H <sub>25</sub> N	Tribenzylamine (b)	267.39	52.5	-292.3	2762.1	2737.9(116)	-	-	55-99



TABLE 112  
ANILINS (liquid)

AGI NO.	FORMULA	NAME	MW	OXYGEN REQUIRED	OXYGEN BALANCE	Q <sup>o</sup> , KCAL/MOLE		Y'	X'	REF.
						OBSERVED	CALCULATED			
	$C_8H_9N_2O_2$	Formylurea	86.07	4.0	- 72.7	287.0	217.8	91.6	2.0	55-133
	$C_8H_9N_2O_2$	D-Tartarimide	125.12	5.0	- 66.4	427.0	419.6	297.9	4.0	55-54
	$C_8H_9N_2O_2$	meso-Tartarimide	"	"	"	426.0	"	297.7	4.0	55-54
	$C_{10}H_{11}N_2O_2$	N-Tartarimidobenzamide	206.22	20.0	-156.7	1064.1	1055.7	512.5	10.0	55-54
	"	N-Tartarimidobenzamide	"	"	"	1066.1	"	519.6	11.0	55-54
	"	meso-Tartarimidobenzamide	"	"	"	1065.1	"	520.1	13.0	55-54
		$\gamma' = -5.47 - 52.68x'$						$1 x' = 40.0$ $1 y' = 2068.4$ $1 x' y' = 17,441.6$ $1 x'^2 = 136.0$ $1 y'^2 = 6$		
	$C_8H_9NO$	Formamide	85.09	2.0	- 50.6	134.9	131.6	-	-	55-209

TABLE 113  
ANILIN (solid)

AGI NO.	FORMULA	NAME	MW	OXYGEN REQUIRED	OXYGEN BALANCE	Q <sup>o</sup> , KCAL/MOLE		Y'	X'	REF.
						OBSERVED	CALCULATED			
	$C_6H_7NO$	Aniline	93.09	2.0	- 11.9	268.4	268.4	127.4	2.5	55-209
	$C_6H_7NO$	Aceamide	89.07	5.5	-189.0	282.8	287.0	276.9	5.5	55-209
	$C_8H_9NO_2$	Methyl oxamate	103.08	"	- 85.4	305.4	300.8	285.9	5.5	55-184
	$C_8H_9NO_2$	Ethyl oxamate	99.09	7.5	-134.7	397.2	401.4	374.6	7.5	55-184
	$C_8H_9NO_2$	Propyl oxamate	73.09	8.5	-185.1	459.9	463.7	434.2	8.5	55-209
	$C_8H_9NO_2$	Ethyl oxamate	117.10	"	-185.1	457.3	462.2	439.1	8.5	55-184
	$C_8H_9NO$	n-Butyramide	87.12	11.5	-213.8	495.0	600.5	593.3	11.5	55-209
	$C_8H_9NO$	Isobutyramide	"	"	"	495.9	597.6	592.9	11.5	55-209
	$C_8H_{11}NO$	1-Valeramide	104.15	14.5	-229.4	551.5	644.8	748.2	14.5	55-209
	$C_8H_{11}NO$	Formanilide	121.13	16.5	-217.9	551.0	645.7	751.3	15.5	55-209
	"	Benzamide	"	"	"	648.0	652.8	652.0	16.5	5
	$C_8H_9NO$	Acetamidide	135.16	19.5	-180.5	1010.4	1012.4	1010.7	19.5	55-209
	$C_8H_9NO_2$	N-Benzoylglycine	179.17	"	-174.1	1012.4	1008.9	1016.2	19.5	55-209
	$C_8H_9NO_2$	N-Phenylpropionamide	144.15	23.5	-256.0	1099.2	1099.2	1051.3	23.5	55-118
	$C_8H_9NO_2$	N-Methylpropionamide	139.19	21.5	-217.1	1150.4	1140.2	1107.4	21.5	55-118
	$C_{10}H_{11}NO_2$	N-(p-Anisyl)glycine	209.20	"	-214.4	1135.2	1128.1	1128.3	21.5	55-209
	$C_{10}H_{11}NO_2$	N-Benzoylalanine	193.20	22.5	-186.3	1168.1	1163.7	1173.8	22.5	55-208
	$C_8H_9NO_2$	Propionamide	116.15	"	-211.3	1167.6	1169.1	1167.9	22.5	55-209
	$C_{10}H_{11}NO_2$	N-(o-Tolyl)glycine	191.20	"	-185.1	1167.7	1163.7	1173.4	22.5	55-208
	"	N-(m-Tolyl)glycine	"	"	"	1167.0	"	1174.7	22.5	55-208
	"	N-(p-Tolyl)glycine	"	"	"	1167.5	"	1173.2	22.5	55-109
	"	N-(Phenylacetyl)glycine	"	"	"	1164.9	"	1170.6	22.5	55-208
	$C_8H_9NO$	N-Benzyloxycarbonylurea	151.22	24.5	-255.8	1394.0	1314.9	1265.0	24.5	55-118
	$C_8H_9NO_2$	Phenacetin	179.17	"	-218.7	1285.2	1279.5	1269.6	24.5	55-160
	$C_{11}H_{13}NO_2$	N-(o-Tolyl)alanine	207.14	25.5	-196.9	1441.7	1418.7	1329.1	25.5	55-208
	"	N-(p-Tolyl)alanine	"	"	"	1441.5	"	1329.7	25.5	55-208
	$C_{10}H_{11}NO$	Benzanilide	197.23	30.5	-217.4	1635.5	1566.9	1576.9	30.5	55-209
	$C_{10}H_{11}NO_2$	N-(Benzoyl)-N-phenylalanine	269.29	36.5	-215.7	1890.1	1876.9	1914.3	36.5	55-66
	$CH_3N_2O$	Urea	60.06	3.0	- 75.9	151.0	150.4	77	1.5	48
	"	"	"	"	"	151.3	"	71.8	1.5	134
	$C_8H_9NO_2$	Formamide	89.07	4.0	- 72.7	205.2	208.6	208.2	2.0	55-184
	$C_8H_9NO_2$	Valeronamide	102.09	7.0	-109.7	354.8	359.2	376.4	3.5	55-184
	$C_8H_9NO_2$	Butylurea	89.11	9.0	-163.1	472.0	474.2	474.7	4.5	55-130
	$C_8H_9NO_2$	Succinamide	116.12	10.0	-137.4	509.2	516.0	525.7	5.0	55-184
	$C_8H_9NO_2$	N,N'-Dimethylsuccinamide	130.15	13.0	-159.0	685.0	677.0	660.1	6.5	55-184
	$C_8H_9NO_2$	N,N'-Dimethylmalonamide	128.20	19.0	-192.2	591.8	666.2	664.6	9.5	55-184
	$C_8H_9NO_2$	1,3-Diphenylurea	212.24	31.0	-233.7	1612.1	1600.9	669.1	15.5	55-184
	$C_8H_9NO_2$	1,1-Diphenylurea	"	"	"	1612.5	1617.9	667.0	15.5	55-184
	$C_{10}H_{14}NO_2$	Succinamide (sym)	268.30	38.0	-226.6	1973.1	1966.8	984.4	19.0	55-184
	$C_8H_9NO_2$	Oxalic acid	132.08	4.0	- 48.5	207.5	191.5	68.7	1.33	55-130
	$C_8H_9NO_2$	Acetylurea	102.09	7.0	-109.7	352.8	352.8	316.6	2.33	55-130
		$\gamma' = -6.03 - 52.21x'$						$1 x' = 660.66$ $1 y' = 31,131.4$ $1 x' y' = 635,623,393$ $1 x'^2 = 12,240,378$ $1 y'^2 = 6$		
	$C_8H_9NO_2$	Dimethylurea	100.11	7.0	- 93.2	384.0	406.1	-	-	121
	$C_8H_9NO_2$	1,3-Dimethyl-1,3-diphenylurea	240.29	37.0	-216.4	1953.0	1948.3	-	-	141
	$C_8H_9NO_2$	1,3-Dimethyl-1,3-diphenylurea	268.35	43.0	-254.4	2266.7	2218.5	-	-	141

TABLE 114  
HYDRAZIDES (contd)

AOL NO.	FORMULA	NAME	MOL. WT.	OXYGEN REQUIRED	OXYGEN BALANCE	O.C. KCAL/MOLF		Y'	X'	REF.
						ORIGIN	CALCULATED			
352	$N_4H_4O_2$	Hydrazine mononitrate	95.06	- 0.5	+ 6.4	117.2	120.9	127.0	2.0	95
356	$C_2H_4N_4O_2$	1-Amino-3-nitroguanidina	119.09	- 2.5	+ 31.6	270.1	265.1	247.5	2.5	77
	$C_2H_4N_4O_2$	Hydrazine hydrogen oxalate	122.09	3.0	- 39.3	259.1	170.4	147.2	4.0	96
355	$C_2H_4N_4O_2$	1,3-Dihydroxy-1,3-tetraole	168.13	8.0	- 57.1	169.7	175.2	135.3	6.0	92
318	$C_2H_4N_4O_2$	2,4-Dinitrophenylhydrazine	174.14	11.0	- 81.8	79.4	78.5	612.2	11.0	56
511	$C_2H_4N_4O_2$	Phenylhydrazine	168.14	15.0	- 236.7	871.7	866.5	849.1	15.0	8
						875.4		871.5	15.0	55-100
490	$C_2H_4N_4O_2$	ac-Methylphenylhydrazine	122.10	19.0	- 249.0	1048.3	1013.0	1025.2	19.0	55-100
693	$C_2H_4N_4O_2$	2,2'-Hydroxybis-1-benzopyr- nitrile	166.22	23.0	- 271.4	1259.2	1261.5	1219.7	23.0	77
	$C_{14}H_{12}O_2$	4,4'-Dinitrohydrosobenzene	274.23	25.0	- 115.9	1166.1	1509.0	1319.4	25.0	26
591	$C_{14}H_{12}N_2$	Hydrosobenzene	184.21	30.0	- 260.5	1530.4	1591.2	1591.1	30.0	99
						1587.5		1588.5	30.0	152
						1591.4		1597.1	30.0	25
						1597.3		1599.0	30.0	55-100
						1598.2		1598.5	30.0	55-147
	$C_{14}H_{12}N_2O_2$	Hydrosobenzene dihydroiodide	440.09	31.0	- 112.7	1614.8	1613.2	1593.5	31.0	99
592	$C_2H_4N_4O_2$	Tetrapropylhydrazine	336.42	50.0	- 275.0	3066.7	3066.2	3066.0	50.0	5
511	$C_6H_4N_4O_2$	1,3-Diaminoguanidina mono- nitrate	152.12	3.0	+ 31.6	329.8	333.1	173.6	2.75	77
	$C_6H_4N_4O_2$	1,2,3-Triaminoquandina mono- nitrate	167.11	3.5	+ 31.5	398.6	411.0	131.3	2.0	61
								$\gamma' = 32.16 + 57.04\beta'$ $2x' = 365.25$ $1y' = 19,666.4$ $1x'y' = 542,346.25$ $1x'' = 10,973.6125$ $n = 19$		
353	$N_4H_4$	Hydrazine (11q)	32.05	2.0	- 99.8	148.5	144.2(e)	-	-	90
691	$C_2H_4N_4O_2$	Hydrazine hydrate (11q)	50.06	2.0	- 31.9	116.9	116.2(e)	-	-	50
1172	$C_2H_4N_4$	Methylhydrazine (11q)	46.07	5.0	- 173.6	311.9	311.4(e)	-	-	20
354	$C_2H_4N_4O_2$	5,5'-Hydrazo-1,3-tetraole	174.13	6.0	- 51.1	159.9	179.0	-	-	77
511	$C_2H_4N_4O_2$	Phenylhydrazine	168.14	15.0	- 236.7	864.4	866.5	-	-	55-147
696	$C_{14}H_{12}N_2O_2$	N,N'-Bisacetamide	240.25	32.0	- 213.1	1615.5	1666.5	-	-	5
695	$C_{14}H_{12}N_2O_2$	N,N'-Diacetamide	260.30	38.0	- 266.6	2006.3	1995.7	-	-	99

TABLE 115  
HYDRAZIDES (contd)

AOL NO.	FORMULA	NAME	MOL. WT.	OXYGEN REQUIRED	OXYGEN BALANCE	O.C. KCAL/MOLF		Y'	X'	REF.
						ORIGIN	CALCULATED			
694	$C_{14}H_{12}N_2O_2$	Acetic 1,3-diphenylhydrazide	266.27	34.0	- 220.6	1792.9	1792.4	1796.3	34.0	99
537	$C_2H_4N_4O_2$	Carbohydrazide hydrogen oxalate	180.13	5.0	+ 44.6	311.6	304.0	169.3	2.5	96
552	$C_2H_4N_4O_2$	Oxalic dihydrazide	116.10	5.0	- 67.7	378.9	325.7	150.0	4.5	57
						325.7		150.0	4.5	52
	$C_2H_4N_4O_2$	Uraminburet	133.12	5.5	- 65.1	345.5	340.5	174.5	- 2.75	96
697	$C_2H_4N_4O_2$	Melonic dihydrazide	132.13	8.0	- 96.9	476.5	481.5	235.3	4.0	57
698	$C_2H_4N_4O_2$	Succinic dihydrazide	166.15	11.0	- 720.4	639.6	597.2	312.5	5.5	57
								$\gamma' = 30.18 + 51.91\beta'$ $2x' = 53.75$ $1y' = 3602.5$ $1x'y' = 45,123.175$ $1x'' = 1,278.5625$ $n = 7$		

TABLE 115  
ASOS (collid)

ADL NO.	FORMULA	NAME	MOL. WT.	OXYGEN EQUIVALENT	OXYGEN BALANCE	G. KCAL/MOLE		Y'	X'	REF.
						OBSERVED	CALCULATED			
	$C_3H_6$	Aso-1-propene (liq)	42.08	19.0	-266.2	1574.4	1550.7	1618.7	12.0	75
	$C_8H_{11}N_3$	2,2'-Azobis-1-butyronitrile	171.21	27.0	-233.8	1211.0	1230.3	1186.0	22.0	77
	$C_{12}H_{10}O_2$	p-Hydroxyacetophenone	198.22	28.0	-276.0	1702.0	1676.3	1515.7	25.0	55-101
	$C_{10}H_{10}O_2$	Asobenzene	162.22	29.0	-254.6	1546.0	1548.9	1557.2	29.0	139
	.	.	.	.	.	1577.2	.	1567.8	28.0	55-147
	.	.	.	.	.	1552.6	.	1565.3	29.0	55-104
	.	.	.	.	.	1575.2	.	1558.5	27.0	55-217
	$C_{12}H_{11}N_3$	p-Aminazobenzene	187.23	27.5	-219.3	1574.0	1576.6	1556.6	27.5	55-101
	$C_{12}H_{11}N_3$	2,2'-Azobis-1-butyronitrile	171.21	30.0	-265.1	1597.4	1666.6	1603.7	30.0	55-101
	$C_8H_8O_2$	p-Azoanisole	142.27	33.0	-217.9	1796.4	1767.6	1777.1	33.0	55-101
	$C_{12}H_{11}N_3$	p-Azophenole	176.22	33.0	-230.8	2104.0	2095.6	1680.2	33.0	55-101
	$C_{22}H_{21}N_3$	Phenylazo(triphenylmethane)	315.43	60.0	-275.5	3171.	3127.4	3203.0	60.0	152
	$C_{22}H_{21}N_3$	Diphenylazo(triphenylmethane)	352.45	63.0	-276.1	3238.	3281.8	3267.0	63.0	152
		$y' = 69.72 + 51.45x'$								
								$\Sigma x' = 139.5$		
								$\Sigma y' = 21,544.7$		
								$\Sigma x'y' = 502,282.0$		
								$\Sigma x'^2 = 16,912.25$		
								$n = 13$		

TABLE 117  
AZOTS (solid)

ADL NO.	FORMULA	NAME	MOL. WT.	OXYGEN EQUIVALENT	OXYGEN BALANCE	G. KCAL/MOLE		Y'	X'	REF.
						OBSERVED	CALCULATED			
	$C_{12}H_{10}O_2$	Azobenzene	162.22	28.0	-266.0	1534.5	1511.7	1547.1	28.0	55-101
	$C_{12}H_{11}N_3$	m-Azobenzidine	256.30	35.0	-215.5	1931.4	1907.4	1909.8	35.0	55-101
	$C_{12}H_{11}N_3$	p-Azobenzidine	256.30	35.0	-215.3	2092.0	2091.0	1975.7	35.0	55-101
	$C_{12}H_{10}O_2$	o-Isoxyphenole	162.22	35.0	-266.0	2068.0		2068.0	35.0	55-101
		$y' = 66.47 + 52.78x'$								
								$\Sigma x' = 167$		
								$\Sigma y' = 611.50$		
								$\Sigma x'y' = 310,413.10$		
								$\Sigma x'^2 = 5681.0$		
								$n = 5$		

TABLE 118  
QUANTITIES (cont'd)

ADL NO	FORMULA	NAME	MOLE WT.	OXYGEN REQUIRED	OXYGEN BALANCE	HEAT OF COMBUSTION		Y'	X'	REF.
						DETERMINED	CALCULATED			
624	C <sub>8</sub> H <sub>10</sub> N <sub>4</sub> O <sub>6</sub>	1-Methyl-2-(2,2,2-trinitroethyl)guanidine	267.17	0.5	- 3.0	432.5	432.6	26.4	0.5	128-1
152	CH <sub>5</sub> N <sub>5</sub> O <sub>5</sub>	Nitroguanidine	164.07	2.0	- 30.7	208.5	206.5	106.0	2.0	77
.	.	.	.	.	.	206.8	.	106.2	2.0	23
.	.	.	.	.	.	210.2	.	107.6	2.0	84
.	.	.	.	.	.	207.8	.	106.2	2.0	124-3
.	.	.	.	.	.	210.3	.	107.7	2.0	74
.	.	.	.	.	.	210.0	.	107.4	2.0	94
114	C <sub>8</sub> H <sub>10</sub> N <sub>4</sub> O <sub>6</sub>	Guanidine nitrate	122.09	.	.	108.2	211.7	229.5	4.5	124
116	C <sub>8</sub> H <sub>10</sub> N <sub>4</sub> O <sub>6</sub>	1-Oxamyl-3-nitrosam	99.10	3.5	5.5	288.9	285.6	184.7	3.5	94
126	C <sub>8</sub> H <sub>10</sub> N <sub>4</sub> O <sub>6</sub>	Oxamylam nitrate	165.12	.	- 33.9	275.1	275.7	181.6	6.0	128-4
130	C <sub>8</sub> H <sub>10</sub> N <sub>4</sub> O <sub>6</sub>	1-Ethyl-3-(2,2,2-trinitropropyl)guanidine dinitrate	268.19	4.0	- 23.9	584.0	585.4	205.8	4.0	128-1
625	C <sub>8</sub> H <sub>10</sub> N <sub>4</sub> O <sub>6</sub>	1-Nitro-3-(1-hydroxyethyl)guanidine nitrate	193.13	4.5	- 37.3	473.4	482.0	224.4	4.5	128-1
671	C <sub>8</sub> H <sub>10</sub> N <sub>4</sub> O <sub>6</sub>	1-Methyl-3-nitroguanidine	173.10	5.0	- 47.7	376.6	377.1	258.3	5.0	84
550	C <sub>8</sub> H <sub>10</sub> N <sub>4</sub> O <sub>6</sub>	Guanoguanidine (diacylamide)	84.04	6.0	- 114.2	330.7	337.9	304.4	6.0	121
.	.	.	.	.	.	.	.	300.7	6.0	92
667	C <sub>8</sub> H <sub>10</sub> N <sub>4</sub> O <sub>6</sub>	1-Ethyl-3-nitroguanidine	172.13	8.0	- 94.9	530.3	532.3	431.7	8.0	84
.	C <sub>4</sub> H <sub>6</sub> N <sub>2</sub> O <sub>2</sub>	Creatine	131.14	10.5	- 123.1	588.1	581.5	516.3	10.5	49
.	C <sub>8</sub> H <sub>10</sub> N <sub>4</sub> O <sub>6</sub>	Oxamidine	113.12	.	- 128.5	688.1	681.1	524.7	10.5	49
668	C <sub>8</sub> H <sub>10</sub> N <sub>4</sub> O <sub>6</sub>	1-(n-Propyl)-3-nitroguanidine	116.15	11.0	- 120.4	684.7	687.3	509.9	11.0	84
669	C <sub>8</sub> H <sub>10</sub> N <sub>4</sub> O <sub>6</sub>	1-(n-Butyl)-3-nitroguanidine	140.15	14.0	- 139.8	844.8	842.4	725.0	14.0	84
670	C <sub>8</sub> H <sub>10</sub> N <sub>4</sub> O <sub>6</sub>	1-(n-Amyl)-3-nitroguanidine	174.20	17.0	- 156.1	1004	997.4	844.8	17.0	84
622	C <sub>8</sub> H <sub>10</sub> N <sub>4</sub> O <sub>7</sub>	1,1'-(2-Hydroxytriethylene)-bis(3-nitroguanidine) mononitrate	309.21	9.5	- 44.0	661.1	662.9	219.2	9.5	128-1
								22		
								127.25		
								653.5		
								58,653.4		
								113,5625		
152	C <sub>8</sub> H <sub>10</sub> N <sub>4</sub> O <sub>5</sub>	Nitroguanidine	164.07	2.0	- 30.7	208.5	206.5	.	.	56
631	C <sub>8</sub> H <sub>10</sub> N <sub>4</sub> O <sub>5</sub>	1-Ethyl-3-(2-hydroxyethyl)guanidine nitrate	193.13	4.5	- 37.3	462.7	462.0	.	.	95-3
543	C <sub>8</sub> H <sub>10</sub> N <sub>4</sub> O <sub>5</sub>	N-Oxamylamino acid	231.09	5.5	- 67.1	338.7	336.3	.	.	56
	C <sub>8</sub> H <sub>10</sub> N <sub>4</sub> O <sub>5</sub>	Bis(guanidine) carbonate	180.17	9.0	- 79.9	499.8	.	.	.	33

TABLE 122

ALTERNATIVE METHODS (solid)

AOI NO.	FORMULA	NAME	MOL. WT.	OXYGEN REQUIRED	OXYGEN BALANCE	D <sub>c</sub> , KCAL/MOLE		Y <sup>1</sup>	X <sup>1</sup>	REF.
						OBSERVED	CALCULATED			
666 192	$C_{12}H_{14}N_2O_6$ $C_{12}H_{14}N_2O_6$ $C_{12}H_{14}N_2O_6$	2-Nitropropene polymer $\alpha, \omega$ -m-Trinitrotoluene m-Trinitrobenzyltoluene	87.06 227.13 257.17	6.5 10.5 21.5	-119.4 -70.0 -255.2	122.2 816.1 1194.3	129.8 699.7 1193.4	519.4 356.9 1197.0	0.4 5.25 21.5	128-3 85 55-243
574 77	$C_6H_8N_2O_4$ $C_6H_8N_2O_4$	2,2-Dinitro-1,1-propanediol 2,4-Dinitro-1-propanol	156.09 150.09	3.0 4.0	-78.9 -42.8	385.6 413.0	381.0 425.9	170.9 200.1	1.5 2.0	128-2 1
57	$C_6H_8N_2O_4$	2,5-Dinitrotoluene	153.07	4.0	-40.5	521.2	427.7	191.4	2.0	65
367	$C_6H_8N_2O_4$	2,2-Dinitropropane	134.09	5.0	-59.7	451.4	446.3	237.9	2.5	81
651	$C_8H_{10}N_2O_4$	1,2-Dinitro-2,1-butanediol	180.12	6.0	-51.3	504.6	514.9	241.7	3.0	124-2
600	$C_8H_{10}N_2O_4$	2-Nitro-2-(2-nitroethyl)-furan	164.11	9.0	-78.2	478.3	693.0	316.8	4.5	45
59	$C_8H_{10}N_2O_4$	2,2-Dimethyl-1,1-dinitropropane	162.15	11.0	-136.4	752.0	752.0	374.5	5.5	86
575	$C_8H_{10}N_2O_4$	4,5-Dinitroisalic acid	260.17	*	*	747.2	777.1	377.1	6.5	79
654	$C_8H_{10}N_2O_4$	3,3-Dinitro-1,5-pentanediamine	192.18	*	-79.1	754.0	749.0	383.0	5.5	124-2
602	$C_8H_{10}N_2O_4$	$\alpha, \omega$ -Dinitrotoluene	182.13	-12.0	-99.9	824.8	815.7	411.8	6.0	111-3
57	$C_{12}H_{14}N_2O_6$ $C_{12}H_{14}N_2O_6$	2,3-Dimethyl-2,3-dinitrobutane	182.13 176.17	13.0 14.0	-114.2 -127.3	831.1 897.7	822.2 936.5	412.9 467.2	6.0 7.0	123-2 128-3
245	$C_8H_{10}N_2O_4$	2-Methyl-2,3-dinitrobutane	207.15	8.5	-65.7	699.1	718.0	395.3	4.833	79
570	$C_8H_{10}N_2O_4$	2-Methyl-2,3-dinitropentane	211.17	11.5	-81.2	811.4	812.4	405.2	3.633	79
343 567	$C_8H_{10}N_2O_4$ $C_8H_{10}N_2O_4$	2,2,3,3-Tetra-nitrotoluene 2,2,3,3-Tetra-nitro-1,5-pentanediol	238.12 234.15	1.0 1.0	-79.2 -77.5	523.0 600.0	531.3 600.0	333.3 446.1	7.76 1.00	71 128-2
653	$C_8H_{10}N_2O_4$	2,2,3,3-Tetra-nitro-1-butanol acetate	296.16	5.0	-32.1	590.1	547.7	124.0	1.5	128-2
555	$C_8H_{10}N_2O_4$	Methyl-2,3,3-Tetra-nitroacrylate	318.16	9.0	-58.4	864.8	850.0	414.4	2.75	128-2
$\chi^2 = 92.83 \pm 51.50n^2$										
								$\chi^2 = 116.915$		
								$\chi^2 = 8237.5$		
								$\chi^2 = 60,269.1678$		
								$\chi^2 = 951.65278$		
								$n = 24$		
179 548	$(C_2H_2NO_2)_n$ $C_6H_6NO_2$	Nitroethylene polymer Tris(hydroxyethyl) nitromethane	121.05 151.12	1.5 4.5	-76.7 -79.4	246.1 609.3	278.4 604.2	-	-	128-3 31a
1676	$C_6H_8NO_4$	2-Nitro-2-methyl-1,1-propanediol	145.12	3.5	-109.7	545.0	556.0	-	-	31a
1771	$C_6H_8NO_4$ $C_6H_8NO_4$	2-Ethyl-2-nitro-1,1-propanediol	144.12 146.15	9.5 11.5	-127.6 -123.4	545.6 709.0	556.0 704.7	-	-	31a 31a
	$C_6H_8NO_4$	2-Nitro-1-propyl-1,1-propanediol	144.12	11.5	-123.4	709.0	704.7	-	-	31a
	$C_6H_8NO_4$	2-Nitro-2-ethyl-1,1-propanediol	146.15	11.5	-123.4	709.0	704.7	-	-	31a
	$C_{10}H_{12}NO_4$	2-Nitro-2-methyl-1-phenyl-1-propanol	195.21	23.5	-192.6	1158.7	1364.5	-	-	31a
		2-Nitro-2-methyl-1-phenyl-1-propanol	"	"	"	1391.5	"	-	-	31a
397	$C_6H_8NO_4$	2,2-Dinitropropene	134.09	5.0	-59.7	427.6	424.3	-	-	65

TABLE 133  
AROMATIC NITRO (solid)

REF. NO.	FORMULA	NAME	MOL. WT.	OXYGEN REQUIRED	OXYGEN BALANCE	G.CAL./MOLE		Y'	X'	REF.
						OBSERVED	CALCULATED			
515	C <sub>6</sub> H <sub>4</sub> NO <sub>2</sub>	2,4-Dinitrobenzene	177.99	10.7	-165.6	671.5	665.1	517.0	15.5	132
		2,6-Dinitrobenzene	177.99	11.7	-137.6	727.7	725.6	522.9	11.5	133
		3,4-Dinitrobenzene	177.99	11.7	-137.6	706.9	706.9	691.7	11.5	134
		Nitrobenzene	123.11	12.5	-167.6	741.7	720.5	711.7	12.5	137
561	C <sub>6</sub> H <sub>5</sub> NO <sub>2</sub>	o-Nitrotoluene	149.11	10.7	-104.7	770.6	771.1	741.5	12.5	136
		m-Nitrotoluene	149.11	10.7	-104.7	770.6	771.1	741.5	12.5	136
562	C <sub>6</sub> H <sub>4</sub> NO <sub>2</sub>	p-Nitrotoluene	149.11	10.7	-104.7	770.6	771.1	741.5	12.5	136
		Nitrotoluene	149.11	10.7	-104.7	770.6	771.1	741.5	12.5	136
560	C <sub>6</sub> H <sub>4</sub> O <sub>2</sub>	o-Nitrobenzoic acid	167.16	17.5	-167.5	827.9	801.2	801.2	15.5	137
		m-Nitrobenzoic acid	167.16	17.5	-167.5	827.9	801.2	801.2	15.5	137
		p-Nitrobenzoic acid	167.16	17.5	-167.5	827.9	801.2	801.2	15.5	137
		2,4-Dinitrobenzoic acid	210.17	19.0	-114.6	915.7	1166.0			56-283

Y' = 135.0  
X' = 755.9  
Y' = 107.65  
X' = 1870.5  
n = 10

TABLE 134  
AROMATIC NITRO (solid)

REF. NO.	FORMULA	NAME	MOL. WT.	OXYGEN REQUIRED	OXYGEN BALANCE	G.CAL./MOLE		Y'	X'	REF.
						OBSERVED	CALCULATED			
529	C <sub>6</sub> H <sub>4</sub> NO <sub>2</sub>	1-Fluoro-2-nitrophenol	157.10	10.5	-106.9	652.5	651.5	615.8	10.5	133
		p-Nitrophenol	139.11	11.5	-132.1	627.1	626.6	637.0	11.5	137
528	C <sub>6</sub> H <sub>5</sub> NO <sub>2</sub>	p-Nitrotoluene	149.11	10.7	-104.7	684.8	684.8	684.1	11.5	137
		o-Nitrotoluene	149.11	10.7	-104.7	684.8	684.8	684.1	11.5	137
527	C <sub>6</sub> H <sub>5</sub> NO <sub>2</sub>	m-Nitrotoluene	149.11	10.7	-104.7	684.8	684.8	684.1	11.5	137
		o-Nitrotoluene	149.11	10.7	-104.7	684.8	684.8	684.1	11.5	137
562	C <sub>6</sub> H <sub>4</sub> NO <sub>2</sub>	p-Nitrobenzoic acid	167.16	17.5	-167.5	827.9	801.2	801.2	15.5	137
		m-Nitrobenzoic acid	167.16	17.5	-167.5	827.9	801.2	801.2	15.5	137
561	C <sub>6</sub> H <sub>5</sub> NO <sub>2</sub>	o-Nitrotoluene	149.11	10.7	-104.7	770.6	771.1	741.5	12.5	136
		m-Nitrotoluene	149.11	10.7	-104.7	770.6	771.1	741.5	12.5	136
560	C <sub>6</sub> H <sub>4</sub> NO <sub>2</sub>	p-Nitrotoluene	149.11	10.7	-104.7	770.6	771.1	741.5	12.5	136
		Nitrotoluene	149.11	10.7	-104.7	770.6	771.1	741.5	12.5	136
572	C <sub>6</sub> H <sub>4</sub> NO <sub>2</sub>	o-Nitrobenzaldehyde	151.12	11.5	-112.9	792.7	774.7	785.9	11.5	137
		m-Nitrobenzaldehyde	151.12	11.5	-112.9	792.7	774.7	785.9	11.5	137
1919	C <sub>6</sub> H <sub>4</sub> NO <sub>2</sub>	p-Nitrobenzaldehyde	151.12	11.5	-112.9	792.7	774.7	785.9	11.5	137
		Nitrobenzaldehyde	151.12	11.5	-112.9	792.7	774.7	785.9	11.5	137
616	C <sub>6</sub> H <sub>4</sub> NO <sub>2</sub>	1-Nitro-2-toluidine	137.13	15.5	-130.9	552.5	627.9	614.0	15.5	139
		2-Nitro-1-toluidine	137.13	15.5	-130.9	552.5	627.9	614.0	15.5	139
660	C <sub>6</sub> H <sub>4</sub> NO <sub>2</sub>	1-Nitro-2-pyridinol	152.15	11.5	-114.1	945.2	932.2	912.3	11.5	137
		2-Nitro-1-pyridinol	152.15	11.5	-114.1	945.2	932.2	912.3	11.5	137
521	C <sub>6</sub> H <sub>4</sub> NO <sub>2</sub>	1-Fluoro-2-nitrobenzotriazole	185.15	16.5	-112.0	980.9	975.1	970.7	16.5	139
		2-Fluoro-1-nitrobenzotriazole	185.15	16.5	-112.0	980.9	975.1	970.7	16.5	139
521	C <sub>6</sub> H <sub>4</sub> NO <sub>2</sub>	o-Nitrophenol	167.16	17.5	-167.5	827.9	801.2	801.2	15.5	137
		m-Nitrophenol	167.16	17.5	-167.5	827.9	801.2	801.2	15.5	137
521	C <sub>6</sub> H <sub>4</sub> NO <sub>2</sub>	o-Nitrobenzoic acid	167.16	17.5	-167.5	827.9	801.2	801.2	15.5	137
		m-Nitrobenzoic acid	167.16	17.5	-167.5	827.9	801.2	801.2	15.5	137
521	C <sub>6</sub> H <sub>4</sub> NO <sub>2</sub>	p-Nitrobenzoic acid	167.16	17.5	-167.5	827.9	801.2	801.2	15.5	137
		Nitrobenzoic acid	167.16	17.5	-167.5	827.9	801.2	801.2	15.5	137
521	C <sub>6</sub> H <sub>4</sub> NO <sub>2</sub>	1-Nitro-2-naphthol	199.20	21.5	-208.2	1028.7	1193.6	1304.4	21.5	139
		2-Nitro-1-naphthol	199.20	21.5	-208.2	1028.7	1193.6	1304.4	21.5	139
71	C <sub>6</sub> H <sub>4</sub> NO <sub>2</sub>	1-Fluoro-2,4-dinitrobenzene	186.10	9.0	-71.1	628.2	660.7	325.9	4.5	133
		2,4-Dinitrophenol	186.10	9.0	-71.1	628.2	660.7	325.9	4.5	133
71	C <sub>6</sub> H <sub>4</sub> NO <sub>2</sub>	2,6-Dinitrophenol	186.10	9.0	-71.1	628.2	660.7	325.9	4.5	133
		2,4-Dinitrophenol	186.10	9.0	-71.1	628.2	660.7	325.9	4.5	133
71	C <sub>6</sub> H <sub>4</sub> NO <sub>2</sub>	2,6-Dinitrophenol	186.10	9.0	-71.1	628.2	660.7	325.9	4.5	133
		2,4-Dinitrophenol	186.10	9.0	-71.1	628.2	660.7	325.9	4.5	133

TABLE 124  
AROMATIC NITROBENZENE (Cont'd)

ADL NO.	FORMULA	NAME	MOL WT	OXYGEN (CALIB)	OXYGEN BALANCE	Q <sub>v</sub> - KCAL/MOLE		Y <sup>a</sup>	X <sup>b</sup>	REF.	
						OBSERVED	ADJUSTED				
147	C <sub>6</sub> H <sub>4</sub> N <sub>2</sub> O <sub>6</sub>	m-Dinitrobenzene	174.11	10.0	-95.2	695.3	697.5	350.0	5.0	110	
		"	"	"	"	"	"	350.5	5.0	15	
		"	"	"	"	"	"	"	351.4	5.0	39
		"	"	"	"	"	"	"	351.6	5.0	9
		"	"	"	"	"	"	"	350.0	5.0	101
		"	"	"	"	"	"	"	352.3	5.0	143
		"	"	"	"	"	"	"	349.3	5.0	23
		"	"	"	"	"	"	"	347.5	5.0	117
		"	"	"	"	"	"	"	353.5	5.0	15
		"	"	"	"	"	"	"	353.4	5.0	37
		"	"	"	"	"	"	"	347.2	5.0	13
148	C <sub>7</sub> H <sub>5</sub> N <sub>2</sub> O <sub>6</sub>	2,4,6-Trinitroanisole	191.13	12.0	-96.9	813.7	814.8	401.4	6.0	23	
		"	"	"	"	"	"	401.5	6.0	10	
24	C <sub>6</sub> H <sub>4</sub> N <sub>2</sub> O <sub>6</sub>	2,4,6-Trinitrotoluene	187.13	11.7	-111.2	817.1	822.3	425.1	5.5	23	
		"	"	"	"	"	"	425.1	6.5	4	
150	C <sub>8</sub> H <sub>7</sub> N <sub>2</sub> O <sub>6</sub>	N-Methyl-2,4-dinitroaniline	197.15	11.5	-139.6	947.2	945.2	469.0	5.75	19	
		2,4,6-Trinitroacetole	212.16	15.0	-133.1	972.4	972.0	489.1	7.5	19	
66	C <sub>7</sub> H <sub>5</sub> N <sub>2</sub> O <sub>6</sub>	2,4,6-Trinitro-m-xylene	186.16	14.6	-125.5	875.2	874.7	434.9	6.0	23	
		"	"	"	"	"	"	434.9	6.0	8	
64	C <sub>10</sub> H <sub>13</sub> N <sub>2</sub> O <sub>6</sub>	2-(2,4,6-Trinitrophenyl)-2-hydroxyethyl-1,3-propanediol	287.23	19.1	-104.6	1222.7	1211.6	591.5	9.75	85	
603	C <sub>10</sub> H <sub>13</sub> N <sub>2</sub> O <sub>6</sub>	N-(2,4,6-Trinitrophenyl)-diethanolamine	271.23	20.5	-123.9	1274.7	1273.2	619.2	10.25	85	
1136	C <sub>10</sub> H <sub>13</sub> N <sub>2</sub> O <sub>6</sub>	2-(2,4,6-Trinitrophenyl)-2-methyl-1,3-propanediol	271.23	"	-120.2	1266.9	1273.9	614.4	10.25	85	
216	C <sub>6</sub> H <sub>3</sub> N <sub>3</sub> O <sub>7</sub>	2,3,5-Trinitrophenol	229.14	6.5	-45.1	616.0	611.3	304.6	2.167	23	
		"	"	"	"	"	"	304.6	2.167	110	
		"	"	"	"	"	"	"	231.7	2.157	110
		"	"	"	"	"	"	"	231.7	1.5	15
		"	"	"	"	"	"	"	231.6	2.167	36
274	C <sub>6</sub> H <sub>3</sub> N <sub>3</sub> O <sub>7</sub>	1,3,5-Trinitrobenzene	213.11	7.5	-56.3	651.7	660.3	269.5	2.59	143	
277	C <sub>6</sub> H <sub>3</sub> N <sub>3</sub> O <sub>7</sub>	1,3,5-Trinitrobenzene	213.11	"	"	657.0	"	225.6	2.59	23	
		"	"	"	"	"	"	221.1	2.50	35	
		"	"	"	"	"	"	221.1	2.833	110	
277	C <sub>6</sub> H <sub>3</sub> N <sub>3</sub> O <sub>7</sub>	2,4,6-Trinitrobenzoic acid	241.12	4.5	-55.1	727.1	727.7	240.7	7.833	110	
		2,3,5-Trinitro-m-cresol	243.13	9.5	-62.5	764.9	768.2	286.1	3.167	10	
277	C <sub>6</sub> H <sub>3</sub> N <sub>3</sub> O <sub>7</sub>	2,4,6-Trinitro-m-xylene	243.13	"	-62.5	764.9	"	287.2	3.167	23	
		"	"	"	"	"	"	287.2	3.167	110	
113	C <sub>6</sub> H <sub>3</sub> N <sub>3</sub> O <sub>7</sub>	2,4,6-Trinitrotoluene	227.13	10.5	-71.0	815.0	815.6	375.1	3.50	143	
		"	"	"	"	"	"	375.1	3.50	110	
		"	"	"	"	"	"	"	375.1	3.50	23
		"	"	"	"	"	"	"	375.1	3.50	86
		"	"	"	"	"	"	"	375.1	3.50	5
292	C <sub>7</sub> H <sub>5</sub> N <sub>3</sub> O <sub>7</sub>	N-Methyl-2,4,6-trinitroaniline	212.15	11.0	-72.7	851.3	851.0	375.1	3.50	37	
294	C <sub>7</sub> H <sub>5</sub> N <sub>3</sub> O <sub>7</sub>	2,4,6-Trinitrophenoxide	257.16	12.5	-75.1	913.7	915.0	375.1	3.50	10	
		"	"	"	"	"	"	375.1	3.50	143	
		"	"	"	"	"	"	"	375.1	3.50	23
318	C <sub>7</sub> H <sub>5</sub> N <sub>3</sub> O <sub>7</sub>	2,4,6-Trinitro-m-xylene	241.12	11.5	-69.6	937.7	968.5	375.1	4.50	6	
		"	"	"	"	"	"	375.1	4.50	110	
		"	"	"	"	"	"	"	375.1	4.50	110
		"	"	"	"	"	"	"	375.1	4.50	23
		"	"	"	"	"	"	"	375.1	4.50	7
325	C <sub>8</sub> H <sub>7</sub> N <sub>3</sub> O <sub>7</sub>	Allyl picrate	259.17	14.5	-56.2	1066.2	1072.5	375.1	4.50	71	
324	C <sub>8</sub> H <sub>7</sub> N <sub>3</sub> O <sub>7</sub>	Propyl picrate	271.17	15.5	-91.4	1131.5	1090.1	375.1	5.167	71	
331	C <sub>8</sub> H <sub>7</sub> N <sub>3</sub> O <sub>7</sub>	2,3,5-Trinitroaniline	271.12	5.5	-32.2	875.0	873.7	375.1	4.375	110	
325	C <sub>7</sub> H <sub>5</sub> N <sub>3</sub> O <sub>7</sub>	2,2',3,3',5,5',6'-Hexanitrodiphenylene	439.22	14.5	-52.6	1877	1321.7	219.5	2.167	58	
		"	"	"	"	"	"	219.5	2.157	110	
		"	"	"	"	"	"	"	219.9	2.167	85
140	C <sub>10</sub> H <sub>6</sub> N <sub>2</sub> O <sub>2</sub>	1-Nitronaphthalene	174.15	71.5	-193.7	1198.5	1190.2	1201.6	21.5	110	
		1-Nitronaphthalene	"	"	"	"	"	1201.2	21.5	7	
141	C <sub>10</sub> H <sub>6</sub> N <sub>2</sub> O <sub>2</sub>	1,5-Dinitronaphthalene	218.16	19.0	-137.3	1156.6	1156.8	261.1	7.5	110	
		"	"	"	"	"	"	261.1	7.5	7	
		"	"	"	"	"	"	"	261.5	7.5	143
		"	"	"	"	"	"	"	261.3	7.5	110
142	C <sub>10</sub> H <sub>6</sub> N <sub>2</sub> O <sub>2</sub>	1,6-Dinitronaphthalene	"	"	"	1156.6	"	261.1	7.5	7	
		"	"	"	"	"	"	261.0	7.5	7	
		"	"	"	"	"	"	"	261.8	7.5	143
		"	"	"	"	"	"	"	261.8	7.5	143
143	C <sub>12</sub> H <sub>8</sub> N <sub>2</sub> O <sub>2</sub>	1,2,5-Trinitronaphthalene	261.16	16.5	-100.3	1117.9	1117.1	375.0	5.5	7	
		"	"	"	"	"	"	375.0	5.5	8	
		"	"	"	"	"	"	"	375.1	5.5	110
144	"	1,2,5-Trinitronaphthalene	"	"	"	1126.3	"	5.5	7		

TABLE I  
AROMATIC NITRO (solid) (Cont'd)

ADL NO	FORMULA	NAME	MOL WT	OXYGEN REQUIRED	OXYGEN BALANCE	$\Delta H_c^\circ$ OF FORMED	KCAL/MOLE CALCULATED	$\Delta V$	$\Delta X$	REF.
90	$C_{10}H_7N_2O_2$	1,3,4,5-Tetraamino-2-naphthalene $\gamma$ = 11.1, $\beta$ = 11.5 <sup>a</sup>	186.15	14.0	- 22.7	1082.3	1003.7	272.9	3.5	85
										$\Delta H_c^\circ$ = 1284.113 $\Delta V^\circ$ = 62,310.2 $\Delta X^\circ$ = 796,817.5372 $\Delta H_c^\circ$ = 13,995.58227 n = 113
92	$C_6H_5NO_2$	p-Nitrophenol	123.11	11.5	2.3	622.7	636.6	-	-	110
	$C_6H_5NO_2$	o-Nitrobenzoic acid	142.12	17.5	-	735.0	738.7	-	-	85-162
97	$C_9H_8N_2O_2$	m-Nitroaniline	135.12	13.0	-	764.1	763.3	-	-	110
	$C_9H_8N_2O_2$	p-Nitroacetophenone	174.14	15.0	-1.6	857.4	950.7	-	-	85-226
98	$C_9H_7N_2O_2$	p-Nitroacetophenone	174.14	15.0	-1.6	822.6	926.9	-	-	85-226
99	$C_9H_7N_2O_2$	m-Nitroacetophenone	174.14	15.0	-1.6	874.1	769.2	-	-	117
100	$C_9H_7N_2O_2$	m-Nitroacetophenone	174.14	15.0	-1.6	909.6	804.7	-	-	117
101	$C_9H_7N_2O_2$	p-Nitroacetophenone	174.14	15.0	-1.6	992.2	887.3	-	-	117
	$C_{12}H_9NO_2$	m-Nitroethylbenzoin-	226.26	17.5	-176.7	1979.9	1551.0	-	-	12
	$C_{12}H_9NO_2$	m-Nitrobenzoinbenzoic	260.27	17.5	-180.3	2686.3	1812.4	-	-	12
	$C_{12}H_9NO_2$	m-Nitrobenzoinbenzoic	260.27	17.5	-180.3	2686.3	1812.4	-	-	12
104	$C_{10}H_7N_2O_2$	4,4'-Dinitrostilbene, cis isomer	270.25	22.0	-171.7	1675.3	920.3	-	-	4
	$C_{10}H_7N_2O_2$	4,4'-Dinitrostilbene, trans	270.25	22.0	-171.7	1670.7	920.3	-	-	4
	$C_{10}H_7N_2O_2$	Trinitrophenylmethane	375.32	37.5	-171.0	2216.7	2216.0	-	-	85-179
	$C_{10}H_7N_2O_2$	Trinitrophenylmethane	379.32	38.5	-162.5	2271.5	2216.3	-	-	85-179
	$C_9H_7N_2O_2$	2-Amino-4-nitrotoluene	174.12	9.5	- 76.3	676.7	677.4	-	-	110
106	$C_8H_6F_2N_2O_2$	2,4-Dinitrofluoro-3,5-difluorobenzene	235.11	10.5	- 67.8	599.2	719.2	-	-	118
	$C_8H_6F_2N_2O_2$	2,6-Dinitrofluoro-3,5-difluorobenzene	235.11	10.5	- 67.8	719.2	599.2	-	-	118
	$C_8H_6F_2N_2O_2$	2,4-Dinitrotoluene	182.13	11.5	-124.2	853.5	852.5	-	-	39
	$C_8H_6F_2N_2O_2$	2,6-Dinitrotoluene	182.13	11.5	-124.2	853.5	852.5	-	-	39
	$C_8H_6F_2N_2O_2$	2,4-Dinitrotoluene	182.13	11.5	-124.2	853.5	852.5	-	-	39
	$C_8H_6F_2N_2O_2$	2,6-Dinitrotoluene	182.13	11.5	-124.2	853.5	852.5	-	-	39
	$C_8H_6F_2N_2O_2$	2,4-Dinitrotoluene	182.13	11.5	-124.2	853.5	852.5	-	-	39
	$C_8H_6F_2N_2O_2$	2,6-Dinitrotoluene	182.13	11.5	-124.2	853.5	852.5	-	-	39
111	$C_{11}H_9NO_2$	m-Nitroacetophenone	210.19	19.0	-144.6	1375.7	1161.3	-	-	85-243
108	$C_8H_6N_2O_2$	m-(4,4'-Dinitrostilbene)- ethanolamine	270.24	22.0	-131.3	1312.6	1369.7	-	-	85
120	$C_8H_6N_2O_2$	2,4,6-Trinitroacetophenone	228.11	9.5	- 85.2	888.8	888.9	-	-	118
	$C_8H_6N_2O_2$	2,4,6-Trinitrophenol	229.11	9.5	- 85.4	871.5	871.8	-	-	118
	$C_8H_6N_2O_2$	2,4,6-Trinitrophenol	229.11	9.5	- 85.4	871.5	871.8	-	-	118
	$C_8H_6N_2O_2$	2,4,6-Trinitrophenol	229.11	9.5	- 85.4	871.5	871.8	-	-	117
	$C_8H_6N_2O_2$	2,4,6-Trinitrophenol	229.11	9.5	- 85.4	871.5	871.8	-	-	117
116	$C_7H_5NO_2$	1,2,4-Trinitrobenzene	153.07	7.5	- 66.9	661.8	660.3	-	-	37
	$C_7H_5NO_2$	1,3,5-Trinitrobenzene	153.07	7.5	- 66.9	675.1	675.1	-	-	119
	$C_7H_5NO_2$	1,2,4-Trinitrobenzene	153.07	7.5	- 66.9	674.0	674.0	-	-	37
113	$C_7H_5NO_2$	2,4,6-Trinitroaniline	229.12	9.5	- 85.1	671.2	669.9	-	-	110
117	$C_8H_7NO_2$	2,4,6-Trinitro-o-cresol	213.13	9.5	- 62.5	779.5	786.2	-	-	113
111	$C_8H_7NO_2$	2,4,6-Trinitro-p-cresol	213.13	9.5	- 62.5	779.5	786.2	-	-	113
112	$C_8H_7NO_2$	2,4,6-Trinitro-m-cresol	213.13	9.5	- 62.5	779.5	786.2	-	-	113
	$C_8H_7NO_2$	2,4,6-Trinitro-m-cresol	213.13	9.5	- 62.5	779.5	786.2	-	-	113
113	$C_8H_7NO_2$	2,4,6-Trinitrotoluene	227.13	10.0	- 71.0	821.9	818.6	-	-	43
	$C_8H_7NO_2$	2,4,6-Trinitrotoluene	227.13	10.0	- 71.0	821.9	818.6	-	-	43
	$C_8H_7NO_2$	2,4,6-Trinitrotoluene	227.13	10.0	- 71.0	821.9	818.6	-	-	113
114	$C_8H_7NO_2$	2,4,6-Trinitrotoluene	227.13	10.0	- 71.0	821.9	818.6	-	-	39
110	$C_8H_7NO_2$	2,4,6-Trinitrotoluene	227.13	10.0	- 71.0	821.9	818.6	-	-	39
111	$C_8H_7NO_2$	2,4,6-Trinitrotoluene	227.13	10.0	- 71.0	821.9	818.6	-	-	39
112	$C_8H_7NO_2$	2,4,6-Trinitrotoluene	227.13	10.0	- 71.0	821.9	818.6	-	-	37
117	$C_8H_7NO_2$	2,4,6-Trinitrotoluene	227.13	10.0	- 71.0	821.9	818.6	-	-	37
118	$C_8H_7NO_2$	2,4,6-Trinitrotoluene	227.13	10.0	- 71.0	821.9	818.6	-	-	37
	$C_8H_7NO_2$	2-Propenyl picrate	267.15	13.5	-	1052.2	1051.9	-	-	71
108	$C_8H_7NO_2$	2,3,4,5,6-Pentanitroaniline	318.12	9.0	- 15.1	682.4	615.9	-	-	71

<sup>a</sup> Ref. 39 gives 1242.5 kcal/mole not a mol. wt. of 60.3. Since the letter is incorrect the heat of combustion was recalculated from the value given, using the proper mol. wt.



CONFIDENTIAL

TABLE 125  
ALTERNATIVE 6-6-TRINITRO (4-114)

ADL NO.	FORMULA	NAME	MOL. WT.	OXYGEN REQUIRED	OXYGEN BALANCE	Q <sub>v</sub> - KCAL/MOLE		Y'	X'	REF.
						OBSERVED	CALCULATED			
377	C <sub>3</sub> H <sub>5</sub> N <sub>3</sub> O <sub>9</sub>	Trinitroethane (Nitroform)	151.04	-5	+ 32.1	109.8	116.9	196.9	-1.5	16
						119.5	111.6		-1.5	31
						122.2		126.5	-1.5	71
574	C <sub>11</sub> H <sub>13</sub> N <sub>3</sub> O <sub>9</sub>	1,1,1-Trinitroethanol acrylate	235.11	+6.5	- 36.6	561.1	502.3	512	+6.5	128-1
579	C <sub>11</sub> H <sub>13</sub> N <sub>3</sub> O <sub>9</sub>	Methyl 1,1,1-trinitrobutyrate	217.13	5.5	- 32.1	517.3	566.7	575.1		128-1
		1,1,1-Trinitro-2-propenol acetate				581.4	566.8	581.3	5.5	128-1
498	C <sub>11</sub> H <sub>13</sub> N <sub>3</sub> O <sub>9</sub>	Vinyl 1,1,1-trinitrobutyrate	210.14		- 32.1	526.5	516.1	513.7	5.5	128-1
616	C <sub>11</sub> H <sub>13</sub> N <sub>3</sub> O <sub>9</sub>	Methyl 1,1,1-trinitrobutyrate isobutyrate	207.16	4.5	- 32.1	511.5	552.0	511.4	4.5	128-1
653	C <sub>11</sub> H <sub>13</sub> N <sub>3</sub> O <sub>9</sub>	2,3,3,4,5-Pentamethyl-1,1,1,1-tetraacetotetroxane	279.21	13.5	- 61.1	1076.1	1054.7	1051.3	13.5	128-1
			336.30	24.0	-116.2	1631.8	1611.5	1616.5	24.0	129-1
								1 x' = 59.5		
								2 x' = 601.0		
								2 x' = 71157.75		
								2 x' = 1037.25		
								n = 11		
210	C <sub>4</sub> H <sub>6</sub> N <sub>2</sub> O <sub>6</sub>	Tetra-nitroethane	196.08	-6.0	+ 30.2	102.4	76.7			116-1
						107.1				61

TABLE 126  
ALTERNATIVE 6-6-TRINITRO (4-114)

ADL NO.	FORMULA	NAME	MOL. WT.	OXYGEN REQUIRED	OXYGEN BALANCE	Q <sub>v</sub> - KCAL/MOLE		Y'	X'	REF.
						OBSERVED	CALCULATED			
340	C <sub>8</sub> H <sub>9</sub> N <sub>3</sub> O <sub>9</sub>	2,2,2-Trinitropropanol	181.07	1.5	+ 33.1	227.0	267.0	212.1	-1.5	120-1
567	C <sub>8</sub> H <sub>9</sub> N <sub>3</sub> O <sub>9</sub>	1,1,1-Trinitroethane	151.04	-0.5	+ 32.0	261.5	288.5	237.6	-0.5	72-1
576	C <sub>8</sub> H <sub>9</sub> N <sub>3</sub> O <sub>9</sub>	1,1,1-Trinitropropene	154.09	0.0	+ 32.0	280.2	354.7	281.7	0.0	126-1
639	C <sub>8</sub> H <sub>9</sub> N <sub>3</sub> O <sub>9</sub>	1,1,1-Trinitrobutyramide	222.12	+1.0	- 26.0	371.8	371.1	375.1	+1.0	128-1
577	C <sub>8</sub> H <sub>9</sub> N <sub>3</sub> O <sub>9</sub>	5,5,5-Trinitro-2-pentanone	221.13	6.5	- 67.1	632.5	634.5	634.5	6.5	120-1
666	C <sub>8</sub> H <sub>9</sub> N <sub>3</sub> O <sub>9</sub>	2,2,2-Trinitroethanol acrylate	220.11	7.5	- 69.1	121.0	101.7	67.5	7.5	128-1
523	C <sub>8</sub> H <sub>9</sub> N <sub>3</sub> O <sub>9</sub>	1,1,1,1,5,5,5-Heptanitro-pentane	387.14	-1.5	+ 6.2	595.5	566.3	239.5	-1.5	128-1
331	C <sub>8</sub> H <sub>9</sub> N <sub>3</sub> O <sub>9</sub>	1,3-Bis(2,2,2-trinitroethyl)-propane	386.16	0.0	0.0	692.5	682.9		0.0	71
110	C <sub>8</sub> H <sub>9</sub> N <sub>3</sub> O <sub>9</sub>	2,2,2-Trinitroethanol	181.07	1.5	+ 33.1	227.0	267.0	212.1	1.5	71
390	C <sub>8</sub> H <sub>9</sub> N <sub>3</sub> O <sub>9</sub>	1,1,1-Trinitrobutyramide (2,2,2-Trinitroethyl)-1,3-bis(2,2,2-trinitroethyl)propane	406.17	2.5	- 16.2	792.5	723.5	392.5	2.5	123-1
								2 x' = 15.5		
								2 x' = 3711.9		
								2 x' = 10,761.75		
								2 x' = 121.625		
								n = 11		
266	C <sub>8</sub> H <sub>9</sub> N <sub>3</sub> O <sub>9</sub>	2-Nitroacetone nitrile	176.05	+2.0	+ 13.2	236.1	187.1			71
567	C <sub>8</sub> H <sub>9</sub> N <sub>3</sub> O <sub>9</sub>	1,1,1-Trinitroethane	151.04	-0.5	+ 32.0	261.5	288.5	237.6	-0.5	117
582	C <sub>8</sub> H <sub>9</sub> N <sub>3</sub> O <sub>9</sub>	4,4-Dimethyl-5,5,5-trinitro-2-pentanone	219.15	-12.5	- 59.2	973.5	973.4			129-1
481	C <sub>8</sub> H <sub>9</sub> N <sub>3</sub> O <sub>9</sub>	Diethyl(2,2,2-trinitroethyl)malonate	321.22	11.5	- 71.6	1162.0	1163.0			123-1
331	C <sub>8</sub> H <sub>9</sub> N <sub>3</sub> O <sub>9</sub>	1,3-Bis(2,2,2-trinitroethyl)-propane	387.16	0.0	0.0	611.3	601.9			124-1
581	C <sub>8</sub> H <sub>9</sub> N <sub>3</sub> O <sub>9</sub>	1,3-Bis(2,2,2-trinitroethyl)-propane dinitroethane	431.23	2.0	- 12.6	874.1	873.0			124-1
581	C <sub>8</sub> H <sub>9</sub> N <sub>3</sub> O <sub>9</sub>	4,4-Dimethyl-5,5,5-trinitro-2-pentanone	219.15	11.5	- 51.7	973.5	973.5			128-1

CONFIDENTIAL

TABLE 127  
NITRATES (liquid)

ADL NO	FORMULA	NAME	MOL WT.	OXYGEN REQUIRED	OXYGEN BALANCE	O <sub>2</sub> - KCAL/MOLE		Y'	X'	REF.
						OBSERVED	CALCULATED			
647	C <sub>8</sub> H <sub>16</sub> N <sub>2</sub> O <sub>4</sub>	Diethyl 3-hydroxyglutarate nitrate	270.22	18.5	-115.8	1327.1	1126.2	1097.5	18.5	128-1
94	C <sub>8</sub> H <sub>14</sub> N <sub>2</sub> O <sub>4</sub>	1,2-Ethandiol dinitrate	192.07	6.0	0.0	268.5	268.5	130.4	0.0	131
37	C <sub>8</sub> H <sub>14</sub> N <sub>2</sub> O <sub>4</sub>	Dimethylene glycol dinitrate	196.12	5.0	-40.9	527.6	513.9	263.0	2.5	95
539	C <sub>8</sub> H <sub>16</sub> N <sub>2</sub> O <sub>4</sub>	1,3-Butanediol dinitrate	188.12	6.0	-52.3	573.6	573.6	283.0	3.0	128-1
157	C <sub>8</sub> H <sub>16</sub> N <sub>2</sub> O <sub>4</sub>	1,3-Butanediol dinitrate	"	"	"	573.6	"	283.0	3.0	128-1
647	C <sub>8</sub> H <sub>16</sub> N <sub>2</sub> O <sub>4</sub>	1,3-Butanediol dinitrate	"	"	"	573.6	"	283.0	3.0	128-1
157	C <sub>8</sub> H <sub>16</sub> N <sub>2</sub> O <sub>4</sub>	1,3-Butanediol dinitrate	"	"	"	573.6	"	283.0	3.0	128-1
532	C <sub>8</sub> H <sub>16</sub> N <sub>2</sub> O <sub>4</sub>	1,5-Pentanediol dinitrate	198.15	7.0	-38.2	732.6	732.6	363.5	4.5	128-1
173	C <sub>8</sub> H <sub>16</sub> N <sub>2</sub> O <sub>4</sub>	2,4-Pentanediol dinitrate	"	"	"	732.6	"	363.5	4.5	128-1
173	C <sub>8</sub> H <sub>16</sub> N <sub>2</sub> O <sub>4</sub>	Tetraethylene glycol dinitrate	280.17	10.0	-66.6	835.4	819.2	409.3	5.0	95
634	C <sub>8</sub> H <sub>12</sub> N <sub>2</sub> O <sub>4</sub>	1,4-Cyclohexanediol dinitrate	280.17	10.0	-92.2	870.1	892.5	436.2	6.0	128-1
972	C <sub>8</sub> H <sub>12</sub> N <sub>2</sub> O <sub>4</sub>	Diethyl carbonate dinitrate	294.17	"	-64.8	937.4	920.0	462.4	6.0	95
151	C <sub>8</sub> H <sub>14</sub> N <sub>2</sub> O <sub>4</sub>	Nitroglycerin	227.09	-0.5	1.5	363.2	357.6	178.8	-0.167	95
"	"	"	"	"	"	363.8	"	179.4	-0.167	151
"	"	"	"	"	"	365.7	"	180.0	-0.167	23
"	"	"	"	"	"	363.3	"	180.9	-0.167	106
"	"	"	"	"	"	365.5	"	179.7	-0.167	122
"	"	"	"	"	"	367.2	"	182.4	-0.167	93
165	C <sub>8</sub> H <sub>18</sub> N <sub>2</sub> O <sub>4</sub>	1,2,3-Butanetriol trinitrate	218.12	9.5	-137.6	517.5	524.7	270.6	0.833	95
157	C <sub>8</sub> H <sub>18</sub> N <sub>2</sub> O <sub>4</sub>	1,2,4-Butanetriol trinitrate	"	"	"	517.5	"	270.6	0.833	95
157	C <sub>8</sub> H <sub>18</sub> N <sub>2</sub> O <sub>4</sub>	2-Methyl-1,2,3-propanetriol trinitrate	"	"	"	517.5	"	270.6	0.833	95
157	C <sub>8</sub> H <sub>18</sub> N <sub>2</sub> O <sub>4</sub>	1,2-Diethoxyethoxy-1,2-propanediol trinitrate	271.15	4.5	-26.6	430.9	615.1	299.5	1.5	95
645	C <sub>8</sub> H <sub>18</sub> N <sub>2</sub> O <sub>4</sub>	1,1,2-Propanetriol trinitrate	255.15	5.5	-34.7	650.9	651.3	325.6	1.5	128-1
764	C <sub>8</sub> H <sub>18</sub> N <sub>2</sub> O <sub>4</sub>	2-Hydroxyethyl-2-methyl-1,1-propanediol trinitrate	"	"	"	671.9	676.3	335.9	1.833	23
111	C <sub>8</sub> H <sub>18</sub> N <sub>2</sub> O <sub>4</sub>	1-Monoethoxy-2-(2-ethoxyethyl)-2-hydroxyethyl-1,1,1-propanediol	313.23	12.5	-55.3	1097.7	1375.4	687.7	4.167	128-1
237	C <sub>8</sub> H <sub>18</sub> N <sub>2</sub> O <sub>4</sub>	2-(Hydroxyethyl)-1,1,1-propanediol	313.22	12.5	-60.7	1107.2	1314.1	662.7	4.5	6
237	C <sub>8</sub> H <sub>18</sub> N <sub>2</sub> O <sub>4</sub>	1,1,3-Propanetriol-2-propanediol tetranitrate	346.17	14.0	-13.7	752.1	760.5	380.7	4.0	95
								$y' = 129.39 + 52.35x'$ $\sum x' = 56.629$ $\sum y' = 526.1$ $\sum x'y' = 63,144.2306$ $\sum x'^2 = 609.444335$ $n = 33$		
150	C <sub>8</sub> H <sub>14</sub> N <sub>2</sub> O <sub>4</sub>	2-Nitroethanol nitrate	192.07	1.0	-1.8	320.1	275.1	"	"	95
160	C <sub>8</sub> H <sub>14</sub> N <sub>2</sub> O <sub>4</sub>	Ethyl nitrate	160.07	3.5	-61.5	326.4	"	"	"	15
"	"	"	"	"	"	322.9	"	"	"	24-273
73	C <sub>8</sub> H <sub>16</sub> N <sub>2</sub> O <sub>4</sub>	1,2,6-Hexanetriol trinitrate	271.19	9.5	-35.0	616.4	614.4	307.2	"	128-1
411	C <sub>8</sub> H <sub>16</sub> N <sub>2</sub> O <sub>4</sub>	All glycerol trinitrate polymer	311.21	12.5	-63.3	934.8	1044.7	"	"	6
527	C <sub>8</sub> H <sub>16</sub> N <sub>2</sub> O <sub>4</sub>	N-Hydroxydiethanolamine dinitrate	266.31	13.0	-51.7	1130.7	1223.8	"	"	95
527	C <sub>8</sub> H <sub>16</sub> N <sub>2</sub> O <sub>4</sub>	2,2-Bis(2-hydroxyethyl)-1,1-propanediol dinitrate	195.27	22.0	-112.9	1517.9	1476.1	"	"	128-1

TABLE 12  
HEAT OF COMBUSTION

APL NO.	FORMULA	NAME	MOL. WT.	OXYGEN REQ (H <sub>2</sub> O)	OXYGEN BALANCE	CAL/MOLE		Y*	X*	RLF.
						OBSERVED	(CALCULATED)			
322	C <sub>4</sub> H <sub>7</sub> N <sub>3</sub> O <sub>6</sub>	Vinyl nitrate polymer	69.06	2.5	47.9	262.5	265.6	750.1	2.0	123.1
326	C <sub>8</sub> H <sub>11</sub> N <sub>3</sub> O <sub>6</sub>	2,4,6-Trinitrobenzyl nitrate	258.13	7.0	44.9	749.1	766.6	499.1	3.2	65
327	C <sub>8</sub> H <sub>9</sub> N <sub>3</sub> O <sub>6</sub>	2-(2,4,6-Trinitrophenyl)-ethanol nitrate	314.34	9.0	50.3	850.9	850.0	570.0	3.0	124
411	C <sub>8</sub> H <sub>9</sub> N <sub>3</sub> O <sub>6</sub>	1,2-Propanediol 2-nitrate 1-(2-nitroacetate)	260.16	12.0	73.9	877.0	890.2	766.5	12.3	35
168	C <sub>8</sub> H <sub>13</sub> N <sub>3</sub> O <sub>6</sub>	2-Methyl-2-nitro-1,3-propanediol dinitrate	275.17	10.5	80.9	931.1	931.8	219.3	1.75	40
43	C <sub>8</sub> H <sub>13</sub> N <sub>3</sub> O <sub>6</sub>	2,2-Dimethyl-1,3-propanediol dinitrate	194.15	9.0	74.9	715.9	731.3	346.6	3.7	66
74	C <sub>8</sub> H <sub>13</sub> N <sub>3</sub> O <sub>6</sub>	2-(3,5-Dinitrophenyl)-2-nitro-1,3-propanediol dinitrate	377.19	9.5	60.3	1022.9	1019.9	179.1	4.75	35
139	C <sub>8</sub> H <sub>13</sub> N <sub>3</sub> O <sub>6</sub>	2-Nitro-2-(o-nitrobenzyl)-1,3-propanediol dinitrate	332.19	12.0	77.4	1071.9	1076.7	621.9	6.0	80
255	C <sub>8</sub> H <sub>13</sub> N <sub>3</sub> O <sub>6</sub>	2-Ethyl-2-(hydroxymethyl)-1,3-propanediol trinitrate	269.17	6.5	60.5	825.5	830.4	274.2	1.033	66
93	C <sub>8</sub> H <sub>13</sub> N <sub>3</sub> O <sub>6</sub>	Erythritol tetrinitrate	300.11	-1.0	5.3	621.3	620.9	110.5	-0.25	124
227	C <sub>8</sub> H <sub>13</sub> N <sub>3</sub> O <sub>6</sub>	Pentaerythritol tetrinitrate	316.15	+2.0	-10.1	615.1	620.6	153.6	+0.5	71
"	"	"	"	"	"	617.5	"	252.0	0.6	23
"	"	"	"	"	"	636.9	"	150.7	0.5	113
231	C <sub>10</sub> H <sub>14</sub> N <sub>4</sub> O <sub>10</sub>	2,2,3,5-Tetraakis(hydroxymethyl)pyroglutazono tetrinitrate	341.22	11.0	65.8	1072.4	1093.1	271.1	2.7	123
"	"	"	"	"	"	1077.1	"	279.0	4.75	38
229	C <sub>10</sub> H <sub>14</sub> N <sub>4</sub> O <sub>10</sub>	2,2,3,5-Tetraakis(hydroxymethyl)cyclohexanono tetrinitrate	359.24	14.0	56.7	1272.7	1251.1	312.9	1.5	124
"	"	"	"	"	"	1210.7	"	194.9	3.6	66
623	C <sub>8</sub> H <sub>13</sub> N <sub>3</sub> O <sub>6</sub>	1-[2,2,2-Tris(hydroxymethyl)ethyl]-1,2-propanediol pentanitrate	339.22	6.5	21.9	1018.9	1020.0	199.2	1.3	124-1
218	C <sub>8</sub> H <sub>13</sub> N <sub>3</sub> O <sub>6</sub>	Tetraakis(2-hydroxyethyl)arsonium pentanitrate	146.26	9.0	30.0	1160.6	1128.6	226.7	1.8	71
230	C <sub>8</sub> H <sub>13</sub> N <sub>3</sub> O <sub>6</sub>	2-Hydroxy-1,1,1,3-cyclohexanetetramethanol pentanitrate	331.23	7.5	56.2	1135.5	1126.0	233.0	1.9	124
228	C <sub>8</sub> H <sub>13</sub> N <sub>3</sub> O <sub>6</sub>	2-Hydroxy-1,1,1,3-cyclohexanetetramethanol pentanitrate	346.26	12.4	41.9	1260.9	1247.0	250.7	-0.5	124
92	C <sub>8</sub> H <sub>13</sub> N <sub>3</sub> O <sub>6</sub>	Dulcitol hexanitrate	352.17	-2.0	7.1	670.9	670.9	110.9	-0.333	67
133	"	Sorbitol hexanitrate	"	"	"	690.5	"	140.5	-0.133	124
152	"	Hamitol hexanitrate	"	"	"	677.1	"	116.8	-0.333	124
"	"	"	"	"	"	691.9	"	112.7	-0.333	124-1
"	"	"	"	"	"	678.9	"	111.5	-0.333	65
"	"	"	"	"	"	714.7	"	116.0	-0.333	73
69	C <sub>10</sub> H <sub>14</sub> N <sub>4</sub> O <sub>10</sub>	1-Pentaerythritol hexanitrate	5	+2.0	-27.5	1271.1	1262.1	206.9	-1.5	124
"	"	"	"	"	"	1250.7	"	206.1	1.5	86
239	C <sub>10</sub> H <sub>14</sub> N <sub>4</sub> O <sub>10</sub>	1,2,3,4,5-Pentakis(hydroxymethyl)hexanitrate	424.26	12.0	35.3	1396.1	1195.7	233.7	2.0	71
"	"	"	"	"	"	"	"	74.35	"	"
"	"	"	"	"	"	"	"	781.40	"	"
"	"	"	"	"	"	"	"	32,465,5912	"	"
"	"	"	"	"	"	"	"	131,666.22	"	"
"	"	"	"	"	"	"	"	31.0	"	"
54	C <sub>8</sub> H <sub>13</sub> N <sub>3</sub> O <sub>6</sub>	Ethanolamine dinitrate	367.13	1.5	14.2	317.5	317.4	-	-	124
323	C <sub>8</sub> H <sub>13</sub> N <sub>3</sub> O <sub>6</sub>	Vinyl nitrate polymer	69.06	2.5	46.5	253.9	265.6	-	-	124-1
75	C <sub>8</sub> H <sub>13</sub> N <sub>3</sub> O <sub>6</sub>	2-(2,4,6-Trinitrobenzyl)-ethanol nitrate	273.16	11.5	67.4	941.6	945.3	-	-	124
604	C <sub>8</sub> H <sub>13</sub> N <sub>3</sub> O <sub>6</sub>	1-(o-Nitrophenyl)-1,2-ethanediol dinitrate	273.15	"	67.4	930.9	945.7	-	-	35
645	C <sub>8</sub> H <sub>13</sub> N <sub>3</sub> O <sub>6</sub>	2,5-Hexanediol dinitrate	208.17	12.0	92.2	871.1	873.9	-	-	124-1
228	C <sub>8</sub> H <sub>13</sub> N <sub>3</sub> O <sub>6</sub>	2-Hydroxy-1,1,1,3-cyclohexanetetramethanol pentanitrate	346.26	12.4	41.9	1260.7	1277.7	-	-	4
133	C <sub>8</sub> H <sub>13</sub> N <sub>3</sub> O <sub>6</sub>	Sorbitol hexanitrate	152.17	-2.0	7.1	773.7	670.9	-	-	124-1
49	C <sub>10</sub> H <sub>14</sub> N <sub>4</sub> O <sub>10</sub>	Dipentaerythritol hexanitrate	524.26	+3.0	-27.5	1175.0	1262.1	-	-	93

TABLE 129  
NITROINES (solid)

ADL NO.	FORMULA	NAME	MOLE WT	OXYGEN RECEIVED	OXYGEN BALANCE	Q <sub>c</sub> , KCAL/MOLE		Y'	X'	REF.
						GENERIG	CALCULATED			
317	C <sub>11</sub> H <sub>16</sub> N <sub>2</sub> O <sub>4</sub>	1,1-(2,2,2-trifluoroethyl)-nitramine (TFN)	359.14	-4.0	-16.5	517.7	491.7	-26.2	-4.3	71
35	C <sub>10</sub> H <sub>14</sub> N <sub>2</sub> O <sub>4</sub>	2,2-Difluoroethyl nitramine	176.7	3.0	-39.1	289.5	303.6	239.9	3.7	1,3
232	C <sub>10</sub> H <sub>14</sub> N <sub>2</sub> O <sub>4</sub>	Dithionit amino dinitrate	240.14	4.0	-76.7	517.7	597.3	291.3	4.2	86
593	C <sub>10</sub> H <sub>14</sub> N <sub>2</sub> O <sub>4</sub>	N-Methyl-1-nitroimidramine	165.11	4.5	-41.6	473.8	487.5	319.6	4.5	86
327	C <sub>10</sub> H <sub>14</sub> N <sub>2</sub> O <sub>4</sub>	Methyl nitramine (11q)	90.66	5.0	-63.9	372.9	370.9	353.7	5.0	136
160	C <sub>10</sub> H <sub>14</sub> N <sub>2</sub> O <sub>4</sub>	N-Methyl-N,2,3,4,5-penta-nitroimidramine	332.15	6.0	-65.9	610.6	776.4	426.6	6.0	71
117	C <sub>10</sub> H <sub>14</sub> N <sub>2</sub> O <sub>4</sub>	2-(4-hydroxymethyl)-2-(N,2,3,4-tetraimidramino)-1,3-propanediol trinitrate	512.23	7.0	-21.9	1160.7	1110.3	486.2	7.0	85
397	C <sub>10</sub> H <sub>14</sub> N <sub>2</sub> O <sub>4</sub>	(Bifluoro)hexazotriazine 2,3,4,5,6-tetra-nitro-2-aminoethanol dinitrate	440.10	5.0	-91.4	555.5	551.6	523.4	5.0	128-3
251	C <sub>10</sub> H <sub>14</sub> N <sub>2</sub> O <sub>4</sub>	Tetrayl(N-methyl-N,2,3,4-tetranitroimidramine)	287.15	8.5	-15.3	935.9	936.9	517.4	8.0	86
.	.	.	.	.	.	816.4	811.6	577.2	8.5	143
.	.	.	.	.	.	816.4	811.6	547.7	8.5	23
.	.	.	.	.	.	816.4	811.6	551.1	8.5	23
.	.	.	.	.	.	816.4	811.6	549.8	8.5	110
.	.	.	.	.	.	816.4	811.6	550.7	8.5	39
.	.	.	.	.	.	816.4	811.6	551.3	8.5	86
260	C <sub>10</sub> H <sub>14</sub> N <sub>2</sub> O <sub>4</sub>	N-Methyl-N,2,3,5-tetranitroimidramine	301.16	11.5	-61.1	1009.8	936.1	727.2	11.5	39
305	C <sub>10</sub> H <sub>14</sub> N <sub>2</sub> O <sub>4</sub>	2-(N,2,3,5-tetranitroimidramine)-1-butanol nitrate	390.23	14.0	-57.4	1250.6	1253.3	813.1	14.0	85
596	C <sub>10</sub> H <sub>14</sub> N <sub>2</sub> O <sub>4</sub>	3,3'-(Nitroamino)dipropionitrile	168.16	.	-133.2	863.2	860.0	819.0	14.0	128-3
323	C <sub>10</sub> H <sub>14</sub> N <sub>2</sub> O <sub>4</sub>	Methylene dinitramine (MENA)	136.07	0.0	0.0	219.5	210.3	489.9	0.0	71
96	C <sub>10</sub> H <sub>14</sub> N <sub>2</sub> O <sub>4</sub>	Methylene dinitramine (MMA)	150.10	3.0	-32.9	370.3	386.7	466.4	1.5	91
594	C <sub>10</sub> H <sub>14</sub> N <sub>2</sub> O <sub>4</sub>	1,3-Dinitroimidramine	162.11	5.0	-69.3	359.7	367.1	466.1	1.5	85
65	C <sub>10</sub> H <sub>14</sub> N <sub>2</sub> O <sub>4</sub>	1,3-bis(methylnitramino)-2-propanol nitrate	253.18	8.5	-53.7	637.1	636.7	322.7	2.5	128-3
65	C <sub>10</sub> H <sub>14</sub> N <sub>2</sub> O <sub>4</sub>	2,7-Dimethyl-3,6-dinitro-3,5-dioxolane-2,7-dicarbonitrile	294.28	24.0	-135.1	1670.5	1526.3	713.1	22.0	128-3
23	C <sub>10</sub> H <sub>14</sub> N <sub>2</sub> O <sub>4</sub>	RM (Hexahydro-1,3,5-triazino-8-triazine)	222.13	3.0	-21.6	500.2	504.7	414.7	1.0	133
.	.	.	.	.	.	504.7	504.7	414.7	1.0	23
.	.	.	.	.	.	504.7	504.7	414.7	1.0	23
.	.	.	.	.	.	504.7	504.7	414.7	1.0	84
.	.	.	.	.	.	504.7	504.7	414.7	1.0	54
98a	C <sub>10</sub> H <sub>14</sub> N <sub>2</sub> O <sub>4</sub>	2,4,6-tris(methylnitramino)-1,3,5-triazinophenol	348.41	21.0	-74.9	1601.7	1511.2	459.0	7.0	95
66a	C <sub>10</sub> H <sub>14</sub> N <sub>2</sub> O <sub>4</sub>	2,10-Dimethyl-3,9-triazino-3,6,9-triazinotriazene-2,10-dicarbonitrile	372.34	28.0	-120.3	1863.9	1819.5	593.4	9.33	128-3
77	C <sub>10</sub> H <sub>14</sub> N <sub>2</sub> O <sub>4</sub>	RM (Hexahydro-1,3,5,7-tetrazino-1,3,5,7-tetrazine)	266.17	0.0	-21.6	495.9	571.3	414.6	1.0	93
.	.	.	.	.	.	507.6	507.6	417.3	1.0	83
.	.	.	.	.	.	63.1	63.1	450.2	1.0	86
								Σ x' = 178.08		
								Σ y' = 12,659.5		
								Σ x'y' = 100,219.444		
								Σ x'^2 = 1,511.8611		
241	C <sub>10</sub> H <sub>14</sub> N <sub>2</sub> O <sub>4</sub>	N-Methyl-N,2,3,4,5-penta-nitroimidramine	332.15	6.0	-65.9	610.6	776.4	-	-	112
253	C <sub>10</sub> H <sub>14</sub> N <sub>2</sub> O <sub>4</sub>	Tetrayl	287.15	8.5	-15.3	935.9	936.9	-	-	117
119	C <sub>10</sub> H <sub>14</sub> N <sub>2</sub> O <sub>4</sub>	N,N'-Bis(4-cyanoethyl)nitramine	272.26	16.0	-44.7	1472.4	1608.4	-	-	84
163	C <sub>10</sub> H <sub>14</sub> N <sub>2</sub> O <sub>4</sub>	2,13-Dimethyl-3,6,9,12-tetranitro-3,6,9,12-tetrazinotriazene-2,13-dicarbonitrile	560.51	17.0	-111.2	2175.0	2175.0	-	-	128-3
67	C <sub>10</sub> H <sub>14</sub> N <sub>2</sub> O <sub>4</sub>	2,16-Dimethyl-1,6,9,12,15-pentazano-2,6,9,12,15-pentazano-2,16-dicarbonitrile	573.52	16.0	-105.2	2293.1	2501.2	-	-	128-3

TABLE 130  
NITRAMIDES (cont'd)

ADI NO.	FORMULA	NAME	MOL. WT.	OXYGEN REQUIRED	OXYGEN BALANCE	O <sub>2</sub> - KCAL/MOLE		Y'	Y*	REF.
						OBSERVED	CALCULATED			
200	C <sub>8</sub> H <sub>12</sub> N <sub>4</sub> O <sub>8</sub>	Nitrourea	165.06	0.0	- 7.6	112.3	127.0	112.5	0.5	124
197	C <sub>10</sub> H <sub>12</sub> N <sub>4</sub> O <sub>8</sub>	N-Methyl-N-nitrosourea nitrate (MISAM)	179.09	2.5	- 22.3	178.2	302.9	227.3	2.5	85
						331.0		230.1	2.5	85
184	C <sub>8</sub> H <sub>8</sub> N <sub>4</sub> O <sub>8</sub>	1-Nitrohydantoin	155.09	3.5	- 38.6	295.0	293.0	285.4	3.5	85
186	C <sub>8</sub> H <sub>8</sub> N <sub>4</sub> O <sub>8</sub>	5-Methyl-1-nitrohydantoin	159.10	6.5	- 65.4	166.8	165.4	165.1	6.5	85
56	C <sub>8</sub> H <sub>8</sub> N <sub>4</sub> O <sub>8</sub>	1,3-Dinitro-2-imidazolidone	175.09	3.0	- 27.3	390.6	396.6	175.6	3.5	128-3
						385.9		174.2	3.5	86
						388.9		175.7	3.5	85
63	C <sub>8</sub> H <sub>8</sub> N <sub>4</sub> O <sub>8</sub>	N,N'-Bis(2-hydroxyethyl)- N,N'-dinitroamide (HSDN)	356.17	4.0	- 18.0	716.2	708.3	210.6	2.0	124
						703.9		205.6	2.0	85
						710.3		206.8	2.0	85
51	C <sub>10</sub> H <sub>8</sub> N <sub>4</sub> O <sub>8</sub>	1,4-Dinitro-2,5-piperazine- dione	204.10	*	- 31.4	452.8	448.0	208.0	2.0	85
58	C <sub>8</sub> H <sub>8</sub> N <sub>4</sub> O <sub>8</sub>	N,N'-Dimethyl-N,N'-dinitro- oxamide	206.12	5.0	- 30.8	506.0	500.0	216.0	2.5	85
60	C <sub>10</sub> H <sub>12</sub> N <sub>4</sub> O <sub>8</sub>	1,1'-Ethylenebis(1-nitrourea)	216.15	6.0	- 40.7	510.4	510.6	250.1	3.0	85
53	C <sub>8</sub> H <sub>8</sub> N <sub>4</sub> O <sub>8</sub>	3,5-Dinitro-1,3,5-dinitro- 2,5-piperazine-dione	212.16	10.0	- 68.9	759.1	755.7	344.8	5.0	85
607	C <sub>8</sub> H <sub>8</sub> N <sub>4</sub> O <sub>8</sub>	Diethyl ethylenebis(N-nitro- carbamate)	294.22	15.0	- 81.6	1035.4	1022.6	490.5	7.5	85
		$y' = 101.05 + 52.1x'$						$x' = 1.500$		
								$x' y' = 3027.7$		
								$\Sigma x' y' = 11,477.9$		
								$\Sigma x'^2 = 156.50$		
								$n = 15$		
612	C <sub>8</sub> H <sub>8</sub> N <sub>4</sub> O <sub>8</sub>	Nitrourea	165.06	2.0	- 21.6	227.6	223.9	-	-	96
	C <sub>10</sub> H <sub>12</sub> N <sub>4</sub> O <sub>8</sub>	N-Methyl-N,5-dinitro-2- furfamide	215.12	4.5	- 63.2	768.0	671.6	-	-	85
55	C <sub>8</sub> H <sub>8</sub> N <sub>4</sub> O <sub>8</sub>	Dinitrobarbit	193.08	-0.5	+ 4.1	703.3	206.0	-	-	96
	C <sub>10</sub> H <sub>12</sub> N <sub>4</sub> O <sub>8</sub>	5,5-Dimethyl-1,3-dinitro- hydantoin	218.13	7.0	- 51.3	604.2	578.5	-	-	85

TABLE 131  
FLUOROS (liquid)

ADL NO	FORMULA	NAME	MOL WT	OXYGEN REQUIRED	OXYGEN BALANCE	C, KCAL/MOLE		Y'	X'	REF.
						OBSERVED	CALCULATED			
				(to HF)	(to HF)					
	C <sub>8</sub> H <sub>17</sub> FO	2-Fluorooctanol	154.26	5.0	-120.9	289.9	289.0	275.3	5.0	138
	C <sub>8</sub> H <sub>17</sub> FO <sub>2</sub>	Ethyl fluorooctanoate	178.10	9.0	-135.7	502.7	503.1	451.7	9.0	138
	C <sub>8</sub> H <sub>17</sub> FO	2-Fluorooctanol acetate	172.10	11.0	-185.4	499.4	499.4	451.4	9.0	138
	C <sub>8</sub> H <sub>17</sub> FO	m-Fluorooctanol	154.26	5.0	-120.9	289.9	289.9	275.9	11.0	138
	C <sub>8</sub> H <sub>17</sub> FO	1-Ethyl 2-Fluoro-2-propyl ether	154.26	5.0	-120.9	289.9	289.9	275.9	11.0	138
	C <sub>8</sub> H <sub>17</sub> FO	Fluorooctane	154.26	5.0	-120.9	289.9	289.9	275.9	11.0	138
	C <sub>8</sub> H <sub>17</sub> FO	o-Fluorooctane	154.26	5.0	-120.9	289.9	289.9	275.9	11.0	138
	C <sub>8</sub> H <sub>17</sub> FO	m-Fluorooctane	154.26	5.0	-120.9	289.9	289.9	275.9	11.0	138
	C <sub>8</sub> H <sub>17</sub> FO	p-Fluorooctane	154.26	5.0	-120.9	289.9	289.9	275.9	11.0	138
	C <sub>8</sub> H <sub>17</sub> FO	2,2-Difluorooctane	170.16	10.0	-146.7	474.7	474.7	421.7	10.0	138
	C <sub>8</sub> H <sub>17</sub> FO	2,2-Difluoroethylamine	154.26	5.0	-120.9	289.9	289.9	275.9	11.0	138
	C <sub>8</sub> H <sub>17</sub> FO	Ethyl difluoroacetate	170.16	10.0	-146.7	474.7	474.7	421.7	10.0	138
	C <sub>8</sub> H <sub>17</sub> FO	2,2-Difluoroethyl acetate	170.16	10.0	-146.7	474.7	474.7	421.7	10.0	138
	C <sub>8</sub> H <sub>17</sub> FO	1-Fluoro-2-naphthene	170.16	10.0	-146.7	474.7	474.7	421.7	10.0	138
	C <sub>8</sub> H <sub>17</sub> FO	1-Fluoro-2-naphthene	170.16	10.0	-146.7	474.7	474.7	421.7	10.0	138
	C <sub>8</sub> H <sub>17</sub> FO <sub>2</sub>	Difluoroacetic acid	98.06	2.0	-31.3	133.4	128.2	66.2	3.0	138
	C <sub>8</sub> H <sub>17</sub> FO <sub>2</sub>	2,2-Difluoroethanol	82.05	1.0	-30.9	115.0	115.0	70.0	2.0	138
	C <sub>8</sub> H <sub>17</sub> FO <sub>2</sub>	2,2-Difluoroethylamine	81.07	5.5	-100.5	121.0	132.2	151.1	2.75	138
	C <sub>8</sub> H <sub>17</sub> FO <sub>2</sub>	Ethyl difluoroacetate	124.09	8.0	-103.2	461.4	461.4	421.4	8.0	138
	C <sub>8</sub> H <sub>17</sub> FO <sub>2</sub>	2,2-Difluoroethyl acetate	124.09	8.0	-103.2	461.4	461.4	421.4	8.0	138
	C <sub>8</sub> H <sub>17</sub> FO <sub>2</sub>	2,2-Difluoroethylamine	111.09	11.0	-122.3	705.2	697.6	677.2	6.5	138
	C <sub>8</sub> H <sub>17</sub> FO <sub>2</sub>	o,p-Difluoroethane	170.16	10.0	-146.7	474.7	474.7	421.7	10.0	138
	C <sub>8</sub> H <sub>17</sub> FO <sub>2</sub>	Bis(2,2-difluoroethyl)amine	170.16	9.5	-103.8	562.3	562.3	522.3	2.375	138
								3 x' = 26.125		
								2 y' = 14,336.6		
								2 x'y' = 213,371.353		
								1 x'' = 3774.29325		
								n = 20		

TABLE 132  
FLUOROS (solid)

ADL NO	FORMULA	NAME	MOL WT	OXYGEN REQUIRED	OXYGEN BALANCE	C, KCAL/MOLE		Y'	X'	REF.
						OBSERVED	CALCULATED			
				(to HF)	(to HF)					
	C <sub>2</sub> H <sub>3</sub> FO <sub>2</sub>	Fluoroacetic acid	76.02	1.0	-61.5	170.3	172.2	169.3	3.0	138
	C <sub>2</sub> H <sub>3</sub> FO <sub>2</sub>	2-Fluoroacetonitrile	77.05	1.0	-61.5	170.3	172.2	169.3	3.0	138
	C <sub>2</sub> H <sub>3</sub> FO	p-Fluorophenol	112.10	11.0	-185.4	499.4	499.4	451.4	11.0	138
	C <sub>2</sub> H <sub>3</sub> FO <sub>2</sub>	o-Fluoroacetic acid	110.11	11.0	-159.9	739.9	739.9	747.0	11.0	138
	C <sub>2</sub> H <sub>3</sub> FO <sub>2</sub>	m-Fluoroacetic acid	110.11	11.0	-159.9	739.9	739.9	747.0	11.0	138
	C <sub>2</sub> H <sub>3</sub> FO <sub>2</sub>	p-Fluoroacetic acid	110.11	11.0	-159.9	739.9	739.9	747.0	11.0	138
	C <sub>2</sub> H <sub>3</sub> FO	Fluoroacetonitrile	151.15	18.5	-151.3	977.9	977.9	981.0	18.5	138
	C <sub>2</sub> H <sub>3</sub> FO	p-Fluoroacetonitrile	151.15	18.5	-151.3	977.9	977.9	981.0	18.5	138
	C <sub>2</sub> H <sub>3</sub> FO <sub>2</sub>	o-Fluoroacetic acid	152.16	19.0	-151.0	1013.5	1013.5	1005.9	19.0	138
	C <sub>2</sub> H <sub>3</sub> FO <sub>2</sub>	Ethyl p-fluoroacetate	180.17	22.0	-195.4	1187.5	1187.1	1257.5	22.0	138
	C <sub>2</sub> H <sub>3</sub> FO <sub>2</sub>	Methyl p-fluoroacetate	133.10	21.0	-176.3	1208.0	1212.0	1211.0	21.0	138
	C <sub>2</sub> H <sub>3</sub> FO <sub>2</sub>	Fluoroacetonitrile	94.05	1.5	-58.9	266.9	212.9	102.7	1.75	138
	C <sub>2</sub> H <sub>3</sub> FO <sub>2</sub>	2,2-Difluoroacetonitrile	171.11	17.5	-161.5	981.4	937.8	873.9	8.75	138
								3 x' = 14.0		
								3 y' = 10,337.0		
								2 x'y' = 176,119.05		
								1 x'' = 3324.375		
								n = 11		

CONFIDENTIAL

TABLE 111  
O<sub>2</sub> TRIFLUOROS (lit.)

ADL NO.	FORMULA	NAME	MOL. WT.	OXYGEN REQUIRED	OXYGEN BALANCE	O <sub>2</sub> KCAL/MOLE		Y'	X'	REF.
						OBSERVED	CALCULATED			
				(lit. ref.)	(lit. ref.)					
	C <sub>7</sub> H <sub>6</sub> F <sub>3</sub> O	o,p,o'-Trifluoro-m-cresol	162.11	11.0	-107.2	761.9	754.6	162.1	11.0	119
	C <sub>7</sub> H <sub>5</sub> F <sub>3</sub>	m,p,o'-Trifluorotoluene	166.11	15.0	-152.1	809.7	807.3	171.1	15.0	119
	C <sub>6</sub> H <sub>5</sub> F <sub>3</sub> O <sub>2</sub>	m,p,o'-Trifluoro-acetic acid (s)	198.12	"	-176.2	855.7	831.1	211.6	15.0	119
	C <sub>7</sub> H <sub>6</sub> F <sub>3</sub> N	o,p,o'-Trifluoro-m-toluidine	161.13	15.0	-101.9	822.7	815.7	167.1	15.0	119
	C <sub>7</sub> H <sub>5</sub> F <sub>3</sub> NO	o,p,o'-Trifluoro-p-acetotolide (s)	209.15	19.5	-151.5	1007.7	1019.9	166.0	19.5	119
		y' = 16.07 + 51.61x'						1 x' = 79.0 2 x' = 40.97 3 x' = 68.251.35 4 x' = 1266.0 n = 5		

TABLE 112  
NITROSAMINES (solid and liquid)

ADL NO.	FORMULA	NAME	MOL. WT.	OXYGEN REQUIRED	OXYGEN BALANCE	O <sub>2</sub> KCAL/MOLE		Y'	X'	REF.
						OBSERVED	CALCULATED			
195	C <sub>2</sub> H <sub>6</sub> N <sub>2</sub> O	Diethylnitrosamine (liq)	72.08	6.0	-129.6	191.3	178.1	171.0	6.0	190
677	C <sub>10</sub> H <sub>18</sub> N <sub>2</sub> O	N-Ethyl-N-nitrosamine (s)	170.16	20.0	-213.1	1138.0	1066.9	1122.2	20.0	190
676	C <sub>12</sub> H <sub>18</sub> N <sub>2</sub> O	Diphenylnitrosamine (s)	198.22	26.0	-226.0	2502.1	1546.1	1531.7	26.0	190
						1530.8		1532.1	26.0	213
776	C <sub>8</sub> H <sub>10</sub> O <sub>2</sub> N <sub>2</sub>	3,7-Dinitroso-3,3,7,7-tetrabicyclo[3,3,1]nonane (s)	186.13	11.0	-111.7	657.7	387.5	191.1	6.5	207
						871.1		395.2	6.5	207
24	C <sub>6</sub> H <sub>8</sub> N <sub>2</sub> O <sub>3</sub>	Hexahydro-1,3,5-trinitroso-oxazine (H.S.S.) (s)	176.13	6.0	-95.1	551.5	525.9	161.5	2.0	201
						556.2		165.2	2.0	15,197
						559.1		166.4	2.0	207
		y' = 56.03 + 53.13x'						1 x' = 101.0 2 x' = 5879.1 3 x' = 117,255.90 4 x' = 2160.5 n = 9		
195	C <sub>8</sub> H <sub>8</sub> N <sub>2</sub> O	Dimethylnitrosamine (liq)	72.08	6.0	-129.6	120.3 <sup>a</sup>	193.1			190
		This value as reported by Tom. Abate is obviously a misprint.								

CONFIDENTIAL

TABLE 145  
 THERM DATA (solid and liquid)

ADL NO.	FORMULA	NAME	MOL. WT.	OXYGEN REQUIRED	OXYGEN BALANCE	Q <sub>c</sub> , KCAL/MOLE		Y'	X'	REF.
						OBSERVED	CALCULATED			
528	C <sub>8</sub> H <sub>8</sub> O <sub>2</sub>	2,2-Dinitrothiophene (a)	174.13	7.0	-54.1	601.7	539.2	195.6	7.0	10
529	C <sub>8</sub> H <sub>8</sub> O <sub>2</sub>	2,4-Dinitrothiophene (a)	174.13	9.5	-177.7	614.7	619.7	620.5	9.5	10
529	C <sub>8</sub> H <sub>8</sub> O <sub>2</sub>	3,5-Dinitrothiophene (a)	174.13	10.6	-95.0	717.1	735.2	571.0	10.6	10
529	C <sub>8</sub> H <sub>8</sub> O <sub>2</sub>	Thiophene (119)	114.13	12.1	-202.7	607.2	601.7	601.7	12.1	79
529	C <sub>8</sub> H <sub>8</sub> O <sub>2</sub>	1-Methyl-2-nitrothiophene (b)	173.16	12.7	-137.7	717.2	777.7	652.5	12.7	53
								$\sum x' = 63.0$ $\sum y' = 371.3$ $\sum x'y' = 17,421.25$ $\sum x'^2 = 63.5$ $n = 6$		
699	C <sub>8</sub> H <sub>7</sub> S	Thiophene (119)	114.13	12.1	-202.7	607.2	603.9	-	-	7-37
	C <sub>8</sub> H <sub>7</sub> S	2-Thiophene carboxylic acid (a)	174.09	12.9	-137.7	616.2	650.0	-	-	7-155
		* Calculated to 50,								

 TABLE 146  
 THERM DATA (solid)

ADL NO.	FORMULA	NAME	MOL. WT.	OXYGEN REQUIRED	OXYGEN BALANCE	Q <sub>c</sub> , KCAL/MOLE		Y'	X'	REF.
						OBSERVED	CALCULATED			
111	C <sub>10</sub> H <sub>13</sub> N	N,N-Dimethyl-p-nitroaniline	195.16	20	-211.1	1126.1	1127.2	1167.0	20	190, 2-212
						1325.8	1376.5	20	211	
110	C <sub>10</sub> H <sub>11</sub> N <sub>2</sub> O	p-Nitro-N,N-diphenylamine	195.22	20	-210.0	1547.2	1517.6	1583.3	20	211
			168.11	10	-95.2	582.5	589.1	287.6	5	190
								$\sum x' = 73.0$ $\sum y' = 1276.3$ $\sum x'y' = 66,126.60$ $\sum x'^2 = 1669.0$ $n = 4$		



CONFIDENTIAL  
SECURITY INFORMATION

TABLE 117  
HEAT OF HYDRATION<sup>a</sup>

$$Q_{\text{Hydration}} = Q_{\text{Formation of hydrate}} - Q_{\text{Formation of anhydrous salt}} - Q_{\text{Formation of water}}$$

Hydrate	$Q_F$ kcal/mole	Anhydrous Compound	$Q_F$ kcal/mole	$\Delta Q_F/H_2O$ kcal/mole
$N_2H_4 \cdot H_2O$ (aq.)	63.9	$N_2H_4$ (aq.)	-8.2	72.1
$N_2H_4 \cdot H_2O$ (liq)	58.0	$N_2H_4$ (liq)	-12.1	70.1
$NaOH \cdot H_2O$ (s)	175.3	$NaOH$ (s)	102.0	73.3
$H_4P_2O_7 \cdot 1.5H_2O$ (s)	637.5	$H_4P_2O_7$ (s)	513.7	82.4
$H_4P_2O_7 \cdot 1.5H_2O$ (liq)	634.4	$H_4P_2O_7$ (liq)	529.4	70.0
$H_2SO_4 \cdot H_2O$ (s)	273.3	$H_2SO_4$ (s)	196.3	70.9
$H_2SO_4 \cdot H_2O$ (liq)	268.7	$H_2SO_4$ (liq)	193.8	74.9
$H_2SeO_4 \cdot H_2O$ (s)	204.5	$H_2SeO_4$ (s)	130.3	74.2
$H_2SeO_4 \cdot H_2O$ (liq)	200.0	$H_2SeO_4$ (liq)	126.8	73.2
$(COOH)_2 \cdot 2H_2O$ (s)	340.2	$(COOH)_2$ (s)	197.5	71.3
$(COOH)_2 \cdot H_2O$ (s)	339.1	$(COOH)_2$ (s)	267.2	71.9
$(NH_4)_2SO_4 \cdot H_2O$ (s)	283.6	$(NH_4)_2SO_4$ (s)	212.3	71.3
$3PbI_2 \cdot PI_3 \cdot 12H_2O$ (s)	977.3	$3PbI_2 \cdot PI_3$ (s)	131.8	70.6
$3PbI_2 \cdot 3I_2 \cdot 12H_2O$ (s)	952.2	$3PbI_2 \cdot 3I_2$ (s)	111.0	70.1
$LiBr \cdot H_2O$ (s)	158.4	$LiBr$ (s)	83.8	74.6
$LiBr \cdot 2H_2O$ (s)	230.0	$LiBr$ (s)	83.8	73.1
$LiBr \cdot 3H_2O$ (s)	302.0	$LiBr$ (s)	83.8	72.4
$Li_2SO_4 \cdot H_2O$ (s)	413.7	$Li_2SO_4$ (s)	342.35	71.2
$KF \cdot 2H_2O$ (s)	277.1	$KF$ (s)	134.5	71.3
$KF \cdot 4H_2O$ (s)	418.0	$KF$ (s)	134.5	70.9

$$\Sigma (\Delta Q_F) = 1455.8$$

$$n = 20$$

$$Av. \Delta Q_F = 72.8$$

$$Q_{\text{Hydration}} = 72.8 - 68.3 = +4.5 \text{ kcal/mole}$$

<sup>a</sup>Data from Ref. 16.

CONFIDENTIAL

TABLE 100

## SALT LINK

$$\Delta C_p (\text{salt link}) = \Delta C_p (\text{salt}) - \Delta C_p (\text{acid} + \text{base}) - \Delta C_p (\text{salt}) = \Delta C_p (\text{acid} + \text{base}) - \Delta C_p (\text{salt})$$

Formula	Name	Mol. wt.	Gr. Needed	Gr. Bal.	Heat Observed, kcal/mole Formation	kcal/mole Combustion	$\Delta C_p$ per salt link kcal/mole	Ref.
$C_2H_5N_3O_2$	Methylamine nitrate (s)	91.07	2.0	-34.0	-	218.4	-	29
$C_2H_5N_3O_2$	Methylamine (liq)	31.06	4.5	-231.0	-	234.1	-	calc.
$HNO_3$	Nitric acid (liq)	63.02	-2.5	+ 63.5	14.4	-7.3	-28.4	82a
$C_2H_5N_3O_2$	Dimethylamine nitrate (s)	106.10	5.0	-74.0	-	372.4	-	29
$C_2H_5N_3O_2$	Dimethylamine (liq)	45.08	7.5	-266.2	-	143.7	-	calc.
$HNO_3$	Nitric acid (liq)	63.02	-2.5	+ 63.5	-	-7.3	-	82a
$C_2H_5N_3O_2$	Trimethylamine nitrate (s)	122.13	6.0	-104.8	-	450.7	-	29
$C_2H_5N_3O_2$	Trimethylamine (liq)	59.11	10.5	-266.2	-	471.9	-	calc.
$HNO_3$	Nitric acid (liq)	63.02	-2.5	+ 63.5	-	-7.3	-	82a
$C_2H_5N_3O_2$	Ethylamine nitrate (liq)	106.10	5.0	-74.0	-	374.5	-	29
$C_2H_5N_3O_2$	Ethylamine (liq)	45.08	7.5	-266.2	-	467.9	-	calc.
$HNO_3$	Nitric acid (liq)	63.02	-2.5	+ 63.5	-	-7.3	-	82a
$C_2H_5N_3O_2$	Diethylamine nitrate (s)	134.15	11.0	-149.3	-	457.4	-	29
$C_2H_5N_3O_2$	Diethylamine (liq)	73.11	13.5	-266.3	-	725.5	-	calc.
$HNO_3$	Nitric acid (liq)	63.02	-2.5	+ 63.5	-	-7.3	-	82a
$C_2H_5N_3O_2$	Triethylamine nitrate (s)	164.20	17.0	-185.7	-	1013.6	-	29
$C_2H_5N_3O_2$	Triethylamine (liq)	101.19	19.5	-300.3	-	1444.5	-	calc.
$HNO_3$	Nitric acid (liq)	63.02	-2.5	+ 63.5	-	-7.3	-	82a
$C_6H_5N_3O_2$	Aniline nitrate (s)	150.11	11.0	-133.2	-	787.9	-	153
$C_6H_5N_3O_2$	Aniline (s)	-	-	-	-	775.1	-	29
$C_6H_5N_3O_2$	Aniline (liq)	91.12	15.5	-266.3	-	809.1	-	calc.
$HNO_3$	Nitric acid (liq)	63.02	-2.5	+ 63.5	-	-7.3	-	82a
$C_{10}H_{10}N_4O_2$	Acridine nitrate (s)	242.23	28.0	-124.9	-	1553.3	-	153
$C_{10}H_{10}N_4O_2$	Acridine (liq)	179.21	30.5	-272.3	-	1577.6	-	2
$HNO_3$	Nitric acid (liq)	63.02	-2.5	+ 63.5	-	-7.3	-	82a
$C_{10}H_{10}N_4O_2$	2-Aminoacridine mono-nitrate (s)	257.24	24.5	-177.3	-	1442.0	-	153
$C_{10}H_{10}N_4O_2$	2-Aminoacridine (s)	194.23	31.0	-255.4	-	1601.7	-	153
$HNO_3$	Nitric acid (liq)	63.02	-2.5	+ 63.5	-	-7.3	-	82a
$C_{10}H_{10}N_4O_2$	9-Aminoacridine mono-nitrate (s)	257.24	24.5	-177.3	-	1553.3	-	153
$C_{10}H_{10}N_4O_2$	9-Aminoacridine (s)	194.23	31.0	-255.4	-	1601.9	-	153
$HNO_3$	Nitric acid (liq)	63.02	-2.5	+ 63.5	-	-7.3	-	82a
$C_{12}H_{12}N_4O_2$	1,6-Diaminocridine mono-nitrate (s)	272.26	29.0	-170.4	-	1594.5	-	153
$C_{12}H_{12}N_4O_2$	1,6-Diaminocridine (s)	209.24	31.5	-249.7	-	1628.4	-	153
$HNO_3$	Nitric acid (liq)	63.02	-2.5	+ 63.5	-	-7.3	-	82a
$C_{12}H_{12}N_4O_2$	3,6-Diaminocridine mono-nitrate (s)	272.26	29.0	-170.4	-	1594.5	-	153
$C_{12}H_{12}N_4O_2$	3,6-Diaminocridine (s)	209.24	31.5	-249.7	-	1628.4	-	153
$HNO_3$	Nitric acid (liq)	63.02	-2.5	+ 63.5	-	-7.3	-	82a
$C_{12}H_{12}N_4O_2$	3,6-Diaminocridine di-nitrate (s)	344.15	3.0	-25.8	-	376.3	-	17b
$C_{12}H_{12}N_4O_2$	3,6-Diaminocridine (liq)	60.10	0.5	-233.0	-	373.2	-	66
$HNO_3$	Nitric acid (liq)	63.02	-2.5	+ 63.5	-	-14.6	-	82a
$C_{12}H_{12}N_4O_2$	5-Aminoacridine nitrate (s)	169.10	1.5	-14.2	-	317.5	-	12f
$C_{12}H_{12}N_4O_2$	5-Aminoacridine (liq)	146.60	2.0	-60.3	-	323.0	-	29
$HNO_3$	Nitric acid (liq)	63.02	-2.5	+ 63.5	-	-7.3	-	82a
$C_{10}H_{10}N_4O_2$	Benzylamine nitrate (s)	176.17	16.0	-140.4	-	912.7	-	29
$C_{10}H_{10}N_4O_2$	Benzylamine (liq)	107.15	19.5	-276.4	-	993.1	-	calc.
$HNO_3$	Nitric acid (liq)	63.02	-2.5	+ 63.5	-	-7.3	-	82a
$C_2H_5N_3O_2$	Glycine nitrate (s)	115.09	2.0	-21.2	-	219.5	-	29
$C_2H_5N_3O_2$	Glycine (s)	75.07	4.8	-94.9	125.1	234.5	-	67a
$HNO_3$	Nitric acid (liq)	63.02	-2.5	+ 63.5	-	-7.3	-	82a
$C_6H_{14}N_4O_2$	Hexamethylenetetramine di-nitrate (s)	226.21	13.5	-91.9	-	943.4	-	86
						977.5	-	33
						952.3	-	35
$C_6H_{14}N_4O_2$	Hexamethylenetetramine (s)	140.19	15.0	-27.4	-	1408.1	-	23
						1037.9	-	13
$HNO_3$	Nitric acid (liq)	63.02	-2.5	+ 63.5	-	1004.7	-	31
						-14.6	-	82a

CONFIDENTIAL

TABLE 11d (cont'd)

Formula	Name	Mol. wt.	Ox.		Heat Observed, kcal/mole Formation	Heat of Combustion kcal/mole	ΔH per salt link kcal/mole	Ref.
			Needed	Ox. Bal.				
$CN_2O_4$	Urea nitrate (s)	121.08	0.5	+ 6.5	-	114.5	-	12L
$CN_2H_4O$	Urea (s)	60.06	1.0	-153.1	-	150.6	-	calc.
$HNO_3$	Nitric acid (liq)	63.02	-2.5	+ 61.5	-	-7.1	-	32a
$C_2H_3N_3O_3$	Thiourea nitrate (s)	119.09	1.0	+ 40.1	74.5	-	- 0.3	32a
$C_2H_3N_3O_3$	Thiourea (s)	76.07	6.0	-160.2	22.1	-	-	32a
$HNO_3$	Nitric acid (liq)	63.02	-2.5	+ 61.5	41.4	-	-	32a
$H_2N_2O$	Hydroxylamine nitrate(s)	76.05	-2.0	+ 22.2	66.2	-	-11.0	16
$H_2NO$	Hydroxylamine (s)	31.31	+0.5	+ 26.2	25.7	-	-	32a
$HNO_3$	Nitric acid (liq)	63.02	-2.5	+ 61.5	41.4	-	-	32a
$H_4N_2O_2$	Ammonium nitrate (s)	80.05	-1.0	+ 20.0	67.1	-	-	32a
$H_4N$	Ammonia (s)	17.01	+1.5	-140.2	16.0	-	-	32a
$HNO_3$	Nitric acid (liq)	63.02	-2.5	+ 61.5	41.4	-	-	32a
$H_2N_2O_2$	Hydrazine mononitrate(s)	95.07	-0.5	+ 6.5	54.4	-	-	71
$H_2N_2$	Hydrazine (aq)	32.05	+2.0	- 77.3	-5.2	-	-	71
$HNO_3$	Nitric acid (aq)	63.02	-2.5	+ 61.5	17.3	-	-	32a
$H_2N_2O_2$	Hydrazine dinitrate (aq)	153.10	-1.0	+ 10.4	101.4	-	-	71
$H_2N_2$	Hydrazine (aq)	32.05	+2.0	- 77.3	-8.2	-	-	71
$HNO_3$	Nitric acid (aq)	63.02	-2.5	+ 61.5	93.8	-	-	32a
$H_4N_2O_4S$	Hydrazine sulfate (aq)	135.08	1.0	- 12.1	219.1	-	-	71
$H_4N_2$	Hydrazine (aq)	32.05	2.0	- 77.3	-5.2	-	-	71
$H_2SO_4$	Sulfuric acid (aq)	98.03	-1.0	+ 16.3	216.7	-	-	32a
$H_2O_4S$	Dihydrazine sulfate (aq)	162.12	1.0	- 29.6	227.1	-	-	71
$H_2N_2$	Hydrazine (aq)	32.05	2.0	- 77.3	-16.4	-	-	71
$H_2SO_4$	Sulfuric acid (aq)	98.03	-1.0	+ 16.3	216.7	-	-	32a
$H_2N_2O_4S$	Hydrazine hydrochloride (aq)	68.52	2.0	+ 46.7	4.6	-	-	71
$H_2N_2$	Hydrazine (aq)	32.05	2.0	- 77.3	-5.2	-	-	71
$HCl$	Hydrochloric acid (aq)	36.47	0.0	0.0	16.0	-	-	32a
$H_2N_2O_4S$	Hydrazine hydrochloride (aq)	104.43	2.0	- 10.5	85.0	-	-	32a
$H_2N_2$	Hydrazine (aq)	32.05	2.0	- 77.3	-3.2	-	-	71
$HCl$	Hydrochloric acid (aq)	36.47	0.0	0.0	80.0	-	-	32a
$CH_5N$	Methylamine hydrochloride (aq)	67.52	4.5	-106.5	-	-	- 6.1	16
$C_2H_7N$	Dimethylamine hydrochloride (aq)	61.55	7.5	-147.1	-	-	-11.3	16
$C_3H_9N$	Etaniline hydrochloride (s)	81.55	7.5	-147.1	-	-	+ 3.4	16
$C_3H_7N$	Guanidine ion (aq)	60.06	7.0	-166.4	-	-	-11.1	16
$C_2H_5NO_2$	Ammonium nitroacetate (aq)	117.07	4.5	- 64.0	-	-	-	15
$C_2H_5NO_2O_3$	Ammonium nitroacetate (aq)	163.03	2.5	- 22.2	-	-	-13.1	16
$C_2H_5NO_2$	Ammonium hydrogen oxalate (aq)	107.07	2.5	- 37.4	-	-	-17.5	16
$C_2H_5NO_2$	Ammonium glycolate (aq)	71.03	4.5	- 77.3	-	-	-14.2	16
$C_2H_5NO_2$	Ammonium glycolate (aq)	159.08	3.5	- 51.1	-	-	-12.2	16
$CH_3NO_2$	Ammonium formate (aq)	61.06	2.5	- 61.4	-	-	-11.9	16
$C_2H_5NO_2$	Diammonium oxalate (s)	124.10	4.0	- 51.5	267.2	-	-	16
$H_3N$	Ammonia (x2) (liq)	17.01	1.5	-140.2	11.2	-	-	32a
$C_2H_5O_4$	Oxalic acid (s)	90.04	1.0	- 17.8	136.2	50.2	-	calc.
$C_2H_5NO_7$	Ammonium picrate (s)	256.14	8.0	- 82.0	-	593.9	-	122
$H_3N$	Ammonia (liq)	17.01	1.5	-140.2	-	197.0	-	32a
$C_2H_5NO_7$	Picric acid (s)	229.11	5.5	- 45.4	-	613.0	-	calc.
$C_2H_5NO_3$	Ammonium carbonate (s)	154.09	1.0	- 60.0	224.6	-	-19.9	16
$H_3N$	Ammonia (x2) (liq)	17.01	1.5	-140.2	14.6	-	-	32a
$CH_3CO_2$	Carbonic acid (aq)	62.03	0.0	0.0	-143.0	-	-	32a
$C_2H_5NO_3$	Methylamine hydrogen carbonate (aq)	71.05	4.5	- 77.4	-	-	+ 2.0	16

TABLE 130 (cont'd)

SALE LINE

Formula	Name		Vol. wt.	Ox. Needed	Ox. Bal.	Heat of Oxidation, kcal/mole	Heat of Combustion, kcal/mole	$\Delta C_p$	$\Delta C_v$	alt link kcal/mole	Ref.
C <sub>4</sub> H <sub>9</sub> NO <sub>2</sub>	Ammonium acetate	(s)	77.08	5.5	-112.2	122.0	272.2	-	-	-	16
N <sub>2</sub> H	Ammonia	(liq)	17.0	1.5	-120.9	-	97.0	-	-	-	82a
C <sub>2</sub> H <sub>4</sub> O <sub>3</sub>	Acetic acid	(liq)	60.06	1.0	-106.6	-	205.6	-	-	-26.4	calc.
C <sub>2</sub> H <sub>3</sub> N <sub>3</sub> O	Ammonium cyanate	(aq)	60.06	1.0	-79.9	68.5	-	-	-	-	82a
N <sub>2</sub> H	Ammonia	(aq)	17.0	1.5	-120.9	19.3	-	-	-	-	82a
C <sub>2</sub> H <sub>3</sub> N <sub>3</sub> O	Cyanic acid	(aq)	43.01	1.5	-55.8	35.1	-	-	-	-	82a
C <sub>2</sub> H <sub>3</sub> N <sub>3</sub> O	Ammonium nitroform	(s)	168.07	-2.0	-19.0	-	195	-	-	-11.1	Pittsiny 62a
N <sub>2</sub> H	Ammonia	(liq)	17.0	1.5	-120.9	-	97.0	-	-	-	calc.
C <sub>2</sub> H <sub>3</sub> N <sub>3</sub> O	Nitroform	(liq)	151.06	-1.5	+17.1	-	111.9	-	-	-	calc.
C <sub>2</sub> H <sub>3</sub> N <sub>3</sub> O <sub>2</sub>	Ethylammonium dinitroform	(s)	271.16	1.5	-34.1	-	55.1	-	-	-	71
C <sub>2</sub> H <sub>3</sub> N <sub>3</sub> O <sub>2</sub>	Nitroammonium dinitroform	(liq)	60.06	1.5	-112.3	-	451.5	-	-	-	calc.
C <sub>2</sub> H <sub>3</sub> N <sub>3</sub> O <sub>2</sub>	Nitroform (x2)	(liq)	151.06	-1.5	+17.1	-	229.0	-	-	-11.1	calc.
N <sub>2</sub> H	Ammonia	(aq)	60.06	2.0	-51.3	-25.9	-	-	-	-	16
N <sub>2</sub> H	Ammonia	(aq)	17.0	1.5	-120.9	19.3	-	-	-	-	82a
H <sub>2</sub>	Hydrochloric acid	(aq)	43.01	0.5	-14.6	-54.5	-	-	-	-	16
C <sub>2</sub> H <sub>3</sub> N <sub>3</sub> O <sub>2</sub>	MEDMA dicyanide salt	(s)	200.17	2.0	-16.0	-	107.9	-	-	-	71
N <sub>2</sub> H	Hydrazine (x2)	(liq)	32.05	2.0	-92.3	-	257.2	-	-	-	73
C <sub>2</sub> H <sub>3</sub> N <sub>3</sub> O <sub>2</sub>	MEDMA	(s)	136.07	0.4	0.0	-	234.3	-	-	-19.9	calc.
C <sub>8</sub> H <sub>11</sub> N <sub>3</sub> OCl	p-Nitrosodimethylam- monium chloride	(s)	156.65	2.0	-171.4	-	1120.5	-	-	-	135
C <sub>8</sub> H <sub>11</sub> N <sub>3</sub> O	p-Nitrosodimethyl- amine	(s)	150.18	20.0	-211.1	-	1120.2	-	-	-	calc.
HCl	Hydrochloric acid	(liq)	36.47	0.0	0.0	-	26.0	-	-	-25.4	82a
C <sub>2</sub> H <sub>3</sub> N <sub>3</sub> O	Ammonium cyanide	(s)	41.05	1.0	-115.3	0.0	-	-	-	-25.4	82a
N <sub>2</sub> H	Ammonia	(liq)	17.0	1.5	-120.9	16.6	-	-	-	-	82a
C <sub>2</sub> H <sub>3</sub> N <sub>3</sub> O	Nytracrynic acid	(liq)	27.01	2.5	-145.0	-25.2	-	-	-	-	82a
C <sub>2</sub> H <sub>3</sub> N <sub>3</sub> O	Ammonium hydrogen carbonate	(s)	79.06	1.5	-30.4	136.9	-	-	-	-	82a
N <sub>2</sub> H	Ammonia	(aq)	17.0	1.5	-120.9	19.3	-	-	-	-	82a
C <sub>2</sub> H <sub>3</sub> N <sub>3</sub> O	Carbonic acid	(cp)	62.03	0.0	0.0	167.0	-	-	-	-10.5	82a
C <sub>2</sub> H <sub>3</sub> N <sub>3</sub> S	Ammonium thiocyanate	(aq)	76.07	4.0	-136.2	15.6	-	-	-	-	82a
N <sub>2</sub> H	Ammonia	(aq)	17.0	1.5	-120.9	19.3	-	-	-	-	82a
C <sub>2</sub> H <sub>3</sub> N <sub>3</sub> O	Thiocyanic acid	(aq)	59.04	4.5	-122.0	-17.7	-	-	-	-12.9	82a
N <sub>2</sub> H <sub>2</sub> OCl	Hydroxylamine mono- hydrochloride	(aq)	69.00	0.5	-11.5	70.7	-	-	-	-	82a
N <sub>2</sub> H <sub>2</sub> O	Hydroxylamine	(aq)	31.03	0.5	-24.2	21.7	-	-	-	-	82a
HCl	Hydrochloric acid	(aq)	36.47	0.0	0.0	10.0	-	-	-	-9.0	82a
						$\Sigma \Delta C_p = -123.2$					
						$n = 52$					
						Av. $\Delta C_p = -16.1$ kcal/mole					
C <sub>12</sub> H <sub>16</sub> N <sub>4</sub> O <sub>12</sub>	Hexanitrodiphenylamine ammonium salt	(s)	456.25	16.0	-56.1	-	1122.6	-	-	-	119
C <sub>2</sub> H <sub>3</sub> N <sub>3</sub> O <sub>2</sub>	Oxallic acid dicyanide dinitrate	(s)	179.15	5.5	-49.1	-	113.5	-	-	-	12b-1
N <sub>2</sub> H <sub>2</sub> O <sub>2</sub> S	Hydroxylamine sulfate	(aq)	131.06	-0.5	+5.1	264.4	-	-	-	-	16
N <sub>2</sub> H <sub>2</sub> OCl <sub>2</sub>	Hydroxylamine dihydro- chloride	(aq)	104.96	-0.5	-7.6	164.0	-	-	-	-	62a
C <sub>12</sub> H <sub>14</sub> N <sub>4</sub> I <sub>2</sub>	Hexidine dihydro- iodide	(s)	444.09	10.0	-109.1	-	1577.0	-	-	-	99

CONFIDENTIAL

TABLE 139  
LIST OF COMPOUNDS

C-

1. Acetylene
2. 1-(Aminoguanyl)-4-(nitroguanyl)-1-tetrazole
3. Ammonium nitrate
4. Nitroform, ammonium salt
5. Picric acid, ammonium salt
6. 4-Hydroxytetrahydropyrazin-3,3,5,5-tetramethanol pentanitrate
7. Dimethylsulfamic acid
8. Sodium azide
9. 2-Azidoethanol
10. 2-Azidoethanol nitrate
11. 5-Azido-1-hydroxy-1H-tetrazole
12. 3-Azido-1,2-propanediol dinitrate
13. 4,4'-Isopropylidenebis(2,6-dinitrophenol)
14. 4,4'-Sulfonylbis(2,6-dinitrophenol)
15. 1,3-Butanediol dinitrate
16. Ethylenediamine dichlorate
17. 2-Chloroethanol nitrate
18. 3-Hydroxypropionitrile nitrate
19. Glycolonitrile nitrate
20. Cyanuric triazide
21. Tetrahydro-3,5-dinitro-1,3,5,2H-oxadiazine
22. HMX
23. RDX
24. Hexahydro-1,3,5-trinitro-2-triazine
25. 2,4,6,8-Tetranitro-2,4,6,8-tetraazanonane-1,9-diol diacetate
26. 2,4,6-Trinitro-2,4,6-triazabenzene-1,7-diol diacetate
27. 5,5'-(Diazamino)di-1H-tetrazole
28. 5,7-Dinitro-1,2,3-benzoxadiazole
29. Diazonium salts
30. 1,9-Dicarboxy-2,4,6,8-tetranitrophenazine N<sup>10</sup>-oxide
31. *o,o'*-Azobis(N'-chloroformamidine)
32. 1,2-Dichloro-3,5-dinitrobenzene
33. Guanylurea perchlorate
34. N,N'-Bis(2,3-dihydroxypropyl)oxamide tetranitrate
35. Diethanolnitramine dinitrate
36. Diethanolnitrosamine dinitrate
37. Diethylene glycol dinitrate
38. N,N'-Dimethylethylenedinitramine
39. 2-(Methoxymethyl)-2-nitro-1,3-propanediol dinitrate
40. 2-Ethyl-2-nitro-1,3-propanediol dinitrate
41. 2,2-Dimethyl-1,3-propanediol dinitrate
42. 1,3-Dinitramino-2-propanol nitrate
43. 2,4-Dinitroaniline
44. 2,4-Dinitroanisole
45. *m*-Dinitrobenzene
46. 2,4-Dinitrobenzyl nitrate
47. 2,2-Dinitro-1-butanol
48. 1-Chloro-2,4-dinitrobenzene
49. 4,6-Dinitro-*o*-cresol
50. N,N'-Diethyl-N,N'-dinitrooxamide

CONFIDENTIAL

C-

51. 1,4-Dinitro-2,5-piperasinedione
52. 2,3-Dimethyl-2,3-dinitrobutane
53. 3,6-Dimethyl-1,4-dinitro-2,5-piperasinedione
54. 3a,6a-Dimethyl-1,6-dinitroglucuril
55. 5,5-Dimethyl-1,3-dinitrohydantoin
56. 4,6-Dinitro-m-xylene- $\alpha,\alpha'$ -diol dinitrate
57. 2,5-Dinitro-p-xylene- $\alpha,\alpha'$ -diol dinitrate
58. N,N-Dimethyl-N,N'-dinitrooxamide
59. 2,2-Dimethyl-1,3-dinitropropane
60. N,N'-Dimethyl-N,N'-dinitrosulfamide
61. N,N'-Dimethyl-N,N'-dinitrotartrazide dinitrate
62. 1,5-Dinitro-2,3-dinitrosobenzene
63. N,N'-Bis(2-hydroxyethyl)-N,N'-dinitrooxamide dinitrate
64. N,N'-Bis(2-hydroxyethyl)-N,N'-dinitrosulfamide dinitrate
65. 1,1-Dinitroethane
66. 1,3-Dinitro-2-imidazolidone
67. 2,5-Dinitrofuran
68. 1,6-Dinitroglucuril
69. Tetrahydro-5-hydroxy-1,3-dinitro-2(1H)-pyrimidone nitrate
70. 2-Methyl-1,2-dinitropropane
71. 2,4-Dinitrophenol
72. 2-(2,4-Dinitrophenoxy)ethanol nitrate
73.  $\alpha,\beta,5$ -Trinitrotoluene
74. 2-(3,5-Dinitrophenyl)-2-nitro-1,3-propanediol dinitrate
75. 1,4-Dinitropiperazine
76. 1,1-Dinitropropane
77. 2,2-Dinitro-1-propanol
78. 2,4-Dinitrosorsorcinol
79. N,N'-Dinitrosobenzamide
80. 2,4-Dinitrosorsorcinol
81. 2,5-Bis(hydroxymethyl)-2,5-dinitro-1,6-hexanediol tetranitrate
82. 2,6-Bis(hydroxymethyl)-2,6-dinitro-1,7-heptanediol tetranitrate
83. 1,10-Dinitro-o-1,4,7,10-tetraepicryl-1,4,7,10-tetraolefane
84. 2,4-Dinitrotoluene
85. Diethanolamine trinitrate
86. 4,6-Dinitro-m-xylene
87. Oxidimethanol dinitrate
88. 2,4,6,8-Tetranitro-2,4,6,8-tetraazanonane-1,9-diol dinitrate
89. Dipentaerythritol hexanitrate
90. N,N'-Dipicryl-1,3-propanedinitramine
91. 3,7-Dinitro-1,3,5,7-tetraazabicyclo[3.3.1]nonane
92. Dulcitol hexanitrate
93. Erythritol tetranitrate
94. Ethanolamine dinitrate
95. Ethylenediamine dinitrate
96. Ethylenedinitramine
97. 1,2-Ethanediol diacetate dinitrate
98. 1,2-Ethanediol dinitrate
99. 1,2-Ethanediol glycolate dinitrate
100. 1,2-Ethanediol lactate dinitrate
101. 3,4-Bis(2-nitroethyl)-1,2,3,4-dioxadiazetidina
102. Ethyl nitrate
103. Galactan trinitrate
104. 1,3(or 2,3)-dichloro-2(or 1)-propanol nitrate

CONFIDENTIAL

TABLE 139 (cont'd)

C-

105. Glycerol 1,2-dinitrate
106. Monocacetin 1,3(or 2,3)dinitrate
107. 3-(2,4-dinitrophenoxy)-1,2-propanediol dinitrate
108. 1,3-Dipicrin 2-nitrate
109. 3-Chloro-1,2-propanediol dinitrate,
110. 1-Monoglycolin trinitrate
111. 1-Monolactin trinitrate
112. 3-Methoxy-1,2-propanediol dinitrate
113. 1-Monopicrin dinitrate
114. Guanidine nitrate
115. Guanidine perchlorate
116. 1-Guanyl-3-nitrourea
117. N,N-Dichloroethylamine
118. 3,4,8,9,12,13-Hexoxa-1,6-diazabicyclo [4.4.4] tetradecane
119. 1,2,3,4,5,6-Benzenohexamethanol hexanitrate
120. Hexamethylenetetramine dinitrate
121. Hexamethylenetetramine diperchlorate
122. Hexamethylenetetramine monopерchlorate
123. 2,2',4,4',6,6'-Hexanitroazobenzene
124. 2,2',4,4',6,6'-Hexanitrobiphenyl
125. 2,2',4,4',6,6'-Hexanitrodiphenylamine
126. 2,2',3,4',6,6'-Hexanitrodiphenylamine
127. N,N-Dipicrylethanolamine nitrate
128. 2,2',4,4',6,6'-Hexanitrodiphenyl ether
129. 2,3',4,4',6,6'-Hexanitrodiphenyl ether
130. N,N'-Dipicrylethylenedinitramine
131. 1,3-Dipicrylguanidine
132. N-Methyl-2,2',4,4',6,6'-hexanitrodiphenylamine
133. 2,2',4,4',6,6'-Hexanitrodiphenyl sulfide
134. 2,2',4,4',6,6'-Hexanitrodiphenyl sulfone
135. 1,3-Dipicrylurea
136. Hexanitroethane
137. 2,2',4,4',6,6'-Hexanitrohydrazobenzene
138. 2,2',4,4',6,6'-Hexanitrooxanilide
139. Hexanitrosobenzene
140. Hydrazine monochlorate
141. N,N'-Bis(azidoformamide)
142. Hydrazine dinitrate
143. Hydrazine monopерchlorate
144. Hydrogen cyanide
145. Inositol hexanitrate
146. Inulin trinitrate
147. Iodose and iodoxy compounds
148. Lactose octanitrate
149. Lead azide
150. Styptic acid, lead salt
151. Maltose octanitrate
152. Mannitol hexanitrate
153. Melamine dinitrate
154. Mercuric fulminate
155. Methazonic acid
156. Methylamine nitrate
157. Nitroform, aminylamine salt

CONFIDENTIAL

TABLE 139 (cont'd)

G-

- 158. Methylamine monoperochlorate
- 159. 4-Methyl-2,6-dinitro-2,4,6-triazahexane
- 160. N,N'-methylenebis(N-nitrosodihydroxyamine)
- 161. (Methylenedioxy)dimethanol dinitrate
- 162. Methanediol dinitrate
- 163. N-Methylethylenedinitramine
- 164.  $\alpha$ -Methylglucoside tetranitrate
- 165. 1,2,3-Butanetriol trinitrate
- 166. Methyl nitramine
- 167. Methyl nitrate
- 168. 2-Methyl-2-nitro-1,3-propanediol dinitrate
- 169. 2,2,4-Tri-(hydroxymethyl)-4-methyl-1,3,5-pentanetriol hexanitrate
- 170. N-Ethylmethylenedinitramine
- 171. 5-Nitrazino-1,2,4,1H-triazole
- 172. Pentaerythritol diglycolate tetranitrate
- 173. Nitroacetone nitrile
- 174. p-Nitrobenzyl nitrate
- 175.  $\alpha$ ,2,4-Trinitrobenzene azoethane
- 176. 5-Nitro-p-toluenediazonium 2-sulfonate
- 177. 5,5-Dimethyl-1-nitrohydantoin
- 178. 2-Nitroethanol
- 179. Nitroethylene polymer
- 180. 2-Nitroethanol nitrate
- 181. Nitroglycerin
- 182. Nitroguanidine
- 183. Nitroguanidine nitrate
- 184. 1-Nitrohydantoin
- 185. Nitrosobenzene
- 186. 5-Methyl-1-nitrohydantoin
- 187. N-Methyl-N-nitroglycolamide nitrate
- 188. 2-Methyl-2-nitro-1-propanol nitrate
- 189. 2-Nitro-2-(m-nitrophenyl)-1,3-propanediol dinitrate
- 190. 1-Nitro-3-(nitrosoamino)guanidine
- 191. 3-Nitro-2-oxazolidone
- 192.  $\alpha$ , $\alpha$ ,m-Trinitrotoluene
- 193. 1-(p-nitrophenyl)-1,2-ethanediol dinitrate
- 194. 2-Nitro-1-(o-nitrophenyl)ethanol nitrate
- 195. Dimethylnitrosamine
- 196. 2,3,5,6-Tetranitroso-1,4-dinitrobenzene
- 197. Nitrosoguanidine
- 198. Urea picrate
- 199. 5-Nitro-1H-tetrazole
- 200. Nitrourea
- 201. Nitrourea picrate
- 202. N-Methylethanolnitramine nitrate
- 203. Dipicryl lead
- 204. Cyclohexene ozonide
- 205. 2,2-Bis(methoxymethyl)-1,3-propanediol dinitrate
- 206. 2-(Hydroxymethyl)-2-(methoxymethyl)-1,3-propanediol trinitrate
- 207. Pentaerythritol tetranitrate
- 208. 2,3,4,5,6-pentanitroaniline
- 209. 1-(2,4-Dinitrophenyl)-2-(2,4,5-trinitrophenyl)ethanol
- 210. 2,2',4,4',6-Pentanitrodiphenyl ether

CONFIDENTIAL



G-

211. 2,2',4,4',6-Pentanitrodiphenyl sulfone  
 212. Ethyl perchlorate  
 213. Acetyl peroxide  
 214. *m*-Phenylenediamine diperchlorate  
 215.  $\alpha$ -Azidoformazidine picrate  
 216. Picric acid  
 217. Picryl chloride  
 218. Picryl fluoride  
 219. 1,2-Propanediol dinitrate  
 220. Propyl nitrate  
 221. Pyridine perchlorate  
 222. Quebrachitol pentanitrate  
 223. Sucrose octanitrate  
 224. Tetraazidoquinone  
 225. 4-Guanyl-1-(nitroguaninoguanyl)-1-tetrazene  
 226. Tetramethylammonium perchlorate  
 227. 1,3,8,10-Tetrazo-5,6,12,13-tetroxacyclotetradecane-2,9-dione  
 228. 2-Hydroxy-1,1,3,3-cyclohexanetetraethanol pentanitrate  
 229. 2,2,6,6-Tetrakis(hydroxymethyl)cyclohexanone tetranitrate  
 230. 2-Hydroxy-1,1,3,3-cyclopentanetetraethanol pentanitrate  
 231. 2,2,5,5-Tetrakis(hydroxymethyl)cyclopentanone tetranitrate  
 232. 2,2-Bis(nitraminomethyl)-1,3-propanedinitramine  
 233. 2,3,4,6-Tetranitroaniline  
 234. 2,3,5,6-Tetranitroanisole  
 235. 3,3',5,5'-Tetranitro-*p,p'*-azoxytoluene  
 236. 1,2,3,5-Tetranitrobenzene  
 237. 3,3'-Oxydi-1,2-propanediol tetranitrate  
 238. 2,2',4,4'-Tetranitrodiphenyl ether  
 239. 1,1,2,2-Tetranitroethane  
 240. Tetranitromethane  
 241. 1,3,6,8-Tetranitronaphthalene  
 242. 2,3,4,6-Tetranitrophenol  
 243. *N*-Methyl-*N*,2,3,4,6-pentanitroaniline  
 244. *N*,2,3,4,6-pentanitroaniline  
 245. 3-nitro-1,2,4,5-tetranitrosobenzene  
 246. 4,4'-Bis(*N,N*-dimethyl-2,6-dinitroaniline)  
 247. 2,3,4,6-Tetranitrotoluene  
 248. Tetrakis(2-hydroxyethyl)ammonium pentanitrate  
 249. *N,N'*-Dipicryl-2,2-bis(*N*,2,4,6-tetranitroanilinosmethyl)-1,3-propanedinitramine  
 250. 1-(Aminoguanyl)-4-guanyl-1-tetrazene  
 251. 1H-Tetrazole  
 252. 5-Azido-1H-tetrazole  
 253. Tetryl  
 254. Triazenes  
 255. 1,2,4,1H-Triazole  
 256. 3,3,6,6,9,9-Hexamethyl-1,2,4,5,7,8-hexoxacyclononane  
 257. Hexahydro-1,3,5-trihydroxy-*s*-triazine trinitrate  
 258. 1,2-Propanedinitramine  
 259. 1,3-Propanediol dinitrate  
 260. 1,3-Propanediol diperchlorate  
 261. 2-Butyl-2-(hydroxymethyl)-1,3-propanediol trinitrate  
 262. 2-(Caloromethyl)-2-(hydroxymethyl)-1,3-propanediol trinitrate  
 263. 2-(Hydroxymethyl)-2-methyl-1,3-propanediol trinitrate  
 264. 2-(Hydroxymethyl)-2-nitro-1,3-propanediol trinitrate

CONFIDENTIAL

TABLE 139 (cont'd)

C-

- 265. 2-Ethyl-2-(hydroxymethyl)-1,3-propanediol trinitrate
- 266. Trinitroacetoneitrile
- 267. 2,4,6-Trinitro-m-anisidine
- 268. 2,4,6-Trinitro-m-phenetidine
- 269. 3-Amino-2,4,6-trinitrophenol
- 270. Picramide
- 271. 2,4,6-Trinitroanisole
- 272. 2,4,6-Trinitrobenzaldehyde
- 273. 1,2,4-Trinitrobenzene
- 274. 1,3,5-Trinitrobenzene
- 275. 2,4,6-Trinitrobenzoic acid
- 276. 2,4,6-Trinitrobenzyl nitrate
- 277. 2,4,6-Trinitro-m-cresol
- 278. N,N-Dimethylpicramide
- 279. N,N'-Dimethyl-N,N',2,4,6-pentanitro-m-phenylenediamine
- 280. 1,2,3-Trinitro-7,8-dinitroso-2-naphthol
- 281. 2,2'-(2,4,6-Trinitro-m-phenylene)diethanol dinitrate
- 282. 3-Ethyl-2,2,3-trinitropentane
- 283. N<sup>2</sup>,N<sup>4</sup>,N<sup>6</sup>-Trinitromelamine
- 284. 2,4,6-Trinitroresitylene
- 285. 2-Methyl-2,3,3-trinitrobutane
- 286. N<sup>1</sup>-Methyl-N<sup>2</sup>,2,4,6-tetranitro-m-phenylenediamine
- 287. N-Methyl-N,2,4,6-tetranitro-m-anisidine
- 288. N-Methyl-N,2,4,6-tetranitro-m-phenetidine
- 289. 3-(Methyltriamino)-2,4,6-trinitrophenol
- 290. N-Methyl-N,2,4,6-tetranitro-m-toluidine
- 291. N-Methyl-2,2,3-trinitropentane
- 292. N-Methylpicramide
- 293. 1,3,8-Trinitronaphthalene
- 294. 2,4,6-Trinitrophenetole
- 295. 2-(2,4,6-Trinitrophenoxy)ethanol nitrate
- 296. Picryl azide
- 297. N-Butyl-N,2,4,6-Tetranitroaniline
- 298. 3-(N,2,4,6-Tetranitroaniline)-1,2-propanediol dinitrate
- 299. 2,4,6-Trinitro-m-phenylenediamine
- 300. N-Ethyl-N,2,4,6-tetranitroaniline
- 301. 2-Picrylethanol nitrate
- 302. Picrylguanidine
- 303. Picrylhydrazine
- 304. N-Methoxy-N,2,4,6-tetranitroaniline
- 305. 2-(N,2,4,6-Tetranitroanilino)-1,2-ethanediol nitrate
- 306. N,2,4,6-Tetranitroaniline
- 307. 2-(N,2,4,6-Tetranitroanilino)ethanol nitrate
- 308. 2-(Hydroxymethyl)-2-(N,2,4,6-tetranitroanilino)-1,3-propanediol trinitrate
- 309. Trinitrochlorogucinol
- 310. Styplinic acid
- 311. 2,2'-(2,4,6-Trinitro-m-phenylenedioxy)diethanol dinitrate
- 312. 2,4,6-Trinitrostilbene
- 313. 2,4,6-Trinitrotoluene
- 314. 2,3,4-Trinitrotoluene
- 315. 1,3,5-Triazido-2,4,6-trinitrobenzene
- 316. N,N',N''-Trimethyl-N,N',N'',2,4,6-hexanitro-1,1,5-benzenetriamine
- 317. Triethanolamine tetranitrate

CONFIDENTIAL

CONFIDENTIAL

TABLE 139 (cont'd)

C-	
318.	2,4,6-Trinitro-m-xylene
319.	1,1',1"-Tris(methylnitramino)trimethylamino
320.	Urea nitrate
321.	4-Hydro-1-butan-3-yne polymer
322.	Vinyl nitrate polymer
323.	$\alpha,\alpha'$ -Diasido-p-xylene
324.	n-Propyl picrate
325.	Allyl picrate
326.	2-Propenyl picrate
327.	Nitroform
328.	Methylenedinitramine
329.	N-Methyl-N,2,3,4,5-pentanitroaniline
330.	2,2,2-Trinitroethanol
331.	1,3-Bis(2,2,2-trinitroethyl)urea
332.	Ethylenediamine dinitroform salt
333.	N,N'-Dichloroethylenedinitramine
334.	(2,4-Dinitrophenyl)hydrazine
335.	1,1-Dinitro-N-(1-nitroethylidene)ethylamine oxide
336.	$\alpha,\alpha,\alpha'$ -Trifluoro-3,5-dinitrotoluene
337.	Bis(2,2,2-trinitroethyl)nitramine
338.	5-Amino-1H-tetrazole hydrate
339.	5-Amino-1H-tetrazole nitrate
340.	2,2,2-Trinitroethanol 4,4,4-trinitrobutyrate
341.	Tris(2,2,2-trinitroethyl)phosphate
342.	5-Amino-1H-tetrazole
343.	2,2,3,3-Tetranitrobutane
344.	E,N'-Methylenebis(N-nitroformamide)
345.	5-(2,2,2-trinitroethylamino)-1H-tetrazole
346.	Methylenedinitramine dihydrazine salt
347.	2,4,6-Trinitro-2,4,6-triazabptans-1,7-diol dinitrate
348.	$\alpha$ -Azido-N-nitroformamide
349.	4,6-Dinitramino-s-triazolo-2-ol
350.	N,N'-Methylenebis(2,2,2-trinitroacetamide)
351.	2,2-Dinitroacetamide
352.	Hydrazine mononitrate
353.	Hydrazine
354.	5,5'-Hydrazodi-1H-tetrazole
355.	By-product from Madina synthesis
356.	2,4-Dinitrothiophene
357.	4-Methyl-3,5-dinitro-1,2,4,4H-triazole
358.	Methylenebis(nitroimino)dimethanol dinitrate
359.	5-Hitramino-1H-tetrazole
360.	Azidodithioformic acid
361.	Azidosulfonic acid
362.	Aniline
363.	p-Azidoaniline perchlorate
364.	2-Azido-4,6-dinitrophenol
365.	Carbonyl azide
366.	p-Azidobenzene
367.	1,2-Diazidethane
368.	1,3-Diazidopropane
369.	$\alpha,\alpha'$ -Diasido-2,5-dinitro-p-xylene
370.	1,3,5-Triazido-2,4-dinitrobenzene

CONFIDENTIAL

CONFIDENTIAL

TABLE 139 (cont'd)

371. Hydrazinium esic  
372.  $\alpha$ -Azido-p-nitrotoluene  
373. 2,4-Diiodo-1,3,5-trinitrobenzene  
374. 1,2,3-Benzoxadiazole  
375. 5,7-Dinitro-1,2,3-benzoxadiazole-4,6-diol  
376. 5,7-Dinitro-1,2,3-benzoxadiazol-4-ol  
377. 5,7-Diethyl-4,6-dinitro-1,2,3-benzoxadiazole  
378. 5-Hydroxy-6-nitro-1,2,3-benzoxadiazole-4,7-dione  
379. 4,5,6,7-Tetranitro-1,2,3-benzoxadiazole  
380. Benzenediazonium nitrate  
381. p-(2,4-Dinitroanilino)benzenediazonium nitrate  
382. 1,5-Dichloro-2,4-dinitrobenzene  
383. 3,5-Dinitroanisole  
384. 1,2-Dinitroethane  
385. 2,5-Dinitrophenol  
386. 2,6-Dinitrophenol  
387. 2,2-Dinitropropane  
388. 1,3-Dinitropropane  
389. 4,6-Dinitroresorcinol  
390. 2,6-Dinitrotoluene  
391. 3,5-Dinitro-o-xylene  
392. Glycerol 1,3-dinitrate  
393. (Nitromethoxy)methanol nitrate  
394. 1,2-Ethandiol polymer dinitrate  
395. 1,2,4-Butanetriol trinitrate  
396. 2,3',4,4',6-Pentanitrodiphenyl ether  
397. Ammonium perchlorate  
398. Ethylenediamine diperchlorate  
399. 1,2-Propanediamine diperchlorate  
400. 1,3-Propanediamine diperchlorate  
401. Bis(2,2,2-trinitroethyl)amine  
402.  $\alpha$ -Azidoformaldimine perchlorate  
403. 2-Methyl-1,3-butanediamine diperchlorate  
404. 1-Naphthylamine perchlorate  
405. 3-(Perchlorato-mercuri)-1,2,4-oxadiazolmercurtan-3-ol  
406. Benzoyl peroxide  
407. Di-2,4-cyclopentadien-1-yl peroxide  
408. Picryl hydroperoxide, sodium salt  
409. 2,3,4,6-Tetranitroanisole  
410. 2,3,5-Trinitrotoluene  
411. 2,3,6-Trinitrotoluene  
412. 2,4,5-Trinitrotoluene  
413. 3,4,5-Trinitrotoluene  
414. 3,4,5-Trinitro-o-xylene  
415. 3,4,6-Trinitro-o-xylene  
416. 2,3,5-Trinitro-p-xylene  
417. (Nitroimino)diacetonitrile  
418. Dimethyltriazine  
419. 1,7-Dinitro-1,4,7-triazahepthane  
420. 1,10-Dinitro-1,4,7,10-tetrazadecane  
421. 1-Methyl-3-nitroguanidine  
422. Nitramide  
423. N-Methyl-N,2,4,5,6-pentanitro-m-toluidine

CONFIDENTIAL

CONFIDENTIAL

TABLE 139 (cont'd)

C-

424.	2-tert-Butoxyethanol nitrate
425.	Tetrahydrofuran-3,4-diol dinitrate
426.	1,3-Diethoxy-2-propanol nitrate
427.	2-Propanol nitrate
428.	Mannitan tetranitrate
429.	Pentaerythritol diacetate dinitrate
430.	Pentaerythritol diformate dinitrate
431.	Pentaerythritol lactate tetranitrate
432.	Pentaerythritol m-nitrobenzoate trinitrate
433.	Sorbitol hexanitrate
434.	2,2,2-Trinitroethanol nitrate
435.	Aminoguanidine nitrate
436.	Guanylurea nitrate
437.	Hydroxylamine nitrate
438.	N-Methylhydroxylamine nitrate
439.	Piperazine dinitrate
440.	Succinonitrile
441.	Malononitrile
442.	1,4-Dibromo-2,3-dinitrobenzene
443.	1-Bromo-2,4-dinitrobenzene
444.	2,4-Dinitroanisylare
445.	3,5-Dinitrosalicylic acid
446.	1,3,5-Tribromo-2,4-dinitrobenzene
447.	2,2',4,4',6,6'-Hexanitroxybenzene
448.	2,2',4,4',6,6'-Hexanitrobiphenyl
449.	3,3'-Azobis(2,4,6-trinitrophenol)
450.	1,3-Dimethyl-1,3-dipicrylurea
451.	2,2',4,4',6,6'-Hexanitrostilbene
452.	2,2',4,4',6,6'-Hexanitro-3,3',5,5'-biphenyltetrrol
453.	2,2',4,4'-Tetranitrobenzophenone
454.	Tetranitrosorsorcinol
455.	2,4,5,6-Tetranitro-m-xylene
456.	1,3,5-Trichloro-2,4,6-trinitrobenzene
457.	Benzene
458.	2-Ethyl-1,3,5-trinitrobenzene
459.	1-Methyl-1,3-dinitrosoguanidine
460.	Benzene trizonide
461.	Biphenyl tetraozonide
462.	Naphthalene diozonide
463.	N,N,N',N'-Tetrabromocethylenediamine
464.	N,N'-Ethylenebis(N-chloro-p-nitrobenzamide)
465.	1,4-Dibromopiperazine
466.	N,N-Dichloroformamide
467.	2,3-Butanediol dinitrate
468.	3-(Chloratomercuri)-1,2,4-oxadimercuretan-3-ol
469.	(Chloratomercuri)mercuriacetaldehyde
470.	Methylenebis(N-nitrosodihydroxylamine) anhydrous sodium salt
471.	Methylenebis(N-nitrosodihydroxylamine) monohydrate sodium salt
472.	Methylenebis(N-nitrosodihydroxylamine) diammonium salt
473.	Methylenebis(N-nitrosodihydroxylamine) monoammonium salt
474.	Methylenebis(N-nitrosodihydroxylamine) anhydrous cadmium salt
475.	Methylenebis(N-nitrosodihydroxylamine) dihydrate cadmium salt
476.	Methylenebis(N-nitrosodihydroxylamine) lead salt
477.	Methylenebis(N-nitrosodihydroxylamine) bis(diethanolamine dinitrate) salt

CONFIDENTIAL

CONFIDENTIAL

TABLE 139 (cont'd)

C-

- 478. Methylenebis(N-nitrosodihydroxylamine) monohydroxylamine salt
- 479. Dipicryl mercury
- 480. 1-Nitronaphthalene
- 481. 1,5-Dinitronaphthalene
- 482. 1,8-Dinitronaphthalene
- 483. 1,3,5-Trinitronaphthalene
- 484. 1,4,5-Trinitronaphthalene
- 485. 2-Methyl-1,2,3-propanetriol trinitrate
- 486. 1,4,5,8-Tetranitronaphthalene
- 487. 1,3,5,8-Tetranitronaphthalene
- 488. 5-[Bis(2,2,2-trinitroethyl)amino]-1H-tetrazole
- 489. 3-(2-Hydroxyethoxy)-1,2-propanediol trinitrate
- 490. 1-Hydroxy-5,5'-azodi-1H-tetrazole
- 491. 5,5'-Azodi-1H-tetrazole
- 492. 5,5'-Bis(1-hydroxy-1H-tetrazole)
- 493. 5,5'-Bi-1H-tetrazole
- 494. Triazato[4,1-a]tetrazole
- 495. 3,4-Benzylidene-4-guanyl-1-(1H-tetrazol-5-yl)-1-tetrazene
- 496. 3-Phenyl-1-(1H-tetrazol-5-yl)-1-tetrazene
- 497. 4-Guanyl-1-(1H-tetrazol-5-yl)-1-tetrazene
- 498. 1-Hydroxy-1H-tetrazole-5-methanol
- 499. 1-Hydroxy-5-nitro-1H-tetrazole
- 500. 1-Hydroxy-1H-tetrazole-5-methanol nitrate
- 501. 1-Hydroxy-1H-tetrazole
- 502. 5-(2,4,6-trinitroanilino)-1H-tetrazole
- 503. 1(or 5)-Amino-5(or 1)-nitramino-1,2,4,4H-triazole
- 504. 5-(Nitrosoamino)-1,2,4,4H-triazole-3-carboxylic acid
- 505. Biacetylenedicarboxylic acid
- 506. 2,6-Dinitro-p-cresol
- 507. m-Nitroaniline
- 508. Phenol
- 509. p-Nitrophenol
- 510. Pentanitrophenol
- 511. Phenylhydrazine
- 512. (p-Nitrophenyl)hydrazine
- 513. (2,3,4,6-Tetranitrophenyl)hydrazine
- 514. (2,3,4,5,6-Pentanitrophenyl)hydrazine
- 515. Nitrobenzene
- 516. Pentanitrobenzene
- 517. Hexanitrobenzene
- 518. 1,1,2,2-Ethanetetranitramine
- 519. 2-Nitrothiophene
- 520. 3-Methyl-2-nitrosothiophene
- 521. 3-Methyl 2,4-dinitrothiophene
- 522. Dinitroacetylene
- 523. 1,1,1,3,5,5,5-Heptanitropentane
- 524. 3,4,5-Triamino-1,2,4,4H-triazole
- 525. Guanazole
- 526. 4-Amino-1,2,4,4H-triazole-3,5-diol
- 527. Urazole
- 528. 1,2,3-Triaminoguanidine
- 529. 1,2,3-Triaminoguanidine dinitrate
- 530. 1,3-Diaminoguanidine
- 531. 1,3-Diaminoguanidine nitrate

CONFIDENTIAL

C-

532.	Oxalic acid, monohydrazine salt
533.	Oxalic acid, bis(hydroxylamine)salt
534.	Oxalic acid, semicarbazide salt
535.	Oxalic acid, aminoguanidine salt
536.	Carbohydrazide
537.	Oxalic acid, monocarbohydrazide salt
538.	1-Carbaalylcarbohydrazide
539.	1,5-Dicarbonylcarbohydrazide
540.	2,4-Diamino-s-triazine
541.	N,N'-Bi(ethyl carbamate)
542.	Biurea
543.	H-Guanidinoxamic acid
544.	5-Amino-1,2,4,1H-triazole-3-carboxylic acid
545.	1,2,3-Trisaminoguanidine monohydrochloride
546.	2,4-Diamino-6-methyl-s-triazine dinitrate
547.	Tetrahydro-1,3,5,7-tetraimino-s-triazolo[a]-s-triazolo
548.	2,4-Diguanidino-s-triazine dinitrate
549.	Cyanuric trihydrazide
550.	1-vanoguanidine
551.	Melamine
552.	Oxalic dihydrazide
553.	Oxamide
554.	Cyanuric acid
555.	6-Nitramino-s-triazine-2,4-diol
556.	m-Nitrotoluene
557.	o-Nitrophenol
558.	n-Nitrophenol
559.	m-Nitrophenetole
560.	o-Nitrophenetole
561.	p-Nitrophenetole
562.	o-Nitroaniline
563.	p-Nitroaniline
564.	o-Nitroacetanilide
565.	m-Nitroacetanilide
566.	p-Nitroacetanilide
567.	1,1,1-Trinitroethane
568.	2,4-Dinitrophenetole
569.	1,1,2-Trinitroethane
570.	2-Methyl-2,3,3-trinitropentane
571.	Picramic acid
572.	p-Nitrobenzaldehyde
573.	1,2,6-Hexanetriol trinitrate
574.	2,2-Dinitro-1,3-propanediol
575.	4,4-Dinitropimelic acid
576.	1,1,1,3-Tetranitropropane
577.	5,5,5-Trinitro-2-pentanone
578.	Methyl 4,4,4-trinitrobutyrate
579.	1,1,1-Trinitro-2-propanol acetate
580.	4,4,4-Trinitro-N-(2,2,2-trinitroethyl)butyramide
581.	1,3-Bis(2,2,2-trinitroethyl)-2-imidazolidone
582.	4,4-Dimethyl-5,5,5-trinitro-2-pentanone
583.	Diethyl (2,2,2-trinitroethyl)malonate
584.	N,N'-Bis(2,2,2-trinitroethyl)-1,4-piperasinedicarboxamide
585.	Urea

CONFIDENTIAL

TABLE 139 (cont'd)

C-

- 586. Guanidinocetic acid
- 587. Ethylurea
- 588. 1,3-Bis(hydroxymethyl)urea
- 589. 2-Imidazolidone
- 590. 1-Methyl-1-phenylhydrazine
- 591. Hydrazobenzene
- 592. Tetraphenylhydrazine
- 593. Ethylnitrazine
- 594. 1,3-Dinitroimidazolidine
- 595. Acetone
- 596. 3,3'-(Nitroimino)dipropionitrile
- 597. 4,4-Dimethyl-5-nitro-2-pentanone
- 598. 1-Picrylaziridine
- 599. 2-(2-Nitrovinyl)furan
- 600. 2-Nitro-5-(2-nitrovinyl)furan
- 601. 2-(2,4-Dinitrophenyl)-5-nitro-2H-benzotriazole
- 602.  $\alpha,\alpha$ -Dinitrotoluene
- 603. N-(2,4-Dinitrophenyl)diethanolamine
- 604. 2-(2,4-Dinitroanilino)-2-(hydroxymethyl)-1,3-propanediol
- 605. 2-(2,4-Dinitroanilino)-1,3-propanediol
- 606. 2-[2-(2,4-Dinitroanilino)ethylamino]ethanol
- 607. Ethylenebis(ethyl nitroacetate)
- 608. 1,1'-Ethylenebis(3-nitrourea)
- 609. 1-(o-Nitrophenyl)-1,2-ethanediol dinitrate
- 610. 1,2-Propanediol 2-nitrate 1-(5-nitro-2-furrate)
- 611. N-Picryldiethanolamine dinitrate
- 612. N-Methyl-N,5-dinitro-2-furamide
- 613. N,N'-Dimethyloxamide
- 614. N-Methylglycolamide
- 615. 5,5-Dimethylhydantoin
- 616. 3,6-Dimethyl-2,5-piperazinedione
- 617. Nitroethane
- 618. 1-Nitropropane
- 619. 2-Nitropropane
- 620. 1-Nitrobutane
- 621. 2-Nitrobutane
- 622. 1,1'-(2-Hydroxytrimethylene)bis(3-nitroguanidines) dinitrate
- 623. 3-[2,2-Tris(hydroxymethyl)ethoxy]-1,2-propanediol pentanitrate
- 624. 2-(Allyloxymethyl)-2-(hydroxymethyl)-1,3-propanediol trinitrate
- 625. 1-Nitro-3-(1-hydroxyethyl)guanidines nitrate
- 626. 2-(Hydroxymethyl)-2-(propoxymethyl)-1,3-propanediol trinitrate
- 627. 2,2-Bis(allyloxymethyl)-1,3-propanediol dinitrate
- 628. 1-Nitro-3-(2,2,2-trinitroethyl)guanidines
- 629. Ethyl 4,4,4-trinitrobutyrate
- 630. 1-Nitro-3-(2,3-dihydroxypropyl)guanidines dinitrate
- 631. 1-Nitro-3-(2-hydroxyethyl)guanidines nitrate
- 632. 1,5-Pentanediol dinitrate
- 633. 2,3,3,4,5-Pentamethyl-1,1,1,5-tetranitrohexane
- 634. 1,6-Hexanediol dinitrate
- 635. 1,4-Butanediol dinitrate
- 636. Bis(3,3,3-trinitropropyl) sulfone
- 637. Oxalimide dihydrazide dinitrate
- 638. Diethyl 3,3,3-trinitro-1-propanephosphonate
- 639. 4,4,4-Trinitrobutyramide

CONFIDENTIAL



CONFIDENTIAL

TABLE 139 (cont'd)

C-

- 640. Dinitroform, dioxane addition compound
- 641. 5,5,5-Trinitro-2-pentanone N-nitroguanylhydrazone
- 642. 5,5,5-Trinitro-2-pentanone semicarbazone
- 643. 2,2-Dinitro-1-propanol acrylate
- 644. Methyl nitroacetate
- 645. 2,5-Hexanediol dinitrate
- 646. n-Butyl 4,4,4-trinitrobutyrate
- 647. Diethyl 3-hydroxyglutarate nitrate
- 648. 1,2,5-Pentanetriol trinitrate
- 649. 1,2-Butanediol dinitrate
- 650. 1,5-Dinitro-2,3-dinitrosobenzene, potassium salt
- 651. 1,4-Dinitro-2,3-butanediol dinitrate
- 652. 3,3-Dinitro-1,5-pentane diisocyanate
- 653. 2,2,4,4-Tetranitro-1-butanol acetate
- 654. 3,3-Dinitro-1,5-pentanediamine
- 655. Methyl 4,4,6,6-tetranitrocaproate
- 656. 2,2,2-Trinitroethanol methacrylate
- 657. 2,2,4,4-Tetranitro-1,5-pentanediol
- 658. 2-Nitramino-2-imidazoline
- 659. 2-Nitramino-1-nitro-2-imidazoline
- 660. 1,4,5,6-Tetrahydro-2-nitraminopyrimidins
- 661. N-Nitroethylensinine polymer diol dinitrate
- 662. 2,16-Dimethyl-3,6,9,12,15-pentanitro-3,6,9,12,15-pentazahptadecane-2,16-dicarbonitrile
- 663. 2,13-Dimethyl-3,6,9,12-tetranitro-3,6,9,12-tetrazatetradecane-,13-dicarbonitrile
- 664. 2,10-Dimethyl-3,6,9-trinitro-3,6,9-triazaundecane-2,10-dicarbonitrile
- 665. 2,7-Dimethyl-3,6-dinitro-3,6-diazaoctane-2,7-dicarbonitrile
- 666. 2-Nitropropene polymer
- 667. 1-Ethyl-3-nitroguanidine
- 668. 1-Nitro-3-propylguanidine
- 669. 1-Butyl-3-nitroguanidine
- 670. 1-Amyl-3-nitroguanidine
- 671. Guanidine
- 672. Pyridine
- 673. Ethyl acetate
- 674. 4,4'-Isopropylidenebis(6-nitro-o-cresol)
- 675. N-Ethyl-N-nitrosoaniline
- 676. N-Nitrosodiphenylamine
- 677. 1,5-Dimethyl-1H-tetrazole
- 678. 1-Phenyl-1H-tetrazole
- 679. 5-Phenyl-1H-tetrazole
- 680. 5-Methyl-1-phenyl-1H-tetrazole
- 681. 5-Methyl-2-phenyl-2H-tetrazole
- 682. 1,5-Diphenyl-1H-tetrazole
- 683. 2,5-Diphenyl-2H-tetrazole
- 684. 1-Phenyl-1H-tetrazol-5-ol
- 685. 2-Phenyl-2H-tetrazole-5-carboxylic acid
- 686. Ethyl N-1H-tetrazol-5-ylcarbonate
- 687. N-1H-tetrazol-5-ylacetamide
- 688. 1H-Tetrazol-5-ylguanidine
- 689. 5-Nitramino-1H-tetrazole monoguanidine salt
- 690. Benzaldehyde, 1H-tetrazol-5-ylhydrazone

CONFIDENTIAL

CONFIDENTIAL

TABLE 139 (cont'd)

C-

- 691. Hydrazine hydrate
- 692. 4,4'-Cyclohexylidenebis(2,6-dinitrophenol)
- 693. 2,2'-Hydrazodisobutyronitrile
- 694. Acetic 1,2-diphenylhydrazide
- 695. N,N'-Biacetanilide
- 696. N,N'-Bibenzamide
- 697. Malonic dihydrazide
- 698. Succinic dihydrazide
- 699. Thiophane
- 700. Catechol

CONFIDENTIAL

SECTION XVII  
Calculation of Power (nRT)  
of Explosives Compounds and Mixtures

A. Abstract

A previously developed method for the calculation of the impulse propellants has been successfully adapted to the calculation of powers of high explosives, relative to TNT. The success of this method is indicated by the correlation achieved between the calculated power (nRT) and the measured ballistic mortar values for 111 pure compounds, 56 mixtures of organic explosives and 30 mixtures of explosives containing metals. In the case of the pure compounds, the average deviation is +1.6 while the standard deviation is 6.7. In the case of the mixture calculations, the average deviation is +0.1 while the standard deviation is 5.8. Corresponding figures for the metallized mixtures are -0.8 for average and 7.8 for standard deviations.

By using the system described in Section XVI for calculating the heat of combustion, and the method described here for calculating nRT, it is possible to evaluate a pure compound or mixture, as to power, by a few simple calculations without the necessity for making and testing the compound.

B. Introduction

This section describes a simplified method for calculating the power of a high explosive (pure compound or mixture) from thermodynamic data which gives results bearing a direct, virtually equivalent, relation to the power of the explosive as measured in the ballistic mortar and the new-type spherical lead block. Power as used here and as measured in the ballistic mortar is that force exercised by an explosive burning in an enclosed space, large enough only to contain the explosive. Thus, it is energy developed at virtually constant volume. All thermodynamic data are therefore referred to constant volume.

Calculation of the PV work product ( $F = nRT$ ) according to the simplified procedure of Hirschfelder has been used for propellant powders for some time (10). The application of this same method to explosives has been described briefly (6). This presentation extends the calculations and data presented previously. Calculations of a large number of pure compounds, mixtures of organic compounds and metallic mixtures indicate the value of this method as a tool for quickly evaluating the power of an explosive.

In this method, one hundred times the quotient of the calculated energy divided by the energy of an equal weight of TNT expresses the power of the explosive in terms of per cent TNT.

CONFIDENTIAL  
SECURITY INFORMATION

Previous simplified methods employed in the calculation of power of high explosives have assumed an overall specific heat for the product gases formed in the detonation (1). The system described in this report, by applying the Hirochfelder method of calculating temperature, takes into account the variation in specific heat for the various product gases formed over the particular temperature range involved. Thus it is believed that explosion-temperatures calculated in the method described in this report are somewhat more realistic, although it is not claimed that they are the true values, since they do not take into account the dissociation of the product gases. The reaction products which are taken into account are governed by the assumptions stated on page 580.

It is also possible to correlate the nRT values obtained with those measured in the new type spherical lead block, a test method which is to be described in a forthcoming report. This correlation is shown in Table 147 and briefly discussed.

One method for calculating power of high explosive used in the past was the so-called "characteristic product" (Berthelot) which gave power as per cent TNT by the formula

$$P (\% \text{ TNT}) = 0.003594 M \cdot H_E$$

in which  $M$  = moles of gas per kg. of explosive

$H_E$  = heat of explosion in kcal/kg.

This system has been applied to a number of compounds and mixtures (4, 5). Comparison of results by the nRT and Berthelot methods of calculation shows distinct advantages of the former as given in the following:

nRT System

- 1) Correlates observed ballistic mortar power values with calculated values, on a 45° line, thus giving identical figures. (See Figures 80, 81 and 82).
- 2) Applicable to primary and secondary phosphores, auxo-plosives, and mixtures including those containing metallic elements.

Berthelot System

- 1) Correlates calculated values with observed lead block values on a straight line, not 45°. (Figure 64 Reference (5)); and with ballistic mortar on a curved line (Figure 57 Reference (5)).
- 2) The same correlations do not apply to mixtures yielding products partly non-gaseous; i.e., metallized mixtures. (Figure 56 Reference (5)).

CONFIDENTIAL

NRT System

Berthelot System

3) Includes calculation of a realistic value for explosion temperature, necessary when gas dissociation is considered.

3) No temperature calculated.

4) Results of power calculation are capable of being expressed in meaningful units, e.g., cal/g; ft.-lbs./g, etc.

4) Results are comparative only.

A series of the Berthelot calculations have been made and compared with the observed Ballistic Mortar values (5). Although there was no direct, straight-line relationship shown, it was possible to formulate an idealized curve.

C. Discussion

The simplified system presented here is based upon the determination of the  $pV$  work product of the gases of detonation. The common expression for this work product is:  $pV = nRT$ , here written as

$$pV = R \cdot T \cdot \ln \quad (1)$$

where  $R$  is the universal gas constant (1.987 cal/°K/mole),  $T$  is the adiabatic flame temperature, °K, and  $\ln$  is the number of moles of gas formed from one mole of explosive. The parameter,  $\ln$ , is obtained by writing the detonation equation. The flame temperature,  $T$ , must be estimated from the equation

$$T = 298 + \frac{Q_V}{\sum n C_V} \times 10^3 \quad (2)$$

where  $Q_V$  is the heat of explosion at constant volume in kcal/mole,  $n$  is the moles of each gas formed, and  $C_V$  is the average heat capacity in cal/mole of each gas, at constant volume, over the range 298 to  $T$ . This temperature,  $T$ , is actually obtained by a process of iteration, using the specific heat values tabulated (Table 149). Examples of detailed calculations are given in Subsection D below.

The substitution of appropriate quantities for  $\ln$  and  $T$  in equation (1) gives a value, which when divided by a corresponding value for TNT, results in a power for the explosive, in terms of TNT, which compares directly with observed ballistic mortar and spherical lead block measurements.

It is fully recognized that in this very simple method of calculation there have been discounted several factors, among which are the possibility of dissociation of the product gases, the effect of  $\gamma$  (the ratio of

CONFIDENTIAL  
SECURITY INFORMATION

specific heats,  $C_p/C_v$ ), and deviations from the perfect-gas law.

Dissociation of the product gases would result in an increase in  $\gamma$  and a corresponding decrease in the adiabatic flame temperature, thus counterbalancing, at least to some extent, the effects described.

It can be shown that in a simplified system such as this,  $\gamma$  has little effect upon the final relative values. This is due to the fact that for all practical purposes only diatomic and triatomic gases are produced in the detonation. A review of some of the more typical high explosives reveals that  $\gamma$  varies over an extremely small range, (Table 140) thus making a refined treatment of this parameter unnecessary in a system where only relative values are involved.

Table 140  
Ratio of Specific Heats,  $\gamma$ , for Product Gases  
for Selected High Explosives

No.	Name	COMPOUND	$\gamma$ for Product Gases, 298°K
22	HMX		1.37
23	RDX		1.37
35	Diethylnitramine dinitrate		1.36
37	Diethylene glycol dinitrate		1.38
208	2,3,4,5,6-Pentanitroaniline		1.36
216	Picric acid		1.39
233	2,3,4,6-Tetranitroaniline		1.38
240	Tetranitromethane		1.39
313	2,4,6-Trinitrotoluene		1.40
328	Methylenedinitramine, MADINA		1.35
331	1,3-Bis(2,2,2-trinitroethyl)urea		1.34
337	Bis(2,2,2-trinitroethyl)nitramine, HOX		1.37
343	2,2,3,3-Tetramitrobutane		1.36
	Average		1.37

Even at high temperatures, and thus lower  $\gamma$ 's, the deviation from an average value is very small.

The heats of combustion used in these calculations were obtained by the method described in Section XVI. It was found that, generally, the use of calculated in place of observed heats of combustion gave nRT values which correlated better with the observed ballistic mortar values. The

CONFIDENTIAL

CONFIDENTIAL  
SECURITY INFORMATION

specific heats,  $C_p/C_v$ ), and deviations from the perfect-gas law.

Dissociation of the product gases would result in an increase in  $\gamma$  and a corresponding decrease in the adiabatic flame temperature, thus counterbalancing, at least to some extent, the effects described.

It can be shown that in a simplified system such as this,  $\gamma$  has little effect upon the final relative values. This is due to the fact that for all practical purposes only diatomic and triatomic gases are produced in the detonation. A review of some of the more typical high explosives reveals that  $\gamma$  varies over an extremely small range, (Table 140) thus making a refined treatment of this parameter unnecessary in a system where only relative values are involved.

Table 140  
Ratio of Specific Heats,  $\gamma$ , for Product Gases  
for Selected High Explosives

No.	Name	$\gamma$ for Product Gases, 298°K
22	HMX	1.37
23	RDX	1.37
35	Diethylenetriamine dinitrate	1.36
37	Diethylene glycol dinitrate	1.38
208	2,3,4,5,6-Pentanitroaniline	1.36
216	Picric acid	1.39
233	2,3,4,6-Tetranitroaniline	1.38
240	Tetranitroethane	1.39
313	2,4,6-Trinitrotoluene	1.40
328	Methylenedinitramine, MMDNA	1.35
331	1,3-Bis(2,2,2-trinitroethyl)urea	1.34
337	Bis(2,2,2-trinitroethyl)nitramine, HOI	1.37
343	2,2,3,3-Tetramitrobutane	1.36
	Average	1.37

Even at high temperatures, and thus lower  $\gamma$ 's, the deviation from an average value is very small.

The heats of combustion used in these calculations were obtained by the method described in Section XVI. It was found that, generally, the use of calculated in place of observed heats of combustion gave NIT values which correlated better with the observed ballistic mortar values. The

CONFIDENTIAL

CONFIDENTIAL  
SECURITY INFORMATION

observed heats of combustion were used in the calculations shown in the original presentation of this method (6).

Other thermodynamic data used were derived from OSRD Report 547 (10), the Bureau of Standard Tables (7), and the open literature (2, 9, 11, 12). These data, as used in these calculations, are given in Table 141, 142 and 149.

Table 141 gives the energy values required for converting H<sub>2</sub>O liquid to H<sub>2</sub>O gas and heats of combustion to heats of explosion.

Table 141  
Thermodynamic Data for Selected Reactions

	<u>Constant + Pressure</u>	<u>Converted to Constant Volume</u>
H <sub>2</sub> O (liq.) → H <sub>2</sub> O (gas)	10.519	9.927 kcal/mol
C + O <sub>2</sub> → CO <sub>2</sub>	94.652	93.460
CO + 0.5 O <sub>2</sub> → CO <sub>2</sub>	67.636	67.341
H <sub>2</sub> + 0.5 O <sub>2</sub> → H <sub>2</sub> O (gas)	57.799	57.502

\* Reference (7)

Table 142 gives the heats of combustion and heats of fusion and vaporization for the metals and their corresponding oxides entering into these calculations.

Table 142  
Thermodynamic Data for Selected Metals  
and their Corresponding Oxides

	<u>Q<sub>c</sub><sup>v</sup> (H<sub>2</sub>O gas)</u>	<u>Q<sub>fusion</sub></u>	<u>Q<sub>vaporization</sub></u>	<u>Q<sub>sublimation</sub></u>	<u>b.p., °K</u>
Al	199.54 a/	2.6 a/	67.9 a/	70.5	2330*
Al <sub>2</sub> O <sub>3</sub>	--	2.6 a/	65.6 a/	91.6	2523 b/
B	151.0 a/	--	--	97.2 b/	2823 b/
B <sub>2</sub> O <sub>3</sub>	--	5.3 a/	77 a/	82.3	>1800 b/
Hg	143.0 a/	2.3 a/	31.5 a/	33.7	1393 a/
H <sub>2</sub> O	--	18.5 a/	100.7 a/	119.2	3823 a/

a/Bureau of Standards value (7); b/Handbook; c/ Estimated; \*Vaporizes at 2073°K

CONFIDENTIAL



CONFIDENTIAL  
SECURITY INFORMATION

In Table 149 are given the specific heats at constant volume, for 10°K intervals, for the assumed products of detonation. In extending these specific heat data to 7000°K it is fully realized that such extrapolation of data, infrequently measured for temperatures above 2000°K, is at best guess-work, but the lack of data, particularly for the metal oxides, forced such extrapolations. It should be noted that in spite of these considerable extrapolations, the degree of accuracy attained in the final values (½ TNT) is good and appears to be in no way related to the temperature range involved, at least up to 5000°K.

Finally, in making these calculations, it was necessary to make certain assumptions as to the products of detonation. These assumptions are:

1. The oxygen present is used first to burn the carbon to CO, then the hydrogen to H<sub>2</sub>O, and finally the CO to CO<sub>2</sub>. This follows the Kistiakowsky-Wilson assumption.
2. If a metal is present, it is oxidized fully before any CO is formed.
3. Any free carbon remaining unburned is considered a solid and is, therefore, not added into the number of moles of gas formed.
4. The metal oxides assumed to be formed remain solids unless the calculated adiabatic flame temperature exceeds the boiling point of the oxide. If the boiling point is exceeded, the heat necessary to vaporize the quantity of oxide is subtracted from the heat of explosion and the adiabatic flame temperature is recalculated. This may result in a calculated temperature less than the boiling point, indicating that only part of the oxide is vaporized.

The accuracy of this method of calculation of power is illustrated in Figures 80, 81 and 82 which show the direct relationship between calculated power and observed ballistic mortar values for 111 pure compounds, 56 organic mixtures and 30 metallized mixtures. Applying simple statistical methods to these data, it may be shown that the average deviation of the observed from the calculated is +1.6 for the pure compounds (eliminating 6 as discussed below), +0.1 for the organic mixtures and -0.8 for the metallized mixtures. The standard (root-mean square) deviations, from the mean, are 6.7 for the pure compounds, 5.8 for the organic mixtures and 7.8 for the metallized mixtures.

Figure 83, drawn to show the statistical scatterings for the pure compounds, mixtures and metallized mixtures, indicate that these scatterings are generally normal. This leads to the conclusion that the deviations are due largely to personal errors and are therefore not inherent in the system or in the compounds.

The summarized results of the calculations are given in Tables 143, 144 and 145. Ballistic mortar data were taken from Parts I and III of this series (3, 5).

CONFIDENTIAL




FIGURE 80  
COMPARISON OF BALLISTIC MORTAR AND CALCULATED nRT FOR  
PURE EXPLOSIVE COMPOUNDS




FIGURE 62  
COMPARISON OF BALLISTIC MORTAR AND CALCULATED TNT FOR  
EXPLOSIVE METALLIC MIXTURES

CONFIDENTIAL  
SECURITY INFORMATION






FIGURE 83  
STATISTICAL DISTRIBUTION OF DEVIATIONS FROM THE MEAN  
COMPARING BALLISTIC MORTAR AND ART FOR EXPLOSIVE  
SUBSTANCES

-50-

CONFIDENTIAL

Arthur D. Little, Inc.

Table 113

Properties of Selected Pure Organic Compounds

No.	Name	O.B.	Calculated		Calc. $T_g, ^\circ F$	Power, kWh		
			Kcal/mol	$T_g, ^\circ C$		Calc.	Exp.	Dev.
6	1,4-dicyclohexyl-2,3,5,5-tetraazabicyclopentane	-30.4	104.1	167.7	3651	120	137	+ 8
22	DMG	-21.6	635.1	355.8	4271	151	150	- 1
23	DMH	-21.6	177.6	275.6	1285	152	150	- 2
24	Hexahydro-1,3,5-triazine-triazine (Bocall)	-85.1	518.9	144.4	2930	223	224	+ 1
25	2,4,6,8-tetraazabicyclo[2,4,6,8]tetraazocane-1,7-diol diacetate	-52.3	1249.5	255.9	2235	96	83	-13
26	2,4,6,8-tetraazabicyclo[2,4,6,8]tetraazocane-1,7-diol diacetate	-88.7	992.6	166.1	1860	81	120	+30
33	Diethylurea perchlorate	-16.2	279.9	145.7	2790	99	92	- 7
35	Dithionitramine dinitrate	-26.7	552.0	282.5	3920	119	110	- 9
37	Diethylurea glycol dinitrate	-10.8	503.7	176.8	3014	171	177	+ 6
41	2,2-Dinitroethyl-1,3-propanediol dinitrate	-70.2	685.5	117.9	2775	103	108	+ 5
42	1,3-Dinitropropane-2-propanol nitrate	-17.8	456.2	289.5	4236	118	112	- 6
43	2,4-Dinitrocellulose	-91.7	701.3	103.7	2110	62	68	+ 7
44	2,4-Dinitrocellulose	-96.9	788.2	92.1	1805	72	67	+15
45	N-Dinitrobenzamide	-95.2	678.8	107.6	2362	86	85	0
47	2,2-Dinitro-1-butanol	-60.2	514.5	72.6	1672	80	94	+14
54	N,N'-bis(2-nitroethyl)-N,N'-dinitroethylenediamine	-61.5	744.7	110.6	1785	78	71	- 7
56	N,N'-bis(2-nitroethyl)-N,N'-dinitroethylenediamine	-39.8	472.3	115.4	2687	103	116	+ 13
63	N,N'-bis(2-nitroethyl)-N,N'-dinitroethylenediamine dinitrate	-18.0	672.7	103.3	4002	128	135	+ 7
65	1,1-Dinitroethane	-26.7	267.2	132.6	3397	111	138	+ 27
66	1,3-Dinitro-2-propanolamine	-27.3	375.8	176.8	3130	130	122	- 8
68	1,6-Dinitrohexanol	-27.6	455.0	185.6	3241	110	101	- 9
71	2,4-Dinitrophenol	-78.2	611.8	86.7	1980	75	52	-23
74	2-(3,5-Dinitrophenyl)-2-nitro-1,3-propanediol dinitrate	-40.3	989.2	351.3	3646	127	126	- 1
75	1,1-Dinitropropane	-72.7	616.6	117.3	2275	111	107	- 4
76	1,1-Dinitropropane	-59.7	412.2	95.2	2435	111	103	+ 8
77	2,2-Dinitro-1-propanol	-42.6	377.3	117.8	2722	111	122	+11
81	2,5-Dinitro-1,3-dioxane-2,6-dinitroethyl dinitrate	-21.4	554.6	554.6	4197	139	147	+8
82	2,6-bis(hydroxyethyl)-2,6-dinitro-1,7-heptanediol tetranitrate	-31.2	1106.6	496.3	3711	131	137	+ 6
84	2,4-Dinitroethane	-114.2	823.9	101.7	1985	76	71	- 5
86	2,4,6,8-tetraazabicyclo[2,4,6,8]tetraazocane-1,7-diol diacetate	-34.7	716.2	658.9	4690	151	255	+ 104
89	Dipentylurea dihydroxide	-27.5	1187.7	581.6	3776	132	144	+12
92	Dinitrocellulose	-71.1	637.1	637.1	4754	130	145	+15
93	Dicyclohexyl tetranitrate	-5.3	460.9	140.9	4968	133	151	+18
95	Ethylurea dinitrate	-25.0	359.8	157.5	2802	119	114	- 5
96	Ethylurea dinitrate, DMA	-32.0	358.1	166.2	3570	116	139	+ 23
102	Diethyl nitrate	-61.5	294.0	73.0	2530	112	123	+ 11
111	1-Hydroxy-1-nitroethane	-29.4	463.7	233.4	2245	117	111	- 6
113	1-Hydroxy-1-nitroethane	-31.5	388.6	196.2	3785	126	120	- 6
115	Diethylurea perchlorate	-5.0	221.3	187.6	4037	133	141	+ 8
116	1-Hydroxy-1-nitroethane	-34.3	382.2	81.3	3755	73	75	+ 2
119	1,2,3,4,5,6-hexanitrohexanetriol tetranitrate	-36.3	1394.5	533.4	3762	130	138	+ 8
125	2,2',4,4',6,6'-hexanitrodiphenylamine	-52.8	1301.3	149.5	3379	121	115	- 6
128	2,2',4,4',6,6'-hexanitrodiphenyl ether	-47.3	1322.3	126.7	3421	116	113	- 3
130	N,N'-dicyclohexylurea dinitrate	-44.5	1575.2	517.4	3674	129	123	- 6
135	1,3-Dicyclopentane	-53.1	1284.7	396.0	2735	101	74	-27
136	Hexanitroethane	-62.7	173.3	173.3	2841	75	102	+27

Table 11) (cont:med)  
Properties of Selected Pure Organic Compounds

No.	Name	M.P.	Calculated		Calc. Mol. Wt.	Power, Stat. Calc.	Stat. Wt.	Error
			U	G				
152	Methyl benzotrate	+7.1	577.1	637.1	4794	130	136	+ 6
163	N-Methyl ethylsulfonamide	-58.5	506.3	131.8	2615	126	170	- 6
165	1,2,3-Substituted triazine	-16.6	492.5	124.1	4368	123	110	- 3
168	2-Methyl-2-nitro-1,3-propanediol dinitrate	-24.8	601.3	265.5	3943	130	136	- 2
170	N-Methyl ethylsulfonamide	-58.5	506.3	130.8	2600	125	118	- 7
180	2-Nitroethanol nitrate	-11.8	577.3	190.0	4470	114	142	- 2
181	Nitroglycerin	+3.5	365.6	365.6	1935	110	140	0
182	Nitroglycerin	+3.5	365.6	365.6	1935	110	140	0
185	Nitroethanol	-19.3	457.2	61.6	3116	134	136	0
187	2-Methyl-2-nitroethylsulfonamide nitrate	-22.3	360.0	181.7	3724	177	133	+ 6
189	2-nitro-2-(n-nitropropyl)-1,3-propanediol dinitrate	-57.8	1440.0	211.4	2938	112	114	+ 2
192	ee, ee, ee-Triethylamine	-74.0	786.8	145.5	2916	97	100	+ 3
194	2-Ethoxy-2-(n-nitropropyl)ethanol nitrate	-77.8	571.2	155.1	2324	95	99	+ 4
200	Nitroethanol	-7.6	433.4	79.8	2810	94	105	+11
202	N-Methyl ethylsulfonamide nitrate	-43.6	454.1	265.8	3448	138	137	- 1
207	Fourmethylol tetraol nitrate	+10.3	544.7	150.0	4431	123	145	+ 4
208	2,3,4,5,6-pentaethylamine	-15.3	410.4	468.4	4977	137	142	+ 5
215	Picric acid	-15.3	601.3	168.6	3130	108	104	+ 1
228	2-Ethoxy-1,1,3,3-cyclohexane tetraethylol pentaol nitrate	-44.9	1227.1	410.0	3205	105	127	+ 1
229	2,2,6,6-Tetraethyl-1,3-cyclohexane tetraethylol pentaol nitrate	-56.3	1184.6	281.2	2575	104	114	+ 6
230	2-Ethoxy-1,1,3,3-cyclohexane tetraethylol pentaol nitrate	-35.3	1525.5	450.7	3640	133	151	+18
231	2,2,5,5-Tetraethyl-1,3-cyclohexane tetraethylol pentaol nitrate	-45.8	1041.5	322.5	3032	117	124	+ 5
232	2,2-bis(nitroethyl)-1,3-propanediol tetraethylol pentaol nitrate	-11.0	812.6	301.4	3247	137	129	- 8
233	2,3,4,6-Tetraethylamine	-32.2	841.5	271.1	4035	127	121	- 8
240	Tetraethylamine	+49.0	74.1	78.1	2253	58	109	-51
241	1,3,6,8-Tetraethylamine	-72.7	1065.0	224.4	2932	100	101	+ 1
243	2-Ethoxy-1,1,3,3-cyclohexane tetraethylol pentaol nitrate	-28.9	160.7	176.6	4432	140	141	+ 1
248	Tetraethyl-2-ethoxyethylamine tetraethylol pentaol nitrate	-33.0	1053.0	456.8	3490	133	138	+ 5
253	Tetraethylamine	-47.3	809.4	251.8	3480	127	130	- 3
258	1,2-Propanediol nitrate	-58.5	506.3	130.2	2512	124	116	- 8
264	2-Ethyl-2-(n-nitropropyl)-1,3-propanediol triethylol pentaol nitrate	-65.0	950.4	200.2	2410	112	117	+ 5
263	2-(n-nitropropyl)-2-(n-nitroethyl)-1,3-propanediol triethylol pentaol nitrate	-34.5	836.3	270.9	3558	134	136	+ 2
265	2-Ethyl-2-(n-nitropropyl)-1,3-propanediol triethylol pentaol nitrate	-50.5	782.7	234.7	2935	124	127	+ 3
270	Picric acid	-56.1	672.4	153.4	2797	107	110	+ 3
271	2,4,6-Triethylamine	-62.5	751.7	135.5	2366	74	106	+12
274	1,3,5-Triethylamine	-56.3	647.4	157.3	3102	115	111	- 4
275	2,4,6-Triethylamine	-60.7	644.4	144.2	2518	67	98	+11
276	2,4,6-Triethylamine nitrate	-38.9	750.0	278.7	3985	130	127	- 3
277	2,4,6-Triethylamine	-62.5	745.4	130.4	2285	92	94	+ 2
282	2-Ethyl-2,2,4,4-tetraethylamine	-94.6	901.3	92.0	1424	70	82	+ 7
283	2,4,6-Triethylamine	-103.5	1020.5	137.1	1735	80	84	+ 4
285	2-Ethyl-1,2,3,3-tetraethylamine	-65.7	674.6	136.7	2318	108	106	- 2
291	3-Ethyl-1,2,3,3-tetraethylamine	-83.2	819.1	98.8	1660	85	86	+ 1
292	1,1,8-Triethylamine	-109.3	1095.2	174.5	2450	82	83	+ 1
294	2,4,6-Triethylamine	-77.8	704.2	136.1	2102	86	80	- 6
305	2-(n-nitropropyl)-2-(n-nitroethyl)-1-butanol nitrate	-57.4	1207.3	303.8	1961	120	117	- 3
307	3-(n-nitropropyl)-2-(n-nitroethyl)-1-butanol nitrate	-35.4	911.1	372.4	2795	115	113	- 2

CONFIDENTIAL  
PROPERTY INFORMATION

Table 11) (continued)

Properties of Selected Pure Organic Compounds

No.	Name	Q <sub>19</sub>	Calculated		Calc. Expt. T <sub>m</sub>	Power, KW		
			C <sub>1</sub>	C <sub>2</sub>		Q <sub>19</sub>	R <sub>19</sub>	ERT
308	2-(4-Glyoxymethyl)-2-(4,2,4,6-tetraoxo-1,3,5-triazine) 1,3-propanediol trinitrate	-21.9	1106.4	635.0	4551	140	143	+ 3
310	Styphnic Acid	-45.7	557.2	186.7	3205	103	103	+ 5
313	2,4,6-Trinitrophenol	-74.0	752.7	151.4	2590	100	100	0
318	2,4,6-Trinitrobenzamide	-89.6	935.4	143.4	2208	86	83	0
319	1,1',1''-Tris(2-methylimidazole)trisethylamide	-55.8	1607.6	170.4	2669	109	107	- 2
320	Urea nitrate	-6.5	105.7	72.0	2163	77	92	+15
321	m-Prop. picrate	-91.5	1047.0	130.0	1845	77	87	+10
325	Allyl picrate	-86.2	1019.7	160.1	2255	68	71	+ 3
326	2-Propargyl picrate	-80.9	1001.3	199.2	2840	102	99	- 3
327	Nitroform	+37.1	112.4	112.4	1280	90	125	+35
328	Methylaltriazine, HPLM	0.0	212.2	212.2	4898	158	158	0
329	5-Methyl-2,3,4,5-tetraoxo-1,2,4-triazine	-28.9	780.7	175.6	4432	140	136	- 4
331	1,3-Bis(2,2-trinitroethyl)urea	0.0	583.4	580.4	5128	110	111	+ 4
337	Bis(2,2,2-trinitroethyl)triazine, HPL	+16.5	478.0	478.0	4649	126	137	+13
340	2,2,2-Trinitroethanol 4,4,4-trinitroethylate	- 4.1	644.6	557.2	4775	75	136	+ 1
343	2,2,3,3-Tetraoxo-1,2,4-triazine	-20.2	504.2	362.2	4335	143	153	+10
348	1,2-Dithiazine	-26.7	267.2	112.6	3720	136	127	- 9
570	2-Methyl-1,3,3-trinitropropane	-83.2	815.1	98.8	1660	85	86	+ 1

Table III

Properties of Selected Explosive Mixtures

No.	Mixture and Composition Ratio	D.S.	Calculated $\rho$ g/cc	Calc. $\rho$ g/cc	Power, KJ/Kg Calcs. Lab. Data	V.V.T. M/s	
M-370	Col (Barite) KEM/AN-2-stylyl cellosol/ F13/612, 71/5.2/2.2/1.6	-45.9	186.4	167.9	3599	137 130	- 1
M-401	Amotal. TRZ/AN, 66/35	-59.8	546.3	117.3	2580	126 106	0
M-402	70/30	-46.8	387.5	115.5	2905	123 114	- 1
M-404	55/45	-31.7	276.6	114.3	3192	122 122	0
M-405	50/50	-27.0	246.7	113.9	3280	121 123	0
M-406	40/60	-17.2	194.9	114.8	2595	128 128	0
M-407	30/70	- 8.2	151.1	114.8	2424	134 134	0
M-408	25/75	- 3.5	132.0	117.9	3132	131 131	0
M-409	20/80	+ 1.2	114.7	114.7	3705	130 130	0
M-410	15/85	+ 5.9	97.5	97.5	3369	121 129	+ 8
M-411	10/90	+10.6	82.1	82.1	3033	122 125	+14
M-412	5/95	+15.3	67.7	67.7	2690	121 117	+16
M-419	KEM/AN, 50/50	- 3.9	116.9	158.5	3455	130 123	- 7
M-420	45/55	- 0.6	113.8	151.0	3550	131 128	- 5
M-421	KEM/TRZ/AN, 20/20/60	- 8.0	157.9	122.4	2530	129 124	- 5
M-422	KEM/AN/AN, 25/30/45	- 8.8	206.3	157.9	3444	125 124	+ 5
M-423	AN/AN, 40/60	- 1.6	154.0	102.7	2630	131 122	- 9
M-430	Comp. A. KEM/AN/AN, 88/12	-57.6	737.5	190.0	2710	123 132	+ 9
M-431	9/9	+4.8	644.3	222.9	3090	122 126	+ 3
M-434	KEM/AN/AN/AN, 45.5/50/4.5	-21.2	209.3	94.5	4897	125 127	+17
M-436	KEM/TRZ/AN, 59.5/39.5/1.0	-45.4	671.9	207.8	3245	132 131	+ 1
M-437	Comp. C. KEM/AN, 68/12	-40.1	524.9	64.3	2810	130 120	+10
M-438	TRZ/TRZ/AN/KEM/AN, 12/2/2/79/5	-40.9	504.4	198.2	2590	117 123	- 4
M-439	TRZ/TRZ/AN/KEM/AN/AN, 10/5/1/17/74/3	-41.8	501.9	206.0	2503	127 126	+11
M-440	Qyalotel. KEM/TRZ, 10/90	-60.2	698.7	147.5	2905	117 123	+ 3
M-441	40/60	-63.0	665.0	143.2	3110	123 127	+ 4
M-442	50/50	-57.5	527.7	228.5	3118	122 125	- 6
M-443	30/70	-42.6	402.0	224.1	3530	125 133	- 2
M-444	70/30	-37.3	370.7	223.7	3715	120 125	- 5
M-447	KEM/TRZ, 45/55	-52.3	576.4	143.4	2519	107 109	+ 2
M-448	Amotal. KEM/TRZ, 50/50	-61.0	521.2	137.8	2809	117 117	0
M-449	55/45	-50.9	500.0	129.7	2850	119 119	0
M-449	64/36	-48.8	482.0	116.7	296	121 121	0
M-451	M-349a. TRZ/AN/TRZ, 31.5/55/13.5	-41.0	575.9	193.7	3399	127 126	- 1
M-452	M-349b. TRZ/AN/TRZ, 28/60/12	-38.7	551.9	209.1	3406	120 127	- 3
M-453	M-349c. TRZ/AN/TRZ, 25/50/25	-42.3	599.5	178.1	3189	122 115	- 7
M-454	Pivonite/AN, 50/50	-38.1	538.8	158.1	3145	126 130	+ 4
M-455	Pivonite/PTR, 50/50	-28.0	771.8	383.4	2643	132 128	0
M-456	Pivonite/PTR, 50/50	-32.7	424.9	227.7	2574	125 124	- 2
M-457	Pivonite/KEM, 53/47	-34.4	700.9	294.1	3024	126 125	+ 1
M-459	Methylite-25. AN/AN/styl pthalate, 75/25	-40.6	516.6	179.9	3035	123 121	- 2
M-460	Methylite-25. AN/AN/styl pthalate, 70/30	-49.5	584.9	118.3	2995	102 102	0
M-463	Pentalite. PTR/TRZ, 25/75	-59.3	752.5	146.7	2700	107 111	+ 4
M-466	50/50	-42.0	705.7	45.1	3475	127 126	- 1
M-467	75/25	-26.2	650.4	334.3	4085	126 126	0
M-468	PEP-1. PTR/TRZ/AN, 21.2/63.7/15	-100.6	799.7	60.1	1990	83 108	+19
M-469	PEP-2. PTR/TRZ/AN, 86/14	-66.6	244.4	65.9	2820	123 123	- 8
M-470	Pivonite. AN/TRZ, 52/48	-58.1	693.4	145.7	2585	103 100	- 3
M-471	PEP-2. PTR/TRZ/AN/TRZ, 28/60/12	-33.1	595.5	280.1	3022	128 128	0
M-472	PEP-1. AN/TRZ/AN/TRZ, 30/50/20	-45.0	652.9	290.7	2572	121 122	+ 1
M-474	PEP-1. KEM/AN, 10/90	-48.5	220.9	39.1	2250	122 122	- 6
M-475	KEM/AN/AN, 55/45	-57.7	277.4	50.1	2730	123 124	0
M-478	Crystal. TRZ/TRZ, 75/25	-54.0	824.4	219.7	3141	120 122	+ 2
M-481	TRZ/TRZ/AN/TRZ/AN/styl pthalate, 22.9/4/64.5/ 4.7/3.3	-42.5	644.4	127.8	3200	126 129	+ 5
M-485	TRZ/TRZ/AN/TRZ/AN/styl pthalate, 24.4/4/64.5/ 3.3/2.4	-43.8	556.5	172.4	3355	133 126	+13
M-486	TRZ/TRZ/AN/TRZ, 14.7/1/79.7/5/5	-31.9	627.0	228.1	2371	137 139	- 2

-508-

CONFIDENTIAL  
Arthur D. Little, Inc.



Table 115

Properties of Selected Metallized Explosives Mixtures

Mixture and Composition Ratio	Metal Oxide State	Qkt	Calculated $\Delta C$ kcal/mole	Calculated $\Delta G$ kcal/mole	Calc. Temp. T, °K	Power, Cal./g.	Heat, Cal./g.	Inst. Temp. T, °K
Unstabilized Comp. A. H <sub>2</sub> O/Al/Al <sub>2</sub> O <sub>3</sub> , 73/16/9	solid	-51.7	373.9	189.3	5980	158	141	-16
	gas			159.1	5071	153		-9
Aluminized Hexamet. H <sub>2</sub> O/TNT/Al	solid	-60.2	319.7	112.1	5056	135	130	-5
	gas			113.7	4377	125		+5
Trinitrol. TNT/Al, 90/10	solid	-75.5	508.6	155.8	3981	118	121	-6
	gas			133.6	3192	110		-13
	solid	-17.0	490.5	157.5	5175	168	117	+9
	gas			125.4	4250	154		+13
80/20	solid	-76.5	223.3	158.5	6117	76	105	+79
	gas			122.6	4925	85		+20
	solid	-83.0	213.1	58.2	4544	21	749	
	gas			44.8	3634	34		
Hexol (Aluminized Hexol). TNT/Al/Al <sub>2</sub> O <sub>3</sub> /BaSO <sub>4</sub>	solid	+4.7	131.6	110.6	3870	129	130	+1
	gas			95.0	3170	146		+4
	solid	+0.7	112.7	112.1	4557	139	136	-3
	gas			100.2	3115	133		+3
	solid	-54.7	335.3	117.2	3147	117	112	-15
	gas			98.1	3180	124		+2
	solid	-55.0	231.7	117.5	3415	137	122	-15
	gas			79.5	2121	124		+2
	solid	-1.4	117.6	113.7	5504	115	134	-11
	gas			79.9	5037	123		+3
	solid	-23.3	196.8	113.3	4670	145	133	-12
	gas			74.4	4082	135		+1
72.5/22.5/5/25	solid	-64.2	359.7	175.6	4180	130	117	-13
	gas			111.0	3790	119		-2
	solid	-71.6	220.8	113.6	4825	119	143	-6
	gas			71.2	4615	135		+8
60/40/18	solid	-37.0	220.2	123.6	4044	119	143	-6
	gas			71.2	4019	125		+8
40/60/20	solid	-39.1	222.0	211.7	5020	151	143	-5
	gas			89.5	4095	146		+7
Dynam. Hex/TNT/Al, 11.5/11.5/17	solid	-54.8	359.9	167.6	5182	142	133	-7
	gas			133.8	4784	132		+1
	solid	-54.8	251.8	162.5	575.8	112	134	-8
	gas			133.0	4130	132		+2
12/88/18	solid	-54.7	152.6	262.7	5506	141	134	-7
	gas			231.1	4166	132		+2
Hex. Hex/TNT/Al/Decomposition, 37.5/37.7/27.1/5.5	solid	-61.0	393.7	162.4	4850	135	146	-9
	gas			133.2	4120	125		0
Hex/TNT/Al/Hex, 11.16/39.20/37.64/1.00	solid	-61.2	370.2	161.2	5045	138	134	-6
	gas			131.7	4245	128		+4
Hex/TNT/Al/Hexol, 39.94/34.04/17.10/5	solid	-66.5	350.6	158.4	4770	133	132	-1
	gas			129.3	4020	123		+9
Hex/TNT/Al/Hexol, 37.80/34.00/16.20/10	solid	-78.2	432.1	153.8	4555	126	126	0
	gas			125.2	3625	116		+10
DNT Dynam. DNT/Hex/Al, 34/18/18	solid	-65.2	304.0	154.3	4751	134	133	-1
	gas			125.0	4168	125		+8
Hex. Hex/TNT/Hex/Al, 22/10/21/10	solid	-65.9	275.8	136.7	4563	131	146	-5
	gas			103.2	4165	137		+9
Bar-tropyx. Hex/TNT/Al, 18.6/16.1/5	solid	-55.9	379.8	138.1	3843	117	127	-10
	gas			115.7	3426	126		+1
	solid	-50.4	321.6	143.9	4745	142	131	-11
	gas			117.5	3792	132		-1
46/46/10	solid	-64.7	395.9	128.5	4450	124	126	-1
	gas			99.8	3751	114		-11
13.5/14.5/15	solid	-73.1	253.6	150.9	6151	113	122	-1
	gas			113.6	5043	122		+0
42/59/10	solid	-78.3	229.0	116.9	3170	74	101	+27
	gas			50.2	1515	45		+36
	8 - BaO <sub>3</sub> gas			47.4	1740	64		+37

\* Duplication of preceding composition; see Appendix in statistical analysis.  
 Note: Ballistic mortar data for additional mixtures explosives in (8) were noted too late for inclusion in this table.

CONFIDENTIAL

SECURITY INFORMATION

1. Organic Compounds

It will be noted that of the 111 pure compounds given in Table 113, only six display deviations from the mean greater than plus or minus 18, that is about three times the standard deviation which is commonly considered to include 99% of the deviation due to chance. These six which were excluded from the statistical analysis, are compound numbers 26, 71, 135, 136, 240 and 327. Data on these are summarized in Table 116.

Table 116  
Pure Compounds Showing Large Differences Between Observed  
Ballistic Mortar and Calculated RKT Values

No.	Name	Power, % TNT			
		O.B.	Calc.	H.M.	Dev.
26	2,4,6-Trinitro-2,4,6-triazahptane- 1,7-diol diacetate	-58.7	82	120	+38
71	2,4-Dinitrophenol	-78.2	75	52	-23
135	1,3-Dipicrylures	-53.1	101	76	+25
136	Hexanitroethane	+42.7	75	108	+33
240	Tetranitromethane	+49.0	58	109	+51
327	Nitroform	+37.1	90	125	+35

Of these compounds, the large negative deviations for numbers 71 and 135 may possibly be explained by the fact that these were not actually the compounds measured or perhaps even that they did not detonate completely in the ballistic mortar. Compounds 136, 240 and 327 have large positive oxygen balances, +42.7, +49.0 and +37.1, respectively, a fact which gives an explanation for the large positive discrepancies found, for it may be reasonably assumed that the excess oxygen reacts with the detonator which itself has a negative oxygen balance. This explanation has been substantiated by two recent ballistic mortar measurements made by the Bureau of Mines. Tetranitromethane 240, and bis(trinitroethyl)trinitramine (hex), 327, were remeasured using in place of the usual electric detonator, a modified No. 6 type detonator free of a combustible seal. The explosive charge of this detonator consisted of 0.29 gram of PETH base charge, and 0.41 gram of 75-25 DDNP-KClO<sub>4</sub> primary charge.

CONFIDENTIAL

The ballistic mortar value for TNT was thereby reduced from 141 to 109% TNT; for NOX from 155 to 139% TNT. Thus the deviation between measured and calculated power for NOX, the only other compound of high positive oxygen balance, was reduced to +13, or within the range of accepted error. The large difference still existing between the measured and calculated values for TNT may well be explained by other thermodynamic characteristics of this material (see page 476). The deviation of +38 for compound number 26 can not be explained except to state that there is the possibility that the compound measured was not what it was thought to be.

Although it does not appear warranted to make any generalizations concerning a correlation between measured and calculated values and the probable degree of purity of the compound or the accuracy of measurement, it should be pointed out that deviations between measured and calculated values are at a minimum for compounds which are best known and hence probably most nearly pure. Among such compounds are HMX (22), RDX (23), ethyl nitrate (102), nitroglycerin (181), nitromethane (185), PETN (207), picric acid (216), Tetryl (253) and MEDINA (328).

## 2. Explosives Mixtures

An even better correlation is indicated for the organic mixtures, as shown in Table 144 and Figure 81. This may be due to the fact that the mixtures are generally made up of the more common, and hence better evaluated, compounds such as TNT, RDX, PETN and EDNA. As in the case of pure compounds, there may be seen the effect of positive oxygen balance (M-410, 411 and 412) upon the ballistic mortar values.

In Figure 84, there are plotted the measured and calculated values for a series of Cyclotols (RDX/TNT). This figure illustrates one of the uses this calculation system has, that is, an independent check on observed values. Here it appears that there are discernable errors in the measured values for there is no explanation for the rather erratic behavior indicated in the range of 40 to 65% RDX (M-441, 442 and 443). Rather, it is to be expected that the power of this mixture increases as the per cent of RDX is increased as shown by the calculated powers.

## 3. Metallized Mixtures

The calculation of metallized mixtures is attended by added complications, including gross assumptions as to the products formed, whether the inorganic oxides assumed to be formed are gaseous or solid, and finally a general insufficiency of thermodynamic data. As pointed out previously, the metals were assumed to oxidize completely (i.e., to  $Al_2O_3$ ,  $B_2O_3$ , or  $MgO$ ). All calculations were made for both solid and gaseous metal oxides and as will be seen in Table 145 and Figure 82, the corresponding "solid" and "gaseous" values generally lie above and below the line, respectively. Hence, it would appear that a true correlation lies between

CONFIDENTIAL  
SECURITY INFORMATION




FIGURE 54  
COMPARISON OF BALLISTIC MORTAR AND CALCULATED nRT  
FOR CYCLOTOLS (RDX/YNT)

CONFIDENTIAL

Arthur D. Little, Inc.

CONFIDENTIAL  
SECURITY INFORMATION

the "solid" and "gaseous" values. This is borne out by the  $\pm 0.8$  average deviation shown when the "solid" and "gaseous" figures are considered together (Figure 83).

Because of the assumptions which had to be made and because some of the temperatures calculated are apparently out of the range which can be handled confidently by such a simplified system, it is not claimed that the calculated temperatures and energies for metallized mixtures represent the true values. They are, however, generally of the same relative order of magnitude as measured by the ballistic mortar and the spherical lead block. (Tables 145 and 147).

There is some reason to believe that the power values for metallized mixtures as measured in the ballistic mortar and lead block to date may be somewhat in error as illustrated by Figure 85. There it is shown that the ballistic mortar and cylindrical lead block measurements for a series of Tritonals (TNT/Al) disagree as to the optimum per cent of aluminum, that is, the location of the peak of the curve. Calculated powers are somewhat lower than measured powers, but again there is general agreement between the observed ballistic mortar and calculated nRT as to the peak and the overall shape of the curve.

Since it is generally believed that the ballistic mortar and the spherical lead block measure shock power, as contrasted to blast power, the use of these two test methods, and hence the nRT calculation method, cannot be considered as a final means of evaluating metallized explosives.

#### 4. Correlation of Spherical Lead Block Measurements and nRT

The new type spherical lead block will be discussed in detail in a forthcoming report. It is sufficient to point out here that results are based upon an equivalent weight, i.e., on a weight of explosive required to give an expansion in the block equivalent to a specified expansion for TNT. The power measurements resulting from the spherical lead block correlate directly with nRT calculations and therefore also with ballistic mortar values. This is shown in Table 147.

Taking all the values given in Table 147, the average deviation is calculated to be  $\pm 0.1$  and the standard deviation 4.9. Thus the correlation (indicated to date on twenty-nine compounds) of calculated nRT with spherical lead block measurements is as good as with ballistic mortar measurements.

CONFIDENTIAL




FIGURE 85  
COMPARISON OF CALCULATED nRT, BALLISTIC MORTAR  
AND SPHERICAL LEAD BLOCK FOR TRITONALS (TNT/AL)

**CONFIDENTIAL**  
**SECURITY INFORMATION**

**Table 147**  
**Correlation Between Calculated nRT and Spherical Lead Block Measurements**

<u>Explosive Compound or Mixture</u>	<u>Power, % TNT</u>			<u>S.M.</u>	
	<u>Calc.</u>	<u>Block</u>	<u>Dev.</u>		
<u>Pure Compounds</u>					
22 HMX	151	146	- 5	150	
23 RDX	152	149	- 3	150	
182 Nitroguanidine	95	98	+ 3	104	
253 Tetryl	127	124	- 3	130	
310 Styphnic acid	103	102	- 1	108	
313 2,4,6-Trinitrotoluene	100	100	0	100	
328 Methylenedinitramine, MEDINA	158	155	- 3	158	
337 Bis(2,2,2-trinitroethyl)nitramine, HOX	125	128	+ 3	139	
342 5-Amino-1H-tetrazole	96	n.d.	-	-	
352 Hydrazine nitrate	122	121	- 1	-	
956 1-Amino-3-nitroguanidine	113	113	0	=	
<u>Organic Mixtures</u>					
TNM/TNT, 56/44	152	160	+ 8	-	
RDX/TNM, 77/23	154	153	- 3	-	
HMX/HOX, 43.3/56.7	153	152	- 1	-	
HOX/TNT, 81.8/18.2	148	153	+ 5	-	
MEDINA/TNT, 80/20 (cast)	151	150	- 1	-	
MEDINA/TNT/Tetryl, 80/10/10	153	147	- 6	-	
MEDINA/Tetryl, 80/20	153	152	- 1	-	
MEDINA/Styphnic acid	149	151	+ 2	-	
RDX/Hydrazine nitrate, 55/45	146	150	+ 4	-	
RDX/TNT, 51.21/48.79	130	124	- 6	-	
<u>Al<sub>2</sub>O<sub>3</sub> State</u>					
<u>Metallized Explosives</u>					
HMX-2H	solid	61	59	- 2	-
	gas	60	-	- 1	-
Torpex II, RDX/TNT/Al, 42/40/18	solid	141	144	+ 3	134
	gas	132	-	+12	-
Torpex II/Nitroguanidine, 61.17/38.83	solid	139	131	- 8	-
	gas	126	-	+ 5	-
Torpex II/Nitraminoguanidine, 57.92/42.08	solid	145	134	-11	-
	gas	132	-	+ 2	-
Aluminized MEDINATOL. MEDINA/TNT/Al, 70/19/5	solid	156	152	- 4	-
	gas	152	-	0	-
	solid	160	157	- 3	-
	gas	153	-	+ 4	-
	solid	163	163	0	-
	gas	154	-	+ 9	-
	solid	164	163	- 1	-
	gas	155	-	+ 8	-

\* New-type spherical lead block.

**CONFIDENTIAL**

CONFIDENTIAL  
SECURITY INFORMATION

5. Application to New Compounds and Mixtures

It is believed, however, that this simplified system has a more important application in predicting the relative power of pure compounds or mixtures as they would be measured by the ballistic mortar. A number of new unusual compounds and mixtures which have been proposed as high explosives have been calculated. A few typical ones are given in Table 118 to illustrate the application of the method for calculating heats of combustion (Section XVI) and the power method here described.

Table 118  
Properties of Selected High-Energy Compounds

<u>Name</u>	<u>O.P.</u>	<u>Calculated</u>		<u>T, °K</u>	<u>Calc. % TNT</u>
		<u>Kcal/mol</u>	<u><math>\frac{Q_v}{Q_E}</math></u>		
5-Trinitroethyl-1H-tetrazole	+11.0	267.7	267.7	6070	164
1,1,2,2-Tetranitroethylbis(5-azo-1H-tetrazole)	- 3.4	670.5	603.1	5849	168
1,1,1,6,6,6-Hexanitro-2,4-hexadiyne	0.0	698.0	698.0	7390	186
1,1,1,4,4,4-Hexanitro-2-butyns	+19.8	655.0	655.0	7750	188

It is recognized that none of the compounds listed in Table 118 have been synthesized and perhaps are not practical explosives, but they point up the need for examining certain structural groups, for example, -C≡C- and the tetrazole ring, which would be desirable from an energy point of view.

It is expected that this system will be extended to sulfur-containing compounds, to some halogenated compounds, and perhaps to mixtures containing metals other than those covered to date.

D. Method of Calculation

A form devised to standardize and expedite the NRT calculations is shown as Figures 86 and 87, the obverse and reverse sides, respectively. The following discussion of procedure is presented in terms of a typical step-by-step calculation for compound number 23, RDX, Figures 86 and 87.

It will be noted that the reverse side contains a small table for calculating the heat of combustion from the values of a' and b', as given in Section XVI of this report (see Table 60). These values are inserted

CONFIDENTIAL



CONFIDENTIAL  
SECURITY INFORMATION

for the various groups present and  $Q_c^D$  ( $H_2O$  liquid) calculated from the general equation

$$Q_c^D (H_2O \text{ liquid}) = a' + x' b'.$$

Having calculated this  $Q_c^D$  ( $H_2O$  liquid) for the compound, there is added the value for  $nRT$  to correct the value at constant pressure to heat of combustion at constant volume,  $Q_c^V$ . These values of 504.7 and 507.4 are then transferred to the "Calc."  $Q_c^V$  and  $Q_c^D$  lines on the front side of the sheet. From the value of 507.4 is subtracted the quantity  $(3.0 \times 9.827)$  to convert to  $H_2O$  gas (See Table 140). From this value for heat of combustion at constant volume are then subtracted the energies required to burn any C to  $CO_2$ , CO to  $CO_2$ , and  $H_2$  to  $H_2O$ . Thus, in the case of RDX, the quantity  $(3 \times 67.341)$  is subtracted from the value of 477.6 to obtain the heat of explosion at constant volume in kcal/mol. The next step involves a series of iterations to arrive at an explosion temperature. This is done by assuming a temperature and setting down on the form the specific heats of the detonation products at this temperature as obtained from Table 149. The summation of the products of these specific heats and the number of moles of each gas present is then divided into the heat of explosion, multiplied by 1000 (to convert from kilogram calories to calories since the specific heats are given in the latter units) and added to 298, the base temperature, to give an approximate temperature. The true temperature lies between these approximate and assumed temperatures. The system is repeated until there results a final calculated temperature corresponding to the assumed temperature. This temperature is next inserted in the  $nRT$  equation along with the value for  $n$ , the total moles of gas formed, which has been obtained from the detonation equation. (See assumption, page 500). The units for  $nRT$  thus found are kcal/mol. Subsequent conversions are made to kcal/kg and then to ft lbs/gram. The latter, when divided by the corresponding figure of 701.0 for TNT and multiplied by 100, results in an  $nRT$  product expressed in terms of per cent TNT.

There is provided on this form a space for calculating the molar and weight oxygen balances, two parameters often required in evaluating a high explosive.

In the case of organic mixtures, it has been found desirable to base all calculation on a total quantity equivalent to one mole of the mixture. Figures 88 and 89 give an example of this type of calculation illustrating the mixture of 60/40 EDNA/TNT (Ednatol). By dividing the weight per cent by the molecular weight of each of the components, a molar


CONFIDENTIAL

Figure 86

CONFIDENTIAL  
SECURITY INFORMATION

## CALCULATION OF POWER BY THE HNT METHOD

Structural Formula

Name RDXNumber 21 Molecular Weight 227.13Oxygen Needed 3.0 atoms/mol cpdOxygen Balance -41.6 gm O<sub>2</sub>/100 gm cpdn = 9 moles gas/mol compound

$$Q_C^v = \underline{477.6} \text{ Kcal/mol (H}_2\text{O gas)}$$

$$\text{Obs. } Q_C^p = \underline{501.8} \text{ Kcal/mol (H}_2\text{O liq.)}$$

$$Q_E^v = \underline{275.6} \text{ Kcal/mol (H}_2\text{O gas)}$$

$$\text{Calc. } Q_C^p = \underline{501.7} \text{ Kcal/mol (H}_2\text{O liq.)}$$

Thermodynamic Data Reference:

$$Q_T = \underline{507.4}$$

No. Mols	Pro- duct	Average C <sub>v</sub> (cal/mol/°K) from 293° to Assumed Flame Temperature									
		1190 °K	1290 °K								
	CO <sub>2</sub>										
3	H <sub>2</sub> O	10.00	10.06								
3	CO	6.50	6.51								
3	N <sub>2</sub>	6.45	6.47								
	H <sub>2</sub>										
	O <sub>2</sub>										
	C										
ΣnC <sub>v</sub>		68.85	69.12								
T=298 + $\frac{Q_C}{\Sigma n C_v} \times 10^3$		4301	4285								

$$F = E \cdot T \cdot \Sigma n = 1.987 \times 4285 \times 9.0 \times 10^{-3}$$

$$= \underline{76.53} \text{ Kcal/mol}$$

$$= \underline{345.0} \text{ Kcal/kg}$$

$$= \underline{1065.0} \text{ ft lbs/g} \quad (3.087 \text{ ft lbs} = 1 \text{ cal})$$

$$= \underline{\hspace{2cm}} \text{ } \left( \underline{\hspace{2cm}} \right)$$

$$\text{Calc. Power } \frac{1065.0}{701.0} = \underline{152} \% \text{ TNT}$$

$$\text{Obs. Power B. H.} = \underline{149} \% \text{ TNT} \quad \left( \frac{\text{Wh Mines}}{\text{ TNT}} \right)$$

$$\text{Equiv. Power, Lead Block} = \underline{149} \% \text{ TNT} \quad \left( \underline{\hspace{2cm}} \right)$$

Compound Number 23

Calculation of Heat of Combustion

Structural Group		a'	b'
No.	Name		
1	Para. Rin	5.7	52.08
2	Nitroline (H)	285.9	+ .81
3	Sec. Amine (1)	54.9	- .12

$$x = \frac{3.0}{\quad}$$

$$\Delta u = -\frac{H}{4} + \frac{2N}{4} + \frac{2O}{4}$$

$$\Delta u = \frac{-6 + 12 + 12}{4}$$

$$\Delta nRT = 4.5 \times 0.5923$$

Total 346.5  
 Ib'x 158.2  
 Q<sub>c</sub> (H<sub>2</sub>O liq) 504.7  
 ΔnRT 2.7  
 Q<sub>c</sub> (H<sub>2</sub>O liq) 507.4

\*\*\*\*\*

Other Calculations and Notes

C	H	N	O	

Figure 88

CONFIDENTIAL  
SECURITY INFORMATION

CALCULATION OF POWER BY THE NBT METHOD

Structural Formula  
  
60/40  
0.3997/0.1761

Name Epibatol  
ECM/TMT

Number 25429 Molecular Weight 173.67

Oxygen Needed +5.2935 atoms/mol cpd

Oxygen Balance -48.77 gm O<sub>2</sub>/100 gm cpd

0.6942 C<sub>2</sub>H<sub>6</sub>N<sub>4</sub>O<sub>4</sub>  
Detonation 0.3058 C<sub>2</sub>H<sub>6</sub>N<sub>4</sub>O<sub>4</sub> → 3.5290 CO + 1.0826 H<sub>2</sub>O + 1.7645 H<sub>2</sub> + 1.8471 N<sub>2</sub>

$\Delta n = 8.2232$  mols gas/mol compound

$Q_C = 180.2$  Kcal/mol (H<sub>2</sub>O gas) Obs.  $Q_C =$  Kcal/mol (H<sub>2</sub>O liq.)

$Q_E = 140.9$  Kcal/mol (H<sub>2</sub>O gas) Calc.  $Q_C =$  Kcal/mol (H<sub>2</sub>O liq.)

Thermodynamic Data Reference: \_\_\_\_\_

No. Mols	Product	Average $C_p$ (cal/mol/°K) from 298° to Assumed Flame Temperature						
		2950°K	3000°K	3100°K	3200°K	3300°K	3400°K	3500°K
	CO <sub>2</sub>							
1.0826	H <sub>2</sub> O	9.17						
3.5290	CO	6.27						
1.8471	N <sub>2</sub>	6.19						
1.7645	H <sub>2</sub>	5.77						
	O <sub>2</sub>							
	C							
$\sum n C_p$		53.56						
$T=298 + \frac{Q_E}{\sum n C_p} \times 10^3$		2929						

$F = R \cdot T \cdot \Delta n = 1.987 \times 2925 \times 8.2232 \times 10^{-3}$

- = 47.79 Kcal/mol
- = 275.2 Kcal/Kg
- = 849.5 ft lbs/g (3.087 ft lbs = 1 cal)
- = / ( )

Calc. Power  $\frac{849.5}{701.5} = 121$  % TNT

Obs. Power E. M. = 121 % TNT (Ref. (5))

Equiv. Power, Lead Block = % TNT ( )

CONFIDENTIAL

Compound Number M-1119

Calculation of Heat of Combustion

Structural Group		a'	b'
No.	Name		

x = \_\_\_\_\_  
 $\Delta n = -\frac{H}{4} + \frac{2N + 2O}{4}$   
 $\Delta n =$  \_\_\_\_\_  
 $\Delta nRT =$  \_\_\_\_\_ x 0.5923

Total \_\_\_\_\_  
 $\sum a'x$  \_\_\_\_\_  
 $Q_c (H_2O liq)$  \_\_\_\_\_  
 $\Delta nRT$  \_\_\_\_\_  
 $Q_c (H_2O liq)$  \_\_\_\_\_

\*\*\*

Other Calculations and Notes

C	H	N	O
1.3884	1.1652	2.7768	2.7768
2.1106	1.5290	0.9171	1.8348
3.5250	5.6942	3.6942	4.6116
	2.8471	1.8471	

1.0625 H<sub>2</sub>O  
 2.7645 H<sub>2</sub>

Figure 90

CONFIDENTIAL  
SECURITY INFORMATION

CALCULATION OF POWER BY THE nRT METHOD

Structural Formula

Name Torpax II  
RDX/TNT/Al

42/40/18  
0.1891/0.1761/0.6674

Number \_\_\_\_\_ Molecular Weight 96.84

Oxygen Needed +3.3022 atoms/mol cpd

Oxygen Balance -54.67 gm O<sub>2</sub>/100 gm cpd

0.1831 C<sub>3</sub>H<sub>6</sub>N<sub>6</sub>O<sub>6</sub>  
0.1705 C<sub>7</sub>H<sub>5</sub>N<sub>3</sub>O<sub>6</sub>  
Detonation 0.2161 Al

→ 1.152 CO + 0.5908 C + 0.9756 H<sub>2</sub> + 0.8050 N<sub>2</sub>  
+ 0.3232 Al<sub>2</sub>O<sub>3</sub>

In = 2.9326 (3.2558) mols gas/mol compound

Q<sub>C</sub> = 351.6 Kcal/mol (H<sub>2</sub>O gas)

Obs. Q<sub>C</sub> = \_\_\_\_\_ Kcal/mol (H<sub>2</sub>O liq.)

Q<sub>E</sub> = 162.7 Kcal/mol (H<sub>2</sub>O gas)  
(133.1)\*

Calc. Q<sub>C</sub> = \_\_\_\_\_ Kcal/mol (H<sub>2</sub>O liq.)

Thermodynamic Data Reference: \_\_\_\_\_

No. Mols	Product	Average C <sub>p</sub> (cal/mol/°K) from 298° to Assumed Flame Temperature				
		5100 °K	5300 °K	(4530) °K	(4480) °K	°K
	CO <sub>2</sub>					
	H <sub>2</sub> O					
1.1520	CO	6.51	6.63	(6.54)	(6.54)	
0.8050	N <sub>2</sub>	6.57	6.60	(6.50)	(6.49)	
0.9756	H <sub>2</sub>	6.42	6.46	(6.30)	(6.29)	
	O <sub>2</sub>					
0.5908	C	6.20	6.26	(5.22)	(5.22)	
0.3232	Al <sub>2</sub> O <sub>3</sub>	29.58	29.70	(29.53)	(29.52)	
	In C <sub>p</sub>	32.36	32.49	(31.96)	(31.93)	
	T=298 + $\frac{Q_E}{In C_p} \times 10^3$	5326	5306	(4463)	(4456)	

$$P = R \cdot T \cdot In - 1.987 \times \frac{(4463)}{5306} \times \frac{(3.2558)}{2.9326} \times 10^{-3}$$

$$= \frac{30.92}{(298.5)} \text{ Kcal/mol}$$

$$= \frac{319.3}{(921.5)} \text{ Kcal/Kg}$$

$$= \frac{985.7}{(921.5)} \text{ ft lbs/g} \quad (3.087 \text{ ft lbs} = 1 \text{ cal})$$

\* Numbers in ( )  
for Al<sub>2</sub>O<sub>3</sub> (g)

$$\text{Calc. Power} = \frac{985.7}{701.0} = \frac{141}{100} \% \text{ TNT}$$

$$\text{Obs. Power} \quad \text{B. M.} = \frac{134.135}{100} \% \text{ TNT} \quad (\text{Ref. (5)})$$

$$\text{Equip. Power, Lead Block} = \frac{143.145}{100} \% \text{ TNT} \quad (\text{Ref. (5)})$$

CONFIDENTIAL

Compound Number \_\_\_\_\_

Calculation of Heat of Combustion

Structural Group		a'	b'
No.	Name		

x = \_\_\_\_\_  

$$\Delta n = -\frac{H}{4} + \frac{2N + 2O}{4}$$

$$\Delta n = \text{_____}$$

$$\Delta nRT = \text{_____} \times 0.5923$$

Total \_\_\_\_\_  
 Eb'x \_\_\_\_\_  
 Q<sub>p</sub> (H<sub>2</sub>O liq) \_\_\_\_\_  
 ΔnRT \_\_\_\_\_  
 Q<sub>v</sub> (H<sub>2</sub>O liq) \_\_\_\_\_

\*\*\*\*\*

Other Calculations and Notes

C	H	N	O	Al
0.5123	1.0286	1.0286	1.0286	
1.1235	0.8625	0.5115	1.0230	0.6161
1.7128	1.9511	1.4101	2.1216	0.4161
1.1520	.9756	.8050		.3232

.5908 C

Table 119  
Average  $C_p$  Over Range from 298 °K to T  
(calories/mole/°K)

T, °K	CO <sub>2</sub>	H <sub>2</sub> O	CO	N <sub>2</sub>	H <sub>2</sub>	O <sub>2</sub>	C(s)	HCl	Al <sub>2</sub> O <sub>3</sub>	H <sub>2</sub> O <sub>g</sub>	H <sub>2</sub> O
1000	9.40	6.88	5.40	5.33	5.06	5.75	4.02	5.15	25.71	26.55	11.07
10	9.43	6.89	5.41	5.33	5.06	5.76	4.04	5.16	25.75	26.63	11.10
20	9.44	6.90	5.42	5.34	5.06	5.76	4.05	5.16	25.81	26.70	11.12
30	9.47	6.92	5.43	5.35	5.06	5.77	4.06	5.17	25.86	26.76	11.14
40	9.49	6.94	5.43	5.36	5.07	5.78	4.07	5.17	25.90	26.80	11.16
50	9.52	6.95	5.44	5.36	5.07	5.79	4.09	5.18	25.94	26.84	11.19
60	9.54	6.96	5.45	5.37	5.07	5.80	4.10	5.18	25.99	26.90	11.21
70	9.56	6.97	5.46	5.38	5.08	5.81	4.11	5.19	26.02	26.95	11.23
80	9.58	7.00	5.47	5.39	5.08	5.81	4.13	5.19	26.04	26.99	11.25
90	9.60	7.02	5.47	5.39	5.08	5.82	4.14	5.20	26.10	27.04	11.26
1100	9.62	7.03	5.48	5.40	5.08	5.83	4.16	5.20	26.13	27.07	11.30
10	9.64	7.04	5.49	5.42	5.09	5.83	4.17	5.21	26.17	27.10	11.33
20	9.64	7.06	5.50	5.42	5.09	5.84	4.18	5.21	26.20	27.14	11.35
30	9.66	7.07	5.50	5.42	5.10	5.85	4.20	5.22	26.24	27.18	11.37
40	9.70	7.08	5.51	5.43	5.10	5.86	4.21	5.22	26.26	27.22	11.40
50	9.72	7.10	5.52	5.44	5.10	5.86	4.23	5.23	26.30	27.25	11.43
60	9.74	7.11	5.52	5.44	5.11	5.87	4.24	5.23	26.32	27.28	11.46
70	9.76	7.13	5.53	5.45	5.11	5.88	4.25	5.24	26.36	27.31	11.48
80	9.78	7.14	5.54	5.46	5.11	5.89	4.26	5.24	26.38	27.35	11.50
90	9.80	7.16	5.55	5.47	5.12	5.89	4.28	5.25	26.42	27.38	11.53
1200	9.82	7.17	5.55	5.47	5.12	5.90	4.30	5.25	26.45	27.41	11.55
10	9.84	7.19	5.56	5.48	5.12	5.90	4.31	5.26	26.48	27.44	11.58
20	9.86	7.20	5.56	5.49	5.12	5.91	4.32	5.27	26.50	27.47	11.57
30	9.87	7.22	5.57	5.49	5.13	5.92	4.33	5.27	26.52	27.50	11.59
40	9.89	7.24	5.58	5.50	5.14	5.93	4.34	5.28	26.56	27.54	11.62
50	9.92	7.24	5.59	5.51	5.14	5.93	4.36	5.28	26.58	27.56	11.64
60	9.92	7.26	5.60	5.51	5.14	5.94	4.37	5.29	26.61	27.60	11.66
70	9.95	7.28	5.60	5.52	5.14	5.95	4.38	5.29	26.64	27.62	11.68
80	9.96	7.29	5.61	5.52	5.15	5.95	4.39	5.30	26.67	27.65	11.70
90	9.98	7.30	5.61	5.53	5.15	5.96	4.40	5.30	26.70	27.68	11.72
1300	10.00	7.32	5.62	5.54	5.16	5.96	4.42	5.30	26.71	27.70	11.75
10	10.01	7.33	5.63	5.54	5.16	5.99	4.43	5.31	26.74	27.73	11.77
20	10.02	7.34	5.64	5.55	5.16	5.99	4.44	5.32	26.77	27.76	11.79
30	10.04	7.36	5.64	5.56	5.16	6.00	4.45	5.32	26.80	27.79	11.81
40	10.06	7.38	5.65	5.56	5.17	6.00	4.46	5.33	26.82	27.82	11.83
50	10.08	7.39	5.65	5.57	5.17	6.01	4.48	5.34	26.85	27.84	11.85
60	10.09	7.40	5.66	5.58	5.18	6.02	4.49	5.34	26.87	27.86	11.87
70	10.11	7.42	5.66	5.58	5.18	6.02	4.50	5.35	26.90	27.89	11.89
80	10.12	7.43	5.67	5.59	5.18	6.03	4.51	5.35	26.92	27.91	11.91
90	10.14	7.45	5.68	5.60	5.19	6.03	4.52	5.36	26.95	27.94	11.93
1400	10.15	7.46	5.68	5.60	5.19	6.03	4.53	5.36	26.97	27.96	11.95
10	10.17	7.48	5.69	5.61	5.20	6.04	4.54	5.37	27.00	27.98	11.97
20	10.18	7.49	5.70	5.61	5.20	6.05	4.55	5.37	27.02	28.00	12.00
30	10.20	7.50	5.70	5.62	5.20	6.05	4.56	5.38	27.05	28.02	12.02
40	10.22	7.52	5.71	5.62	5.21	6.06	4.58	5.38	27.07	28.04	12.04
50	10.23	7.53	5.72	5.63	5.21	6.06	4.59	5.39	27.09	28.06	12.06
60	10.24	7.54	5.72	5.64	5.21	6.07	4.60	5.40	27.11	28.08	12.08
70	10.26	7.56	5.72	5.64	5.22	6.07	4.60	5.40	27.13	28.10	12.10
80	10.28	7.58	5.73	5.65	5.22	6.08	4.61	5.41	27.16	28.12	12.12
90	10.29	7.59	5.73	5.65	5.22	6.08	4.62	5.41	27.18	28.14	12.14
1500	10.30	7.60	5.74	5.66	5.23	6.09	4.63	5.42	27.20	28.16	12.16
10	10.32	7.62	5.75	5.66	5.23	6.09	4.64	5.43	27.22	28.18	12.18
20	10.33	7.63	5.75	5.67	5.24	6.10	4.65	5.43	27.24	28.20	12.20
30	10.34	7.65	5.76	5.68	5.24	6.10	4.66	5.44	27.26	28.22	12.22
40	10.36	7.66	5.76	5.68	5.24	6.11	4.67	5.45	27.28	28.24	12.24
50	10.37	7.68	5.77	5.69	5.25	6.11	4.68	5.45	27.30	28.26	12.26
60	10.38	7.69	5.78	5.69	5.25	6.12	4.69	5.46	27.32	28.27	12.28
70	10.40	7.70	5.78	5.70	5.25	6.12	4.70	5.46	27.34	28.29	12.30
80	10.41	7.72	5.79	5.71	5.26	6.13	4.70	5.47	27.36	28.30	12.32
90	10.42	7.73	5.79	5.71	5.27	6.14	4.71	5.48	27.38	28.32	12.34
1600	10.44	7.75	5.80	5.72	5.27	6.14	4.72	5.48	27.40	28.34	12.36
10	10.45	7.76	5.80	5.72	5.28	6.15	4.73	5.49	27.42	28.36	12.38
20	10.46	7.77	5.81	5.73	5.28	6.15	4.74	5.49	27.44	28.37	12.40
30	10.48	7.79	5.81	5.73	5.28	6.16	4.75	5.50	27.46	28.39	12.42
40	10.49	7.80	5.82	5.74	5.29	6.16	4.76	5.50	27.48	28.40	12.44
50	10.50	7.82	5.82	5.74	5.29	6.17	4.76	5.51	27.50	28.42	12.46
60	10.52	7.83	5.83	5.75	5.30	6.17	4.77	5.52	27.52	28.43	12.48
70	10.53	7.84	5.83	5.75	5.30	6.18	4.78	5.52	27.54	28.45	12.50
80	10.54	7.85	5.84	5.76	5.30	6.18	4.78	5.53	27.57	28.47	12.52
90	10.55	7.87	5.84	5.76	5.31	6.19	4.79	5.53	27.57	28.47	12.53



Table 11. (continued)

Average  $C_p$  Over Range from 298 °K to T  
 (calories/mole/°E)

T, °K	CO <sub>2</sub>	H <sub>2</sub> O	CO	N <sub>2</sub>	H <sub>2</sub>	O <sub>2</sub>	C(s)	HCl	Al <sub>2</sub> O <sub>3</sub>	B <sub>2</sub> O <sub>3</sub>	MgO
1700	10.56	7.58	5.85	5.77	5.31	6.19	4.80	5.54	27.58	26.19	12.55
10	10.58	7.69	5.86	5.77	5.32	6.20	4.81	5.54	27.60	26.50	12.57
20	10.59	7.91	5.86	5.78	5.32	6.20	4.82	5.55	27.62	26.52	12.59
30	10.60	7.92	5.87	5.78	5.32	6.21	4.82	5.55	27.64	26.53	12.61
40	10.61	7.94	5.87	5.79	5.33	6.21	4.83	5.56	27.65	26.54	12.63
50	10.62	7.95	5.88	5.79	5.33	6.22	4.84	5.56	27.67	26.56	12.65
60	10.64	7.98	5.88	5.80	5.34	6.22	4.84	5.57	27.69	26.57	12.67
70	10.65	7.98	5.89	5.80	5.34	6.23	4.85	5.57	27.70	26.58	12.69
80	10.66	7.99	5.89	5.81	5.35	6.23	4.86	5.58	27.72	26.60	12.71
90	10.67	8.00	5.89	5.81	5.35	6.24	4.87	5.58	27.73	26.62	12.73
1800	10.68	8.01	5.90	5.82	5.36	6.24	4.88	5.59	27.75	26.63	12.75
10	10.69	8.07	5.90	5.82	5.36	6.24	4.88	5.59	27.77	26.64	12.77
20	10.70	8.04	5.91	5.83	5.37	6.25	4.89	5.60	27.78	26.65	12.78
30	10.71	8.05	5.91	5.83	5.37	6.25	4.89	5.60	27.80	26.66	12.80
40	10.72	8.07	5.92	5.83	5.38	6.26	4.90	5.61	27.81	26.68	12.82
50	10.73	8.08	5.92	5.84	5.38	6.26	4.91	5.61	27.82	26.69	12.84
60	10.74	8.09	5.93	5.84	5.38	6.27	4.91	5.62	27.83	26.70	12.86
70	10.76	8.10	5.93	5.85	5.39	6.27	4.92	5.62	27.84	26.71	12.88
80	10.77	8.12	5.94	5.85	5.39	6.28	4.93	5.63	27.85	26.72	12.90
90	10.78	8.13	5.94	5.86	5.40	6.28	4.93	5.64	27.86	26.74	12.92
1900	10.75	8.14	5.94	5.86	5.40	6.29	4.94	5.64	27.89	26.75	12.94
10	10.80	8.16	5.95	5.86	5.41	6.29	4.94	5.65	27.90	26.76	12.96
20	10.81	8.17	5.95	5.87	5.41	6.30	4.95	5.65	27.91	26.77	12.98
30	10.82	8.20	5.96	5.88	5.42	6.30	4.96	5.66	27.92	26.78	13.00
40	10.83	8.20	5.96	5.88	5.42	6.30	4.96	5.66	27.94	26.79	13.02
50	10.84	8.21	5.97	5.88	5.43	6.31	4.97	5.67	27.95	26.80	13.04
60	10.85	8.22	5.97	5.89	5.43	6.31	4.98	5.67	27.97	26.81	13.06
70	10.86	8.23	5.98	5.89	5.44	6.32	4.98	5.68	27.98	26.82	13.08
80	10.87	8.24	5.98	5.90	5.44	6.32	4.99	5.68	27.99	26.83	13.10
90	10.88	8.25	5.98	5.90	5.45	6.32	4.99	5.69	28.01	26.84	13.11
2000	10.89	8.26	5.99	5.90	5.45	6.33	5.00	5.70	28.02	26.85	13.13
10	10.90	8.28	5.99	5.91	5.46	6.33	5.01	5.70	28.03	26.86	13.15
20	10.91	8.29	6.00	5.91	5.46	6.34	5.01	5.71	28.04	26.87	13.17
30	10.92	8.30	6.00	5.92	5.47	6.34	5.02	5.71	28.06	26.88	13.18
40	10.93	8.31	6.00	5.92	5.47	6.34	5.02	5.71	28.07	26.89	13.20
50	10.94	8.32	6.01	5.92	5.48	6.35	5.03	5.72	28.08	26.90	13.22
60	10.94	8.33	6.01	5.93	5.48	6.35	5.04	5.72	28.09	26.90	13.24
70	10.95	8.34	6.02	5.93	5.49	6.36	5.04	5.72	28.10	26.91	13.26
80	10.96	8.35	6.02	5.94	5.49	6.36	5.05	5.73	28.11	26.92	13.28
90	10.97	8.36	6.02	5.94	5.50	6.36	5.05	5.73	28.13	26.93	13.29
2100	10.98	8.38	6.03	5.94	5.50	6.37	5.06	5.73	28.14	26.94	13.31
10	11.00	8.39	6.03	5.95	5.50	6.37	5.06	5.74	28.15	26.94	13.33
20	11.01	8.40	6.04	5.95	5.51	6.38	5.07	5.74	28.16	26.95	13.34
30	11.01	8.41	6.04	5.96	5.51	6.38	5.08	5.74	28.17	26.96	13.36
40	11.02	8.42	6.04	5.96	5.52	6.38	5.08	5.75	28.18	26.96	13.38
50	11.02	8.43	6.05	5.96	5.52	6.39	5.09	5.75	28.20	26.97	13.40
60	11.03	8.44	6.05	5.97	5.52	6.39	5.09	5.76	28.21	26.98	13.41
70	11.04	8.45	6.05	5.97	5.53	6.40	5.10	5.76	28.22	26.99	13.42
80	11.05	8.46	6.06	5.97	5.53	6.40	5.10	5.76	28.23	26.99	13.44
90	11.06	8.47	6.06	5.98	5.54	6.40	5.11	5.77	28.24	26.99	13.46
2200	11.06	8.49	6.07	5.98	5.54	6.41	5.11	5.77	28.25	26.99	13.47
10	11.07	8.50	6.07	5.99	5.54	6.41	5.12	5.77	28.26	26.99	13.49
20	11.08	8.51	6.07	5.99	5.55	6.42	5.12	5.78	28.27	27.00	13.50
30	11.09	8.52	6.08	6.00	5.55	6.42	5.13	5.78	28.28	29.01	13.52
40	11.10	8.53	6.08	6.00	5.56	6.42	5.13	5.78	28.29	29.01	13.54
50	11.10	8.54	6.08	6.00	5.56	6.43	5.14	5.79	28.30	29.01	13.56
60	11.11	8.55	6.09	6.01	5.56	6.43	5.14	5.79	28.31	29.02	13.57
70	11.12	8.56	6.09	6.01	5.57	6.44	5.15	5.80	28.32	29.02	13.58
80	11.13	8.57	6.09	6.02	5.57	6.44	5.15	5.80	28.33	29.03	13.60
90	11.14	8.58	6.10	6.02	5.58	6.44	5.16	5.80	28.34	29.03	13.62
2300	11.15	8.59	6.10	6.02	5.58	6.45	5.16	5.81	28.35	29.04	13.63
10	11.15	8.60	6.10	6.03	5.58	6.45	5.17	5.81	28.36	29.04	13.65
20	11.16	8.62	6.11	6.03	5.59	6.46	5.18	5.82	28.37	29.05	13.67
30	11.17	8.63	6.11	6.04	5.59	6.46	5.18	5.82	28.38	29.05	13.68
40	11.18	8.64	6.12	6.04	5.60	6.46	5.19	5.82	28.39	29.05	13.70
50	11.19	8.65	6.12	6.04	5.60	6.46	5.20	5.83	28.40	29.06	13.72
60	11.20	8.66	6.12	6.05	5.60	6.47	5.20	5.83	28.41	29.06	13.73
70	11.20	8.67	6.13	6.05	5.61	6.47	5.20	5.83	28.42	29.07	13.75
80	11.21	8.68	6.13	6.05	5.61	6.48	5.21	5.84	28.43	29.07	13.76
90	11.22	8.69	6.13	6.05	5.62	6.48	5.21	5.84	28.44	29.08	13.78

Table 11.0 (continued)

Average  $C_p$  Over Range from 798 °K to T  
(calories/mole/°K)

T, °K	CO <sub>2</sub>	H <sub>2</sub> O	CO	H <sub>2</sub>	N <sub>2</sub>	O <sub>2</sub>	C(s)	HCl	Al <sub>2</sub> O <sub>3</sub>	Fe <sub>2</sub> O <sub>3</sub>	MgO
2400	11.23	8.70	6.14	6.04	5.62	6.48	5.22	5.84	28.45	29.14	13.60
10	11.24	8.71	6.14	6.05	5.63	6.49	5.22	5.85	28.46	29.15	13.62
20	11.24	8.72	6.14	6.06	5.63	6.49	5.23	5.85	28.47	29.15	13.63
30	11.25	8.73	6.15	6.06	5.63	6.49	5.23	5.86	28.48	29.16	13.64
40	11.26	8.74	6.15	6.07	5.64	6.50	5.24	5.86	28.48	29.16	13.66
50	11.27	8.75	6.15	6.07	5.64	6.50	5.24	5.86	28.49	29.17	13.68
60	11.27	8.76	6.16	6.07	5.64	6.50	5.25	5.87	28.50	29.18	13.90
70	11.28	8.77	6.16	6.08	5.65	6.51	5.25	5.87	28.51	29.19	13.92
80	11.29	8.78	6.16	6.08	5.65	6.51	5.26	5.88	28.52	29.19	13.93
90	11.30	8.79	6.17	6.08	5.66	6.52	5.26	5.88	28.52	29.19	13.94
2500	11.30	8.79	6.17	6.09	5.66	6.52	5.27	5.88	28.53	29.20	13.96
10	11.31	8.80	6.17	6.09	5.66	6.52	5.27	5.89	28.54	29.20	13.98
20	11.32	8.82	6.17	6.09	5.67	6.52	5.28	5.89	28.55	29.21	14.00
30	11.32	8.83	6.18	6.09	5.67	6.53	5.28	5.90	28.56	29.21	14.02
40	11.33	8.84	6.18	6.10	5.67	6.53	5.28	5.90	28.57	29.22	14.04
50	11.34	8.84	6.18	6.10	5.68	6.54	5.29	5.90	28.58	29.22	14.06
60	11.35	8.85	6.19	6.10	5.68	6.54	5.29	5.90	28.58	29.23	14.07
70	11.36	8.86	6.19	6.10	5.69	6.54	5.30	5.91	28.59	29.23	14.08
80	11.36	8.87	6.19	6.11	5.69	6.54	5.30	5.91	28.60	29.24	14.10
90	11.37	8.88	6.19	6.11	5.69	6.55	5.30	5.91	28.61	29.24	14.12
2600	11.38	8.89	6.20	6.12	5.70	6.55	5.31	5.92	28.62	29.25	14.14
10	11.38	8.90	6.20	6.12	5.70	6.56	5.31	5.92	28.62	29.25	14.16
20	11.39	8.91	6.20	6.12	5.71	6.56	5.32	5.93	28.63	29.26	14.18
30	11.40	8.92	6.21	6.12	5.71	6.56	5.32	5.93	28.64	29.26	14.20
40	11.40	8.93	6.21	6.13	5.72	6.57	5.32	5.93	28.64	29.27	14.21
50	11.41	8.94	6.21	6.13	5.72	6.57	5.33	5.94	28.65	29.27	14.22
60	11.42	8.95	6.21	6.13	5.72	6.57	5.33	5.94	28.66	29.28	14.24
70	11.42	8.96	6.22	6.14	5.73	6.58	5.34	5.94	28.67	29.28	14.26
80	11.43	8.97	6.22	6.14	5.73	6.58	5.34	5.95	28.68	29.29	14.27
90	11.44	8.98	6.22	6.14	5.74	6.58	5.34	5.95	28.68	29.29	14.29
2700	11.44	8.98	6.22	6.14	5.74	6.59	5.35	5.95	28.69	29.30	14.30
10	11.45	8.99	6.23	6.14	5.74	6.59	5.35	5.96	28.70	29.30	14.32
20	11.46	9.00	6.23	6.15	5.75	6.60	5.36	5.96	28.71	29.31	14.34
30	11.46	9.01	6.23	6.15	5.75	6.60	5.36	5.96	28.71	29.31	14.35
40	11.46	9.02	6.23	6.15	5.75	6.60	5.36	5.97	28.72	29.32	14.36
50	11.47	9.03	6.24	6.15	5.76	6.61	5.37	5.97	28.73	29.32	14.38
60	11.48	9.04	6.24	6.16	5.76	6.61	5.37	5.98	28.73	29.33	14.40
70	11.48	9.05	6.24	6.16	5.76	6.61	5.38	5.98	28.74	29.33	14.41
80	11.49	9.06	6.24	6.16	5.77	6.62	5.38	5.98	28.75	29.34	14.42
90	11.50	9.07	6.25	6.17	5.77	6.62	5.38	5.99	28.76	29.34	14.44
2800	11.50	9.08	6.25	6.17	5.77	6.62	5.39	5.99	28.76	29.35	14.45
10	11.51	9.09	6.25	6.17	5.78	6.63	5.39	6.00	28.77	29.35	14.47
20	11.51	9.10	6.25	6.17	5.78	6.63	5.40	6.00	28.77	29.36	14.48
30	11.52	9.11	6.26	6.18	5.79	6.63	5.40	6.00	28.78	29.36	14.50
40	11.53	9.11	6.26	6.18	5.79	6.64	5.40	6.01	28.79	29.36	14.51
50	11.53	9.12	6.26	6.18	5.79	6.64	5.41	6.01	28.80	29.37	14.52
60	11.54	9.13	6.26	6.18	5.80	6.65	5.41	6.02	28.81	29.38	14.55
70	11.54	9.14	6.26	6.19	5.80	6.65	5.42	6.02	28.82	29.38	14.56
80	11.54	9.15	6.27	6.19	5.80	6.65	5.42	6.03	28.82	29.38	14.58
90	11.55	9.16	6.27	6.19	5.81	6.65	5.42	6.03	28.82	29.38	14.58
2900	11.56	9.17	6.27	6.19	5.81	6.66	5.42	6.03	28.83	29.39	14.59
10	11.56	9.17	6.27	6.20	5.81	6.66	5.43	6.03	28.84	29.39	14.60
20	11.56	9.18	6.28	6.20	5.82	6.66	5.43	6.04	28.84	29.40	14.62
30	11.57	9.19	6.28	6.20	5.82	6.66	5.44	6.04	28.85	29.40	14.63
40	11.57	9.20	6.28	6.20	5.83	6.67	5.44	6.05	28.85	29.40	14.64
50	11.57	9.21	6.28	6.21	5.83	6.67	5.44	6.05	28.86	29.41	14.65
60	11.57	9.22	6.29	6.21	5.83	6.67	5.45	6.05	28.86	29.41	14.67
70	11.58	9.23	6.29	6.21	5.84	6.68	5.45	6.06	28.87	29.42	14.68
80	11.59	9.23	6.29	6.21	5.84	6.68	5.45	6.06	28.88	29.42	14.69
90	11.59	9.24	6.29	6.22	5.84	6.68	5.46	6.06	28.89	29.42	14.70
3000	11.60	9.25	6.29	6.22	5.85	6.69	5.46	6.07	28.90	29.43	14.71
10	11.61	9.26	6.30	6.22	5.85	6.69	5.46	6.07	28.90	29.43	14.73
20	11.62	9.27	6.30	6.22	5.86	6.69	5.47	6.07	28.91	29.44	14.75
30	11.62	9.28	6.30	6.23	5.86	6.70	5.47	6.08	28.92	29.44	14.75
40	11.63	9.29	6.30	6.23	5.86	6.70	5.47	6.08	28.92	29.44	14.76
50	11.63	9.29	6.30	6.23	5.87	6.70	5.48	6.08	28.93	29.45	14.78
60	11.64	9.30	6.31	6.23	5.87	6.71	5.48	6.09	28.94	29.45	14.79
70	11.64	9.30	6.31	6.23	5.87	6.71	5.48	6.09	28.94	29.45	14.80
80	11.65	9.31	6.31	6.24	5.88	6.71	5.49	6.10	28.95	29.46	14.81
90	11.65	9.31	6.31	6.24	5.88	6.72	5.49	6.10	28.95	29.46	14.82

Table 11.2 (continued)  
Average  $C_p$  Over Range from 298 °K to T  
(calories/mole/°K)

T, °K	CO <sub>2</sub>	H <sub>2</sub> O	CO	H <sub>2</sub>	N <sub>2</sub>	O <sub>2</sub>	C(s)	HCl	Al <sub>2</sub> O <sub>3</sub>	B <sub>2</sub> O <sub>3</sub>	MgO
1100	11.66	9.32	6.32	6.24	5.83	6.72	5.49	6.10	28.96	29.46	14.63
10	11.66	9.33	6.32	6.24	5.89	6.72	5.50	6.11	28.96	29.47	14.64
20	11.66	9.34	6.32	6.24	5.89	6.73	5.50	6.11	28.97	29.47	14.66
30	11.67	9.34	6.32	6.25	5.89	6.73	5.50	6.11	28.98	29.48	14.67
40	11.68	9.35	6.32	6.25	5.90	6.73	5.51	6.12	28.98	29.48	14.68
50	11.68	9.36	6.32	6.25	5.90	6.74	5.51	6.12	28.99	29.48	14.69
60	11.69	9.37	6.33	6.25	5.90	6.74	5.52	6.12	29.00	29.49	14.90
70	11.69	9.38	6.33	6.26	5.91	6.74	5.52	6.13	29.00	29.49	14.91
80	11.70	9.38	6.33	6.26	5.91	6.74	5.52	6.13	29.01	29.50	14.92
90	11.70	9.39	6.33	6.26	5.91	6.75	5.52	6.14	29.01	29.50	14.93
3700	11.71	9.40	6.34	6.26	5.92	6.75	5.53	6.14	29.02	29.50	14.94
10	11.72	9.41	6.34	6.26	5.92	6.75	5.53	6.14	29.02	29.50	14.95
20	11.72	9.41	6.34	6.26	5.92	6.76	5.54	6.15	29.03	29.51	14.96
30	11.72	9.42	6.34	6.27	5.93	6.76	5.54	6.15	29.03	29.51	14.97
40	11.73	9.42	6.34	6.27	5.93	6.76	5.54	6.16	29.04	29.51	14.98
50	11.73	9.43	6.34	6.27	5.94	6.76	5.54	6.16	29.04	29.52	14.99
60	11.74	9.44	6.35	6.28	5.94	6.77	5.55	6.16	29.05	29.52	15.00
70	11.74	9.44	6.35	6.28	5.94	6.77	5.55	6.17	29.05	29.52	15.01
80	11.75	9.45	6.35	6.28	5.95	6.77	5.55	6.17	29.06	29.53	15.02
90	11.75	9.46	6.35	6.28	5.95	6.78	5.56	6.17	29.06	29.53	15.03
3300	11.76	9.47	6.36	6.28	5.95	6.78	5.56	6.18	29.07	29.53	15.04
10	11.76	9.48	6.36	6.29	5.95	6.78	5.56	6.18	29.07	29.54	15.05
20	11.77	9.48	6.36	6.29	5.96	6.78	5.56	6.18	29.08	29.54	15.06
30	11.77	9.49	6.36	6.29	5.96	6.79	5.57	6.19	29.08	29.54	15.07
40	11.78	9.49	6.36	6.29	5.96	6.79	5.57	6.19	29.09	29.54	15.08
50	11.78	9.50	6.36	6.29	5.97	6.80	5.57	6.20	29.09	29.55	15.09
60	11.79	9.51	6.37	6.30	5.97	6.80	5.58	6.20	29.10	29.55	15.10
70	11.80	9.52	6.37	6.30	5.98	6.80	5.58	6.21	29.10	29.55	15.11
80	11.80	9.52	6.37	6.30	5.98	6.80	5.58	6.21	29.11	29.56	15.12
90	11.80	9.53	6.37	6.30	5.98	6.81	5.59	6.21	29.11	29.56	15.13
3100	11.81	9.53	6.37	6.30	5.98	6.81	5.59	6.22	29.12	29.56	15.14
10	11.82	9.54	6.38	6.31	5.99	6.82	5.59	6.22	29.12	29.56	15.15
20	11.82	9.54	6.38	6.31	5.99	6.82	5.60	6.22	29.13	29.57	15.16
30	11.82	9.55	6.38	6.31	5.99	6.82	5.60	6.23	29.13	29.57	15.17
40	11.83	9.56	6.38	6.31	6.00	6.82	5.60	6.23	29.14	29.57	15.18
50	11.83	9.56	6.38	6.32	6.00	6.82	5.61	6.24	29.14	29.58	15.19
60	11.84	9.57	6.39	6.32	6.01	6.83	5.61	6.24	29.14	29.58	15.20
70	11.84	9.58	6.39	6.32	6.01	6.83	5.62	6.24	29.15	29.58	15.21
80	11.84	9.58	6.39	6.32	6.01	6.83	5.62	6.24	29.15	29.58	15.22
90	11.85	9.59	6.39	6.32	6.02	6.84	5.62	6.25	29.16	29.59	15.23
3500	11.85	9.60	6.39	6.32	6.02	6.84	5.62	6.25	29.16	29.59	15.24
10	11.86	9.60	6.39	6.33	6.02	6.84	5.62	6.26	29.16	29.59	15.25
20	11.86	9.61	6.40	6.33	6.03	6.84	5.63	6.26	29.17	29.60	15.26
30	11.86	9.62	6.40	6.33	6.03	6.85	5.63	6.26	29.17	29.60	15.27
40	11.87	9.62	6.40	6.33	6.03	6.85	5.63	6.27	29.18	29.60	15.28
50	11.87	9.63	6.40	6.34	6.03	6.85	5.64	6.27	29.18	29.60	15.28
60	11.88	9.64	6.40	6.34	6.04	6.86	5.64	6.28	29.19	29.60	15.29
70	11.88	9.64	6.41	6.34	6.04	6.86	5.64	6.28	29.19	29.60	15.30
80	11.88	9.65	6.41	6.34	6.04	6.86	5.64	6.28	29.20	29.61	15.31
90	11.89	9.66	6.41	6.34	6.05	6.86	5.65	6.29	29.20	29.61	15.32
3600	11.89	9.66	6.41	6.34	6.05	6.86	5.65	6.29	29.20	29.61	15.33
10	11.90	9.67	6.42	6.35	6.05	6.87	5.65	6.30	29.21	29.62	15.34
20	11.90	9.68	6.42	6.35	6.06	6.87	5.66	6.30	29.21	29.62	15.35
30	11.90	9.68	6.42	6.35	6.06	6.88	5.66	6.30	29.22	29.62	15.36
40	11.91	9.69	6.42	6.36	6.06	6.88	5.66	6.31	29.22	29.62	15.36
50	11.91	9.70	6.42	6.36	6.06	6.88	5.66	6.31	29.22	29.63	15.37
60	11.92	9.70	6.42	6.36	6.07	6.88	5.67	6.31	29.23	29.63	15.38
70	11.92	9.71	6.42	6.36	6.07	6.89	5.67	6.32	29.23	29.63	15.39
80	11.92	9.72	6.42	6.36	6.07	6.89	5.67	6.32	29.24	29.63	15.40
90	11.93	9.72	6.42	6.36	6.08	6.89	5.68	6.33	29.24	29.64	15.41
3700	11.93	9.73	6.43	6.37	6.08	6.89	5.68	6.33	29.24	29.64	15.42
10	11.94	9.73	6.43	6.37	6.08	6.90	5.68	6.33	29.25	29.64	15.42
20	11.94	9.74	6.43	6.37	6.09	6.90	5.69	6.34	29.25	29.64	15.43
30	11.94	9.74	6.43	6.37	6.09	6.90	5.69	6.34	29.26	29.65	15.44
40	11.95	9.75	6.43	6.37	6.09	6.90	5.69	6.34	29.26	29.65	15.45
50	11.95	9.76	6.43	6.38	6.10	6.91	5.70	6.35	29.27	29.66	15.46
60	11.96	9.76	6.43	6.38	6.10	6.91	5.70	6.35	29.27	29.66	15.47
70	11.96	9.77	6.43	6.38	6.10	6.91	5.70	6.35	29.28	29.66	15.48
80	11.97	9.77	6.43	6.38	6.10	6.91	5.70	6.35	29.28	29.66	15.48
90	11.97	9.78	6.43	6.38	6.11	6.92	5.71	6.36	29.28	29.66	15.49

Table 11. (continued)  
Average  $C_p$  Over Range from 250 to T  
(calories/mole/°K)

T, °K	CO <sub>2</sub>	H <sub>2</sub> O	CO	N <sub>2</sub>	F <sub>2</sub>	O <sub>2</sub>	C(s)	HCl	Li <sub>2</sub> O <sub>2</sub>	B <sub>2</sub> O <sub>3</sub>	MgO
3800	11.98	9.78	6.45	6.38	6.11	6.92	5.71	6.37	29.28	29.66	15.50
10	11.98	9.79	6.45	6.38	6.11	6.93	5.71	6.37	29.29	29.67	15.51
20	11.98	9.80	6.45	6.39	6.12	6.93	5.72	6.38	29.29	29.67	15.52
30	11.99	9.80	6.45	6.39	6.12	6.93	5.72	6.38	29.30	29.67	15.52
40	11.99	9.81	6.45	6.39	6.12	6.93	5.72	6.38	29.30	29.68	15.53
50	12.00	9.82	6.45	6.39	6.12	6.94	5.72	6.39	29.30	29.68	15.54
60	12.00	9.82	6.46	6.39	6.13	6.94	5.73	6.39	29.31	29.68	15.55
70	12.00	9.83	6.46	6.40	6.13	6.94	5.73	6.40	29.31	29.68	15.56
80	12.01	9.83	6.46	6.40	6.14	6.94	5.73	6.40	29.32	29.68	15.56
90	12.01	9.84	6.46	6.40	6.14	6.94	5.74	6.40	29.32	29.68	15.57
3900	12.02	9.84	6.46	6.40	6.14	6.95	5.74	6.41	29.32	29.69	15.58
10	12.02	9.85	6.46	6.40	6.14	6.95	5.74	6.41	29.32	29.69	15.59
20	12.02	9.86	6.46	6.41	6.15	6.95	5.75	6.41	29.33	29.69	15.60
30	12.03	9.86	6.47	6.41	6.15	6.96	5.75	6.42	29.33	29.69	15.61
40	12.03	9.87	6.47	6.41	6.15	6.96	5.75	6.42	29.33	29.70	15.62
50	12.03	9.88	6.47	6.41	6.16	6.96	5.76	6.42	29.34	29.70	15.62
60	12.04	9.88	6.47	6.41	6.16	6.96	5.76	6.43	29.34	29.70	15.63
70	12.04	9.89	6.47	6.42	6.16	6.97	5.76	6.43	29.34	29.70	15.64
80	12.04	9.89	6.47	6.42	6.16	6.97	5.76	6.44	29.34	29.70	15.64
90	12.04	9.90	6.48	6.42	6.17	6.97	5.77	6.44	29.35	29.71	15.65
4000	12.05	9.90	6.48	6.42	6.17	6.98	5.77	6.45	29.35	29.71	15.66
10	12.05	9.91	6.48	6.42	6.17	6.99	5.77	6.45	29.35	29.71	15.66
20	12.05	9.92	6.48	6.42	6.17	6.99	5.78	6.45	29.36	29.72	15.67
30	12.06	9.92	6.48	6.42	6.18	6.98	5.78	6.46	29.36	29.72	15.68
40	12.06	9.93	6.48	6.42	6.18	6.98	5.78	6.46	29.37	29.72	15.69
50	12.06	9.93	6.48	6.42	6.18	6.99	5.78	6.46	29.37	29.72	15.70
60	12.07	9.94	6.48	6.43	6.18	6.99	5.79	6.46	29.38	29.72	15.70
70	12.07	9.94	6.48	6.43	6.18	6.99	5.79	6.47	29.38	29.72	15.71
80	12.07	9.95	6.49	6.43	6.19	6.99	5.79	6.47	29.38	29.73	15.72
90	12.08	9.95	6.49	6.43	6.19	7.00	5.80	6.47	29.39	29.73	15.74
4100	12.08	9.96	6.49	6.43	6.19	7.00	5.80	6.48	29.39	29.73	15.73
10	12.08	9.96	6.49	6.44	6.20	7.00	5.80	6.48	29.39	29.73	15.74
20	12.08	9.97	6.49	6.44	6.20	7.00	5.80	6.49	29.40	29.74	15.74
30	12.09	9.97	6.49	6.44	6.20	7.01	5.81	6.49	29.40	29.74	15.75
40	12.10	9.98	6.50	6.44	6.20	7.01	5.81	6.50	29.40	29.74	15.76
50	12.10	9.98	6.50	6.44	6.21	7.01	5.81	6.50	29.41	29.74	15.76
60	12.10	9.99	6.50	6.44	6.21	7.02	5.82	6.50	29.41	29.74	15.77
70	12.11	9.99	6.50	6.45	6.21	7.02	5.82	6.51	29.42	29.75	15.78
80	12.12	10.00	6.50	6.45	6.21	7.02	5.82	6.51	29.42	29.75	15.78
90	12.12	10.00	6.50	6.45	6.22	7.02	5.82	6.52	29.42	29.75	15.79
4200	12.12	10.01	6.50	6.45	6.22	7.02	5.83	6.52	29.43	29.75	15.79
10	12.12	10.02	6.50	6.45	6.22	7.03	5.83	6.52	29.43	29.76	15.80
20	12.12	10.02	6.51	6.45	6.22	7.03	5.83	6.53	29.43	29.76	15.80
30	12.12	10.03	6.51	6.46	6.23	7.03	5.84	6.53	29.44	29.76	15.81
40	12.13	10.03	6.51	6.46	6.23	7.04	5.84	6.53	29.44	29.76	15.82
50	12.13	10.04	6.51	6.46	6.23	7.04	5.84	6.54	29.44	29.76	15.82
60	12.13	10.04	6.51	6.46	6.24	7.04	5.84	6.54	29.45	29.76	15.83
70	12.14	10.05	6.51	6.46	6.24	7.04	5.85	6.54	29.45	29.77	15.84
80	12.14	10.05	6.51	6.46	6.24	7.05	5.85	6.55	29.46	29.77	15.84
90	12.14	10.06	6.51	6.47	6.24	7.05	5.85	6.55	29.46	29.77	15.85
4300	12.14	10.06	6.52	6.47	6.25	7.05	5.85	6.56	29.46	29.77	15.86
10	12.15	10.07	6.52	6.47	6.25	7.05	5.86	6.56	29.46	29.78	15.86
20	12.15	10.07	6.52	6.47	6.25	7.06	5.86	6.56	29.47	29.78	15.87
30	12.16	10.08	6.52	6.47	6.25	7.06	5.87	6.57	29.47	29.78	15.87
40	12.16	10.08	6.52	6.47	6.26	7.06	5.87	6.57	29.47	29.78	15.88
50	12.17	10.09	6.52	6.48	6.26	7.07	5.88	6.58	29.48	29.78	15.89
60	12.17	10.09	6.52	6.48	6.26	7.07	5.88	6.58	29.48	29.78	15.89
70	12.17	10.10	6.52	6.48	6.27	7.07	5.88	6.59	29.48	29.79	15.90
80	12.17	10.10	6.52	6.48	6.27	7.07	5.88	6.59	29.48	29.79	15.90
90	12.18	10.11	6.53	6.48	6.27	7.08	5.89	6.60	29.49	29.79	15.91
4400	12.18	10.11	6.53	6.48	6.27	7.08	5.89	6.60	29.49	29.79	15.92
10	12.18	10.12	6.53	6.49	6.27	7.08	5.89	6.60	29.50	29.79	15.92
20	12.19	10.12	6.53	6.49	6.28	7.08	5.90	6.60	29.50	29.80	15.92
30	12.19	10.13	6.53	6.49	6.28	7.08	5.90	6.61	29.50	29.80	15.93
40	12.20	10.14	6.53	6.49	6.28	7.09	5.90	6.61	29.51	29.80	15.94
50	12.20	10.14	6.53	6.49	6.28	7.09	5.91	6.62	29.51	29.80	15.94
60	12.20	10.15	6.54	6.49	6.28	7.09	5.91	6.62	29.51	29.80	15.95
70	12.21	10.15	6.54	6.49	6.29	7.09	5.91	6.62	29.51	29.80	15.95
80	12.21	10.16	6.54	6.49	6.29	7.09	5.91	6.62	29.51	29.80	15.95
90	12.21	10.16	6.54	6.50	6.29	7.10	5.92	6.63	29.52	29.80	15.96

Table III (continued)  
Average  $C_p$  Over Range from 275 °K to T  
(calories/mole/°K)

T, °K	CO <sub>2</sub>	H <sub>2</sub> O	CO	H <sub>2</sub>	H <sub>2</sub>	O <sub>2</sub>	C(s)	HCl	Al <sub>2</sub> O <sub>3</sub>	H <sub>2</sub> O <sub>2</sub>	H <sub>2</sub> O
1500	12.22	10.17	6.54	6.50	6.30	7.10	5.92	6.63	29.52	29.80	15.96
10	12.22	10.17	6.54	6.50	6.30	7.10	5.92	6.64	29.52	29.80	15.97
20	12.22	10.18	6.54	6.50	6.30	7.10	5.93	6.64	29.53	29.81	15.97
30	12.23	10.18	6.54	6.50	6.30	7.10	5.93	6.64	29.53	29.81	15.98
40	12.23	10.19	6.54	6.50	6.30	7.10	5.93	6.65	29.53	29.81	15.98
50	12.23	10.19	6.55	6.50	6.31	7.11	5.94	6.65	29.54	29.82	15.99
60	12.23	10.20	6.55	6.51	6.31	7.11	5.94	6.66	29.54	29.82	15.99
70	12.24	10.20	6.55	6.51	6.31	7.11	5.94	6.66	29.54	29.82	16.00
80	12.24	10.20	6.55	6.51	6.31	7.12	5.94	6.66	29.54	29.82	16.00
90	12.24	10.21	6.55	6.51	6.32	7.12	5.95	6.67	29.55	29.82	16.01
1600	12.25	10.21	6.55	6.51	6.32	7.12	5.95	6.67	29.55	29.82	16.01
10	12.25	10.22	6.55	6.51	6.32	7.12	5.95	6.68	29.55	29.82	16.02
20	12.25	10.22	6.55	6.51	6.32	7.12	5.96	6.68	29.55	29.82	16.02
30	12.26	10.23	6.56	6.52	6.33	7.12	5.96	6.68	29.56	29.82	16.03
40	12.26	10.23	6.56	6.52	6.33	7.13	5.96	6.68	29.56	29.82	16.03
50	12.26	10.24	6.56	6.52	6.33	7.13	5.96	6.69	29.56	29.82	16.04
60	12.27	10.24	6.56	6.52	6.33	7.13	5.97	6.69	29.56	29.82	16.04
70	12.27	10.25	6.56	6.52	6.34	7.14	5.97	6.70	29.57	29.83	16.05
80	12.27	10.25	6.56	6.52	6.34	7.14	5.97	6.70	29.57	29.83	16.05
90	12.27	10.26	6.56	6.52	6.34	7.14	5.97	6.70	29.57	29.83	16.06
1700	12.28	10.26	6.56	6.52	6.34	7.14	5.98	6.71	29.57	29.83	16.06
10	12.28	10.26	6.56	6.52	6.34	7.14	5.98	6.71	29.58	29.83	16.06
20	12.28	10.27	6.56	6.53	6.35	7.14	5.98	6.72	29.58	29.83	16.07
30	12.28	10.27	6.57	6.53	6.35	7.15	5.99	6.72	29.58	29.83	16.07
40	12.29	10.27	6.57	6.53	6.35	7.15	5.99	6.72	29.58	29.83	16.08
50	12.29	10.28	6.57	6.53	6.35	7.15	5.99	6.72	29.59	29.83	16.08
60	12.29	10.29	6.57	6.53	6.36	7.15	5.99	6.73	29.59	29.83	16.09
70	12.30	10.29	6.57	6.53	6.36	7.16	6.00	6.74	29.59	29.83	16.09
80	12.30	10.30	6.57	6.54	6.36	7.16	6.00	6.74	29.59	29.83	16.10
90	12.30	10.30	6.57	6.54	6.36	7.16	6.00	6.74	29.60	29.83	16.10
1800	12.31	10.31	6.57	6.54	6.36	7.16	6.00	6.75	29.60	29.83	16.10
10	12.31	10.31	6.57	6.54	6.37	7.16	6.00	6.75	29.60	29.83	16.11
20	12.31	10.31	6.58	6.54	6.37	7.17	6.01	6.75	29.60	29.83	16.11
30	12.32	10.32	6.58	6.54	6.37	7.17	6.01	6.76	29.60	29.85	16.12
40	12.32	10.32	6.58	6.54	6.37	7.17	6.02	6.76	29.61	29.85	16.12
50	12.32	10.33	6.58	6.55	6.37	7.17	6.02	6.76	29.61	29.85	16.13
60	12.32	10.33	6.58	6.55	6.37	7.17	6.02	6.77	29.61	29.85	16.13
70	12.32	10.34	6.58	6.55	6.38	7.18	6.03	6.77	29.62	29.85	16.14
80	12.33	10.34	6.58	6.55	6.38	7.18	6.03	6.78	29.62	29.85	16.14
90	12.33	10.34	6.58	6.55	6.38	7.18	6.03	6.78	29.62	29.85	16.14
1900	12.33	10.35	6.58	6.55	6.38	7.18	6.04	6.78	29.62	29.85	16.15
10	12.34	10.35	6.58	6.55	6.38	7.18	6.04	6.79	29.62	29.86	16.15
20	12.34	10.36	6.59	6.55	6.39	7.19	6.04	6.79	29.63	29.86	16.16
30	12.34	10.36	6.59	6.56	6.39	7.19	6.05	6.80	29.63	29.86	16.16
40	12.34	10.36	6.59	6.56	6.39	7.19	6.05	6.80	29.63	29.86	16.16
50	12.35	10.37	6.59	6.56	6.39	7.19	6.05	6.81	29.64	29.86	16.17
60	12.35	10.37	6.59	6.56	6.39	7.20	6.05	6.81	29.64	29.86	16.17
70	12.36	10.38	6.59	6.56	6.40	7.20	6.06	6.82	29.64	29.86	16.18
80	12.36	10.38	6.59	6.56	6.40	7.20	6.06	6.82	29.64	29.86	16.18
90	12.36	10.38	6.59	6.56	6.40	7.20	6.06	6.82	29.64	29.86	16.18
5000	12.36	10.39	6.59	6.56	6.40	7.20	6.06	6.82	29.64	29.86	16.18
10	12.36	10.40	6.60	6.56	6.40	7.21	6.07	6.83	29.64	29.86	16.19
20	12.37	10.40	6.60	6.56	6.41	7.21	6.07	6.83	29.64	29.86	16.19
30	12.37	10.40	6.60	6.57	6.41	7.21	6.08	6.84	29.65	29.86	16.20
40	12.38	10.41	6.60	6.57	6.41	7.22	6.08	6.84	29.65	29.86	16.20
50	12.38	10.41	6.60	6.57	6.41	7.22	6.08	6.84	29.65	29.86	16.20
60	12.38	10.42	6.60	6.57	6.41	7.22	6.09	6.85	29.66	29.87	16.21
70	12.38	10.42	6.60	6.57	6.42	7.22	6.09	6.85	29.66	29.87	16.21
80	12.39	10.42	6.60	6.57	6.42	7.22	6.09	6.86	29.66	29.87	16.22
90	12.39	10.42	6.60	6.57	6.42	7.22	6.09	6.86	29.66	29.87	16.22
2100	12.39	10.43	6.61	6.57	6.42	7.22	6.10	6.86	29.66	29.87	16.22
10	12.40	10.43	6.61	6.58	6.43	7.22	6.10	6.87	29.67	29.87	16.23
20	12.40	10.44	6.61	6.58	6.43	7.23	6.10	6.87	29.67	29.87	16.23
30	12.40	10.45	6.61	6.58	6.43	7.23	6.11	6.88	29.67	29.87	16.24
40	12.40	10.45	6.61	6.58	6.43	7.23	6.11	6.88	29.68	29.88	16.24
50	12.41	10.46	6.61	6.58	6.43	7.24	6.11	6.88	29.68	29.88	16.25
60	12.41	10.46	6.61	6.58	6.44	7.24	6.12	6.89	29.68	29.88	16.25
70	12.41	10.46	6.61	6.58	6.44	7.24	6.12	6.89	29.68	29.88	16.25
80	12.42	10.47	6.61	6.59	6.44	7.24	6.12	6.89	29.68	29.88	16.25
90	12.42	10.47	6.62	6.59	6.44	7.24	6.12	6.90	29.68	29.88	16.25

Table 149 (continued)

Average  $C_p$  Over Range  $T_1 = 225$  °F to  $T_2$   
(calorie./mole/°K)

$T_2$ , °K	CO <sub>2</sub>	H <sub>2</sub> O	CO	N <sub>2</sub>	H <sub>2</sub>	O <sub>2</sub>	C(s)	HCl	Al <sub>2</sub> O <sub>3</sub>	SiO <sub>2</sub>	K <sub>2</sub> C
5200	12.42	10.48	6.62	6.59	6.44	7.25	6.13	6.90	29.68	29.88	16.26
10	12.42	10.48	6.62	6.59	6.45	7.25	6.13	6.90	29.68	29.88	16.27
20	12.43	10.49	6.62	6.59	6.45	7.25	6.13	6.91	29.69	29.88	16.27
30	12.43	10.49	6.62	6.59	6.45	7.25	6.14	6.91	29.69	29.88	16.28
40	12.44	10.49	6.62	6.59	6.45	7.26	6.14	6.92	29.69	29.88	16.28
50	12.44	10.50	6.62	6.59	6.45	7.26	6.14	6.92	29.69	29.88	16.28
60	12.44	10.50	6.62	6.59	6.45	7.26	6.14	6.92	29.69	29.88	16.28
70	12.44	10.51	6.62	6.59	6.46	7.26	6.15	6.93	29.70	29.88	16.29
80	12.45	10.51	6.62	6.60	6.46	7.26	6.15	6.93	29.70	29.88	16.29
90	12.45	10.52	6.63	6.60	6.46	7.27	6.15	6.94	29.70	29.88	16.30
5300	12.45	10.52	6.63	6.60	6.46	7.27	6.16	6.94	29.70	29.88	16.30
10	12.46	10.53	6.63	6.60	6.46	7.27	6.16	6.94	29.70	29.89	16.30
20	12.46	10.53	6.63	6.60	6.47	7.28	6.16	6.94	29.71	29.89	16.31
30	12.46	10.54	6.63	6.60	6.47	7.28	6.16	6.95	29.71	29.89	16.31
40	12.46	10.54	6.63	6.60	6.47	7.28	6.16	6.95	29.71	29.89	16.32
50	12.47	10.54	6.63	6.60	6.47	7.28	6.17	6.96	29.71	29.89	16.32
60	12.47	10.55	6.63	6.61	6.47	7.28	6.17	6.96	29.71	29.89	16.32
70	12.47	10.55	6.63	6.61	6.48	7.28	6.18	6.96	29.72	29.89	16.33
80	12.48	10.56	6.64	6.61	6.48	7.29	6.18	6.97	29.72	29.89	16.33
90	12.48	10.56	6.64	6.61	6.48	7.29	6.19	6.97	29.72	29.90	16.33
5400	12.48	10.56	6.64	6.61	6.48	7.29	6.19	6.97	29.72	29.90	16.34
10	12.48	10.57	6.64	6.61	6.48	7.29	6.19	6.98	29.72	29.90	16.34
20	12.49	10.57	6.64	6.61	6.49	7.29	6.20	6.98	29.72	29.90	16.34
30	12.49	10.58	6.64	6.61	6.49	7.29	6.20	6.98	29.73	29.90	16.35
40	12.49	10.58	6.64	6.62	6.49	7.30	6.20	6.99	29.73	29.90	16.35
50	12.50	10.59	6.64	6.62	6.49	7.30	6.20	6.99	29.73	29.90	16.36
60	12.50	10.59	6.64	6.62	6.49	7.30	6.21	7.00	29.73	29.90	16.36
70	12.50	10.59	6.64	6.62	6.50	7.30	6.21	7.00	29.74	29.90	16.36
80	12.50	10.59	6.64	6.62	6.50	7.30	6.21	7.00	29.74	29.90	16.37
90	12.51	10.60	6.65	6.62	6.50	7.31	6.22	7.01	29.74	29.90	16.37
5500	12.51	10.61	6.65	6.62	6.50	7.31	6.22	7.01	29.74	29.90	16.37
10	12.51	10.61	6.65	6.62	6.50	7.31	6.22	7.02	29.74	29.92	16.38
20	12.52	10.62	6.65	6.63	6.51	7.31	6.23	7.02	29.74	29.91	16.38
30	12.52	10.62	6.65	6.63	6.51	7.32	6.23	7.02	29.75	29.91	16.38
40	12.52	10.62	6.65	6.63	6.51	7.32	6.23	7.03	29.75	29.91	16.39
50	12.52	10.63	6.65	6.63	6.51	7.32	6.24	7.03	29.75	29.91	16.39
60	12.53	10.63	6.65	6.63	6.51	7.32	6.24	7.04	29.75	29.91	16.40
70	12.53	10.64	6.65	6.63	6.52	7.32	6.24	7.04	29.76	29.91	16.40
80	12.53	10.64	6.65	6.63	6.52	7.33	6.24	7.04	29.76	29.91	16.40
90	12.54	10.65	6.66	6.63	6.52	7.33	6.25	7.04	29.76	29.91	16.41
5600	12.54	10.65	6.66	6.63	6.52	7.33	6.25	7.05	29.76	29.92	16.41
10	12.54	10.66	6.66	6.63	6.53	7.33	6.25	7.05	29.76	29.92	16.42
20	12.54	10.66	6.66	6.64	6.53	7.33	6.25	7.06	29.76	29.92	16.42
30	12.55	10.66	6.66	6.64	6.53	7.34	6.26	7.06	29.76	29.92	16.42
40	12.55	10.67	6.66	6.64	6.53	7.34	6.26	7.06	29.77	29.92	16.42
50	12.55	10.67	6.66	6.64	6.53	7.34	6.27	7.07	29.77	29.92	16.43
60	12.56	10.68	6.66	6.64	6.54	7.34	6.27	7.07	29.77	29.92	16.43
70	12.56	10.68	6.66	6.64	6.54	7.34	6.27	7.06	29.77	29.92	16.44
80	12.56	10.69	6.67	6.64	6.54	7.35	6.28	7.08	29.78	29.92	16.44
90	12.56	10.69	6.67	6.64	6.54	7.35	6.28	7.08	29.78	29.92	16.44
5700	12.57	10.69	6.67	6.65	6.54	7.35	6.28	7.09	29.78	29.92	16.45
10	12.57	10.70	6.67	6.65	6.55	7.35	6.29	7.10	29.78	29.92	16.45
20	12.57	10.70	6.67	6.65	6.55	7.35	6.29	7.10	29.78	29.92	16.46
30	12.58	10.71	6.67	6.65	6.55	7.36	6.29	7.10	29.78	29.93	16.46
40	12.58	10.71	6.67	6.65	6.55	7.36	6.30	7.10	29.78	29.93	16.46
50	12.58	10.72	6.67	6.65	6.55	7.36	6.30	7.11	29.79	29.93	16.46
60	12.59	10.72	6.67	6.65	6.56	7.36	6.30	7.11	29.79	29.93	16.47
70	12.59	10.72	6.68	6.65	6.55	7.36	6.30	7.12	29.79	29.93	16.47
80	12.59	10.73	6.68	6.66	6.56	7.36	6.31	7.12	29.80	29.93	16.48
90	12.60	10.73	6.68	6.66	6.56	7.36	6.31	7.12	29.80	29.93	16.48
5800	12.60	10.74	6.68	6.66	6.56	7.37	6.31	7.12	29.80	29.93	16.48
10	12.60	10.75	6.68	6.66	6.57	7.37	6.31	7.13	29.80	29.93	16.48
20	12.60	10.75	6.68	6.66	6.57	7.37	6.32	7.13	29.80	29.93	16.49
30	12.61	10.75	6.68	6.66	6.57	7.38	6.32	7.14	29.80	29.93	16.49
40	12.61	10.76	6.68	6.67	6.57	7.38	6.33	7.14	29.81	29.93	16.50
50	12.62	10.76	6.69	6.67	6.57	7.38	6.33	7.14	29.81	29.93	16.50
60	12.62	10.76	6.69	6.67	6.58	7.38	6.33	7.15	29.81	29.94	16.50
70	12.62	10.77	6.69	6.67	6.58	7.38	6.33	7.15	29.82	29.94	16.51
80	12.62	10.77	6.69	6.67	6.58	7.38	6.34	7.16	29.82	29.94	16.51
90	12.62	10.78	6.69	6.67	6.58	7.39	6.34	7.16	29.82	29.94	16.51

Table 149 (continued)

Average  $C_p$  Over Range from 290 °K to T  
(calories/mole/°K)

T, °K	CO <sub>2</sub>	H <sub>2</sub> (l)	CO	H <sub>2</sub>	H <sub>2</sub>	O <sub>2</sub>	C(s)	HCl	Al <sub>2</sub> O <sub>3</sub>	B <sub>2</sub> O <sub>3</sub>	H <sub>2</sub> O
5900	12.63	10.78	6.69	6.67	6.59	7.39	6.34	7.16	29.82	29.94	16.52
10	12.63	10.78	6.69	6.67	6.59	7.39	6.34	7.17	29.82	29.94	16.52
20	12.64	10.79	6.69	6.67	6.59	7.39	6.35	7.17	29.82	29.94	16.52
30	12.64	10.79	6.69	6.67	6.59	7.40	6.35	7.18	29.82	29.94	16.53
40	12.64	10.80	6.69	6.67	6.59	7.40	6.36	7.18	29.83	29.94	16.53
50	12.64	10.80	6.70	6.68	6.59	7.40	6.36	7.18	29.83	29.94	16.53
60	12.65	10.81	6.70	6.68	6.60	7.40	6.36	7.19	29.83	29.94	16.54
70	12.65	10.81	6.70	6.68	6.60	7.40	6.36	7.19	29.83	29.94	16.54
80	12.65	10.82	6.70	6.68	6.60	7.41	6.37	7.20	29.84	29.95	16.54
90	12.66	10.82	6.70	6.68	6.60	7.41	6.37	7.20	29.84	29.95	16.55
6000	12.66	10.82	6.70	6.68	6.60	7.41	6.37	7.20	29.84	29.95	16.55
10	12.66	10.83	6.70	6.68	6.61	7.42	6.38	7.21	29.84	29.95	16.55
20	12.67	10.83	6.70	6.68	6.61	7.42	6.38	7.21	29.84	29.95	16.56
30	12.67	10.84	6.70	6.69	6.61	7.42	6.39	7.22	29.84	29.95	16.56
40	12.67	10.84	6.71	6.69	6.61	7.42	6.39	7.22	29.85	29.95	16.57
50	12.68	10.85	6.71	6.69	6.62	7.42	6.39	7.22	29.85	29.95	16.57
60	12.68	10.85	6.71	6.69	6.62	7.43	6.40	7.23	29.85	29.95	16.57
70	12.68	10.85	6.71	6.69	6.62	7.43	6.40	7.23	29.85	29.96	16.58
80	12.69	10.86	6.71	6.69	6.62	7.43	6.40	7.24	29.86	29.96	16.58
90	12.69	10.86	6.71	6.69	6.62	7.43	6.40	7.24	29.86	29.96	16.58
6100	12.69	10.87	6.71	6.69	6.62	7.43	6.40	7.24	29.86	29.96	16.58
10	12.70	10.87	6.71	6.70	6.63	7.44	6.41	7.25	29.86	29.96	16.59
20	12.70	10.88	6.71	6.70	6.63	7.44	6.41	7.25	29.86	29.96	16.59
30	12.70	10.88	6.71	6.70	6.63	7.44	6.41	7.25	29.86	29.96	16.59
40	12.70	10.88	6.72	6.70	6.63	7.44	6.42	7.26	29.87	29.96	16.60
50	12.71	10.89	6.72	6.70	6.63	7.44	6.42	7.26	29.87	29.96	16.60
60	12.71	10.89	6.72	6.70	6.64	7.44	6.42	7.26	29.87	29.96	16.60
70	12.71	10.90	6.72	6.70	6.64	7.45	6.43	7.27	29.87	29.96	16.60
80	12.72	10.90	6.72	6.71	6.64	7.45	6.43	7.27	29.88	29.96	16.61
90	12.72	10.91	6.72	6.71	6.64	7.45	6.43	7.28	29.88	29.96	16.61
6200	12.72	10.91	6.72	6.71	6.65	7.45	6.44	7.28	29.88	29.97	16.62
10	12.72	10.92	6.72	6.71	6.65	7.45	6.44	7.28	29.88	29.97	16.62
20	12.73	10.92	6.72	6.71	6.65	7.46	6.44	7.29	29.88	29.97	16.62
30	12.73	10.93	6.73	6.71	6.65	7.46	6.45	7.30	29.88	29.97	16.62
40	12.73	10.93	6.73	6.71	6.65	7.46	6.45	7.30	29.88	29.97	16.63
50	12.74	10.93	6.73	6.71	6.65	7.46	6.45	7.30	29.88	29.97	16.63
60	12.74	10.94	6.73	6.72	6.66	7.47	6.45	7.31	29.89	29.97	16.64
70	12.74	10.94	6.73	6.72	6.66	7.47	6.45	7.31	29.89	29.97	16.64
80	12.75	10.94	6.73	6.72	6.66	7.47	6.46	7.31	29.89	29.97	16.64
90	12.75	10.95	6.73	6.72	6.66	7.47	6.46	7.31	29.89	29.98	16.64
6300	12.75	10.95	6.73	6.72	6.67	7.47	6.46	7.32	29.90	29.98	16.65
10	12.76	10.96	6.73	6.72	6.67	7.47	6.47	7.32	29.90	29.98	16.65
20	12.76	10.96	6.73	6.72	6.67	7.47	6.47	7.32	29.90	29.98	16.65
30	12.76	10.97	6.74	6.72	6.67	7.48	6.47	7.33	29.90	29.98	16.66
40	12.76	10.98	6.74	6.73	6.67	7.48	6.48	7.33	29.91	29.98	16.66
50	12.77	10.98	6.74	6.73	6.68	7.48	6.48	7.34	29.91	29.98	16.67
60	12.77	10.98	6.74	6.73	6.68	7.48	6.48	7.34	29.91	29.98	16.67
70	12.77	10.99	6.74	6.73	6.68	7.49	6.49	7.35	29.91	29.98	16.67
80	12.78	10.99	6.74	6.73	6.68	7.49	6.49	7.35	29.91	29.98	16.67
90	12.78	11.00	6.74	6.73	6.68	7.49	6.49	7.35	29.91	29.98	16.67
6400	12.78	11.00	6.74	6.73	6.69	7.49	6.50	7.35	29.92	29.98	16.68
10	12.79	11.01	6.74	6.73	6.69	7.49	6.50	7.36	29.92	29.99	16.68
20	12.79	11.01	6.75	6.73	6.69	7.49	6.50	7.36	29.92	29.99	16.69
30	12.79	11.01	6.75	6.74	6.70	7.50	6.51	7.37	29.92	29.99	16.69
40	12.80	11.02	6.75	6.74	6.70	7.50	6.51	7.37	29.93	29.99	16.70
50	12.80	11.02	6.75	6.74	6.70	7.50	6.51	7.37	29.93	29.99	16.70
60	12.80	11.03	6.75	6.74	6.70	7.50	6.52	7.38	29.93	29.99	16.70
70	12.80	11.03	6.75	6.74	6.70	7.50	6.52	7.38	29.93	29.99	16.70
80	12.81	11.03	6.75	6.74	6.70	7.50	6.52	7.39	29.93	29.99	16.70
90	12.81	11.04	6.75	6.74	6.70	7.50	6.52	7.39	29.93	29.99	16.70
6500	12.82	11.04	6.75	6.74	6.71	7.51	6.52	7.39	29.94	29.99	16.71
10	12.82	11.05	6.75	6.74	6.71	7.51	6.53	7.40	29.94	30.00	16.71
20	12.82	11.05	6.76	6.75	6.71	7.51	6.53	7.40	29.94	30.00	16.72
30	12.82	11.06	6.76	6.75	6.71	7.52	6.54	7.41	29.94	30.00	16.72
40	12.83	11.06	6.76	6.75	6.71	7.52	6.54	7.41	29.94	30.00	16.72
50	12.83	11.07	6.76	6.75	6.72	7.52	6.54	7.42	29.95	30.00	16.73
60	12.83	11.07	6.76	6.75	6.72	7.52	6.54	7.42	29.95	30.00	16.73
70	12.84	11.07	6.76	6.75	6.72	7.52	6.55	7.42	29.95	30.00	16.74
80	12.84	11.08	6.76	6.75	6.72	7.52	6.55	7.43	29.95	30.00	16.74
90	12.84	11.08	6.76	6.75	6.72	7.53	6.56	7.43	29.95	30.00	16.74

Table 149 (continued)  
Average  $C_p$  Over Range from 298 °K to T  
(caloria/mole/°K)

T, °K	CO <sub>2</sub>	H <sub>2</sub> O	CO	H <sub>2</sub>	H <sub>2</sub>	O <sub>2</sub>	C(s)	NaCl	Al <sub>2</sub> O <sub>3</sub>	B <sub>2</sub> O <sub>3</sub>	MgO
6500	12.84	11.09	6.77	6.76	6.73	7.53	6.56	7.43	29.96	30.00	16.74
10	12.85	11.09	6.77	6.76	6.73	7.53	6.56	7.43	29.96	30.00	16.75
20	12.85	11.10	6.77	6.76	6.73	7.54	6.56	7.44	29.96	30.00	16.75
30	12.85	11.10	6.77	6.76	6.73	7.54	6.57	7.44	29.96	30.00	16.76
40	12.86	11.10	6.77	6.76	6.74	7.54	6.57	7.44	29.96	30.00	16.76
50	12.86	11.11	6.77	6.76	6.74	7.54	6.57	7.45	29.97	30.01	16.77
60	12.86	11.11	6.77	6.76	6.74	7.54	6.57	7.45	29.97	30.01	16.77
70	12.86	11.12	6.77	6.76	6.74	7.55	6.58	7.46	29.97	30.01	16.77
80	12.87	11.12	6.77	6.77	6.74	7.55	6.58	7.46	29.97	30.01	16.77
90	12.87	11.12	6.77	6.77	6.75	7.55	6.58	7.46	29.97	30.01	16.78
6700	12.88	11.13	6.77	6.77	6.75	7.55	6.58	7.47	29.98	30.01	16.76
10	12.88	11.13	6.78	6.77	6.75	7.56	6.58	7.47	29.98	30.01	16.78
20	12.88	11.14	6.78	6.77	6.75	7.56	6.58	7.47	29.98	30.02	16.79
30	12.88	11.14	6.78	6.77	6.75	7.56	6.59	7.48	29.98	30.02	16.79
40	12.89	11.15	6.78	6.77	6.76	7.56	6.60	7.48	29.98	30.02	16.79
50	12.89	11.15	6.78	6.77	6.76	7.56	6.60	7.48	29.99	30.02	16.80
60	12.89	11.16	6.78	6.77	6.76	7.56	6.60	7.49	29.99	30.02	16.80
70	12.90	11.16	6.78	6.77	6.76	7.56	6.60	7.49	29.99	30.02	16.80
80	12.90	11.16	6.78	6.77	6.76	7.57	6.61	7.50	29.99	30.02	16.80
90	12.90	11.17	6.78	6.77	6.77	7.57	6.61	7.50	29.99	30.02	16.81
6800	12.91	11.17	6.78	6.78	6.77	7.57	6.61	7.50	30.00	30.02	16.81
10	12.91	11.18	6.78	6.78	6.77	7.57	6.62	7.51	30.00	30.02	16.81
20	12.91	11.18	6.78	6.78	6.77	7.58	6.62	7.51	30.00	30.02	16.82
30	12.92	11.19	6.79	6.78	6.77	7.58	6.62	7.52	30.00	30.02	16.82
40	12.92	11.19	6.79	6.78	6.78	7.58	6.62	7.52	30.00	30.02	16.82
50	12.92	11.20	6.79	6.79	6.78	7.58	6.63	7.52	30.01	30.03	16.82
60	12.92	11.20	6.79	6.79	6.78	7.58	6.63	7.53	30.01	30.03	16.83
70	12.93	11.20	6.79	6.79	6.78	7.58	6.63	7.53	30.01	30.03	16.83
80	12.93	11.21	6.79	6.79	6.79	7.59	6.64	7.53	30.01	30.03	16.83
90	12.93	11.21	6.79	6.79	6.79	7.59	6.64	7.54	30.01	30.03	16.84
6900	12.94	11.22	6.79	6.79	6.79	7.59	6.64	7.54	30.02	30.03	16.84
10	12.94	11.22	6.80	6.79	6.79	7.59	6.64	7.54	30.02	30.03	16.84
20	12.94	11.23	6.80	6.80	6.79	7.60	6.65	7.55	30.02	30.03	16.84
30	12.95	11.23	6.80	6.80	6.79	7.60	6.65	7.55	30.02	30.03	16.85
40	12.95	11.24	6.80	6.80	6.80	7.60	6.65	7.56	30.02	30.03	16.85
50	12.95	11.24	6.80	6.80	6.80	7.60	6.66	7.56	30.03	30.04	16.86
60	12.96	11.24	6.80	6.80	6.80	7.60	6.66	7.56	30.03	30.04	16.86
70	12.96	11.25	6.80	6.80	6.80	7.61	6.66	7.57	30.03	30.04	16.86
80	12.96	11.25	6.80	6.80	6.80	7.61	6.67	7.57	30.03	30.04	16.86
90	12.96	11.26	6.80	6.80	6.80	7.61	6.67	7.57	30.04	30.04	16.87



CONFIDENTIAL  
SECURITY INFORMATION

ratio,  $\frac{a}{M_1}$ , is obtained. The molecular weight of the mixture may be calculated from the equation

$$\frac{M_w}{\text{mixture}} = \frac{100}{\frac{a}{M_1} + \frac{b}{M_2}} \quad (3)$$

where a and b are the weight per cents of the components and  $M_1$  and  $M_2$  their respective molecular weights. The molar ratios are then multiplied by the  $\frac{M_w}{\text{mixture}}$  to convert to the basis of one mole of mixtures as shown.

The total number of carbon, hydrogen, nitrogen and oxygen atoms are added in the form provided on the reverse side and the detonation equation is written according to the rules set forth on page 580. The heat of combustion at constant volume ( $H_2O$  gas) is calculated as the sum of the heat of combustion of each component times its molar quantity. The heat of explosion, flame temperature, number of moles,  $n_{H_2O}$ , and final power are then calculated in the same manner as that described for a pure compound.

Figures 90 and 91 represent a similar calculation for a metallized mixture, Torpex, illustrating Rule 4, page 580. This calculation shows the effect of considering the  $Al_2O_3$  as solid and gaseous.

### E. Conclusions and Recommendations

The application of this system to the calculation of some two hundred high explosives (pure compounds and mixtures) for which ballistic mortar values have been determined shows a reasonably good correlation between observed and calculated power values as expressed in per cent TNT.

This method of calculation has two major uses: (1) the preliminary evaluation of new compounds and mixtures without the necessity for making and testing them, and (2) the checking of observed data. These uses have been illustrated in the discussion.

The future extension of this system depends upon the availability of additional thermodynamic data, upon further studies of the reaction mechanisms involved in the explosive mixtures particularly those containing metals, and upon the retesting of those explosives for which the correlation between measured and calculated power is unsatisfactory.

The somewhat less satisfactory agreement achieved for the metallized mixtures (Table 115) leads to the conclusion that the observed values bear re-determination either in the ballistic mortar or the new-type spherical lead block. Rather than merely repeating the measurements already made, it is recommended that a program be laid out to include the investigation of

CONFIDENTIAL

CONFIDENTIAL  
SECURITY INFORMATION

1. The optimum amount of aluminum for any one mixture;
2. The relative contribution to the power of high explosives of a number of metals, including among others aluminum, boron, beryllium, magnesium and lithium and possibly some of their hydrides;
3. The effect of the addition of high-nitrogen or other high energy compounds to metallized explosives; and
4. The relationship between the various test procedures including ballistic mortar, lead block, closed-blast and open-air blast.

The details of such a program must await the completion of a study on metallized explosives which is now underway.

In Table 116 there are listed pure compounds calculated to have powers in excess of 100% TNT, the highest so far measured or calculated for a known pure compound, MEDINA. It is recommended that if feasible, one or more of these compounds be synthesized and measured in the ballistic mortar and spherical lead block.

The large discrepancies existing between observed and calculated power for compounds of large positive oxygen balance, numbers 136, 240, 327 and 337 (Table 116) should probably be investigated further by a re-measurement of compounds 136 and 327 and any others of high positive oxygen balance which can be conveniently obtained.

CONFIDENTIAL

CONFIDENTIAL  
SECURITY INFORMATION

F. References

1. Aerojet Engineering Corp., Report No. 512, Contract N7 onr-46208 by M. H. Gold and A. O. Dekker (14 June 1951). CONFIDENTIAL
2. J. S. Arthur, J. App. Phys., 21, 732 (1950). "The specific heats of MgO, TiO<sub>2</sub> and ZrO<sub>2</sub> at high temperatures".
3. Arthur D. Little, Inc., Contract W-19-020-ORD-6436 (2 January 1947). "Study of pure explosive compounds. Part I. Correlation of organic structure with explosive properties from existing data". CONFIDENTIAL
4. Arthur D. Little, Inc., Contract W-19-020-ORD-6436 (2 April 1947). "Study of pure explosive compounds. Part II. Correlation of thermal quantities with explosive properties". CONFIDENTIAL
5. Arthur D. Little, Inc., Contract W-19-020-ORD-12 (1 May 1950). "Study of pure explosive compounds. Part III. Correlation of composition of mixtures with performance". CONFIDENTIAL
6. Arthur D. Little, Inc., Contract W-19-020-ORD-6436 and DA-19-020-ORD-12 (1 March 1951). "Second report on synthesis and testing of high explosives". pp. 367-371. CONFIDENTIAL See also Arthur D. Little, Inc., "Fundamental research on high explosives. Abstract of work performed under contract DA-19-020-ORD-12 with Office of the Chief of Ordnance". (11 October 1950) CONFIDENTIAL
7. National Bureau of Standards, "Selected values of chemical thermodynamic properties. Series I and II". U. S. Dept. of Commerce, Washington, D. C. (30 June 1947).
8. Naval Ordnance Laboratory, "Report on the program to develop an improved HXI-type explosive". NavOrd Report 1502, 26 July 1950. CONFIDENTIAL
9. W. Nernst and K. Wohl, Zeitschr. f. techn. physik. 608-614 (1929). "Specific heats at high temperature".
10. ONRD 547. J. O. Hirschfelder, F. T. McClure, C. F. Curties, and D. W. Osborne, "Thermodynamic properties of propellant gases". (23 November 1942) RESTRICTED
11. J. C. Southard, J. Am. Chem. Soc., 63, 3147 (1941). "The thermal properties of crystalline and glassy boron trioxide".
12. A. Thuret, J. Soc. Glass Tech., 20, 680-84 (1936). "Calculation of the specific heats of vitreous silica and components of glasses as functions of the temperature".

CONFIDENTIAL

CONFIDENTIAL  
SECURITY INFORMATION

INITIAL DISTRIBUTION LIST  
OF  
"FUNDAMENTAL STUDY OF HIGH EXPLOSIVES"

PART IV

<u>Copy No's.</u>		<u>No. of copies</u>
1-6	Chief, Bureau of Ordnance, Department of the Navy, Washington 25, D. C.	
	Technical Library, Ad3	1
	Chief, High Explosives and Amphibious Munitions, Re2o	2
	Chief, Propellants, Re2d	1
	Chief, Guided Missiles Jet Propulsion and Counter- measures, Re9	1
	U. S. Naval Attache, Technical Section, London, England	1
7,8	Commander, Naval Ordnance Laboratory, White Oak, Silver Spring 19, Md.	
9,10	Commander, Naval Ordnance Test Station, China Lake, Inyokern, Calif.	
11	Supt. U. S. Naval Postgraduate School, Monterey, Calif.	
12	Commander, Naval Powder Factory, Indian Head, Md.	
13	Commander, Naval Proving Ground, Dahlgren, Va.	
14,15	Chief of Naval Research, Department of the Navy, Washington 25, D. C.	
16	Director, Naval Research Laboratory, Bellevue, Washington 25, D. C.	
17	Chief of Naval Research, Technical Section, Navy Dept., Washington 25, D. C.	
18	Chief, Bureau of Aeronautics, Department of Navy, Washington 25, D. C.	
19	Chief, Bureau of Ships, Department of Navy, Washington, 25, D. C.	
20	Commandant, U. S. Marine Corps, Washington 25, D. C.	
21	Aerojet Engineering Corp., Azusa, Calif. Attn: Mr. H. H. Gold	
22	George Washington University, Dept. of Chemistry, Washington, D. C. Attn: Mr. John Farago	
23,24	Director, Applied Physics Laboratory, Johns Hopkins University, Silver Spring, Md.	
25	National Fireworks, Inc., West Hanover, Mass. Attn: J. J. Porter	
26	Naugatuck Chemical, Naugatuck, Conn. Attn: Mr. F. J. Foster	
27	Dr. Ralph Conner, Rohm and Haas Co., 5000 Richmond St., Philadelphia 37, Pa.	
28	U. S. Rubber Co., General Laboratories, Passaic, N. J. Attn: Dr. P. O. Towney	

CONFIDENTIAL

CONFIDENTIAL  
SECURITY INFORMATION

Copy No's.

No. of Copies

29	University of Louisville, Dept. of Chemistry, Louisville, Ky. Attn: Prof. R. H. Wiley	
30	University of Maryland, Dept. of Chemistry, Colleges Park, Md. Attn: Dr. N. L. Drake	
31	Dr. J. C. Holts, Bureau of Mines, 4800 Forbes St., Pittsburgh, Pa.	
32	Dr. Bernard Lewis, Bureau of Mines, 4800 Forbes St., Pittsburgh, Pa.	
33-37	Office, Chief of Chemical Corps, Res. and Dev. Div., Washington 25, D. C.	
38	National Bureau of Standards, Ord. Dev. Div., Washington 25, D. C. Attn: Mr. H. Lipnick	
39, 40	Director, Research and Development, GSUSA, Washington 25, D. C.	
41	War Dept. Library, Cataloguing and Accessions, Pentagon, Washington 25, D. C. Attn: V. A. Schaefer	
42	OSI, Armed Services Explosive Safety Board, Washington 25, D. C.	
43	Atomic Energy Commission, 1901 Constitution Ave., Washington 25, D. C. Attn: DMA	
44	Atomic Energy Commission, Santa Fe Operations Office, P.O. Box 5100, Albuquerque, N. M.	
45-49	Armed Services Tech Info Agency, Document Service Center, U. B. Bldg., Dayton 2, Ohio Attn: DSC-SD	
50	Commanding General, U. S. Air Force, Washington 25, D. C. Attn: AFIRD-AR, MD Tippelito	
51	Office, Chief of Engineers, Gravelly Point, Washington, D.C. Attn: EROHP, Mr. H. Lieske	
52	Engineer Research and Development Laboratories, Fort Belvoir, Va. Attn: R. L. Thornton	
53	Commanding General, Army Field Forces, Fort Monroe, Va. Attn: Asst. Chief for R and D	
54-58	Commanding General, Aberdeen Proving Ground, Maryland Attn: Col. A. P. Taber Dr. W. E. Sterne Mr. R. Kent Ordnance Board Ordnance School	1 1 1 1 1
59	Commanding Officer, Frankford Arsenal, Philadelphia 37, Pa.	
60, 61	Commanding Officer, Picatinny Arsenal, Dover, N. J. Attn: Tech. Div.	

CONFIDENTIAL

CONFIDENTIAL  
SECURITY INFORMATION

Copy No's.

No. of copies

62	Commanding Officer, Redstone Arsenal, Huntsville, Ala.	
	Attn: Tech Library, ORDDWR	
63	Bohn and Mann Co., Redstone Arsenal, Huntsville, Ala.	
64	Dr. Duncan McDougall, Los Alamos Scientific Laboratory, Los Alamos, N. M.	
65	Professor of Ordnance and Gunnery, U. S. Military Academy, West Point, N. Y.	
66	Office of Ordnance Research, 2127 Kyrtle Drive, Duke University, Durham, N. C.	
67	Atlas Powder Company, Reynolds Exp. Lab., Tamaqua, Pa.	
	Attn: Dr. W. J. Taylor, Jr.	
68	Dr. E. M. Pugh, Carnegie Institute of Technology, Pittsburgh 13, Pa.	
69	Dr. R. L. Evans, Evans Res. and Dev. Corp., 250 East 43rd St., New York, N. Y.	
70	Dr. G. B. Kistlikowsky, Harvard University Dept. of Chemistry, Cambridge, Mass.	
71	Dr. A. M. Ball, Hercules Powder Co., Wilmington 99, Del.	
72	Office, Chief of Ordnance, Washington 25, D. C., Attn: ORDTM-Dr. C. M. Salb	
73	Office, Chief of Ordnance, Washington 25, D. C., Attn: ORDTB-Mr. G. Stetson	
74	Office, Chief of Ordnance, Washington 25, D. C., Attn: ORDTX-AR, Mrs. G. R. Myers	
75	Office, Chief of Ordnance, Washington 25, D. C., Attn: ORDDU-CA Col. C. S. Reed	
76-102	OO, Chief, Ammunition Br, ORDTA, Res. and Dev. Div., Washington 25, D. C.	
	Dr. L. R. Littleton	2
	Military Attache, London, England	1
	British Joint Services Mission, Technical Service, 1800 H St., N.W., Washington, D. C.	
	Attn: Miss Mary O. Scott	5
	British Joint Services Mission, Naval Staff, Box 165, Benjamin Franklin Station, Washington, D. C., Attn: Com. Carew Hunt	2
	British Ministry of Supply, ARE, Main Bldg., Royal Arsenal, Woolich, SE18, England	
	Attn: Dr. H. J. Poole	4
	British Ministry of Supply, Shell-Mex House, Strand, London WC2, England	
	Attn: DGRD, Mr. C. S. Bryant	1
	Dr. W. B. Shringall	1
	Director, Nat'l Physics Laboratory, Engineering Section, Teddington, Middlesex, England	1

CONFIDENTIAL

NOTIFICATION OF MISSING PAGES

INSTRUCTIONS: THIS FORM IS INSERTED INTO ASTIA CATALOGED DOCUMENTS TO DENOTE MISSING PAGES.

DOCUMENT	CLASSIFICATION (CHECK ONE)		
	UNCLASSIFIED	CONFIDENTIAL	SECRET
AD 302 890		✓	
ATI			

THE PAGES, FIGURES, CHARTS, PHOTOGRAPHS, ETC., MISSING FROM THIS DOCUMENT ARE: BLANK

**DO NOT REMOVE**