

MEMORANDUM
RM-4088-NIH
MAY 1964

AD-600 161

NUMERICAL SOLUTION OF
FUNCTIONAL EQUATIONS BY
MEANS OF LAPLACE TRANSFORM-IV:
NONLINEAR EQUATIONS

R. E. Bellman, R. E. Kalaba and J. Lockett

This investigation was supported in part by Public Health Service Research Grant Number GM-09608-02, from the Division of General Medical Sciences, National Institutes of Health.

PREFACE

Mathematical studies of biological processes have many nonlinear functional equations. This study shows that the method of successive approximations, coupled with the numerical inversion of Laplace transforms, provides an effective computational approach.

copy → For illustrative purposes a nonlinear differential equation, a nonlinear differential-difference equation and a nonlinear diffusion equation are considered.

See also ↑ AD-428 074.

SUMMARY

↙ It is shown that Laplace transform can be effective in the
numerical solution of nonlinear functional equations. → to p iii

CONTENTS

PREFACE	iii
SUMMARY	v
Section	
I. INTRODUCTION	1
II. DIFFERENTIAL EQUATIONS	2
III. NUMERICAL RESULTS	3
IV. DIFFERENTIAL-DIFFERENCE EQUATIONS	3
V. NUMERICAL RESULTS	4
VI. HEAT EQUATION	4
VII. NUMERICAL RESULTS	5
VIII. CONVERGENCE ASPECTS	6
REFERENCES	14

I. INTRODUCTION

In the previous Memoranda in this series, ⁽¹⁻³⁾ we have demonstrated the effectiveness of the Laplace transform in obtaining the numerical solution of linear functional equations. In I, we considered the renewal equation, in II, differential-difference equations, and in III, the heat equation.

Here we wish to consider the possibility of using numerical inversion of the Laplace transform plus successive approximations to treat nonlinear functional equations. To illustrate our techniques, we consider the nonlinear differential equation

$$u' = -u - u^2, \quad u(0) = c, \quad (1)$$

the nonlinear differential-difference equation

$$u'(t) = -u(t-1) - u^2(t), \quad u(t) = c, \quad -1 \leq t \leq 0, \quad (2)$$

and the nonlinear heat equation

$$k(x)u_t = u_{xx} + bu^2 \quad (3)$$

$$u(x,0) = c \sin \pi x, \quad u(0,t) = u(1,t) = 0 .$$

Numerical results are given, as well as times of execution.

II. DIFFERENTIAL EQUATIONS

To illustrate the general approach in its simplest form, let us take the nonlinear differential Eq. (1). Taking Laplace transforms, we have

$$L(u') = -L(u) - L(u^2), \quad (4)$$

$$(s+1)L(u) = c - L(u^2), \quad (5)$$

$$L(u) = \frac{c - L(u^2)}{(s+1)}.$$

We solve this equation numerically by means of successive approximations plus numerical quadratures in the following fashion. The function $u_0(t)$ is taken as the solution of the equation

$$L(u) = \frac{c}{s+1}. \quad (6)$$

Ignoring the fact that we can find $u(t)$ explicitly, we use the quadrature techniques presented in I, II, and III to obtain the values $\{u_0(r_i)\}$.

Next we write

$$L(u_1) = \frac{c - L(u_0^2)}{s+1}. \quad (7)$$

To evaluate $L(u_0^2)$, we employ the same quadrature formula

$$L(u_0^2) = \sum_{i=1}^N w_i u_0^2(r_i) r_i^{s-1}. \quad (8)$$

We now determine u_1 using quadrature techniques and the explicit inversion formula.

Continuing in this fashion, we generate a sequence of functions $\{u_n(t)\}$ evaluated at the points $t_i = -\log r_i$, $i=1,2,\dots,N$, with

$$L(u_n) = \frac{c - L(u_{n-1}^2)}{(s+1)} \quad (9)$$

Naturally, the procedure will converge only if $|c|$ is sufficiently small. Subsequently, we shall discuss what techniques can be used in the general case.

III. NUMERICAL RESULTS

Ten successive approximations for each value of c can be accomplished in 6 sec of execution time on the IBM 7090. Graphs of the solutions are given in Figs. 1 and 2.

IV. DIFFERENTIAL-DIFFERENCE EQUATIONS

Similarly, starting with Eq. (2), we have

$$L(u'(t)) = -L(u(t-1)) - L(u^2(t)), \quad (10)$$

whence

$$L(u) (s+e^s) = c - \int_0^1 ce^{-st} dt - L(u^2). \quad (11)$$

Hence, we compute u_0 by means of

$$L(u_0) = \frac{c - \int_0^1 ce^{-st} dt}{s + e^{-s}} \quad (12)$$

and then u_1 from

$$L(u_1) = \frac{c - \int_0^1 ce^{-st} dt - L(u_0^2)}{(s+e^{-s})} \quad (13)$$

We see that despite the great increase in complexity of a differential-difference equation over an ordinary differential equation, the computational time and effort using this procedure is virtually the same for both types of equations. Once again, we have convergence of this iteration procedure only if $|c|$ is sufficiently small.

V. NUMERICAL RESULTS

Fifteen seconds of execution time are required for ten successive approximations. Graphs of the numerical results are given in Fig. 3. For some reasons which we have not as yet understood, all of the values obtained following the procedure given above are in excellent agreement with the actual solution, except for the very last value of t . This will be investigated at a later time.

VI. HEAT EQUATION

Consider next the nonlinear heat equation

$$\begin{aligned} k(x)u_t &= u_{xx} + b g(u), \\ u(0,t) &= u(1,t) = 0, \\ u(x,0) &= h(x). \end{aligned} \tag{14}$$

Using the Laplace transform, we readily obtain

$$L(u)'' - s k(x) L(u) = - k(x)h(x) - b L(g(u)), \tag{15}$$

with the two-point boundary condition

$$L(u) = 0, \quad x = 0,1. \quad (16)$$

Here ' denotes d/dx .

As above, we employ successive approximations. Let u_0 be the solution of

$$\begin{aligned} L(u_0)'' - s k(x)L(u_0) &= -k(x)h(x), \\ L(u_0) &= 0, \quad x = 0,1. \end{aligned} \quad (17)$$

First we solve the differential equation numerically for a set of s -values, $s = 1, 2, \dots, N$, and then we use the Laplace inversion technique. Storing the values $\{u_0(x, r_1)\}$ for $x = 0, \Delta, 2\Delta, \dots, M\Delta=1$, we then solve the equation

$$\begin{aligned} L(u_1)'' - s k(x)L(u_1) &= -k(x)h(x) - b L(g(u_0)), \\ L(u_1) &= 0, \quad x = 0,1. \end{aligned} \quad (18)$$

Continuing in this way, we obtain a sequence of values $\{u_n(x, r_1)\}$.

VII. NUMERICAL RESULTS

Five successive approximations require 6 min 10 sec of IBM 7090 execution time, and the numerical agreement was quite satisfactory. Results of this method are shown in Figs. 4 and 5.

VIII. CONVERGENCE ASPECTS

From the standpoint of rapidity of convergence, we would prefer to use quasilinearization procedures (cf. Refs. 4-6). For example, in treating the differential equation of Section II, we would prefer to use successive approximations in the form

$$L(u'_n) = -L(u_n) - L(u_{n-1}^2 + 2u_{n-1}(u_n - u_{n-1})). \quad (19)$$

This, however, complicates the matrix inversion aspects. Consequently, we feel that it is better to use the simpler, but less efficient approximation technique.

There is also the problem that the initial function, the solution of the linear equation, may have too large a norm for convergence. Provided that the solution of the nonlinear problem exists, we can often use extrapolation techniques (cf. Ref. 7).

For example, we can replace

$$k(x)u_t = u_{xx} + u^2 \quad (20)$$

by

$$k(x)u_t = u_{xx} + bu^2, \quad (21)$$

and use the new procedures for small b . The solution $u(x,t) = u(x,t,b)$ can then be determined for larger values of b by using extrapolation formulas.




Fig. 1




Fig. 2




Fig. 3




Fig. 4




Fig. 5

```

*      XEQ
*      LIST
*      LABEL
CNONLIN
C      NON LINEAR DIFFERENTIAL EQUATION
D      DIMENSION R(15),W(15),A(15,15),T(15),U(15),B(15)
      10 FORMAT(I12)
      11 FORMAT(1H1,4X11HDIMENSION =,I3)
      13 FORMAT(/5X32HROOTS / (5X6E20.8))
      15 FORMAT(/5X32HWEIGHTS / (5X6E20.8))
      16 FORMAT(/5X16HEXPlicit INVERSE//)
      17 FORMAT(5X6E20.8)
      101 FORMAT(/5X32HT = -LOG R / (5X6E20.8))
      102 FORMAT(E12.8)
      103 FORMAT(/10X4HC = ,F5.2)
      32 FORMAT(/5X32HLAPLACE TRANSFORM / (5X6E20.8))
      21 FORMAT(/5X32HINITIAL APPROXIMATION OF U(I) / (5X6E20.8))
      42 FORMAT (/5X23HVALUE OF U(I) AT STAGE , I2 / (5X6E20.8))

C
C      I INPUT
C
      READ 10, NA
      READ 10,N
      PRINT 11,N
      DO 12 I=1,N
D      CALL DBLRED(X)
D      12 R(I)=X
      PRINT 13, (R(I),I=1,N)
      DO 14 I=1,N
D      CALL DBLRED(X)
D      14 W(I)=X
      PRINT 15, (W(I),I=1,N)
D      CALL EXPINV (N,R,W,A)
      PRINT 16
      DO 18 I=1,N
      18 PRINT 17, (A(I,J),J=1,N)
      DO 19 I=1,N
D      19 T(I)=-LOGF(R(I))
      PRINT 101, (T(I),I=1,N)
      100 READ 102, C
      PRINT 103, C

C
C      II INITIAL APPROXIMATION
C
      DO 20 I=1,N
D      20 U(I)=C*R(I)
      PRINT 21, (U(I),I=1,N)

C
C      III TRANSFORM
C
      DO 30 K=1,NA
      DO 31 I=1,N
D      S=I

```


REFERENCES

1. Bellman, R. E., R. E. Kalaba, and Jo Ann Lockett, Numerical Solution of Functional Equations by Means of Laplace Transform - I: Renewal Equation, The RAND Corporation, RM-3948-NIH, December 1963.
2. Bellman, R. E., R. E. Kalaba, and Jo Ann Lockett, Numerical Solution of Functional Equations by Means of Laplace Transform - II: Differential-Difference Equations, The RAND Corporation, RM-3952-NIH, December 1963.
3. Bellman, R. E., R. E. Kalaba, and Jo Ann Lockett, Numerical Solution of Functional Equations by Means of Laplace Transform - III: The Diffusion Equation, The RAND Corporation, RM-3974-NIH, January 1964.
4. Bellman, R. E., Adaptive Control Processes, Princeton University Press, Princeton, New Jersey, 1961.
5. Kalaba, R. E., "On Nonlinear Differential Equations, the Maximum Operation and Monotone Convergence," J. of Math. and Mech., Vol. 8, 1959, pp. 519-574.
6. Bellman, R. E., and R. E. Kalaba, Quasilinearization and Boundary Value Problems, American Elsevier Publishing Co., New York, 1964.
7. Bellman, R. E., and R. E. Kalaba, "A Note on Nonlinear Summability Techniques in Invariant Imbedding," J. of Math. Anal. and Applications, Vol. 6, 1963, pp. 465-472.

```
B(I)=C/(S+1.)  
DO 31 J=1,N  
31 B(I)=B(I)-(W(J)*R(J)**(I-1)*U(J)**2)/(S+1.)  
PRINT 32, (B(I),I=1,N)
```

IV NEW APPROXIMATION

```
DO 41 I=1,N  
U(I)=0.0  
DO 41 J=1,N  
41 U(I)=U(I)+A(I,J)*B(J)  
PRINT 42, K, (U(I),I=1,N)  
30 CONTINUE  
GO TO 100  
END
```