

CLEARINGHOUSE FOR FEDERAL SCIENTIFIC AND TECHNICAL INFORMATION CFSTI
DOCUMENT MANAGEMENT BRANCH 410.11

LIMITATIONS IN REPRODUCTION QUALITY

ACCESSION # AD605075

- 1. WE REGRET THAT LEGIBILITY OF THIS DOCUMENT IS IN PART UNSATISFACTORY. REPRODUCTION HAS BEEN MADE FROM BEST AVAILABLE COPY.
- 2. A PORTION OF THE ORIGINAL DOCUMENT CONTAINS FINE DETAIL WHICH MAY MAKE READING OF PHOTOCOPY DIFFICULT.
- 3. THE ORIGINAL DOCUMENT CONTAINS COLOR, BUT DISTRIBUTION COPIES ARE AVAILABLE IN BLACK-AND-WHITE REPRODUCTION ONLY.
- 4. THE INITIAL DISTRIBUTION COPIES CONTAIN COLOR WHICH WILL BE SHOWN IN BLACK-AND-WHITE WHEN IT IS NECESSARY TO REPRINT.
- 5. LIMITED SUPPLY ON HAND: WHEN EXHAUSTED, DOCUMENT WILL BE AVAILABLE IN MICROFICHE ONLY.
- 6. LIMITED SUPPLY ON HAND: WHEN EXHAUSTED DOCUMENT WILL NOT BE AVAILABLE.
- 7. DOCUMENT IS AVAILABLE IN MICROFICHE ONLY.
- 8. DOCUMENT AVAILABLE ON LOAN FROM CFSTI (TT DOCUMENTS ONLY).
- 9.

NBS 9/64

PROCESSOR: *pm*

605 075

T
N

x

**Best
Available
Copy**

560509

605075

NOTES ON MATRIX THEORY—XIII:

SLIGHTLY INTERTWINED
LINEAR PROGRAMMING MATRICES

Richard Bellman

✓
P-918 *ptt*

Revised
January 23, 1957

Approved for OIS release

COPY _____ OF 6 *ptt*

HARD COPY	\$. 1.00
MICROFICHE	\$. 0.50

The **RAND** Corporation 1964
 1700 MAIN ST. • SANTA MONICA • CALIFORNIA

DDC RECEIVED
RECEIVED
 DDC-IRA E

SUMMARY

In this paper the functional-equation approach of dynamic programming is used to treat a linear programming problem involving a "slightly intertwined" matrix—i.e., one that is almost block diagonal.

SLIGHTLY INTERTWINED LINEAR PROGRAMMING MATRICES

1. INTRODUCTION

Consider the problem of maximizing the linear form

$$(1) \quad L_N(x) = \sum_{i=1}^{3N} x_i$$

over all x_i satisfying the constraints

$$(2) \quad a_{11}x_1 + a_{12}x_2 + a_{13}x_3 \leq c_1,$$

$$a_{21}x_1 + a_{22}x_2 + a_{23}x_3 \leq c_2,$$

$$a_{31}x_1 + a_{32}x_2 + a_{33}x_3 + b_1x_4 \leq c_3,$$

$$a_{44}x_4 + a_{45}x_5 + a_{46}x_6 \leq c_4,$$

$$a_{54}x_4 + a_{55}x_5 + a_{56}x_6 \leq c_5,$$

$$a_{64}x_4 + a_{65}x_5 + a_{66}x_6 + b_2x_7 \leq c_6,$$

⋮

$$a_{3N-2, 3N-2}x_{3N-2} + a_{3N-2, 3N-1}x_{3N-1} + a_{3N-2, 3N}x_{3N} \leq c_{3N-2},$$

$$a_{3N-1, 3N-2}x_{3N-2} + a_{3N-1, 3N-1}x_{3N-1} + a_{3N-1, 3N}x_{3N} \leq c_{3N-1},$$

$$a_{3N, 3N-2}x_{3N-2} + a_{3N, 3N-1}x_{3N-1} + a_{3N, 3N}x_{3N} \leq c_{3N},$$

and

$$(3) \quad x_i \geq 0.$$

It is assumed throughout that $a_{ij} \geq 0$, $b_i > 0$, $c_i \geq 0$, with a sufficient set of the a_{ij} positive so that the maximum of $L_N(x)$ is not infinite.

We wish to attack this problem—which arises from the study of weakly coupled economic systems, or alternatively from the study of multistage processes with almost-independent stages—by means of the techniques of dynamic programming. Specifically, we shall show that the computational solution can be obtained by means of a sequence of functions of one variable.

It will be clear that a number of similar problems involving almost block-diagonal matrices can be treated by means of the same general method. In another paper, [2], we have illustrated the application of this idea to mechanical and electrical systems in which the matrices are symmetric.

2. DYNAMIC PROGRAMMING FORMULATION

Let us define the sequence of functions of z ,

$$(1) \quad f_0(z) = 0, \quad f_K(z) = \text{Max}_{x_1} L_K(x), \quad K = 1, 2, \dots, N,$$

where the x_1 are subject to the constraints given above, with the exception that the last constraint is now

$$(2) \quad a_{3K, 3K-2} x_{3K-2} + a_{3K, 3K-1} x_{3K-1} + a_{3K, 3K} x_{3K} \leq z.$$

Employing the principle of optimality, cf. [1], we see that the sequence $\{f_K(z)\}$ satisfies the recurrence relation

$$(3) \quad f_K(z) = \text{Max}_{[x_{3K-2}, x_{3K-1}, x_{3K}]} \left[x_{3K-2} + x_{3K-1} + x_{3K} + f_{K-1} \right. \\ \left. (c_{3K-3} - b_{K-1} x_{3K-2}) \right], \quad K \geq 1,$$

with the variables $x_{3K-2}, x_{3K-1}, x_{3K}$ subject to the constraints

$$\begin{aligned}
 (4) \quad & a_{3K-2, 3K-2} x_{3K-2} + a_{3K-2, 3K-1} x_{3K-1} + a_{3K-2, 3K} x_{3K} \leq c_{3K-2}, \\
 & a_{3K-1, 3K-2} x_{3K-2} + a_{3K-1, 3K-1} x_{3K-1} + a_{3K-1, 3K} x_{3K} \leq c_{3K-1}, \\
 & a_{3K, 3K-2} x_{3K-2} + a_{3K, 3K-1} x_{3K-1} + a_{3K, 3K} x_{3K} \leq z, \\
 & b_{K-1} x_{3K-2} \leq c_{3K-3}, \\
 & x_{3K-2}, x_{3K-1}, x_{3K} \geq 0.
 \end{aligned}$$

3. SIMPLIFICATION

We can write the recurrence relation of (2.3) in the form

$$\begin{aligned}
 (1) \quad f_K(z) &= \text{Max}_{x_{3K-2}} \left[\text{Max}_{x_{3K-1}, x_{3K}} [\dots] \right] \\
 &= \text{Max}_{x_{3K-2}} \left[\text{Max}_{R_K} (x_{3K-1} + x_{3K}) + x_{3K-2} \right. \\
 &\quad \left. + f_{K-1}(c_{3K-3} - b_{K-1} x_{3K-2}) \right],
 \end{aligned}$$

where R_K is the region in (x_{3K-1}, x_{3K}) space defined by

$$\begin{aligned}
 (2) \quad & a_{3K-2, 3K-1} x_{3K-1} + a_{3K-2, 3K} x_{3K} \leq c_{3K-2} - a_{3K-2, 3K-2} x_{3K-2}, \\
 & a_{3K-1, 3K-1} x_{3K-1} + a_{3K-1, 3K} x_{3K} \leq c_{3K-1} - a_{3K-1, 3K-2} x_{3K-2}, \\
 & a_{3K, 3K-1} x_{3K-1} + a_{3K, 3K} x_{3K} \leq z - a_{3K, 3K-2} x_{3K-2}, \\
 & x_{3K-1}, x_{3K} \geq 0.
 \end{aligned}$$

Thus we can write

$$(3) f_K(z) = \text{Max}_{x_{3K-2}} \left[g_K(x_{3K-2}, z) + f_{K-1}(c_{3K-3} - b_{K-1}x_{3K-2}) \right],$$

where x_{3K-2} is constrained by

$$(4) 0 \leq x_{3K-2} \leq \text{Min} \left[\frac{c_{3K-3}}{b_{K-1}}, \frac{c_{3K-2}}{a_{3K-2, 3K-2}}, \frac{c_{3K-1}}{a_{3K-1, 3K-2}}, \frac{z}{a_{3K, 3K-2}} \right]$$

The foregoing function $g_K(y, z)$ is readily determined, since the maximum over R_K is attained at a vertex of the region; indeed, for the present problem at most four such vertices need to be considered for any particular assignment of the parameters y, z .

REFERENCES

1. Bellman, R., "The Theory of Dynamic Programming," Bull. Amer. Math. Soc., Vol. 60, 1954, pp. 503-516.
2. _____, Notes on Matrix Theory-XII: Slightly Intertwined Matrices, The RAND Corporation, Paper P-917, 1956.