

AD 607027

1 (1)
for


FUNCTIONAL EQUATIONS AND MAXIMUM RANGE

Richard Bellman

✓
P-1494

September 24, 1958

Approved for OTS release



COPY	1	OF	DDC-IRA G
HARD COPY	\$. 1 00		
MICROFICHE	\$. 0.50		

The RAND Corporation

1700 MAIN STREET, SANTA MONICA, CALIFORNIA

ARCHIVE COPY

H

Summary

The current interest in rockets and space travel has aroused a corresponding interest in the determination of maximum range, minimum time, and so on, for various types of trajectories.

A variety of questions of this type have been treated by means of the theory of dynamic programming. Here we wish to show how to use functional equations to determine the range, the maximum elevation, and similar quantities, as functions of initial position and velocities.

FUNCTIONAL EQUATIONS AND MAXIMUM RANGE

Richard Bellman

1. Introduction

The current interest in rockets and space travel has aroused a corresponding interest in the determination of maximum range, minimum time, and so on, for various types of trajectories.

A variety of questions of this type have been treated by means of the theory of dynamic programming, see [1,2,4]. Here we wish to show how to use functional equations to determine the range, the maximum elevation, and similar quantities, as functions of initial position and velocities.

2. Vertical Motion--I

Consider an object, subject only to the force of gravity and the resistance of the air, which is propelled straight up. In order to illustrate the technique we shall employ, let us treat the problem of determining the maximum altitude.

Let the defining equation be

$$(1) \quad u'' = -g - h(u'),$$

with the initial conditions $u(0) = 0$, $u'(0) = v$. Here $v > 0$, and $h(u') \geq 0$ for all u' .

Since the maximum altitude is a function of v , let us introduce the function

- (2) $f(v)$ = the maximum altitude attained starting
with initial velocity v .

From the definition of the function it follows that

$$(3) \quad f(v) = v\Delta + f(v - [g + h(v)]\Delta) + o(\Delta),$$

for Δ an infinitesimal. Verbally, this states that the maximum altitude is the altitude gained over an initial time Δ , plus the maximum altitude attained starting with a velocity $v - [g + h(v)]\Delta$, the velocity of the object at the end of time Δ , to within $o(\Delta)$.

Expanding both sides and letting $\Delta \rightarrow 0$, we see that

$$(4) \quad f'(v) = \frac{v}{g + h(v)}.$$

Since $f(0) = 0$, this yields

$$(5) \quad f(v) = \int_0^v \frac{v_1 dv_1}{g + h(v_1)}.$$

In the particular case where $h(v) = 0$, we obtain the standard result $v^2/2g$.

3. Vertical Motion--II

Consider the more general case where motion is through an inhomogeneous medium. Let the defining equation be

$$(1) \quad u'' = h(u, u'), \quad u(0) = c_1, \quad u'(0) = c_2.$$

Assume that $h(u, u') \leq 0$ for all u and u' , so that $c_2 = 0$ implies no motion.

The maximum altitude is now a function of both c_1 and c_2 . Introduce

(2) $f(c_1, c_2)$ = the maximum altitude attained starting with the initial position c_1 and initial velocity c_2 .

Then, as above,

$$(3) \quad f(c_1, c_2) = c_2 \Delta + f(c_1 + c_2 \Delta, c_2 + h(c_1, c_2) \Delta) + o(\Delta),$$

which yields in the limit the partial differential equation

$$(4) \quad c_2 + c_2 \frac{\partial f}{\partial c_1} + h(c_1, c_2) \frac{\partial f}{\partial c_2} = 0.$$

By virtue of our assumptions, $f(c_1, 0) \equiv 0$, for $c_1 \geq 0$.

4. Computational Aspects

One can, of course, use the method of characteristics, or standard difference methods, to solve (3.4). Let us present another method which reduces the solution to the tabulation of a sequence of functions of one variable.

In place of (3.4), let us use the discrete approximation of (3.3),

$$(1) \quad f(c_1, c_2) = c_2 \Delta + f(c_1 + c_2 \Delta, c_2 + h(c_1, c_2) \Delta).$$

Since c_2 is monotone decreasing, it can be used to play the role of time. Let us write $c_2 = N\delta$, where δ is a positive quantity, and $f(c_1, c_2) = f_N(c_1)$. We consider then only values of c_2 which are multiples of δ . To overcome the fact that $c_2 + h(c_1, c_2)\Delta$ in general will not be a multiple of δ , we can either replace it by $[(c_2 + h(c_1, c_2)\Delta)/\delta]$, or use interpolation. Although use of an interpolation formula slows up the computation, it greatly improves the accuracy. For an application of the foregoing techniques to a more complicated partial differential equation, see [3].

5. Maximum Altitude

Consider now the case where motion takes place in a plane. Let the equations be

$$(1) \quad x'' = g(x', y'), \quad x(0) = 0, \quad x'(0) = c_1,$$

$$y'' = h(x', y'), \quad y(0) = 0, \quad y'(0) = c_2.$$

Introducing, as before, the function $f(c_1, c_2)$ equal to the maximum altitude, we see that

$$(2) \quad f(c_1, c_2) = (c_1^2 + c_2^2)^{1/2} \Delta + f(c_1 + g(c_1, c_2)\Delta, c_2 + h(c_1, c_2)\Delta) + o(\Delta).$$

Hence,

$$(3) \quad (c_1^2 + c_2^2)^{1/2} + g(c_1, c_2) \frac{\partial f}{\partial c_1} + h(c_1, c_2) \frac{\partial f}{\partial c_2} = 0.$$

Once again, let us assume that $c_2 = 0$ implies no vertical motion. Then $f(c_1, 0) = 0$ for $c_1 \geq 0$. It follows that we can again compute the solution by means of a sequence of functions of one variable.

6. Maximum Range

To tackle the problem of maximum range directly requires the introduction of another state variable, the initial altitude. It can also be broken up into two problems, corresponding to the ascent to maximum altitude, and the descent.

References

1. Bellman, R., Dynamic Programming, Princeton University Press, Princeton, New Jersey, 1957.
2. Bellman, R., and S. Dreyfus, An Application of Dynamic Programming to the Determination of Optimal Satellite Trajectories, The RAND Corporation, Paper P-1463, August 19, 1958.
3. Bellman, R., I. Cherry, and G. M. Wing, "A note on the numerical integration of a class of nonlinear hyperbolic equations," Q. Appl. Math., vol. 16, 1958, pp. 181-183.
4. Cartaino, T., and S. Dreyfus, "Application of dynamic programming to the airplane minimum time-to-climb problem," Aeronautical Engineering Review, vol. 16, 1957, pp. 74-77.