

**CLEARINGHOUSE FOR FEDERAL SCIENTIFIC AND TECHNICAL INFORMATION, CFSTI
INPUT SECTION 410.11**

**LIMITATIONS IN REPRODUCTION QUALITY OF TECHNICAL ABSTRACT BULLETIN
DOCUMENTS, DEFENSE DOCUMENTATION CENTER (DDC)**

AD 613524

- 1. AVAILABLE ONLY FOR REFERENCE USE AT DDC FIELD SERVICES. COPY IS NOT AVAILABLE FOR PUBLIC SALE.
- 2. AVAILABLE COPY WILL NOT PERMIT FULLY LEGIBLE REPRODUCTION. REPRODUCTION WILL BE MADE IF REQUESTED BY USERS OF DDC.
 - A. COPY IS AVAILABLE FOR PUBLIC SALE.
 - B. COPY IS NOT AVAILABLE FOR PUBLIC SALE.
- 3. LIMITED NUMBER OF COPIES CONTAINING COLOR OTHER THAN BLACK AND WHITE ARE AVAILABLE UNTIL STOCK IS EXHAUSTED. REPRODUCTIONS WILL BE MADE IN BLACK AND WHITE ONLY.

TSL-121-2/64

DATE PROCESSED:

PROCESSOR:

4-19-65
B. Lee

AD 613524

Technical Report No. 51

to the

Office of Naval Research

ARPA Order No. 26-60, Task 2, Item 3

Contract No.: Nonr 477 (16)

PREPARATION AND PROPERTIES OF BISTRIFLUOROMETHYLSULFURYL PEROXIDE AND TRIFLUOROMETHYL TRIFLUOROMETHANE SULFONATE

by

Ronald E. Noftle and George H. Cady

Department of Chemistry

University of Washington

Seattle, Washington

98105

1965

DDC

APR 13 1965

Thom B

13-p
COPY 2 OF 3
HARD COPY \$.
MICROFICHE \$.
Reference only

Note: The junior author (R.E.N.) was employed under Contract Nonr 477 (16) prior to September 1964 and will resume such employment in May 1965. During the period while the work described in this report was in progress, he held an Allied Chemical Co. Fellowship. Since his future work under the contract will extend that covered here, it is desirable that this paper be distributed as a technical report.

ARCHIVE COPY

Reproduction in whole or in part is permitted for any purpose United States Government.

Contribution From the Department of Chemistry
University of Washington, Seattle, Washington

PREPARATION AND PROPERTIES OF BISTRIFLUOROMETHYLSULFURYL
PEROXIDE AND TRIFLUOROMETHYL TRIFLUOROMETHANESULFONATE

By Ronald E. Nofle and George H. Cady

Electrolysis of trifluoromethanesulfonic acid at -23° yields the unstable compound bistrifluoromethylsulfuryl peroxide which decomposes explosively to perfluoroethane, sulfur trioxide, and the ester, trifluoromethyl trifluoromethanesulfonate.

The synthesis of peroxydisulfuryl difluoride, $S_2O_6F_2$, by electrolysis of fluorosulfuric acid¹ suggests that other new

(1) F. B. Dudley, J. Chem. Soc., 3407 (1963).

peroxides may be prepared by electrolysis of their corresponding strong acids. When trifluoromethanesulfonic acid, CF_3SO_3H , was electrolyzed under high vacuum at about -23° in a cell similar to that described by Dudley¹, the products obtained at the anode included bistrifluoromethylsulfuryl peroxide, $CF_3SO_2OOSO_2CF_3$. As a cold sample of this liquid compound was allowed to warm up, it decomposed suddenly* and with evolution of heat when its

* Although the authors experienced no violent or destructive explosions due to the decomposition of $(CF_3SO_3)_2$ during the course of this work (possibly due to the fact that only small samples were prepared), larger samples might be expected to explode with considerable force, and caution should be exercised when working with this material.

temperature reached about 10° . The products formed were perfluoroethane, sulfur trioxide, and the new ester, trifluoromethyl trifluoromethanesulfonate, $CF_3SO_3CF_3$. Although esters of trifluoromethanesulfonic acid containing hydrocarbon^{2,3,4} and polyfluoro hydrocarbon⁵ groups have been reported, this ester appears to be

(2) T. Gramstad and R. N. Haszeldine, *ibid*, 173 (1956).


(3) T. J. Erice and P. W. Trott, U. S. Patent 2,732,398 (1956).

(4) T. Gramstad and R. N. Haszeldine, *ibid*, 4069 (1957).

(5) J. Burdon, *Tetrahedron*, 1 (1965).

the first instance of a perfluorocarbon trifluoromethanesulfonate. Trifluoromethyl trifluoromethanesulfonate is resistant to hydrolysis by water but does hydrolyze at 100° in 0.1N sodium hydroxide.

The products observed from the decomposition of $(CF_3SO_3)_2$ can be explained on the basis of the following hypothetical mechanism:



Recently Haszeldine, Heslop and Lethbridge have reported the synthesis of the peroxide $(CF_3SO_3)_2$ by the electrolysis of CH_3SO_3H .⁶

(6) R. N. Haszeldine, R. B. Heslop and J. W. Lethbridge, J. Chem. Soc., 4901 (1964).

EXPERIMENTAL

Materials.— Trifluoromethanesulfonic acid was prepared by distillation from a mixture of barium trifluoromethanesulfonate⁷ and concentrated sulfuric acid.⁸ The product was redistilled and

(7) Gift from the Minnesota Mining and Manufacturing Co.

(8) J. M. Kidd and R. N. Haszeldine, J. Chem. Soc., 4228 (1956).

the fraction boiling near 56° under 15 mm pressure was used.

Electrolysis.— The electrolysis cell employed for this work was similar in design to that described previously by Dudley.¹

It was a small glass cell which could be cooled in a Dewar vessel and operated under vacuum with separate collection of gaseous or volatile products from the anode and cathode. The cathode was a 1 cm. square piece of platinum foil, and the anode consisted of two 26 gauge platinum wires of 3 cm. length wound about each other. Since the conductivity of the pure trifluoromethanesulfonic acid was very low, a 0.1 molal solution of sodium trifluoromethanesulfonate in trifluoromethanesulfonic acid was used. This was prepared by adding 0.202 g. of dry sodium chloride and 33.1 g. of trifluoromethanesulfonic acid to the cell, care being taken to exclude moisture from the system. The hydrogen chloride produced was pumped away. The cell was held at about -25°C . (to minimize transfer of the acid under vacuum) by a carbon tetrachloride slush bath, and the traps for collecting products were held at the boiling point of oxygen. When 36 volts was applied to the cell, a current of 0.014 amp. passed. A copper coulometer was used to measure the total charge passed. While the cell was operating, a non-condensable gas was evolved at the cathode, and mass spectral analysis showed it to be hydrogen. No oxygen could be detected as a product at the anode. After about 10 hours of steady operation, a blue color, also observed by Dudley, developed in the liquid occupying the cathode compartment of the cell. Liquid in the anode compartment remained colorless. In some runs the material in the anode trap was allowed to warm to room temperature while transferring the vapor to another trap at -183°C . When the mixture of materials including trifluoromethanesulfonic acid approached about 10° , a sudden evolution of gas and heat was observed. The volatile products were separated by fractional

co-distillation⁹ and each cut was analyzed by infra-red spectroscopy.

(9) G. H. Cady and P. P. Siegarth, Anal. Chem., 31, 618 (1959).

The products were C_2F_6 , $(CF_3)_2SO_3$, SO_3 , and small amounts of COF_2 , SO_2 , trifluoromethanesulfonic anhydride,^{3,4,10} and tri-

(10) J. Burdon, I. Farazmand, M. Stacey and J. C. Tatlow, J. Chem. Soc., 2574 (1957).

fluoromethanesulfonic acid. At the cathode, the same products were observed by the same method of analysis, but were present in much lower yield except for SO_2 and CF_3SO_3H . Hydrogen was also liberated at the cathode but was not collected. It appeared to be the principal substance escaping from the cathode chamber of the cell. Assuming a two electron charge to produce $(CF_3SO_3)_2$ at the anode, the current efficiency for a selected run was about 60%. This value was determined by measuring the total charge passed ($0.91 \times 10^{-2}F$) during electrolysis of a sample by means of a copper coulometer. The volatile products were collected, separated and weighed. The total number of millimoles of $(CF_3)_2SO_3$ (1.81) plus the total number of millimoles of C_2F_6 (0.89) were taken to equal the number of millimoles of $(CF_3SO_3)_2$ (2.70) originally produced.

Bistrifluoromethylsulfuryl peroxide.— Since the peroxide, $(CF_3SO_3)_2$, decomposed easily and was difficult to separate completely from CF_3SO_3H , no pure sample was obtained. In some runs decomposition of the material collected in the anode trap was avoided by handling it when in the solid or liquid state at

temperatures below 0°. In spite of these difficulties, a chemical analysis was made in the manner now to be described: Before electrolysis was begun, 3 ml. of a 67% solution of sulfuric acid in water was placed in the anode trap to absorb SO₃ and CF₃SO₃II. After electrolyzing the acid under vacuum for 10 hours, the anode trap was held at about -70 and pumped to remove C₂F₆ and (CF₃)₂SO₃ as well as other volatile products. The trap was then warmed slowly to -10° and the product volatile at this temperature was pumped through a length of connecting glass tubing of 22 cm. into a weighed bulb held at -183°. The bulb was then closed and allowed to warm slowly. The small amount of liquid present was observed to explode mildly when the temperature of the bulb reached about 10°. The bulb was again weighed, (net wt. 0.3745g. of product) the material was cooled to -183° (no O₂ was present),* and 5 ml.

* On one occasion O₂ was observed as a product when a very impure sample of (CF₃SO₃)₂ decomposed. This result was not observed in subsequent experiments.

of distilled water was added. The volatile C₂F₆ and (CF₃)₂SO₃ were pumped off, separated from any water and each weighed after separation by trap to trap distillation. The solution left behind was analyzed for SO₄[•] by precipitation as BaSO₄. The acid equivalents present were determined by titration with standard base. The analysis indicated a sample before decomposition to consist of 0.3332 g. of (CF₃SO₃)₂ and 0.0413 g. of CF₃SO₃II.

	Calcd. for above mixture	Found
Wt. $(CF_3)_2SO_3$	0.1648*	0.1648
Wt. C_2F_6	0.0500*	0.0500
Wt. SO_3	0.1184*	0.1204
total g. equiv. of acid	3.24	3.22

* Various proportions of these three substances could result from the decomposition of $(CF_3SO_3)_2$. The set given here was chosen to correspond to the observed amounts of C_2F_6 and $(CF_3)_2SO_3$ and from this the amount of SO_3 was calculated.

While this analysis shows the explosive material to be $(CF_3SO_3)_n$, it does not show n to equal 2. The conclusion that n equals 2 comes from the fact that the substance is colorless, apparently not a free radical, and of the correct volatility. By comparison with other similar fluorides including SO_3F , $(SO_3F)_2$, SO_2F_2 , $S_2O_5F_2$, $S_3O_8F_2$, CF_3SO_3F , $(CF_3SO_2)_2O$ etc., one would expect $(CF_3SO_3)_2$ to boil at about 100° while CF_3SO_3 would boil much lower than this and the hypothetical $(CF_3SO_3)_3$ or $(CF_3SO_3)_4$ considerably higher. Since the material in question has approximately the same volatility as $(CF_3SO_2)_2O$ (b.p. 84°),^{3,4,10} it is reasonable to conclude that n = 2. The fact that the substance gives a good n.m.r. spectrum showing all fluorine atoms to be equivalent supports this conclusion.

Trifluoromethyl trifluoromethanesulfonate.— A purified weighed sample of $(CF_3)_2SO_3$ was fused with potassium overnight at 450° . The excess potassium was destroyed with a small amount of ethanol, and the aqueous solution analyzed for sulfide by

adding an aliquot to an acidic standard potassium iodate - potassium iodide mixture. The excess iodine was back titrated with standard sodium thiosulfate solution. Fluoride was determined by steam distillation and subsequent titration with standard thorium nitrate solution. Anal. Calcd. for $(CF_3)_2SO_3$: S, 14.70; F, 52.27. Found: S, 15.1; F, 51.8.

The molecular weight obtained from vapor density measurements assuming ideal gas behavior was 217.8 ± 0.8 g./g.m.v. (avg. 4 determinations) Calcd: 218.1 g./g.m.v.

The melting point was determined by freezing a portion of a sample held in a 17 mm. o.d. glass tube. The tube was allowed to warm in an n-butyl chloride bath at the rate of $0.4^\circ/\text{min.}$, the temperature being measured by means of a calibrated thermocouple. Four determinations were made, each on a fresh portion of the sample. The solid was observed to start to melt at about -109° . The melting point was close to -108.2° .

Vapor pressures of $(CF_3)_2SO_3$ were measured in Pyrex glass apparatus similar to that described by Kellogg and Cady.¹¹ In

(11) K. B. Kellogg and G. H. Cady, J. Am. Chem. Soc., 70, 3986 (1948).

this method the liquid was allowed to boil under various constant and measured pressures while the vapor was condensed in a reflux condenser at -78° and returned to the pot as a stream of liquid. The temperature corresponding to the equilibrium between liquid and vapor was measured by a copper-constantan thermocouple located about 10 cm. above the boiling liquid and 8 cm. below the condenser.

Vapor pressures are shown in Table I. The approximate molar heat of vaporization at the boiling point was calculated from the slope of the vapor pressure curve with the aid of the Clausius-Clayron equation. The boiling point was estimated to be 21.1° by extrapolation of the vapor pressure curve. These values lead to a Trouton Constant of 22.7 e.u.

TABLE I
Vapor Pressures of $(CF_3)_2SO_3$

<u>P, mm.</u>	<u>T, °K.</u>	<u>P, mm.</u>	<u>T, °K.</u>	<u>P, mm.</u>	<u>T, °K.</u>
46	237.8	241	267.1	471	282.1
107	251.2	290	271.4	565	286.6
144	256.8	362	277.5	645	290.2
182	261.8	412	278.6	682	291.6
				732	293.5

Nuclear Magnetic Resonance Spectra.— Nuclear magnetic resonance spectra were recorded using a Varian Model 4311B high resolution spectrometer with a 40 mc. oscillator. Trichlorofluoromethane (Freon 11) was used as an external standard. Chemical shifts in ppm. relative to this standard were obtained for trifluoromethanesulfonic acid, +78.48 ppm., and trifluoromethanesulfonic anhydride, +74.02 ppm. The spectrum of trifluoromethyl trifluoromethanesulfonate first appeared as two separate peaks; one at +55.61 ppm. due to the CF_3 group attached to oxygen and one at +76.22 ppm. due to the CF_3 group attached to sulfur. Under high resolution these two peaks each split into a similar quartet having relative intensities 1:3:3:1. A spacing of 3.4 cycles/sec. in the sharp quartets indicated this to be the spin-spin coupling constant between the two types of fluorine atoms in the F-C-S-O-C-F bond and space system. The decomposition of the peroxide, $(CF_3SO_3)_2$, was followed by n.m.r. spectrometry. When an impure sample of

the peroxide which had been kept cold was scanned at low temperature in the C-F region, two large peaks were observed. The larger one at +77.19 ppm. could be ascribed to $\text{CF}_3\text{SO}_3\text{H}$ while the smaller one at +72.36 ppm. could be ascribed to $(\text{CF}_3\text{SO}_3)_2$. Two small peaks were also observed for $(\text{CF}_3)_2\text{SO}_3$ at +54.66 ppm. and +75.30 ppm. as well as a trace peak for the anhydride of trifluoromethanesulfonic acid at +74.05 ppm. When the sample was taken out of the spectrometer and allowed to warm to room temperature, a reaction occurred in the tube with the evolution of heat. When the sample was again scanned in the C-F region, the spectrogram revealed that the peroxide peak had disappeared, the ester peaks had grown, and a new peak at +89.06 ppm. (attributable to perfluoroethane) had appeared. A rerun of the same sample when cold did not produce the original spectrum, but the ester and perfluoroethane peaks were enlarged still further due to the gas phase being more highly dissolved in the liquid phase at lower temperatures.

Mass Spectrum.— A Consolidated Engineering Corporation Type 21-103 mass spectrometer was used to record the mass spectrogram for $(\text{CF}_3)_2\text{SO}_3$ which showed peaks corresponding to the following ions in order of decreasing intensity: CF_3^+ , SO^+ , SO_2^+ , CF^+ , CF_2^+ , COF^+ , S^+ , CF_3SO^+ , COF_2^+ , and CF_3SO_2^+ . In addition, small peaks which could have been due to CF_3O^+ , SO_3^+ , and CF_3SO_3^+ were observed at about the same low intensity level as some impurity peaks in the mass spectrometer background. No peak corresponding to the molecule-ion was noted.


Infrared Spectrum.— A Perkin Elmer Model 21 infrared spectrometer with a sodium chloride prism was used to obtain the

spectrum of $(CF_3)_2SO_3$ in the gas phase at 5 mm. pressure in a 10 cm. cell. Strong absorptions were found at 1461, 1258, 1230, 1134, and 954 $cm.^{-1}$. Two medium absorptions were noted at 736 and 766 $cm.^{-1}$. At 10 mm. pressure a weak band appeared at 858 $cm.^{-1}$. Some of these bands are close to those reported by Gramstad and Haszeldine for the S=O asymmetric vibration (1435 $cm.^{-1}$) and the S=O symmetric vibration (either 1227 or 1156 $cm.^{-1}$) in $CF_3SO_3CH_3$.²

Chemical Reactions.— An impure sample of $(CF_3SO_3)_2$ was held at -70° and pumped to remove volatile impurities. Then the sample was warmed until material began to distil and condense in a weighed bulb containing frozen KI solution at -183° . Iodine was liberated at once. The bulb was weighed again, and the iodine was titrated with standard sodium thiosulfate solution. The oxidizing equivalents found corresponded to about 20% of the distilled mixture being present as $(CF_3SO_3)_2$.

A sample of $(CF_3)_2SO_3$ was allowed to remain in contact with water at room temperature for 16 hours. The compound was immiscible and did not appear to react. Sample wt: 0.7496 g. Recovered: 0.7291 g. Infrared analysis showed the recovered material to be the unchanged ester.

A sample of $(CF_3)_2SO_3$ was held in contact with 0.1N. sodium hydroxide at 100° for about 12 hours (with occasional shaking). Then 5 ml. of a 5% $BaCl_2$ solution was added to precipitate barium carbonate. The excess base was titrated to the phenolphthalein end point and the milliequivalents of base used per milliequivalent of $(CF_3)_2SO_3$ calculated. Calcd. meq. base used per mM. $(CF_3)_2SO_3$: 6.0. Found: 5.7. Apparently the basic hydrolysis proceeded:



but did not go to completion under these conditions.

Acknowledgement.— This work was performed in part with the support of a fellowship from the Allied Chemical Company. The authors thank the Minnesota Mining and Manufacturing Company for furnishing the varium salt of trifluoromethanesulfonic acid. They also thank Mr. B. J. Nist for the N.M.R. spectra.