

AD617001

ON A CLASS OF NONLINEAR DIFFERENTIAL EQUATIONS
WITH NONUNIQUE SOLUTIONS

Richard Bellman

June 1965

COPY	OF	196
HARD COPY	\$.	1.00
MICROFICHE	\$.	0.50

DDC
JUL 2 1965
TISA B

P-3147

ARCHIVE COPY

ON A CLASS OF NONLINEAR DIFFERENTIAL EQUATIONS
WITH NONUNIQUE SOLUTIONS

Richard Bellman*

The RAND Corporation, Santa Monica, California

1. INTRODUCTION

As we know, nonlinear differential equations subject to initial-value conditions possess unique solutions, under reasonable assumptions, whereas the same equations subject to two-point or multipoint boundary-value conditions can easily possess an infinite number of solutions. The purpose of this note is to describe a class of nonlinear differential equations subject to conditions analogous but not equivalent to multipoint conditions which can possess an infinite number of solutions. These equations are suggested by some work in respiratory control theory which will be described elsewhere.

2. A SCALAR EXAMPLE

Consider the equation

$$(2.1) \quad \frac{du}{dt} = (a + bu(t_1))u(t), \quad u(0) = c,$$

* Any views expressed in this paper are those of the author. They should not be interpreted as reflecting the views of The RAND Corporation or the official opinion or policy of any of its governmental or private research sponsors. Papers are reproduced by The RAND Corporation as a courtesy to members of its staff.

This paper will be submitted for publication in the Bolletino dell'Unione Matematica Italiana.

valid for $0 \leq t \leq t_2$, where $t_2 \geq t_1$. Using the explicit form of the solution of (2.1),

$$(2.2) \quad u(t) = ce^{(a+bu(t_1))t},$$

we obtain the following transcendental equation for $u(t_1)$,

$$(2.3) \quad u(t_1) = ce^{at_1} e^{bt_1 u(t_1)}.$$

Depending upon the values of a , b , c , and t_1 , this equation can have no, one, or two real roots, and it always possesses an infinite set of complex roots. To each root corresponds a solution of (2.1).

3. DISCUSSION

It is easy to see that the corresponding situation will persist for the general vector differential equation

$$(3.1) \quad \frac{dx}{dt} = g(x(t), x(t_1), \dots, x(t_M)), \quad x(0) = c,$$

valid for $0 \leq t \leq t_0$, where $0 < t_1 < t_2 < \dots < t_M < t_0$.

An interesting question is that of determining what additional conditions single out a unique solution.