

USNRDL-TR-943

12 November 1965

AD627127

# $\text{Pd}^{115}$ YIELD IN THERMAL NEUTRON FISSION OF $\text{U}^{235}$

by

H. V. Weiss

W. L. Reichert\*

CLEARINGHOUSE FOR FEDERAL SCIENTIFIC AND TECHNICAL INFORMATION		
Hardcopy	Microfilm	
\$1.00	\$0.50	12.00
ARCHIVE COPY		

Code 1

DDC  
RECEIVED  
FEB 2 1966  
DDC-IRA F

U.S. NAVAL RADIOLOGICAL  
DEFENSE LABORATORY

SAN FRANCISCO • CALIFORNIA • 94135

NUCLEAR CHEMISTRY BRANCH  
N. E. Ballou, Head

CHEMICAL TECHNOLOGY DIVISION  
R. Cole, Head

---

ADMINISTRATIVE INFORMATION


This work is part of a project sponsored  
by the Atomic Energy Commission under Contract  
No. AT(49-2)-1167.

\* University of Washington, Seattle,  
Washington.

DDC AVAILABILITY NOTICE

Distribution of this document is unlimited.

  
Eugene P. Cooper  
Scientific Director

  
D.C. Campbell, CAPT USN  
Commanding Officer and Director

**BLANK PAGE**

# ABSTRACT

The formation of  $\text{Pd}^{115}$  in thermal neutron fission of  $\text{U}^{235}$  was studied quantitatively. A procedure was developed which rapidly separated palladium from daughter elements silver and cadmium. Separations were made from 2.8-120 seconds after fission. The growth of  $\text{Cd}^{115}$  from palladium separated at various times constituted a decay curve for  $\text{Pd}^{115}$ . The half-life thus determined was  $40.5^{+4.0}_{-3.0}$  seconds, a value which agrees with that previously reported. The cumulative fractional yield of  $\text{Pd}^{115}$  compared to the total  $\text{Cd}^{115}$  yield was  $90.5 \pm 7.2\%$ . Estimation of the most probable nuclear charge ( $Z_p$ ) must await information on the contribution of other members to the yield of the 115 mass chain.

## SUMMARY

### Problem

As part of a general study of nuclear charge distribution in the region of symmetric fission of uranium, information on the formation of a palladium isotope (mass 115) was required.

### Findings

A rapid radiochemical separation method was developed to study the formation of this palladium isotope. The half-life of this fission product was confirmed and its cumulative contribution to the mass 115 chain yield was determined.

## INTRODUCTION

The distribution of nuclear charge in near-symmetric thermal neutron fission of  $U^{235}$  is being studied. For this study, information is needed on the cumulative fission yield of  $Pd^{115}$ .

Alexander and co-workers<sup>1</sup> reported the formation of  $Pd^{115}$  upon bombardment of natural uranium with deuterons. Palladium was isolated free from silver and cadmium within about 3 minutes after fission. By the periodic extraction of silver decay products and purification and counting of 2.2-day  $Cd^{115}$  after decay of 21.1-minute  $Ag^{115}$ , a half-life of  $45 \pm 3$  seconds was determined for  $Pd^{115}$ .

The present work was aimed at quantitatively examining the formation of  $Pd^{115}$  as quickly after fission as possible. Speed was essential to reduce the probability of failure to observe the contribution to the yield by any as yet unidentified short-lived isomer.

A rapid method was developed for the separation of palladium from its descendants, silver and cadmium. The procedure was based upon the reduction and simultaneous removal from solution of palladium by copper powder.<sup>2</sup> The reduction of silver was inhibited by complexation with bromide; the oxidation potential of cadmium is unfavorable for its reduction by copper. The unreduced elements remained essentially quantitatively in solution.

With this method the quantity of descendant  $Cd^{115}$  was measured in palladium samples isolated at various times after irradiation. The curve which resulted from a plot of  $Cd^{115}$  activity as a function of separation time constitutes a decay curve for  $Pd^{115}$ . From this data together with a separate measurement of the total  $Cd^{115}$  yield, the cumulative fractional yield of  $Pd^{115}$  was computed. The difficulty of using this cumulative fractional yield to estimate the most probable nuclear charge ( $Z_p$ ) is described.

## EXPERIMENTAL

### Chemical Solutions

The weights given in the solutions listed below refer to the metallic element. All reagents used were of analytical grade.

Uranyl bromide was prepared from 93.17 % enriched  $U^{235}$  metal. The metal was dissolved in conc.  $HNO_3$ , and excess  $HNO_3$  was removed by several evaporations with conc.  $HBr$ . The dried salt was finally dissolved in conc.  $HBr$  at a concentration of 400 mg/ml.

Ruthenium, rhodium and palladium carriers were prepared from the respective metals. After dissolution they were converted to chlorides and finally dissolved in 3 N  $HCl$  at concentrations of 10, 6.1 and 10.0 mg/ml, respectively.

Silver carrier was prepared from  $AgNO_3$  in conc.  $HBr$  at a concentration of 10 mg/ml.

The conc.  $HBr-Br_2$  solution contained 1 drop of  $Br_2$ /ml.

### Procedure for $Pd^{115}$ Yield Determination

A solution of enriched uranium (250  $\lambda$ ), of ruthenium, rhodium, silver, and palladium carriers (100, 150, 100 and 200  $\lambda$ , respectively), and of conc.  $HBr-Br_2$  (500  $\lambda$ ) was contained in a pneumatically-driven sample carrier (rabbit). Each rabbit was irradiated for 5 seconds in the Vallecitos Nuclear Test Reactor in a flux of  $\sim 10^{12}$  neutrons  $cm^{-2} sec^{-1}$ . The cadmium ratio for gold was determined to be 2.7. A gold foil taped to the rabbit served as a monitor of the number of fissions in the sample as before.<sup>3</sup>

At the end of the irradiation the rabbit was transferred in about 1 second a distance of 50 feet to the laboratory. The irradiated solution was transferred by suction to a tube containing 10 ml of hot conc.  $HBr$ , and in most cases the rabbit was washed with 1 ml of conc.  $HBr$ . The combined solution was passed through 2 grams of copper powder in a filtration apparatus<sup>2</sup> at a definite time after the end of the irradiation. Passage of the solution through the copper was complete in less than 1 second. For separations which occurred more than several seconds after irradiation the copper bed was washed immediately with 5 ml of hot conc.  $HBr$ .

After separation the copper was transferred to a centrifuge tube which contained a definite volume of standardized cadmium carrier. Several hours after irradiation the copper was dissolved with 10 ml conc.  $\text{HNO}_3$ , evaporated to dryness and then brought twice more to dryness after the addition of 5-ml volumes of conc.  $\text{HCl}$ . The residue was dissolved in 125 ml of 4 N  $\text{HCl}$ , and both copper and palladium were quantitatively precipitated by bubbling  $\text{H}_2\text{S}$  gas for 10 minutes through the previously warmed solution. The insoluble sulfides were collected on sintered glass and saved for the palladium-yield determination.

The clear filtrate containing the cadmium was alkalized with  $\text{NaOH}$  pellets.  $\text{H}_2\text{S}$  gas was further bubbled through the filtrate for several minutes, and the precipitated cadmium sulfide was separated by centrifugation. This precipitate was dissolved in 1 ml of conc.  $\text{HCl}$  and the solution was analyzed for  $\text{Cd}^{115}$  by the radiochemical procedure of Hicks.<sup>4</sup> The beta-ray activity was measured on a gas-flow proportional counter for 1 to 2 months and the decay curve was resolved into the 2.3- and 43-day isomers of  $\text{Cd}^{115}$ . The count rate of the shorter-lived isomer was corrected for decay from the time of irradiation, for cadmium carrier yield in the radiochemical procedure, and for palladium yield (see below), and was normalized to  $10^{12}$  fissions.

The palladium carrier yield in the initial separation was determined spectrophotometrically<sup>5</sup> with corrections by the isotopic dilution technique for losses incurred in the purification process. The purification was performed as follows: The copper-palladium insoluble sulfide fraction was dissolved from the sintered glass filter with aqua regia after the addition of a known quantity of  $\text{Pd}^{109}$ . This solution was evaporated to dryness and the residue was dissolved in 20 ml of water.

Two ml of 1 % dimethylglyoxime in ethyl alcohol were added and the solution was extracted two times with 10-ml portions of chloroform. The chloroform extract was washed three times with 5 ml of 0.3 N  $\text{HCl}$  and the palladium was then extracted into 10 ml  $\text{NH}_4\text{OH}$ . After the ammonia extract was heated to a boil,  $\text{H}_2\text{S}$  gas was bubbled through it, and conc.  $\text{HCl}$  was added dropwise until the palladium sulfide precipitate appeared. After 3 minutes more of  $\text{H}_2\text{S}$  bubbling, the precipitate was separated from the solution. The sulfide was dissolved in 2 or 3 drops of aqua regia and carefully evaporated to complete dryness. The residue was dissolved and brought to a definite volume with  $1.5 \times 10^{-3}$  N  $\text{HCl}$ . Colorimetric<sup>5</sup> and  $\text{Pd}^{109}$  analysis of aliquots of this solution provided the values for the palladium yield determination.

#### Total $\text{Cd}^{115}$ Yield Determination

Two rabbits with contents as above including the various carriers were irradiated for 40 seconds. Several hours after the irradiation



they were analyzed for  $\text{Cd}^{115}$ . The count rate was corrected for decay and cadmium carrier yield, and again was normalized to  $10^{12}$  fissions.

## RESULTS AND DISCUSSION

In the rapid separation procedure of palladium from silver and cadmium, 75-80 % of the palladium was reduced and retained by the copper bed. Preliminary experiments with radioactive tracers had indicated that contamination by cadmium and silver was  $4.2 \times 10^{-2}$  % and 0.95 %, respectively. Ruthenium and rhodium, potential palladium precursors, are associated with the copper bed to the extent of 3.0 % and 14.3 %. As discussed below, error from contamination by ruthenium and rhodium was not evident in the decay measurements of  $\text{Pd}^{115}$ . Half-lives of isotopes of these elements with a mass number of 115 are probably too short for them to be contaminants in this procedure.

The counting rate of  $\text{Cd}^{115}$  for 16 separations extending in time from 2.8-120 seconds after fission appears in Fig. 1. The time of separation was the time between the end of irradiation and filtration through the copper bed. The relationship between the logarithm of the counting rate and the separation time is linear. Analysis by the method of least squares gives a half-life of  $40.5^{+4.0}_{-3.0}$  sec for  $\text{Pd}^{115}$  which agrees with the previously reported half-life.<sup>1</sup> No shorter-lived isomer of  $\text{Pd}^{115}$  was detectable, nor was interference from precursors evident.

The counting rate of  $\text{Cd}^{115}$  extrapolated to the end of the irradiation was corrected for the formation and decay of  $\text{Pd}^{115}$  during the course of the 5-second irradiation time. This corrected value was compared with the counting rate for the total  $\text{Cd}^{115}$  formed. The cumulative fractional chain yield of  $\text{Pd}^{115}$  thereby determined was  $90.6 \pm 7.2$  %. Formerly, under the assumption that the distribution of nuclear charge is Gaussian and that the width parameter  $\sigma$  is 0.62, values of  $Z_p$  were computed from cumulative fractional yields.<sup>6,7</sup> However, recent information points to the unreliability of this approach; in the 131-136 mass chains the values of  $\sigma$  were determined to vary over the range of 0.28-0.69.\* To illustrate the sensitivity of  $Z_p$  over this range of  $\sigma$  values, the above treatment gives values of  $Z_p$  which vary from 45.58-46.08.

\*Strom, P. O., et al., to be published.




Fig. 1 The Growth of  $\text{Cd}^{115}$  From  $\text{Pd}^{115}$  at Various Times of Pd Separation

More adequate characterization of  $Z_p$  for mass 115 will depend upon a separate evaluation of the contribution of other chain members to the total yield. Currently experiments are underway to measure the independent yield of the two  $\text{Ag}^{115}$  isomers; subsequently, if practicable, the independent yield of  $\text{Cd}^{115}$  will be determined.

#### REFERENCES

1. Alexander, J. M., Schindewolf, U., and Coryell, C. D. Phys. Rev. 111:228 (1958).
2. Weiss, H. V., and Reichert, W. L., "Rapid Isolation of Radiosilver and Certain Other Radioelements From Solution. Application to a Search for  $\text{Ag}^{121}$  in  $\text{U}^{235}$  Fission." U. S. Naval Radiological Defense Laboratory, USNRDL-TR-844, 11 April 1965.
3. Weiss, H. V., and Ballou, H. E., J. Inorg. Nucl. Chem., 27:1917 (1965).
4. Hicks, H. G., University of California Radiation Laboratory, UCRL-4377 (1954).
5. Cheng, K. L., Anal. Chem., 26:1894 (1954).
6. Weiss, H. V., Phys. Rev., 139:B304 (1965).
7. Wahl, A. C., Ferguson, D. R., Nethaway, D. R., Troutner, D. E. and Wolfsberg, K., Phys. Rev. 126:112 (1962).

UNCLASSIFIED

Security Classification.

DOCUMENT CONTROL DATA - R&D		
(Security classification of title, body of abstract and indexing annotation must be entered when the overall report is classified)		
1. ORIGINATING ACTIVITY (Corporate author) U. S. Naval Radiological Defense Laboratory San Francisco, California 94135		2a. REPORT SECURITY CLASSIFICATION UNCLASSIFIED
		2b. GROUP
3. REPORT TITLE Pd <sup>115</sup> YIELD IN THERMAL NEUTRON FISSION OF U <sup>235</sup>		
4. DESCRIPTIVE NOTES (Type of report and inclusive dates)		
5. AUTHOR(S) (Last name, first name, initial) Weiss, Herbert V. Reichert, W. L.		
6. REPORT DATE 7 February 1966	7a. TOTAL NO. OF PAGES 10	7b. NO. OF REFS 7
8a. CONTRACT OR GRANT NO. AT(49-2)-1167	9a. ORIGINATOR'S REPORT NUMBER(S) USNADL-TR-943	
b. PROJECT NO.		
c.	9b. OTHER REPORT NO(S) (Any other numbers that may be assigned this report)	
d.		
10. AVAILABILITY/LIMITATION NOTICES Distribution of this document is unlimited.		
11. SUPPLEMENTARY NOTES	12. SPONSORING MILITARY ACTIVITY Atomic Energy Commission Washington, D.C. 20545	
13. ABSTRACT <p>The formation of Pd<sup>115</sup> in thermal neutron fission of U<sup>235</sup> was studied quantitatively. A procedure was developed which rapidly separated palladium from daughter elements silver and cadmium. Separations were made from 2.8-120 seconds after fission. The growth of Cd<sup>115</sup> from palladium separated at various times constituted a decay curve for Pd<sup>115</sup>. The half-life thus determined was 40.5 <math>\pm</math> 3.0 seconds, a value which agrees with that previously reported. The cumulative fractional yield of Pd<sup>115</sup> compared to the total Cd<sup>115</sup> yield was 90.5 <math>\pm</math> 7.2 %. Estimation of the most probably nuclear charge (Z<sub>p</sub>) must await information on the contribution of other members to the yield of the 115 mass chain.</p>		

UNCLASSIFIED

Security Classification

14. KEY WORDS	LINK A		LINK B		LINK C	
	ROLE	WT	ROLE	WT	ROLE	WT
Pd <sup>115</sup> Fission of U <sup>235</sup> Separation of Pd from Ag and Cd Nuclear charge distribution						

## INSTRUCTIONS

1. **ORIGINATING ACTIVITY:** Enter the name and address of the contractor, subcontractor, grantee, Department of Defense activity or other organization (corporate author) issuing the report.

2a. **REPORT SECURITY CLASSIFICATION:** Enter the overall security classification of the report. Indicate whether "Restricted Data" is included. Marking is to be in accordance with appropriate security regulations.

2b. **GROUP:** Automatic downgrading is specified in DoD Directive 5200.10 and Armed Forces Industrial Manual. Enter the group number. Also, when applicable, show that optional markings have been used for Group 3 and Group 4 as authorized.

3. **REPORT TITLE:** Enter the complete report title in all capital letters. Titles in all cases should be unclassified. If a meaningful title cannot be selected without classification, show title classification in all capitals in parenthesis immediately following the title.

4. **DESCRIPTIVE NOTES:** If appropriate, enter the type of report, e.g., interim, progress, summary, annual, or final. Give the inclusive dates when a specific reporting period is covered.

5. **AUTHOR(S):** Enter the name(s) of author(s) as shown on or in the report. Enter last name, first name, middle initial. If military, show rank and branch of service. The name of the principal author is an absolute minimum requirement.

6. **REPORT DATE:** Enter the date of the report as day, month, year, or month, year. If more than one date appears on the report, use date of publication.

7a. **TOTAL NUMBER OF PAGES:** The total page count should follow normal pagination procedures, i.e., enter the number of pages containing information.

7b. **NUMBER OF REFERENCES:** Enter the total number of references cited in the report.

8a. **CONTRACT OR GRANT NUMBER:** If appropriate, enter the applicable number of the contract or grant under which the report was written.

8b, 8c, & 8d. **PROJECT NUMBER:** Enter the appropriate military department identification, such as project number, subproject number, system numbers, task number, etc.

9a. **ORIGINATOR'S REPORT NUMBER(S):** Enter the official report number by which the document will be identified and controlled by the originating activity. This number must be unique to this report.

9b. **OTHER REPORT NUMBER(S):** If the report has been assigned any other report numbers (either by the originator or by the sponsor), also enter this number(s).

10. **AVAILABILITY/LIMITATION NOTICES:** Enter any limitations on further dissemination of the report, other than those

imposed by security classification, using standard statements such as:

- (1) "Qualified requesters may obtain copies of this report from DDC."
- (2) "Foreign announcement and dissemination of this report by DDC is not authorized."
- (3) "U. S. Government agencies may obtain copies of this report directly from DDC. Other qualified DDC users shall request through \_\_\_\_\_."
- (4) "U. S. military agencies may obtain copies of this report directly from DDC. Other qualified users shall request through \_\_\_\_\_."
- (5) "All distribution of this report is controlled. Qualified DDC users shall request through \_\_\_\_\_."

If the report has been furnished to the Office of Technical Services, Department of Commerce, for sale to the public, indicate this fact and enter the price, if known.

11. **SUPPLEMENTARY NOTES:** Use for additional explanatory notes.

12. **SPONSORING MILITARY ACTIVITY:** Enter the name of the departmental project office or laboratory sponsoring (paying for) the research and development. Include address.

13. **ABSTRACT:** Enter an abstract giving a brief and factual summary of the document indicative of the report, even though it may also appear elsewhere in the body of the technical report. If additional space is required, a continuation sheet shall be attached.

It is highly desirable that the abstract of classified reports be unclassified. Each paragraph of the abstract shall end with an indication of the military security classification of the information in the paragraph, represented as (TS), (S), (C), or (U).

There is no limitation on the length of the abstract. However, the suggested length is from 150 to 225 words.

14. **KEY WORDS:** Key words are technically meaningful terms or short phrases that characterize a report and may be used as index entries for cataloging the report. Key words must be selected so that no security classification is required. Identifiers, such as equipment model designation, trade name, military project code name, geographic location, may be used as key words but will be followed by an indication of technical context. The assignment of links, roles, and weights is optional.

END

DATE

FILMED

2-19-66