

Distribution of this  
document is unlimited.

AD 632360

Technical Report

66 - 26 FD

CALCULATED INTERNAL RADIATION DOSE  
FROM INGESTION OF MEAT STERILIZED  
BY ELECTRON IRRADIATION

by

THOMAS G. MARTIN, III  
Radiation Sources Branch

Project Reference:  
1-C-O-25601-A-033

Series: FD-38

April 1966

Food Division  
U. S. ARMY NATICK LABORATORIES  
Natick, Massachusetts

**Best  
Available  
Copy**

APPROVED FOR							
WRITE SECTION <input checked="" type="checkbox"/>	WRITE SECTION <input type="checkbox"/>						
DATE	DATE						
<i>Paul J. [unclear]</i>							
DATE SECTION/AVAILABILITY CODES							
<table border="1"> <tr> <th>DATE</th> <th>AVAIL. CODE</th> <th>REASON</th> </tr> <tr> <td>/</td> <td></td> <td></td> </tr> </table>	DATE	AVAIL. CODE	REASON	/			
DATE	AVAIL. CODE	REASON					
/							

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED.

The findings in this report are not to be construed as an official Department of the Army position unless so designated by other authorized documents.

Citation of trade names in this report does not constitute an official endorsement or approval of the use of such items.

Destroy this report when no longer needed. Do not return it to the originator.

## FOREWORD

This report presents a means of estimating the magnitude of internal radiation hazard which may result from the ingestion of meat sterilized by irradiation with high energy electrons.

Beef is used as a basis for the calculations but the treatment can be employed for other foods for which radiation sterilization has been proposed.

From the results of this report, it can be stated that the amount of internal exposure is relatively small even for the highest energy considered. The calculated values represented approximately 0.1 to 2% of the average mean radiation exposure of humans, 130 mrem/year due to natural environmental radiation<sup>(12)</sup>. It points out, however, the need for further study in this area to confirm and improve the accuracy of such estimations.

EDWARD S. JOSEPHSON  
Associate Director for Food Radiation  
Food Division

### APPROVED:

FERDINAND P. MEHRLICH, Ph.D.  
Director Food Division

DALE H. SIELING, Ph.D.  
Scientific Director

W. M. MANTZ  
Colonel, QMC  
Commanding

## CONTENTS

	<u>Page</u>
List of tables/figures	iii
Abstract	iv
1. Introduction	1
2. Basis of Calculations	1
3. Calculations	2
4. Sources of Error	5
5. Discussion	6
6. References	8
7. Appendix I	13

## LIST OF TABLES/FIGURES

<u>Table</u>	<u>Page</u>
I. Data used in calculation of radionuclide production	9
II. Data used for calculation of Dose values	10
III. Annual dose (mrem) calculated for several energies	11
<u>Figure</u>	
I. Annual dose (mrem) vs. electron energy (MeV)	12

## ABSTRACT

The production of radionuclides in food irradiated with high energy electrons can be predicted reasonably well with the equation presented by R. A. Meyer (1). Using this equation and others derived from the reports of the International Commission on Radiological Protection (3) it is possible to estimate the amount of radiation received by humans consuming meat irradiated by high energy electrons.

The estimated annual internal doses decrease with storage time. At the maximum energy treated the dose would decrease from a maximum of 2.36 mrem/year to a minimum of 0.113 mrem/year with a storage time of 7 days and two years, respectively. The calculated values represented approximately 0.1 to 2% of the average mean radiation exposure of humans, 130 mrem/year due to natural environmental radiation (12).

CALCULATED INTERNAL RADIATION DOSE FROM INGESTION  
OF MEAT STERILIZED BY ELECTRON IRRADIATION

1. Introduction

The use of electron accelerators as sources of high intensity radiation for food sterilization has shown great promise because of the relatively high dose rate obtainable as well as economic and operational efficiency.

One disadvantage of the electron accelerator is the limited penetration of the electron beam. This penetration is related to the energy of the beam electrons, i.e., in order to penetrate a thicker package it requires an increase in energy, and from this standpoint, the most efficient operation of the machine should be at the highest energy obtainable. There are costs associated with an increase of efficiency. With the accelerator, one of these costs is the production of radioactive nuclides in the irradiated sample. As the energy is increased to enable the irradiation of thicker packages the probability of inducing radioactivity is also increased.

Below some threshold energy, the reaction which causes the induced activity cannot occur. However, it appears reasonable that even above this threshold energy, there is a range which should be of little concern since the amount of induced radioactivity may not produce significant internal radiation exposure to the population eating the food.

During the years that accelerators have been considered for the sterilization of food, several reports concerning the amount of radioactivity induced in food have been published (1, 2, 5, 6, 7). Meyer (1) has suggested an equation for predicting concentrations of radionuclides produced in food as a function of incident electron energy and dose. This equation agrees reasonably well with available experimental data. This present report attempts to predict the internal absorbed dose to humans who eat meat sterilized by electron irradiation at several energies. Using the equation of Meyer and those of the International Commission in Radiological Protection (3), calculations are made for beef sterilized by electrons of energies 12, 14, 16, 18, 20, and 24 MeV and a dose of five megarads.

2. Basis of Calculations

In order to predict the dose of radiation which will be received by persons eating electron sterilized beef, it is necessary to make a series of assumptions. With each assumption made, there is further error introduced when an individual case is to be considered; however, in order to establish a

reference point from which to work, it is necessary to set conditions which may vary in their validity. At this point, the assumptions will be stated, but not defended. Under Sources of Error, the effect of the assumptions will be discussed.

In the calculations, it was assumed that:

1. Meyer's equation was valid.
2. Beef received five megarad dose at the stated energies.
3. Beef was stored for the stated times prior to ingestion.
4. 200 grams of irradiated beef were ingested per day.
5. A state of equilibrium was reached in the body with reference to the radionuclide in question, which may be expressed by equation number (2).
6. Total dose received was sum of doses from each of the radionuclides considered.
7. Radionuclides were uniformly distributed throughout the body.

### 3. Calculations

Meyer, after reviewing previously published information, arrives at the following equation for predicting radioactivity in food irradiated with electrons:

$$R = KAnDT^{-1} (E - E_0)^3 \quad (1)$$

where:

- R = activity in pc/gm food/D Mrads  
 K =  $4 \times 10^{-3}$   
 A = atomic number of the target isotope  
 n = fractional abundance of the target isotope in the food  
 $\frac{\text{atoms of target isotope}}{\text{atoms of target element}} \times \frac{\text{grams of element}}{\text{grams of beef}}$   
 D = dose in Megarads  
 T = half life of product activity in years  
 E = initial electron energy in MeV  
 E<sub>0</sub> = threshold energy for the reaction producing the product activity in MeV.

Tables I and II give the data used for the calculations in this report.

In order to establish values for an equilibrium concentration within the body of the radionuclide in question, the following form derived from the ICRP report (3) was used:

$$C = \frac{T_{eff} I f}{0.693 m} \quad (2)$$

where:

(2a)

$$I = a \cdot R \cdot e^{-\frac{0.693 \cdot t}{T}}$$



$C$  = the equilibrium concentration of the element in the body of radionuclide pCi/g of body  
 $T_{eff}$  = the effective half period for the radionuclide in the body in days  
 $I$  = rate of ingestion of the element (radionuclide) pCi/day  
 $f$  = the fraction of the ingested material remaining in the body  
 $m$  = the mass of the body (taken as 70 Kg for the "standard man")  
 $a$  = 200 grams of beef (see assumption 4)  
 $t$  = time between radiation and ingestion

The absorbed dose in rem/year is then calculated from:

$$(AD) = \frac{C f \epsilon}{53.6} \quad (3)$$

where:

$(AD)$  = absorbed dose received by the body in rem/year  
 $C$  = concentration in pCi/g of body  
 $f$  = fraction reaching critical organ (taken as 1 for total body)  
 $\epsilon$  = energy absorbed by tissue in MeV/disintegration  
 53.6 = constant value such that units are properly converted i.e.

$$53.6 = \frac{100 \text{ ergs}}{\left(\frac{\text{g rads}}{3.7 \times 10^4 \frac{\text{dps}}{\text{uCi}}}\right) \left(\frac{10^6 \text{ pCi/uCi}}{1.6 \times 10^6 \frac{\text{ergs}}{\text{MeV}}}\right) \left(\frac{3.15 \times 10^7 \text{ sec}}{\text{year}}\right)}$$

Two sample calculations are shown in order to show the general form used in the calculation. For  $\text{Na}^{22}$ , the 2.6 year half-lived isotope of sodium which is produced by a  $(\gamma, n)$  reaction, the following example is presented: (20 MeV electrons: 1 year storage)

$$\begin{aligned}
 R &= K \lambda n D T^{-1} (E - E_0)^3 \\
 R &= \frac{(4 \times 10^{-3}) (23) (5 \times 10^{-4}) (5)}{2.6} (20 - 12.4)^3 \\
 R &= 3.88 \times 10^{-2} \text{ pCi/g of beef} \\
 C &= \frac{(I) (f)}{(T_{eff}) (m)} \\
 C &= \frac{(11) (5.92) (1)}{(0.693) (7 \times 10^4)} \\
 C &= 1.34 \times 10^{-3} \text{ pCi/g of tissue} \\
 AD &= \frac{C f \epsilon}{53.6} \\
 AD &= \frac{(1.34 \times 10^{-3}) (1) (1.6)}{53.6} \\
 AD &= 4.00 \times 10^{-5} \text{ rem/year}
 \end{aligned}
 \quad
 \begin{aligned}
 I &= 200 \cdot R \cdot e^{-\frac{0.693 \cdot 365}{T}} - 253 \\
 I &= (2 \times 10^2) (3.88 \times 10^{-2}) e^{-\frac{253}{949}} \\
 I &= 7.76 e^{-0.266} \\
 I &= 5.92 \text{ pCi/day}
 \end{aligned}$$

For  $P^{32}$  the day isotope of phosphorus which is produced by a ( $\gamma$ , pn) reaction with  $S^{34}$ , the following example is presented: (20 MeV, 1 month storage)

$$R = K \Delta DT^{-1} (E - E^0)^3$$

$$R = \frac{(4 \times 10^{-3})(34)(0.92 \times 10^{-4})(5)}{3.9 \times 10^{-2}} (20 - 10.9)^3$$

$$R = 1.21 \text{ pCi/g of beef}$$

$$I' = I R$$

$$I' = (2 \times 10^2)(1.21)(e^{-(.693)(30)})$$

$$I' = (2.42 \times 10^2)(e^{-1.46}) \quad 14.2$$

$$I = 5.62 \times 10 \text{ pCi/day}$$

$$C = \frac{(T_{eff})(I')(f)}{0.693 \text{ m}}$$

$$C = \frac{(13.5)(56.2)(0.75)}{(0.693)(7 \times 10^4)}$$

$$C = 1.17 \times 10^{-2} \text{ pCi/g of tissue}$$

$$AD = \frac{C f \epsilon}{53.6}$$

$$AD = \frac{(1.17 \times 10^{-2})(1)(0.69)}{53.6}$$

$$AD = 1.51 \times 10^{-4} \text{ rem/year}$$

#### 4. Sources of Error

The assumptions made in paragraph 2 are considered in some detail in the order in which they were listed.

##### 1. Validity of Meyer's Equation

Meyer's equation can only be expected to predict the activity produced to within approximately a factor of 2, since the fractional abundance of the parent isotope in a given food often varies. Further, the equation is based on a simple approximation of both the photonuclear cross section and the bremsstrahlung spectra. Meyer and Burkhardt (7) have shown, however, that this equation and a graphical integration are in reasonable agreement with each other and available experimental data.

##### 2. Beef received five Megarad dose at the stated energies

Dose to beef may vary throughout the product up to 25% under present irradiation requirements. The energy of the electrons is not monochromatic, therefore, electrons up to the maximum energy will be present in the beam. This condition will tend to lower actual production of radionuclides as compared to the theoretical production.

##### 3. Beef was stored for the stated times prior to ingestion

No problem arises from this assumption.

##### 4. 200 grams of irradiated beef are ingested per day

This is a relatively conservative consumption since it assumes all of the beef eaten is irradiated and that beef is eaten every day. Beef was chosen, however, since it is the most likely candidate for such an assumption.

##### 5. A state of equilibrium is reached in the body with reference to the radionuclides in question, which may be expressed by equation (2).

This assumption is one that has been made by the ICRP in their calculations of maximum permissible concentrations of radionuclides in the human body, water, and air.

6. Total dose received is the sum of doses from each of the radionuclides considered.

There are other possible sources of induced radioactivity. One of the most significant reactions is that which produces isomeric radioculides. However, according to Meyer (5) no isomeric activation has been found in foods. Another source of error might be the gamma-triton reaction which produces tritium. Meyer states that tritium has not been found experimentally above background levels in food.

Other elements than those considered in Table I and which might be present in foods could also produce radioactivity via the reactions considered. However, these elements are generally in very small concentrations and would produce negligible amounts of the radioactive products.

7. Whole body dose assumes uniform distribution of each of the radionuclides in the body

This assumption is one which leaves much to discuss. Actually, no element is truly uniformly distributed throughout the entire body. When one considers the localization of a particular radionuclide in a specific organ, then the dose to that organ will be significantly higher than the dose to the whole body. The two most significant radionuclides  $\text{Na}^{22}$  and  $\text{P}^{32}$  are relatively uniform in distribution, therefore, the dose calculations are acceptable.

For the short storage time,  $\text{I}^{126}$  represents a relatively low total body exposure  $8.02 \times 10^{-5}$  mrem/year, however, iodine is selectively absorbed by the thyroid gland. If the thyroid gland is considered alone, the dose would be about  $5.63 \times 10^{-2}$  mrem/year or nearly 1,000 times the dose to the whole body contributed by Iodine-126.

The major contributor to the internal dose to the body during short storage times, 7 days and 30 days, is phosphorus-32; after longer storage times Sodium-22 becomes the major contributor.

5. Discussion

Table I indicates the data used in solution of Meyer's equation for each of the nuclides listed.

Table II indicates the data used in the internal dose equations. Appendix I shows calculations of the effective energy per disintegration for these radionuclides for which data are not available in the TRP report.

Table III indicates the total annual internal absorbed dose in mrem for various energies and storage times. Figure 1 shows graphically the same

information allowing for some interpolation of energies and storage times.

It is the intention of this report to present an estimate of internal dose received by persons eating electron sterilized meat and thereby demonstrating that this dose is relatively low. It is not intended to minimize the need for further experimental studies to confirm these data.

## REFERENCES

1. R. A. Meyer, "Induced Radioactivity in Food and Electron Sterilization" U. S. Army Natick Laboratories Technical Report, FD-6, 1964
2. W. H. Newkirk, H. D. Smith, and R. A. Glass, "Radioactivities Produced in Foods by High-Energy Electrons: Theoretical Analysis and Computer Calculations," Stanford Research Institute Project SU-242, Report No. 11, 14 April 1960 (DA 19-129-QM-1100)
3. Report of International Commission on Radiological Protection Committee II on Permissible Dose for Internal Radiation (1959), Health Physics 3 (June 1960)
4. K. T. Bainbridge, and A. O. Mier, "Relative Isotopic Abundances of the Elements," Nuclear Science Series Report No. 9, National Research Council
5. E. Haddad, G. D. Trimble, and W. R. Mowry, "Electron Beam Irradiation Services," General Atomic Final Report (13 Nov 1961 to 13 Nov 1963) Project No. 7-94-01-002, July 1, 1963 (DA 19-129-QM-1924)
6. M. Stein, "A Comparison of the Doses Produced from Ingestion of Electron Irradiated Food to Doses from Several Radiation Sources", Quartermaster Food & Container Institute for the Armed Forces Research Project No. 15-59, 1 May 1959
7. R. A. Meyer and J. L. Burkhardt, "Induced Radioactivity in Food and Electron Sterilization; Calculations of Induced Radioactivity II", To be published.
8. D. O. Hunt, A. Brynjolfsson, and R. D. Cooper, "Handbook of Photonuclear Reactions," U. S. Army Natick Laboratories Technical Report, FD-1 (1963)
9. Radiological Health Data, U. S. Department of Health, Education, and Welfare, Vol. V (1964)
10. Radiation Quantities and Units, International Commission on Radiological Units and Measurements, Report 10a, National Bureau of Standards Handbook 84, (1962)
11. C. F. Miller and S. L. Brown, "Models for Estimating the absorbed Dose from Assimilation of Radionuclides in Body Organs of Humans", Stanford Research Institute Project No. IMU-4021 May 1963 (OCD-OS-62-135)
12. Report of the United Nations Scientific Committee on the Effects of Atomic Radiation, New York: United Nations, General Assembly Official Records 17th Session, Supplement 16, p.442. (1962)

TABLE I

Data Used in Calculation of  
Radionuclide Production

Nuclide	Parent/reaction	Fraction of parent nuclide in beef	Half life of product Nuclide (years)	Threshold energy for reaction (MeV)
Na <sup>22</sup>	Na <sup>23</sup> (γ-n)	5 x 10 <sup>-4</sup>	2.6	12.4
Na <sup>24</sup>	Mg <sup>24</sup> (γ-p)	3.1 x 10 <sup>-5</sup>	1.71 x 10 <sup>-3</sup>	12.1
P <sup>32</sup>	S <sup>34</sup> (γ-pn)	0.92 x 10 <sup>-4</sup>	3.9 x 10 <sup>-2</sup>	10.9
P <sup>33</sup>	S <sup>34</sup> (γ-p)	0.92 x 10 <sup>-4</sup>	6.6 x 10 <sup>-2</sup>	18.8
S <sup>35</sup>	Cl <sup>37</sup> (γ-pn)	1.47 x 10 <sup>-4</sup>	0.24	16.1
Ca <sup>45</sup>	Ca <sup>46</sup> (γ-n)	3.3 x 10 <sup>-9</sup>	0.45	10.4
Cr <sup>51</sup>	Cr <sup>52</sup> (γ-n)	2.5 x 10 <sup>-9</sup>	7.4 x 10 <sup>-2</sup>	12.0
Mn <sup>54</sup>	Mn <sup>55</sup> (γ-n)	2.0 x 10 <sup>-7</sup>	0.822	10.2
Fe <sup>55</sup>	Fe <sup>56</sup> (γ-n)	3.8 x 10 <sup>-5</sup>	2.6	11.2
Zn <sup>65</sup>	Zn <sup>66</sup> (γ-n)	4.17 x 10 <sup>-6</sup>	0.671	11.0
Rb <sup>84</sup>	Rb <sup>85</sup> (γ-n)	0.94 x 10 <sup>-6</sup>	9.0 x 10 <sup>-2</sup>	10.5
I <sup>126</sup>	I <sup>127</sup> (γ-n)	3.5 x 10 <sup>-8</sup>	3.6 x 10 <sup>-2</sup>	9.2
Cs <sup>132</sup>	Cs <sup>133</sup> (γ-n)	9.2 x 10 <sup>-9</sup>	1.7 x 10 <sup>-2</sup>	9.0

TABLE II  
Data for the Dose Calculations

Nuclide	Effective Half Life (days)	Fraction of Ingested Nuclide Remaining in Body	Effective Energy Absorbed per Disintegration + (MeV/dis.)
Na <sup>22</sup>	11	1.0	1.6
Na <sup>24</sup>	0.6	1.0	3.6
P <sup>32</sup>	13.5	0.75	0.69
P <sup>33</sup>	22.8	0.75	0.086
S <sup>35</sup>	76.4	1.0	0.056
Ca <sup>45</sup>	162	1.0	0.43
Cr <sup>51</sup>	26.6	$5 \times 10^{-3}$	0.025
Mn <sup>54</sup>	5.6	0.1	0.23
Fe <sup>55</sup>	388	0.1	$6.5 \times 10^{-3}$
Zn <sup>65</sup>	194	0.1	0.32
Rb <sup>84</sup>	0.047	1.0	1.63
I <sup>126</sup>	12.1	1.0	0.16
Cs <sup>132</sup>	0.168	1.0	0.47


+ See Appendix I for calculations of  $\epsilon$  not given in reference (3).



TABLE III

Annual Dose (mrem) Calculated for Several Energies

Electron Energy	Storage Time (days)				
	7	30	180	365	720
12	8.11 (-4)	2.64 (-4)	-	-	-
14	1.92 (-2)	6.44 (-3)	4.42 (-4)	3.83 (-4)	2.94 (-4)
16	9.02 (-2)	3.18 (-2)	4.97 (-3)	4.32 (-3)	3.31 (-3)
18	2.53 (-1)	9.26 (-2)	1.87 (-2)	1.61 (-2)	1.24 (-3)
20	5.54 (-1)	2.13 (-1)	4.86 (-2)	4.11 (-2)	3.14 (-2)
22	1.29	6.18 (-1)	1.67 (-1)	9.85 (-2)	6.41 (-2)
24	2.36	1.21	3.40 (-1)	1.84 (-1)	1.13 (-1)



**FIG 1, ANNUAL DOSE (mrem) vs ELECTRON ENERGY (MeV)**

APPENDIX I

Calculations of effective energy per disintegration based on the techniques described in the report of the International Commission on Radiological Protection (1959) are made for the following radionuclides:

Na<sup>24</sup> RBE = 1 (More recent terminology in personnel dosimetry uses QF (quality factor) to effectively describe the relative effectiveness of various types and energies of radiation)

$$n = 1$$

$$F = 1$$

$$E_{\beta} = 0.33 E_m f(1 - \frac{Z^{\frac{1}{2}}}{50})(1 - \frac{E_m^{\frac{1}{2}}}{4})$$

$$E_{\beta} = (0.33)(1.40)(1)(1 - \frac{(11)^{\frac{1}{2}}}{50})(1 + \frac{(1.40)^{\frac{1}{2}}}{4})$$

$$E_{\beta} = 0.559$$

$$E_{\gamma_1} = E_m (1 - e^{-\sigma r})$$

$$E_{\gamma_1} = (1.37)(1)(1 - e^{-(0.060)(30)})$$

$$E_{\gamma_1} = 1.14$$

$$E_{\gamma_2} = (2.75)(1)(1 - e^{-(0.040)(30)})$$

$$E_{\gamma_2} = 1.93$$

$$E = E F(RBE) = 0.569 + 1.14 + 1.93$$

$$E_{Na^{24}} = 3.6$$

Rb<sup>84</sup>

$$RBE = 1$$

$$n = 1$$

$$F = 1$$

$$E_{\beta} = (0.33)(0.91)(0.025)(1 - \frac{6.08}{50})(1 - \frac{.954}{4})$$

$$E_{\beta} = 5.02 \times 10^{-3}$$

$$E_{\beta} = 0.33 E_m f(1 - \frac{E_m^{\frac{1}{2}}}{4}) + 2f(0.51)(1 - e^{-\sigma r})$$

$$E_{\beta_3} = (0.33)(1.64)(0.56)\left(1 - \frac{1.28}{4}\right) + (2)(0.51)(0.56)(1 - e^{-(0.04)(30)})$$

$$E = 0.604$$

$$E_{\beta_1} = (0.33)(0.79)(0.44)\left(1 - \frac{0.89}{4}\right) + (2)(0.51)(0.44)(1 - e^{-(0.04)(30)})$$

$$E_{\beta_1} = 0.404$$

$$E_{\beta_2} = (0.88)(1)(1 - e^{-(0.04)(30)})$$

$$E_{\beta_2} = 0.615$$

$$\epsilon_{Rb 84} = 1.63$$

Cs 132

$$E = (0.67)(1)(1 - e^{-(0.04)(30)})$$

$$E = 0.468$$

$$\epsilon = 0.468$$

Unclassified

Security Classification

**DOCUMENT CONTROL DATA - R&D**

(Security classification of title, body of abstract and indexing annotation must be entered when the overall report is classified)

1. ORIGINATING ACTIVITY (Corporate author) U. S. Army Natick Laboratories Natick, Mass. 01760		2a. REPORT SECURITY CLASSIFICATION Unclassified	
		2b. GROUP	
3. REPORT TITLE CALCULATED INTERNAL RADIATION DOSE FROM INGESTION OF MEAT STERILIZED BY ELECTRON IRRADIATION			
4. DESCRIPTIVE NOTES (Type of report and inclusive dates)			
5. AUTHOR(S) (Last name, first name, initial) Martin, Thomas G. III			
6. REPORT DATE April 1966		7a. TOTAL NO. OF PAGES 14	7b. NO. OF REFS 12
8a. CONTRACT OR GRANT NO.		9a. ORIGINATOR'S REPORT NUMBER(S) 66-26 FD	
b. PROJECT NO. 1-C-O-25601-A-033			
c.		9b. OTHER REPORT NO(S) (Any other numbers that may be assigned this report)	
d.		FD-38	
10. AVAILABILITY/LIMITATION NOTICES Distribution of this document is unlimited. Release to CFSTI is authorized.			
11. SUPPLEMENTARY NOTES		12. SPONSORING MILITARY ACTIVITY Food Division U. S. Army Natick Laboratories Natick, Massachusetts 01760	
13. ABSTRACT <p>The production of radionuclides in food irradiated with high energy electrons can be predicted reasonably well with the equation presented by Dr. R. A. Meyer. Using this equation and others derived from the reports of the International Commission on Radiological Protection it is possible to estimate the amount of radiation received by humans consuming meat irradiated by high energy electrons.</p> <p>The estimated annual internal doses decrease with storage time. At the maximum energy treated the dose would decrease from a maximum of 2.36 mrem/year to a minimum of 0.113 mrem/year with storage times of 7 days and two years respectively. The calculated values represented approximately 0.1 to 2% of the average mrem radiation exposure of humans, 130 mrem/year due to natural environmental radiation. (12)</p>			

KEY WORDS	LINK A		LINK B		LINK C	
	ROLE	WT	ROLE	WT	ROLE	WT
Measurement	8					
Estimation	8				8	
Predictions	8					
Radiation	9		8			
Dosage	9		8			
Internal	0		0		0	
Irradiated Food			9		9	
Beef			9		9	
Meat			9		9	
Irradiated			0		0	
Radiation hazard			0		9	

INSTRUCTIONS

1. **ORIGINATING ACTIVITY:** Enter the name and address of the contractor, subcontractor, grantee, Department of Defense activity or other organization (*corporate author*) issuing the report.

2a. **REPORT SECURITY CLASSIFICATION:** Enter the overall security classification of the report. Indicate whether "Restricted Data" is included. Marking is to be in accordance with appropriate security regulations.

2b. **GROUP:** Automatic downgrading is specified in DoD Directive 5200.10 and Armed Forces Industrial Manual. Enter the group number. Also, when applicable, show that optional markings have been used for Group 3 and Group 4 as authorized.

3. **REPORT TITLE:** Enter the complete report title in all capital letters. Titles in all cases should be unclassified. If a meaningful title cannot be selected without classification, show title classification in all capitals in parenthesis immediately following the title.

4. **DESCRIPTIVE NOTES:** If appropriate, enter the type of report, e.g., interim, progress, summary, annual, or final. Give the inclusive dates when a specific reporting period is covered.

5. **AUTHOR(S):** Enter the name(s) of author(s) as shown on or in the report. Enter last name, first name, middle initial. If military, show rank and branch of service. The name of the principal author is an absolute minimum requirement.

6. **REPORT DATE:** Enter the date of the report as day, month, year, or month, year. If more than one date appears on the report, use date of publication.

7a. **TOTAL NUMBER OF PAGES:** The total page count should follow normal pagination procedures, i.e., enter the number of pages containing information.

7b. **NUMBER OF REFERENCES:** Enter the total number of references cited in the report.

8a. **CONTRACT OR GRANT NUMBER:** If appropriate, enter the applicable number of the contract or grant under which the report was written.

8c, & 8d. **PROJECT NUMBER:** Enter the appropriate military department identification, such as project number, subproject number, system numbers, task number, etc.

9a. **ORIGINATOR'S REPORT NUMBER(S):** Enter the official report number by which the document will be identified and controlled by the originating activity. This number must be unique to this report.

9b. **OTHER REPORT NUMBER(S):** If the report has been assigned any other report numbers (*either by the originator or by the sponsor*), also enter this number(s).

10. **AVAILABILITY/LIMITATION NOTICES:** Enter any limitations on further dissemination of the report, other than those imposed by security classification, using standard statements such as:

- (1) "Qualified requesters may obtain copies of this report from DDC."
- (2) "Foreign announcement and dissemination of this report by DDC is not authorized."
- (3) "U. S. Government agencies may obtain copies of this report directly from DDC. Other qualified DDC users shall request through \_\_\_\_\_."
- (4) "U. S. military agencies may obtain copies of this report directly from DDC. Other qualified users shall request through \_\_\_\_\_."
- (5) "All distribution of this report is controlled. Qualified DDC users shall request through \_\_\_\_\_."

If the report has been furnished to the Office of Technical Services, Department of Commerce, for sale to the public, indicate this fact and enter the price, if known.

11. **SUPPLEMENTARY NOTES:** Use for additional explanatory notes.

12. **SPONSORING MILITARY ACTIVITY:** Enter the name of the departmental project office or laboratory sponsoring (*paying for*) the research and development. Include address.

13. **ABSTRACT:** Enter an abstract giving a brief and factual summary of the document indicative of the report, even though it may also appear elsewhere in the body of the technical report. If additional space is required, a continuation sheet shall be attached.

It is highly desirable that the abstract of classified reports be unclassified. Each paragraph of the abstract shall end with an indication of the military security classification of the information in the paragraph, represented as (TS), (S), (C), or (U).

There is no limitation on the length of the abstract. However, the suggested length is from 150 to 225 words.

14. **KEY WORDS:** Key words are technically meaningful terms or short phrases that characterize a report and may be used as index entries for cataloging the report. Key words must be selected so that no security classification is required. Identifiers, such as equipment model designation, trade name, military project code name, geographic location, may be used as key words but will be followed by an indication of technical context. The assignment of links, rules, and weights is optional.