

AD 041217

Technical Report 56

to the

Office of Naval Research

ARPA Order No. 26-60, Task 2, Item 3

Contract No. Nonr 477(16)

"A New Method for the Preparation of Fluorodisulfate Salts"

By Darryl D. Des Marteau and George H. Cady

University of Washington

October 14, 1966

D D C
RECEIVED
NOV 4 1966
C

Reproduction in whole or in part is permitted for any purpose of the United States Government.

CLEARINGHOUSE FOR FEDERAL SCIENTIFIC AND TECHNICAL INFORMATION			
Hardcopy	Microfiche	8	72
\$ 1.00	\$.50	pp	
/ ARCHIVE COPY			


**BEST
AVAILABLE COPY**

Contribution from the Department of Chemistry
University of Washington, Seattle, Washington

"A New Method for the Preparation
of Fluorodisulfate Salts"

By Darryl D. Des Marteau and George H. Cady

The only reported fluorodisulfate salt is that of potassium. It was prepared by the reaction of potassium fluoride with excess sulfur trioxide.¹



1. H. A. Lehmann and E. Kolditz, Z. Anorg. Allgem. Chem., 272, 69 (1953).

The white crystalline compound was unstable and decomposed completely to potassium fluorosulfate and sulfur trioxide when heated to 100° for 4 hours. Calcium fluorodisulfate has been suggested as an intermediate in the reaction of calcium fluoride with sulfur trioxide.²

2. E. L. Meutterties and D. D. Coffman, J. Am. Chem. Soc., 80, 5914 (1958).

Potassium, sodium and, probably, ammonium fluorodisulfates have now been produced by reaction of peroxydisulfuryl difluoride with the corresponding peroxydisulfates, the potassium and sodium

$$\text{M}_2\text{S}_2\text{O}_8 + \text{S}_2\text{O}_6\text{F}_2 = 2\text{MS}_2\text{O}_6\text{F} + \text{O}_2 = 2\text{MOSO}_2\text{F} + 2\text{SO}_3 \quad (\text{M} = \text{K}, \text{Na or NH}_4)$$

salts being obtained in high yield. The ammonium salt was impure and was obtained in approximately 15% yield.

Experimental

Reagents.— Peroxydisulfuryl difluoride was prepared and purified by the method of Shreeve and Cady.³ The peroxydisulfate salts

3. J. M. Shreeve and G. H. Cady, *Inorg. Chem.*, 2, 466 (1963).

were obtained from commercial sources and were dried by heating in vacuo.

Preparation and Analysis.—The reactions were run in a Pyrex glass bulb fitted with a Fischer-Porter glass valve closed by a Teflon stem. $S_2O_6F_2$ was condensed onto the dried, powdered peroxydisulfate at -183° and the mixture was then allowed to warm to room temperature. A slow reaction took place near room temperature liberating oxygen. An approximate 10:1 mole ratio of $S_2O_6F_2$ to $M_2S_2O_8$ was used in each case. The reactions were very slow and became slower with time, probably due to surface effects. Oxygen was removed periodically by pumping on the reaction vessel at -183° . After no further oxygen was observed to form (21 days for 1.7384g $K_2S_2O_8$, 4 days for 1.033g $Na_2S_2O_8$, 8 days for 1.0356g $(NH_4)_2S_2O_8$), the excess $S_2O_6F_2$ was removed by pumping at room temperature until the rate of weight loss was only a few mg. per hr. The products could not be pumped to constant weight because the unstable fluorodisulfate salts dissociated slowly into SO_3 and MO_2SO_2F at room temperature. For the above samples of $K_2S_2O_8$ and $Na_2S_2O_8$ the weights of product were 2.743g and 1.716g respectively (Theory: 2.807g

and 1.456g). For the sample of $(\text{NH}_4)_2\text{S}_2\text{O}_8$ the weight of product was 1.169g (1.789g theor) and corresponded to approximately 15% $(\text{NH}_4)_2\text{S}_2\text{O}_5\text{F}$ mixed with 85% $\text{NH}_4\text{SO}_3\text{F}$. The products were then heated to 155° maximum and pumped to constant weight. After this treatment the above samples weighed 1.760, 1.073 and 1.035 g. respectively, corresponding closely to the theoretical weights, 1.770, 1.060 and 1.063 g. respectively for KSO_3F , NaSO_3F and $\text{NH}_4\text{SO}_3\text{F}$.

Other runs were similar to those just described. The KOSO_2F , NaOSO_2F and $\text{NH}_4\text{OSO}_2\text{F}$ were then hydrolyzed in 1N. sodium hydroxide for 10 hours at 100° and the resulting solutions were analyzed for sulfur and fluorine. Sulfur was determined gravimetrically as barium sulfate and fluorine volumetrically by the method of Willard and Winter.⁴ The sulfur trioxide was identified qualita-

4. H. H. Willard and O. B. Winter, Ind. Eng. Chem., Anal. Ed.,
5, 7 (1933).

tively by its physical properties, by tendency to liberate iodine from aqueous potassium iodide and by giving a positive test for sulfate after hydrolysis. Oxygen was identified by its mass spectrum and volatility at -183° . The infrared spectrum of the excess peroxydisulfuryl difluoride remaining after reaction corresponded to $\text{S}_2\text{O}_6\text{F}_2$ containing a trace of SiF_4 . Analytical data for the runs given above are summarized in table I.

X-ray powder photographs were taken for samples of KOSO_2F and $\text{KS}_2\text{O}_6\text{F}$ prepared by the method given above and by the method of Lehmann and Kolditz.¹ The two methods gave materials having essentially the same patterns. These are given in the

Table I
Analytical Data

Cation	Na ⁺	K ⁺	NH ₄ ⁺
Moles O ₂ ^a	0.98	0.99	1.02
Moles SO ₂ ^b	0.99	1.00	1.01
Wt % C ^c found	16.2	13.6	16.0
calc.	15.6	13.8	16.2
Wt % S ^c found	26.2	22.5	27.6
calc.	26.3	23.2	27.3

a per mole of M₂S₂O₈ consumed

b per mole MS₂O₆F decomposed by heating

c analysis of MOSO₂F

Ph.D. thesis of the junior author.⁵ When the patterns were com-

S. Darryl D. Des Marteau, Thesis, University of Washington, 1966.

pared with those reported by Lehmann and Kolditz¹ the extent of agreement was incomplete. Prof. Dr. Kolditz has also compared the data and powder patterns (private correspondence) and agrees with us that the materials obtained by the two methods must be the same even though the reported powder patterns are not exactly the same.

DOCUMENT CONTROL DATA - R & D

(Security classification of title, body of abstract and indexing annotation must be entered when the overall report is classified)

ORIGINATING ACTIVITY (Corporate author)

University of Washington
Seattle, Washington

20. REPORT SECURITY CLASSIFICATION

None

21. GROUP

REPORT TITLE

"A New Method for the Preparation of Fluorodisulfate Salts"

DESCRIPTIVE NOTES (Type of report and inclusive dates)

Technical Report

AUTHOR(S) (First name, middle initial, last name)

Darryl D. Des Marteau and George H. Cady

REPORT DATE

October 14, 1966

22. TOTAL NO OF PAGES

5

23. NO OF REFS

5

CONTRACT OR GRANT NO

Nonr 477(16)

24. ORIGINATOR'S REPORT NUMBER(S)

56

PROJECT NO

25. OTHER REPORT NO(S) (Any other numbers that may be assigned this report)

None

DISTRIBUTION STATEMENT

Distribution of this document is unlimited.


SUPPLEMENTARY NOTES

12. SPONSORING MILITARY ACTIVITY


Office of Naval Research

ABSTRACT

Potassium and sodium fluorodisulfates are produced by the reaction of peroxydisulfuryl difluoride with the corresponding peroxydisulfates:



When M is ammonium, some $NH_4S_2O_6F$ is obtained together with much NH_4SO_3F , apparently formed by the reaction:



REV 00000

LINE A		LINE B		LINE C	
ROLE	WT	ROLE	WT	ROLE	WT

Fluorodisulfate salts