

ARL 66-0215
OCTOBER 1966



AD 647318

Aerospace Research Laboratories

A NOTE ON RECURRENCE RELATIONS BETWEEN EXPECTED
VALUES OF FUNCTIONS OF ORDER STATISTICS


(Reprint from The Annals of Mathematical Statistics, Vol. 37,
No. 3, June 1966)

P. R. KRISHNAIAH

APPLIED MATHEMATICS RESEARCH LABORATORY


M. HASEEB RIZVI

THE OHIO STATE UNIVERSITY
COLUMBUS, OHIO



Distribution of this document is unlimited

OFFICE OF AEROSPACE RESEARCH
United States Air Force



ARCHIVE COPY

NOTICES

When Government drawings, specifications, or other data are used for any purpose other than in connection with a definitely related Government procurement operation, the United States Government thereby incurs no responsibility nor any obligation whatsoever; and the fact that the Government may have formulated, furnished, or in any way supplied the said drawings, specifications, or other data, is not to be regarded by implication or otherwise as in any manner licensing the holder or any other person or corporation, or conveying any rights or permission to manufacture, use, or sell any patented invention that may in any way be related thereto.

Qualified requesters may obtain copies of this report from the Defense Documentation Center, (DDC), Cameron Station, Alexandria, Virginia.
(Reproduction in whole or in part is permitted for any purpose of the U.S. Gov't)

Distribution of this document is unlimited

Copies of ARL Technical Documentary Reports should not be returned to Aerospace Research Laboratories unless return is required by security considerations, contractual obligations or notices on a specified document.

| | | | | |
|---|------------------------------|--------------|------|------|
| 1 | DEFENSE DOCUMENTATION CENTER | REPRODUCTION | DATE | 1968 |
| 1 | DEFENSE DOCUMENTATION CENTER | REPRODUCTION | DATE | 1968 |
| 1 | DEFENSE DOCUMENTATION CENTER | REPRODUCTION | DATE | 1968 |
| 1 | DEFENSE DOCUMENTATION CENTER | REPRODUCTION | DATE | 1968 |

**A NOTE ON RECURRENCE RELATIONS BETWEEN EXPECTED
 VALUES OF FUNCTIONS OF ORDER STATISTICS**

By P. R. KRISHNAIAH AND M. HASEEB RIZVI¹

Aerosol Research Laboratories, Wright-Patterson AFB

1. Summary. In this note, some recurrence relations are derived between the expected values of functions of order statistics from any arbitrary distribution with continuous cumulative distribution function (cdf). These recurrence relations are closely related to some results obtained by Srikantan [2].

2. Some recurrence relations. Suppose X has an arbitrary distribution with a continuous cdf $F(x)$ and $h(\cdot)$ is a specified function such that $Eh(X)$ exists. Let $X_{k,n}$ denote the k th ($1 \leq k \leq n$) order statistic in a random sample of size n from the distribution with cdf $F(x)$. Then we have the following theorem:

THEOREM. For $1 \leq k \leq m \leq n$,

$$(2.1) \quad E\{h(X_{k,m})\} = \binom{m}{k} \sum_{i=0}^k (-1)^i (k/(k-i)) \binom{i}{i} / \binom{m-i+i}{k-i+i} E\{h(X_{k-i,m-i+i})\},$$

$$0 \leq i \leq k-1,$$

and also

$$(2.2) \quad E\{h(X_{k,m})\} = \binom{m}{k} \sum_{s=0}^j (-1)^s (k/(k+s)) \binom{j}{s} / \binom{m-j+s}{k+s} E\{h(X_{k+s,m-j+s})\},$$

$$0 \leq j \leq m-k.$$

PROOF. Let $G(u) = 1 - F(u)$ for all u and let \int denote the integral over the entire real line. Then for $1 \leq k \leq m \leq n$ and $0 \leq i \leq k-1$,

$$(2.3) \quad E\{h(X_{k,m})\} = k \binom{m}{k} \int h(u) \{1 - G(u)\}^i \{1 - G(u)\}^{k-i-1} G^{m-k}(u) dF(u)$$

$$= k \binom{m}{k} \sum_{i=0}^k (-1)^i \binom{i}{i} \int h(u) F^{k-i-1}(u) \{1 - F(u)\}^{m-k+i} dF(u)$$

$$= \binom{m}{k} \sum_{i=0}^k (-1)^i (k/(k-i)) \binom{i}{i} / \binom{m-i+i}{k-i+i} E\{h(X_{k-i,m-i+i})\}.$$

Also, for $1 \leq k \leq m \leq n$ and $0 \leq j \leq m-k$,

$$(2.4) \quad E\{h(X_{k,m})\} = k \binom{m}{k} \int h(u) F^{k-1}(u) \{1 - F(u)\}^j \{1 - F(u)\}^{m-k-j} dF(u)$$

$$= k \binom{m}{k} \sum_{s=0}^j (-1)^s \binom{j}{s} \int h(u) F^{k+s-1}(u) \{1 - F(u)\}^{m-k-j} dF(u)$$

$$= \binom{m}{k} \sum_{s=0}^j (-1)^s (k/(k+s)) \binom{j}{s} / \binom{m-j+s}{k+s} E\{h(X_{k+s,m-j+s})\}.$$

It should be pointed out that formulae (5) and (6) of Srikantan [2] are equivalent respectively to (2.1) with $i = k-1$ and (2.2) with $j = m-k$. The recurrence relations between the moments, between the moment generating functions and between the characteristic functions of the order statistics can be obtained

Received 22 June 1964; revised 15 September 1965.

¹ Now at Ohio State University.

from (2.1) and (2.2) by setting $h(u) = u^r$ ($r = 1, 2, \dots$), $h(u) = \exp\{tu\}$, (t real) and $h(u) = \exp\{(-1)^j tu\}$, (t real) respectively. Similarly by letting $h(u) = 1$ if $u \in (-\infty, x)$ and $h(u) = 0$ otherwise, (2.1) and (2.2) yield recurrence relations between cdf's of order statistics. These in turn lead to results connecting pdf's (provided densities exist). Srikantan [2] makes some similar remarks for his results. He also discusses the interrelation between the results like (2.1) and (2.2) and those available in the literature. For the problem of gamma order statistics, (2.2) was proved in [1] when $h(u) = u^r$.

REFERENCES

- [1] GUPTA, S. S. (1960). Order statistics from the gamma distribution. *Technometrics* **2** 243-262.
- [2] SRIKANTAN, K. S. (1962). Recurrence relations between the pdf's of order statistics, and some applications. *Ann. Math. Statist.* **33** 169-177.