۱Y

CH'

AD 64761

# RESEARCH INVESTIGATION OF LASER LINE PROFILES

## REPORT F-920479-2 SEMI-ANNUAL REPORT



PERIOD COVERED: 1 AUGU 1 1966 TO 31 JANUARY 1967

PREPARED UNDER CONTRACT N00014-66-CO344 ARPA ORDER NO. 306, PROJECT COST CODE NO. 6E30K21

United Aircraft Research Laboratories



EAST HARTFORD, CONNECTICUT

ABGEINVE USUPY



UNITED AIRCRAFT CORPORATION RESEARCH LABORATORIES East Hartford, Connecticut

F-920479-2 Semiannual Report Under Contract NO0014-66-C0344 1 August 1966 through 31 January 1967

Project Title: RESEARCH INVESTIGATION OF LASER LI PROFILES Name of Contractor: United Aircraft Corporation Research Laboratories Project Code No. 6E30K21 ARPA Order No 306

This research is part of project DEFENDER under the joint sponsorship of the Advanced Research Projects Agency, the Office of Naval Research, and the Department of Defense.

> Reported By: <u>A. L. Lamb</u>, Jr. G. L. Lamb, Jr. Senior Theoretical Physicist

tetur

D. A. Stetser Research Scientist

J. De Maria J. DeMaria

Principal Scientist, Quantum Physics

Approved By: X H. D. Taylor.

Manager, Physics Department

Date: February 27, 1967

"Reproduction in whole or in part is permitted for any purpose of the United States Government."

## Report F-920479-2

Semiannual Report Under Contract NOOO14-66-CO344 for the Period 1 August 1966 through 31 January 1967

## RESEARCH INVESTIGATION OF LASER LINE PROFILES

ARPA Order No. 306, Project Code No. 6E30K21

## TABLE OF CONTENTS

	rage
SUMMARY	. 1
INTRODUCTION	. 2
KINETIC THEORY FOR AN ION LASER	. 2
Introduction	. 2 . 3
The Polarization of the Medium and Reduced Distribution Functions	. 6 . 9
Wigner Distribution Function	11
Long Range Forces	12 14 21 23
MEASUREMENTS OF ARGON ION LASER LINE PROFILE	28
Introduction	28 28
SIX-MONTH STATUS EVALUATION	30
REFERENCES	31
LIST OF FIGURES	32
FIGURES	

DISTRIBUTION LIST

Highly Street

#### Report F-920479-2

Semiannual Report Under Contract NO0014-66-C0344 for the Period 1 August 1966 through 31 January 1967

Research Investigation of Laser Line Profiles

ARPA Order No. 306, Project Code No. 6E30K21

#### SUMMARY

Some aspects of the theory of a gas laser recently developed by W. E. Lamb, Jr. are recast in a form which more fully displays the role played by the particle dynamics. The Wigner distribution function is used to derive kinetic equations which govern the external center of mass motion of the two-level systems as well as their internal dynamics. The effect of long range forces is discussed by treating the collision integral in a manner similar to that employed in plasma kinetic theory. A modification in the criterion for the existence of a dip in the output is obtained. It is also shown that effects due to long range forces are most noticeable at long optical wavelengths and when there is a large difference between the lifetimes of the two laser levels.

The experimental system for measuring the line profile of a dc excited argon laser has been designed, constructed, and made operative. Preliminary data on the "Lamb Dip" of an argon ion laser has been obtained as a function of pressure and excitation. The results of the experimental portion of this program and attempts to interpret the experiments in terms of this theory will be reported in later reports.

60

#### INTRODUCTION

The objective of this program is to conduct an experimental and theoretical investigation of laser line profiles. The theoretical analysis will be based on a discussion of many particle distribution functions which provide a single unified theory for treating both the long range nature of the coulomb forces and the quantum mechanical interaction of the two-level system with the optical field. The effects of such long range forces on the steady state conditions in a single mode gas laser will be determined. The experimental program will determine the homogeneous line width in an argon ion laser and relate experimental effects to the prediction from the theoretical portion of the investigation.

#### KINETIC THEORY FOR AN ION LASER

#### Introduction

A satisfactory description of the operating conditions for a gas laser including effects due to rectilinear particle motion is presently available. <sup>(1)</sup> A modification of this theory to include the effect of short range collisions has also been developed. <sup>(2)</sup> However, a theoretical framework permitting the systematic calculation of corrections associated with particle interactions has yet to be provided. The purpose of this work is to show that such refinements appear quite naturally when the dynamics of the laser medium is described in terms of the Wigner distribution function.

Following the technique which has been employed recently in both classical and quantum plasmas, the Liouville equation

-2-

for an N-particle Wigner distribution function is first obtained. This distribution function contains coordinates describing both the external center of mass motion of each particle (and is treated classically) as well as the internal motion which is quantized. From the Liouville equation, an equation for a one particle distribution function is obtained which contains an integral over a two particle function. This integral contains all effects associated with particle interactions and is exact. To treat the long range coulomb forces that are present in the ionized gas laser, the collision integral is treated by methods developed recently in classical plasma physics. Implicit in this treatment is the standard assumption of plasma physics that one is dealing with a completely ionized system so that short range collisions among neutral atoms may be neglected. Unfortunately, this idealization is not realizable in practice. The effect of long range collisions on the steady state conditions of an ion laser operating in a single mode is then derived from a third order perturbation analysis.

#### The N-Particle Wigner Distribution Function

We consider a cavity of volume V containing N two-level systems interacting with an electromagnetic field. Each particle is assumed to have a mass m and the velocity distributions of particles in each of the two levels are assumed to be Maxwellian at the same temperature T. The i<sup>th</sup> such system has internal coordinates  $p_i$ ,  $q_i$ , and external center of mass coordinates  $P_i$ ,  $Q_i$ .

The Hamiltonian operator governing the dynamics of the N two-level systems is assumed to be composed of four types of energy operators. They are the kinetic energy operator  $K_i$  for the center of mass motion of the i<sup>th</sup> particle,  $H_i$  the operator associated with the internal energy of an isolated two-level system,  $V_i$  the

-3-

the potential energy of a particle interacting with the electromagnetic field and finally,  $V_{ij}$  the two-body potential expressing the interaction between pairs of two-level systems in the cavity. In actual laser systems, the scattering will be predominantly with particles other than those in either of the two laser levels. Hence, to describe the scattering, it will be neces ary to consider a three-level system. Since optical transitions to this third level will be ignored, only a minor extension of the two-level analysis will be required.

The Hamiltonian is, then,

$$\mathcal{F} = \sum_{i=1}^{N} (\kappa_i + H_i + V_i) + \sum_{i,j} V_{ij}$$
 (1)

¥,

The two-body potential affects the internal dynamics of the two-level systems as well as their translational motion. It will be assumed that these two effects are additive so that one may decompose the potential into the form

$$\sum_{i,j} V_{ij} = \sum_{i < j} V_{ij} (IQ_i - Q_j I) + \sum_{i,j} W_{ij} (q_i, IQ_i - Q_j I) , \qquad (2)$$

where  $U_{ij}$  is the two-body potential influencing the center of mass motion and  $W_{ij}$  represents the effect of the external field of the j<sup>th</sup> particle upon the internal dynamics of the i<sup>th</sup> particle.

The first and third types of terms in the Hamiltonian are, respectively,

$$K_{i} = -\frac{\hbar^{2}}{2m} \frac{\partial^{2}}{\partial Q_{i}^{2}}$$
(3)

$$V_i = eE(Q_i, t) \cdot q_i$$
(4)

The operator  $H_i$  and the second term in Eq. (2) are defined in terms of the Schroedinger equation

$$(H_i + \sum_j W_{ij})\psi_i(q_i, t) = i + \frac{\partial}{\partial t}\psi(q_i, t) .$$

$$(5)$$

The spatial dependence of  $W_{ij}$  may be replaced by a stochastic time dependence. Furthermore, since the time required for a collision to take place  $(10^{-13} \text{ sec})$  is much longer than the period of the optical field  $(10^{-15} \text{ sec})$ , Eq. (5) may be solved in the adiabatic approximation yielding wave functions of the form

$$\psi_{in}(q_i,t) = \chi_{in}(q_i) \exp\left[-i\widetilde{E}_n t/\hbar - (i/\hbar) \int_{-\infty}^{t} dt' \Delta E_n(t')\right], \quad n = a,b, \quad (6)$$

where

$$\Delta E_n(i) = \Delta E_n(Q_i, \dots Q_N) = \sum_j W_{ij} , \qquad (7)$$

and the x in and  $\tilde{E}_n$  are defined by the eigenvalue equation

$$H_1 X_{in} = \widetilde{E}_n X_{in}$$
 (8)

The constants <u>a</u> and <u>b</u> refer, respectively, to the upper and lower levels of an individual two-level system. It is assumed that the  $\tilde{E}_n$  contain an imaginary part which accounts for decay to the ground state labelled by the subscript <u>c</u>, i.e.,

$$\widetilde{\mathsf{E}}_{\mathsf{n}} = \mathsf{E}_{\mathsf{n}} - i\hbar \gamma_{\mathsf{n}}/2 \,. \tag{9}$$

As a result of this damping, the number of particles in states <u>a</u> and <u>b</u> decreases with increasing time. It will be assumed that these excited states are replenished by an external pumping mechanism to be introduced later.

$$\mathcal{P}\Psi(\eta,t) = i\hbar \frac{\partial \Psi}{\partial t}$$
, (10)

where  $\eta$  represents the set of all position coordinates  $\eta = (Q,q) = (Q_1, \ldots, Q_N, q_1, \ldots, q_N)$ . The N-particle Wigner distribution function may now be introduced through the definition

$$f_{N}(\xi,\eta,t) = \int \frac{d^{6N}\lambda}{(2\pi)^{6N}} e^{-i\lambda\cdot\xi} \Psi^{*}(\eta-\frac{\hbar}{2}\lambda,t)\Psi(\eta+\frac{\hbar}{2}\lambda,t), \quad (11)$$

-5-

where  $\xi$  represents the set of all momentum coordinates  $\xi = (P,p) = (\sum_1, \dots, P_N, p_1, \dots, p_N)$ , and the 6N dimensional transform variable  $\lambda$  has components  $(T_1, \dots, T_N, \tau_1, \dots, \tau_N)$  corresponding to the 3N external and 3N internal coordinates respectively.

The Polarization of the Medium and Reduced Distribution Functions

As a result of using the Wigner distribution function, quantum statistical averages are obtained from expressions closely resembling those used in classical statistical mechanics. In particular, the polarization density of the medium due to the i<sup>th</sup> two-level system is

$$\mathcal{P}_{i}(Q_{i},t) = \int d^{3(N-1)}Q \int d^{3N}p \int d^{3N}p \int d^{3N}p (-eq_{i}) f_{n}(\xi,\eta,t)$$
(12)  
=  $-\frac{e}{\sqrt{2}} \int d^{3}p_{i} d^{3}p_{i} d^{3}q_{i} q_{i} f_{i}(\xi_{i},\eta_{i},t) ,$ (13)

where  $f_1$  is the reduced distribution function obtained by integrating  $f_N$  over all phase space except that of the i<sup>th</sup> particle, i.e.,

$$f_{1}(\xi_{1},\eta_{1},t) \equiv \bigvee \int d^{6(N-1)} \xi \, d^{6(N-1)} \eta \, f_{N}(\xi,\eta,t) \quad . \tag{14}$$

For later use, a two-particle reduced distribution function for a pair of particles i and j is defined in a similar way as

$$f_{2}(\xi_{i},\xi_{j},\eta_{i},\eta_{j},t) \equiv \sqrt{2} \int d^{6(N-2)} \xi \, d^{6(N-2)} \eta \, f_{N}(\xi,\eta,t) \,. \tag{15}$$

The one-particle Wigner distribution function also has a representation in terms of one-particle wave functions as

$$f_{1}(\xi_{1},\eta_{1},t) = \sqrt{\int} \frac{d^{6} \lambda}{(2\pi)^{6}} e^{-i\lambda \cdot \xi_{1}} \psi_{1}^{*}(\eta_{1} - \frac{\hbar}{2} \lambda_{1},t) \psi(\eta_{1} + \frac{\hbar}{2} \lambda_{1},t) , \quad (16)$$

-6-

where

$$\psi_{1}(\eta_{1}, t) = \sum_{\nu, n} a_{\nu n}(t) \phi_{\nu}(Q_{1}, t) \chi_{n}(q_{1}). \qquad (17)$$

The sum on <u>n</u> extends over <u>a</u>, <u>b</u> and <u>c</u> while that on v is actually over the continuous values of translational momentum. With the expansion given in Eq. (17), the one-particle Wigner distribution function may be written

$$f_{1}(\xi_{1},\eta_{1},t) = \sum_{m,n} F_{lnm}(P_{1},Q_{1},t) f_{lmn}(p_{1},q_{1},t), \quad m,n = \pi, \Sigma_{\eta}$$
(18)

where

$$F_{\rm inm}(\tilde{P}_{1},Q_{1},t) = \bigvee \sum_{\mu,\nu} \sigma_{\mu m}^{*}(t) q_{\nu n}(t) \int \frac{d^{3}T_{1}}{(2\pi)^{3}} e^{-iT_{1}\cdot P_{1}} \phi_{\mu}^{*}(Q_{1}-\frac{\hbar}{2}T_{1},t) \phi_{\nu}(Q_{1}+\frac{\hbar}{2}T_{1},t) \qquad (15)$$

$$f_{\rm inn}(p_{1},q_{1},t) = \int \frac{d^{3}\tau_{1}}{(2\pi)^{3}} e^{-i\tau_{1}\cdot p_{1}} \chi_{m}^{*}(q_{1}-\frac{\hbar}{2}\tau) \chi_{n}(q_{1}+\frac{\hbar}{2}\tau) \qquad (20)$$

From these definitions it follows that

$$F_{\text{inm}} = F_{\text{imn}}^{*}$$
(21)  
$$f_{\text{inm}} = f_{\text{imn}}^{*}$$

Now  $P_i(Q_i,t)$  is the polarization density due to the i<sup>th</sup> particle at point  $Q_i$  in the cavity. The polarization density at some point Q in the cavity due to the entire assemblage of N two-level systems is

$$\mathcal{P}(Q,t) = \sum_{i=1}^{N} \mathcal{P}_{1}(Q_{i},t) \Big|_{Q_{i}=Q} = -n^{(0)} e \int d^{3}P d^{3}p d^{3}q q f_{1}(P,Q,p,q,t), \quad (22)$$

where  $n^{(0)}$  is the sum of the average number density of particles in all levels, i.e.,  $n^{(0)} = n_a^{(0)} + n_b^{(0)} + n_c^{(0)}$ .

With the form of  $f_1$  given in Eq. (18), the macroscopic

-7-

polarization density becomes

$$\mathcal{C}(Q,t) = -n^{(0)} \left[ \mathscr{B}_{ba} \int d^{3} PF_{lab}(P,Q,t) + \mathscr{E}_{ca} \int d^{3} PF_{lab}(P,Q,t) + \mathscr{E}_{cb} \int d^{3} PF_{lab}(P,Q,t) \right] + c.c., (23)$$

where

$$\mathscr{C}_{mn} = \vartheta \int d^3 q \chi^{*}_{m}(q) q \chi_{n}(q) = \mathscr{C}^{*}_{nm} . \qquad (24)$$

In obtaining Eq. (23) it has been assumed that the off diagonal matrix elements are real and hence equal. Also, it is assumed that there are no permanent dipole moments so that the diagonal matrix elements vanish.

The optical field and the associated polarization of the medium may be expanded in terms of the eigenmodes of the cavity  $U_n(Q)$  which satisfy a vector Helmholtz equation.  $(\frac{1}{2})$  One then has

$$E(Q,t) = \sum_{n} E_{n}(t) U_{n}(Q)$$

$$\mathcal{P}(Q,t) = \sum_{n} \mathcal{P}_{n}(t) U_{n}(Q) .$$
(25)

Assuming that the optical field in the cavity varies in only one direction (labeled by a Z axis) and that the length of the cavity in this direction is L, a set of eigenfunctions adequate for the present work are

$$U_n(Q) = e \sin k_n Z$$
, (26)

where e is a unit vector perpendicular to the Z axis,

$$k_n = n\pi/L$$
, (27)

and n is a large positive integer. It will also be assumed that the polarization of the individual two-level system takes place along e so the q = eq| and  $\mathscr{C}_{mn} = e|\mathscr{C}_{mn}|$ . Finally, since we are only concerned with the contribution of level <u>c</u> to the particle dynamics and not to laser action between levels <u>a</u> and <u>c</u> or <u>b</u> and <u>c</u>,

-8-

the dipole moments  $\varphi_{ac}$  and  $\varphi_{bc}$  will be set equal to zero. From Eqs. (23) and (25) we then have

$$\mathcal{P}_{n}(t) = - \mathcal{E}n^{(0)} \frac{2}{L} \int_{0}^{L} dZ U_{n}(Z) \int d^{3}P F_{1ab}(P,Q,t) + c.c.,$$
 (28)

where  $\varphi = \varphi_{ab} = \varphi_{ba}$ .

As has been shown,  $(\frac{1}{2})$  the steady state operating conditions for the laser may be obtained once  $\mathcal{P}_n(t)$  is known as a function of the cavity field beyond the linear approximation.

#### Quantum Mechanical Liouville Equation

The time derivative of Eq. (11) along with the Schroedinger equation (10) lead to

$$\frac{\partial f_{\rm N}}{\partial t} = \frac{i}{i\hbar} \int \frac{d^{6} \lambda}{(2\pi)^{6} N} e^{-i\lambda\xi} \left[ \mathcal{H}(\eta + \frac{\hbar}{2}\lambda) - \mathcal{H}(\eta - \frac{\hbar}{2}\lambda) \right] \Psi^{*}(\eta - \frac{\hbar}{2}\lambda, t) \Psi(\eta + \frac{\hbar}{2}\lambda, t) .$$
(29)

The contribution to  $\partial f_N / \partial t$  which is made by each of the four types of terms in  $\frac{1}{N}$  will now be considered separately.

The kinetic energy term given in Eq. (13) is readily shown (3) to contribute the convective term

$$\frac{\partial f_{N}}{\partial t}\Big|_{I} = -\frac{1}{m}\sum_{i} P_{i} \cdot \frac{\partial f_{N}}{\partial Q_{i}} \qquad (30)$$

The contribution from the internal dynamics will merely be rewritten in the form

$$\frac{\partial f_{N}}{\partial t}\Big|_{\Pi} = \frac{1}{ih} \int \frac{d^{6N}\lambda}{(2\pi)^{6N}} \int d^{6N} \xi e^{-i(\xi \cdot \xi)\lambda} \Big[ \sum_{i} (H_{i} + \sum_{j} W_{ij})_{+} - \sum_{i} (H_{i} + \sum_{j} W_{ij})_{-} \Big] f_{N}(\xi, \eta, t), \quad (31)$$

where the subscripts ± refer to the displaced arguments of the type introduced in Eq. (29), and use has been made of the inverse Fourier

transform relation associated with Eq. (11). The rate of change of  $f_N$  resulting from the third term in  $\frac{4}{3}$  is

$$\frac{\partial f_{N}}{\partial t} \Big|_{\mathbf{III}} = i \mathbf{e} \sum_{i} \mathbf{E}(\mathbf{Q}_{i}, t) \cdot \int \frac{d^{\mathbf{e}N} \lambda}{(2\pi)} \mathbf{e}^{-i\lambda \cdot \mathbf{\zeta}} \tau_{i} \Psi^{*}(\eta - \frac{\hbar}{2}\lambda, t) \Psi(\eta + \frac{\hbar}{2}\lambda, t)$$

$$= \mathbf{e} \sum_{i} \mathbf{E}(\mathbf{Q}_{i}, t) \cdot \frac{\partial f_{N}}{\partial \mathbf{p}_{i}} .$$
(32)

ā.

Finally, the part of the Hamiltonian containing the interaction due to translational motion leads to

$$\frac{\partial f_{N}}{\partial t} \Big|_{III} = \frac{I}{ih} \int \frac{d^{6N} \lambda}{(2\pi)^{6N}} \int d^{6N} \xi e^{-i(\xi-\zeta)\cdot\lambda} \sum_{i < j} \left[ U_{ij}(IQ_{i} + \frac{\hbar}{2}T_{i} - Q_{j} - \frac{\hbar}{2}T_{j}I) - U_{ij}(IQ_{i} - \frac{\hbar}{2}T_{i} - Q_{j} + \frac{\hbar}{2}T_{j}I) \right] f_{N}(\xi, \eta, t) .$$

$$(33)$$

The center of mass motion will be treated classically. This limit is readily obtained by expanding the integrand in Eq. (33) in powers of Planck's constant and retaining only the first nonvanishing term. One recovers the standard expression of kinetic theory,

$$\frac{\partial f_{N}}{\partial t} \bigg|_{III} = \sum_{i < j} \left( \frac{\partial U_{ij}}{\partial Q_{i}} \cdot \frac{\partial}{\partial P_{i}} - \frac{\partial U_{ij}}{\partial Q_{j}} \cdot \frac{\partial}{\partial P_{j}} \right) f_{N}(\zeta, \eta, t) .$$
(34)

As mentioned previously,  $f_N$  also changes due to an external pumping process which replenishes the excited states <u>a</u> and <u>b</u>. This change in  $f_N$  will be accounted for by including a phenomenological source term  $\mathcal{A}(\xi,\eta,t)$  in the quantum Liouville equation satisfied by  $f_N$ . The combination of Eqs. (30), (31), (32) and (34) and the source term yields the quantum Liouville equation in the form

$$\frac{\partial f_{N}}{\partial t} + \frac{1}{m} \sum_{i=1}^{n} P_{i} \cdot \frac{\partial f_{N}}{\partial Q} + \frac{1}{ih} \int \frac{d^{6N} \lambda}{(2\pi)^{6N}} \int d^{6N} \sum_{j=1}^{n} (\xi,\zeta) \lambda \left[ \sum_{i=1}^{n} (H_{i} + \sum_{j=1}^{n} W_{ij}) + (35) \right]$$

$$(35)$$

$$\sum_{i=1}^{n} (H_{i} + \sum_{j=1}^{n} W_{ij}) = \int_{i=1}^{n} f_{N} (\xi,\eta,t) - e \sum_{i=1}^{n} E(Q_{n},t) \cdot \frac{\partial f_{N}}{\partial P_{n}} = \sum_{i=1}^{n} \left( \frac{\partial U_{ij}}{\partial Q_{i}} \cdot \frac{\partial}{\partial P_{i}} - \frac{\partial U_{ij}}{\partial Q_{j}} \cdot \frac{\partial}{\partial P_{j}} \right) f_{N} + \mathcal{Q}(\xi,\eta,t) .$$

-10-

## Kinetic Equation Satisfied by the One-Particle Wigner Distribution Function

It has been shown previously that the physically interesting quantity, the macroscopic polarization density, may be expressed in terms of a one-particle Wigner distribution function. One obtains the equation determining this function by integrating Eq. (35) over the phase space of all particles except one. Employing the definitions given in Eqs. (19) and (20) this procedure yields

$$\frac{\partial f_{1}}{\partial t} + \frac{P_{1}}{m} \cdot \frac{\partial f_{1}}{\partial Q_{1}} + \sum_{n,m=a,b,c} (i \omega_{n,m} + \gamma_{n,m}) F_{lnm} f_{lmn} - e E(Q_{1},t) \cdot \frac{\partial f_{1}}{\partial p}$$

$$= n^{(0)} \int d^{6} \xi_{2} d^{6} \eta_{2} \frac{\partial U_{12}}{\partial Q_{1}} \cdot \frac{\partial f_{12}}{\partial P_{1}} + \mathcal{Q}_{1}(\xi_{1},\eta_{1},t) , \qquad (36)$$

where

$$\mathcal{Q}_{1}(\xi_{1},\eta_{1},t) = \int d^{6(N-1)} \xi d^{6(N-1)} \eta \mathcal{Q}(\xi,\eta,t),$$
 (37)

and

$$\gamma_{mn} = \gamma_{nm} = (\gamma_m + \gamma_n)/2$$

$$\omega_{mn} = (\varepsilon_m - \varepsilon_n)/\hbar .$$
(38)

It will prove useful to rewrite the term containing the cyternal field by using the identity

$$eE(Q_{I},t)\cdot\frac{\partial f_{I}}{\partial p_{I}} = -ieE(Q_{I},t)\cdot\int\frac{d^{6}\lambda_{I}}{(2\pi)^{6}}e^{-i\lambda_{I}\cdot\xi_{I}}\tau_{I}\psi_{I}^{*}(\eta_{I}-\frac{\hbar}{2}\lambda_{I},t)\psi(\eta_{I}+\frac{\hbar}{2}\lambda_{I},t), \quad (39)$$

where  $d^6\lambda_1 = d^3T_1 d^3\tau_1$ . In terms of the functions  $F_{1nm}$  introduced previously,

$$e E(Q_{I},t) \cdot \frac{\partial f_{I}}{\partial p_{I}} = -ie E(Q_{I},t) \cdot \int \frac{d^{3}\tau}{(2\pi)^{3}} e^{-i\tau_{I} \cdot p_{I}} \tau_{I} \sum F_{Inm} \chi_{m}^{*} (q_{I} - \frac{\hbar}{2}\tau_{I}) \chi_{n} (q_{I} + \frac{\hbar}{2}\tau_{I}) \cdot (40)$$

Modification of the Collision Term for Long Range Forces

The one term in Eq. (36) requiring further elucidation is the collision integral. Since the distribution function  $f_2$ satisfies an equation containing integrals over  $f_3$  which describes three particle effects etc., a means of closing this hierarchy must be introduced. Long-range forces may be treated by truncating Eq. (36) in a manner similar to that employed in plasma kinetic theory. Neglecting particle correlation entirely and writing

$$f_{2}(\xi_{1},\xi_{2},\eta_{1},\eta_{2},t) = f_{1}(\xi_{1},\eta_{1},t) f_{1}(\xi_{2},\eta_{2},t) , \qquad (41)$$

a pair of Vlasov-like equations may be derived for the distribution functions  $F_{aa}$  and  $F_{ab}$  introduced previously. It should be emphasized that no expansion in any plasma parameter is employed; spatial inhomogeneity in the assemblage of two-level systems is assumed to be due solely to the presence of the externally applied optical field. One finds that the introduction of Eq. (40) into Eq. (36) yields

$$\frac{\partial f_{I}}{\partial t} + \frac{P_{I}}{m} \cdot \frac{\partial f_{I}}{\partial Q_{I}} + \sum_{n,m=a,b,c} (i\omega_{nm} + \gamma_{nm}) F_{nm} f_{mn} - eE \cdot \frac{\partial f_{I}}{\partial P_{I}} + eE \cdot \frac{\partial f_{I}}{\partial P_{I}} = \mathcal{Q}(P_{I},Q_{I},t), \quad (42)$$

where

$$e \mathcal{E} = -n^{(0)} \frac{\partial}{\partial Q_1} \int d^3 p_2 d^3 Q_2 U_{12} \left[ F_{aa} + F_{bb} + F_{cc} \right] .$$
 (43)

Introducing a potential through the definition

$$\mathcal{E} = -\frac{\partial \Phi}{\partial Q_1} \tag{44}$$

-1.2-

F-9 0479-2

Name of Street

and employing the orthogonality properties of the fij, namely,

$$(2\pi\hbar)^{3}\int d^{3}pd^{3}qf_{ij}(p,q)f_{ab}(p,q) = \delta_{ja}\delta_{bi}$$
, (45)

one finds

$$\frac{\partial F_{aa}}{\partial t} + \frac{P_{I}}{m} \cdot \frac{\partial F_{aa}}{\partial Q_{I}} + \gamma_{a} F_{aa} - i \frac{E \cdot \delta^{a} ab}{\hbar} (F_{ab} - F_{ba}) - e \frac{\partial \phi}{\partial Q_{I}} \cdot \frac{\partial F_{aa}}{\partial P_{I}} = \lambda_{a} \quad (46)$$

$$\frac{\partial F_{ab}}{\partial t} + \frac{P_i}{m} \cdot \frac{\partial F_{ab}}{\partial Q_i} + i\Omega_{ab}F_{ab} - i\frac{E \cdot \delta^2 ab}{\hbar} (F_{aa} - F_{bb}) - e\frac{\partial \phi}{\partial Q_i} \cdot \frac{\partial F_{ab}}{\partial P_i} = 0 \qquad (47)$$

$$\frac{\partial F_{cc}}{\partial t} + \frac{P_{I}}{m} \cdot \frac{\partial F_{cc}}{\partial Q_{I}} - e \frac{\partial \phi}{\partial Q_{I}} \cdot \frac{\partial F_{cc}}{\partial P_{I}} = 0 , \qquad (48)$$

where

$$\lambda_{a}(P_{1},Q_{1},t) \equiv (2\pi h)^{3} \int d^{3}p_{1}d^{3}q_{1}f_{1ca} \mathcal{Q}_{1}(\xi_{1},\eta_{1},t) , \qquad (19)$$

When these equations are expanded to third order in the optical field strength, it is found that the equations governing  $F_{aa}^{(1)}$  and  $F_{ab}^{(1)}$  are the same as those in the collisionless theory, namely,

$$\frac{\partial F_{aa}^{(0)}}{\partial t} + \frac{P}{m} \cdot \frac{\partial F_{aa}^{(0)}}{\partial Q} + \gamma_{a} F_{aa}^{(0)} = \lambda_{a} (P,Q,t)$$
(50)

$$\frac{\partial F_{ab}^{(1)}}{\partial t} + \frac{P}{m} \frac{\partial F_{ab}^{(1)}}{\partial Q} + i \Omega_{ab} F_{ab}^{(1)} = \frac{i}{\hbar} E \cdot \mathscr{B}_{ab} \left[ F_{au}^{(0)} - F_{bb}^{(0)} \right]$$
(51)

$$\frac{\partial F_{aa}^{(2)}}{\partial t} + \frac{P}{m} \cdot \frac{\partial F_{aa}^{(2)}}{\partial Q} + \gamma_{a} F_{aa}^{(2)} = \frac{i}{\hbar} E \cdot \delta_{ab}^{2} \left[ F_{ab}^{(1)} - F_{ba}^{(1)} \right]$$
(52)

-13-

----

Pake 4- -----

$$\frac{\partial F_{ab}^{(3)}}{\partial t} + \frac{P}{n} \cdot \frac{\partial F_{ab}^{(3)}}{\partial Q} + i\Omega_{ab}F_{ab}^{(3)} = \frac{i}{\hbar} E \cdot \mathscr{C}_{ab} \left[ F_{ac}^{(2)} - F_{bb}^{(2)} \right], \quad (53)$$

where

$$\Omega_{ab} = \omega_{ab} - i \gamma_{ab} . \qquad (54)$$

1

One also obtains

$$\frac{\partial F_{oa}^{(2)}}{\partial t} + \frac{P_{I}}{m} \cdot \frac{\partial F_{aa}^{(2)}}{\partial Q_{I}} + \gamma_{a} F_{aa}^{(2)} = \frac{\partial \phi}{\partial Q_{I}} \cdot \frac{\partial F_{aa}^{(0)}}{\partial P_{I}} = \frac{iE \cdot \delta^{2} ab}{\hbar} \left[ F_{ab}^{(1)} - F_{ba}^{(1)} \right]$$
(55)

$$\frac{\partial F_{ab}}{\partial t} + \frac{P_{I}}{m} \cdot \frac{\partial F_{ab}}{\partial Q_{I}} + i\Omega_{ab}F_{ab} = \frac{iE \cdot \delta_{ab}}{\hbar} \left[ F_{aa}^{(2)} - F_{bb}^{(2)} \right] + e \frac{\partial \phi^{(2)}}{\partial Q_{I}} \cdot \frac{\partial F_{ab}^{(1)}}{\partial P_{I}} , (56)$$

It is seen that  $F_{aa}^{(2)}$  is goverened by an inhomogeneous Vlasov equation and hence may be obtained through the use of standard integral transform techniques.

## A Review of Collisionless Results

Before considering the effect of collisions, the collisionless results of Ref. (1) will first be recovered. From Eq. (50) one obtains

$$F_{aa}^{(00)}(P,Q,t) = \int_{0}^{\infty} d\tau e^{-\gamma_{a}\tau} \lambda_{a} (P,Q-P\tau/m,t-\tau) . \qquad (57)$$

Assuming that

$$\lambda_{\alpha}(P,Q,t) = \Lambda_{\alpha} \Phi^{(0)}(P) , \qquad (58)$$

where  $\Lambda_{a}$  is constant or varies slowly compared to the other length

and time scales of interest in the problem, one may write

$$F_{aa}^{(00)}(P,Q,t) = \Lambda_{a} \Phi^{(0)}(P) / \gamma_{a}$$
 (59)

For use in Eq. (51), it is useful to write

$$F_{aa}^{(00)} - F_{bb}^{(00)} = \Phi^{(0)} N / n^{(0)}$$
(60)

where

$$N \equiv \frac{\Lambda_{o}}{\gamma_{c}} - \frac{\Lambda_{b}}{\gamma_{b}}$$
(61)

and is referred to  $\binom{1}{2}$  as the "excitation density". The equation obtained by interchanging <u>a</u> and <u>b</u> in Eq. (51) is

$$F_{ab}^{(0)}(P,Q,t) - F_{ba}^{(0)}(P,Q,t) = i \delta (N/n^{(0)}) \Phi^{(0)}(P) \int_{0}^{\infty} d\tau (e^{-i\Omega_{ab}\tau} + e^{-i\Omega_{ba}\tau})$$

$$(62)$$

$$\times U_{n}(Q - P\tau/m)A_{n}(t - \tau) ,$$

where the electromagnetic field has been written

$$\mathbf{E}(\mathbf{Q}, \mathbf{t}) = \mathbf{e} \mathbf{E}_{\mathbf{n}} \mathbf{A}_{\mathbf{n}}(\mathbf{t}) \mathbf{U}_{\mathbf{n}}(\mathbf{Q}) \tag{63}$$

and

$$\delta = E_n \, \delta / \hbar \,. \tag{64}$$

Obtaining a similar solution for  $F_{aa}$  (20) -  $F_{bb}$  (20) from Eq. (52) and substituting the result into Eq. (53) leads to

$$F_{ab}^{(30)}(P,Q,t) = (i8)^{3} (N/n^{(0)}) \Phi^{(0)}(P) \sum_{c=a,b} \int_{0}^{\infty} d\tau d\tau' d\tau'' e^{-i\Omega_{ab}\tau - \gamma'_{c}\tau'} \times (e^{-i\Omega_{ab}\tau''} + e^{-i\Omega_{ba}\tau''}) A_{n}(t-\tau) A_{n}(t-\tau-\tau') A_{n}(t-\tau-\tau'-\tau'') \times U_{n}(Q-P\tau/m) U_{n} [Q-(P\tau+P\tau')/m] U_{n} [Q-(P\tau+P\tau'+P\tau'')/m].$$
(65)

-15-

From Eq. (28), the first-order contribution to the polarization is

$$\mathcal{P}_{n}^{(10)}(t) = -n^{(0)} \, \& \, \frac{2}{L} \int d^{3}P \int_{0}^{L} dZ U_{n}(Z) F_{ab}^{(10)}(P,Q,t) + c.c \quad . \tag{66}$$

Introducing  $F_{ab}^{(10)}$ , carrying out the spatial integration and neglecting the second harmonic contribution yields

$$\mathcal{O}_{n}^{(10)}(t) = -i8 \,^{\otimes} N \int d^{3}P \Phi^{(0)}(P) \int_{0}^{\infty} d\tau e^{-i\Omega_{ab}\tau} A_{n}(t-\tau) \cos k_{n}P\tau/m + c.c.$$
(67)  
=  $-i8 \,^{\otimes} N \int d\tau e^{-\Omega_{o}^{2}\tau^{2} - i\Omega_{ab}\tau} A_{n}(t-\tau) + c.c.$ 

where

$$\Omega_{o} = k_{n}p_{o}/2m \quad . \tag{68}$$

\$

Setting

$$A_n(t) = \cos(\omega_n t + \phi) \tag{69}$$

and using the rotating wave approximation (1) one obtains

$$\left(\widehat{\Gamma}_{n}^{(10)}(t)\right) = -\frac{88N}{2k_{n}\nu_{o}}e^{-1(\omega_{n}t+\phi)}Z\left(\frac{\omega_{n}-\Omega_{ab}}{k_{n}\nu_{o}}\right) + c.c.$$
(70)

where

$$Z(\zeta) \equiv i \int_{0}^{\infty} dt e^{i\zeta t - t^{2}/4} = Z_{r} + i Z_{i}$$
 (71)

In Ref. (1), it is shown that the steady state operating conditions follow from a knowledge of that part of  $\mathcal{P}_n$  that varies as  $\sin(\omega_n t + \phi)$ .

Writing

$$\mathcal{P}_{n}(t) = C_{n} \cos(\omega_{n} t + \phi) + S_{n} \sin(\omega_{n} t + \phi)$$
(72)

there results

$$S_{n}^{(10)} = -\frac{g^{2}NE_{n}}{\hbar k_{n}v_{o}} Z_{i}\left(\frac{\omega_{n}-\Omega_{ob}}{k_{n}v_{o}}\right).$$
(73)

The third-order contribution is obtained from Eqs. (28) and (65) by a similar but somewhat more lengthy procedure. The spatial integration is first carried out by employing the formula

$$\frac{2}{L} \int_{0}^{L} dZ U_{n}(Z) U_{n}(Z - \alpha_{1}) U_{n}(Z - \alpha_{2}) U_{n}(Z - \alpha_{3})$$

$$= \frac{1}{4} \left[ \cos (\alpha_{1} + \alpha_{2} - \alpha_{3}) + \cos (\alpha_{2} + \alpha_{3} - \alpha_{1}) + \cos (\alpha_{3} + \alpha_{1} - \alpha_{2}) \right] .$$
(74)

In the present case

$$a_{1} + a_{2} - a_{3} = k_{n} \cdot P(\tau - \tau^{*}) / m$$

$$a_{2} + a_{3} - a_{1} = k_{n} \cdot P(\tau + 2\tau' + \tau^{*}) / m \qquad (75)$$

$$a_{3} + a_{1} - a_{2} = k_{n} \cdot P(\tau + \tau^{*}) / m \qquad .$$

As has been pointed out,  $\binom{1}{2}$  the second and third of these terms give contributions with higher inverse powers of  $k_n v_0$  and hence in the "extreme Doppler limit" may be neglected. Carrying

-17-

out the momentum integration then yields

$$\mathcal{P}_{n}^{(30)}(t) = -\frac{1}{4} \delta^{2}(i\delta)^{3} N \sum_{c=0,b} \int_{0}^{\infty} d\tau \ d\tau' \ d\tau'' \ e^{-i\Omega_{0}b^{T} - \gamma_{c}'\tau' - \Omega_{0}(\tau - \tau'')^{2}}$$

$$(76)$$

$$x(e^{-i\Omega_{0}b^{T''}} + e^{-i\Omega_{0}b^{T''}}) A_{n}(t - \tau) A_{n}(t - \tau - \tau') A_{n}(t - \tau - \tau'') + c.c.$$

4

When the three time factors are written in exponential form, it is found that only the terms  $(1/8)e^{-i(\omega_n t + \psi) + i\omega_n(\tau + \tau'')}$  and  $(1/8)e^{-i(\omega_n t + \psi) + i\omega_n(\tau - \tau'')}$  give rise to denominators that will have a resonance when the various time integrations in Eq. (76) are performed. Retaining only these terms, Eq. (76) becomes

$$\left( P_{n}^{(30)}(t) = -\frac{1}{32} \, \mathscr{C}(i\,8)^{3} \, N \, e^{-i(\omega_{n} t + \Psi)} \sum_{c=a,b} \frac{1}{\gamma_{c}} \int_{0}^{\infty} d\tau \, d\tau' d\tau'' \, e^{-\Omega_{0}^{2}(\tau - \tau'')^{2}} \right)$$

$$\times e^{-i(\Omega_{0}b^{-}\omega_{n})\tau} \left[ e^{-i(\Omega_{0}b^{-}\omega_{n})\tau''} \div e^{-i(\Omega_{0}b_{0} + \omega_{n})\tau''} \right] + c.c.$$
(77)

$$=-\frac{1}{32} \left\{ \left(i \right)^{3} \operatorname{Ne}^{-i(\omega_{n} t + \psi)} \left( \frac{1}{\gamma_{a}} + \frac{1}{\gamma_{b}} \right) \left[ I \left( \Omega_{c}, \omega_{1}, \omega_{1} \right) + I \left( \Omega_{o}, \omega_{1}, -\omega_{1}^{*} \right) \right] + c.c.,$$

where

$$\boldsymbol{\omega}_{1} = \boldsymbol{\omega}_{n} - \boldsymbol{\Omega}_{ob}, \qquad (78)$$

and

$$I(\Omega_{0},\omega,\omega') \equiv \int_{0}^{\infty} d\tau d\tau' e^{-\Omega_{0}^{2}(\tau-t')^{2}+i\omega\tau+i\omega'\tau'}.$$
 (79)

One integration in this expression is readily performed after introducing the transformation

$$\begin{array}{l} x = \tau_1 + \tau_2 \\ y = \tau_1 - \tau_2 \end{array}$$
 (80)

The remaining integrals are expressible in terms of the function introduced in Eq. (71). One finds

$$I(\Omega_{o} \mu, \mu') = \left[ Z\left(\frac{\omega}{2\Omega_{o}}\right) + Z\left(\frac{\omega'}{2\Omega_{o}}\right) \right] / \left[ 2\Omega_{o}(\omega + \omega') \right].$$
(81)

Since  $Z(-z^*) = -Z^*(z)$ , it follows that

$$I(\Omega_{o}, \omega_{I}, \omega_{I}) = (k_{n} v_{o} \omega_{I})^{T} Z(\omega_{I} / 2 \Omega_{o}), \qquad (82)$$

and

$$I(\Omega_{0},\omega_{1},\omega_{1}) = (k_{n}v_{0}Im\omega_{1})^{-1}Z_{i}(\omega_{i}/2\Omega_{0}).$$
(83)

In terms of these results, the third-order polarization becomes

$$\mathcal{P}_{n}^{(30)}(t) = -\frac{\mathscr{C}(i8) N}{32 k_{n} v_{o}} \left(\frac{1}{\gamma_{o}} - \frac{1}{\gamma_{b}}\right) e^{-i(\omega_{n} t + \phi)} \left[\omega_{1}^{-1} Z(\omega_{1} / k_{n} v_{o}) + (I_{m} \omega_{1})^{-1} Z_{1}(\omega_{1} / k_{n} v_{o})\right].$$
(84)

From which one finds

$$S_{n}^{(30)} = \frac{\mathscr{B}^{4}E^{-3}N}{8n^{3}\gamma_{a}\gamma_{b}k_{n}v_{o}}Z_{i}(\omega_{1}/k_{n}v_{o})\left[1+\gamma_{a}b\mathcal{Z}(\omega_{n}-\omega_{a}b)\right], \qquad (85)$$

where

$$\mathcal{L}(\omega) \equiv \frac{1}{\omega^2 + \gamma_{ob}^2} . \tag{86}$$

-19-

Now the operating condition for a laser is obtained from 2 steady state solution of the equation  $(\frac{1}{2})$ 

$$\dot{\mathbf{E}}_{n} = \boldsymbol{\alpha}_{n} \mathbf{E}_{n} - \boldsymbol{\beta}_{n} \mathbf{F}_{n}^{3} \qquad (37)$$

where

$$\alpha_{n} = \frac{1}{2} \left( \frac{\omega_{n}}{Q_{L}} \right) - \frac{1}{2} \left( \frac{\omega_{n}}{\epsilon_{0}} \right) \frac{S_{n}^{(1)}}{E_{n}} , \qquad (88)$$

$$\beta_n = \frac{1}{2} \left( \frac{\omega_n}{\epsilon_0} \right) \frac{S_n^{(3)}}{E_n^3} , \qquad (89)$$

while  $Q_n$  is the Q of the n<sup>th</sup> cavity mode and  $\epsilon_o$  is the permittivity of free space.

The threshold excitation density for laser operation is obtained by setting  $\alpha_n = 0$  in Eq. (38). Neglecting effects that are  $O(\epsilon)$ , one finds that the threshold density at resonance  $(\omega_n = \omega_{ob})$  is

$$N_{T} = \frac{\epsilon_{o} h k_{n} v_{o}}{Q_{n} \mathcal{E}^{2} \sqrt{\pi}} \qquad (90)$$

Setting  $\dot{E}_n = 0$  in Eq. (87) and introducing Eqs. (73), (85) and (90), the equilibrium field amplitude is given by

$$\frac{g^{2}E_{n}^{2}}{g^{2}\pi^{2}\gamma_{a}\gamma_{b}} = \frac{1-\gamma_{i}^{-1}e^{(\omega_{n}-\omega_{ab})^{2}/(k_{n}v_{o})^{2}}}{1+\gamma_{ab}^{2}\mathcal{L}(\omega_{n}-\omega_{ab})}, \qquad (91)$$

where

$$\gamma = N/N_{\rm T} \qquad (92)$$

In obtaining this result the approximation

$$Z_i(x) = \pi^{1/2} e^{-x^2} , \qquad (93)$$

has been used in the numerator to yield the Doppler profile. In the denominator this exponential has been replaced by unity since the dip in the response curve occurs in the region  $|\omega_n - \omega_{ab}| \le \gamma_{ab}$ for which the exponential is essentially constant in the extreme Doppler limit.

Solution of Vlasov Equation for  $F_{aa}^{(2)}$ 

The solution of Eq. (55) may be carried out by introducing the transforms

$$F_{aa}^{(2)}(P,Q,t) = \sum_{\ell} e^{ik_{\ell} \cdot Q} \int \frac{dS}{2\pi i} e^{tt} f_{a\ell}^{(2)}(P,s) , \qquad (94)$$

$$\phi^{(2)}(Q,t) = \sum_{\ell} e^{ik_{\ell} \cdot Q} \int \frac{ds}{2\pi i} e^{st} \phi_{\ell}^{(2)}(s) , \qquad (95)$$

$$h_{o}^{(2)}(P,Q,t) = \sum_{\ell} e^{ik_{\ell}Q} \int \frac{ds}{2\pi i} e^{st} H_{o\ell}^{(2)}(P,s) ,$$
 (96)

where  $h_c^{(2)}(P,Q,\tau)$  is the inhomogeneous term in Eq. (55). The transformed version of Eq. (55) is

$$(s+\gamma_{\alpha}+ik_{\ell}\cdot P_{l}/m)\mathcal{F}_{\sigma\ell}^{(2)}+iek_{\ell}\cdot\frac{\partial F_{\alpha\alpha}^{(0)}}{\partial P_{l}}\dot{\phi}_{\ell}^{(2)}=H_{\sigma\ell}^{(2)}.$$
 (97)

-21-

-----

From Eq. (43) it follows that

$$\phi_{\ell}^{(2)}(s) = \frac{4\pi n_{n}^{(0)}e}{k_{\ell}^{2}} \int d^{3} P \left[ f_{\alpha\ell}^{(2)} + f_{h}^{(2)} + f_{c\ell}^{(2)} \right] .$$
(98)

Equation (97), along with a similar equation for  $F_{bl}^{(2)}$  plus a homogeneous equation for  $F_{cl}^{(2)}$  (since  $\mathscr{P}_{ac} = \mathscr{P}_{bc} = 0$ ) may now be solved simultaneously. One finds

$$f_{o_{\ell}}^{(2)}(P,s) = f_{o_{\ell}}^{(20)}(P,s) + \frac{4\pi n^{(0)}e^{2}}{k_{\ell}^{2}} \frac{n_{\ell}^{(20)}(s)}{n^{(c)}\epsilon_{\ell}(s)} \frac{ik_{\ell}}{s + \gamma_{\alpha} + ik_{\ell} \cdot P/m}, \quad (99)$$

where

$$\mathcal{F}_{a,r}^{(20)}(P,s) = \frac{H_{a,\ell}(P,s)}{s + \gamma_a + ik_r P/m} , \qquad (100)$$

and

$$n_{\ell}^{(20)}(s) = \int d^{3}P \left[ f_{0\ell}^{(20)} + f_{0\ell}^{(20)} + f_{0\ell}^{(20)} \right]$$
(101)

Also,  $\epsilon_{\mathcal{L}}(s)$  is defined by

$$\epsilon_{\ell}(s) = \epsilon_{\alpha_{\ell}}(s) + \epsilon_{b\ell}(s) + \epsilon_{c\ell}(s) - 2 , \qquad (102)$$

where

$$\epsilon_{n\ell}(s) = 1 - \frac{4\pi n_n^{(0)} e^2}{k_\ell^2} \int d^3 P \frac{i k_\ell}{s + \gamma_n + i k_\ell} \frac{\partial \Phi^{(0)}}{\partial P} , \quad n = a, b, c . \quad (103)$$

Third-Order Polarization for the Ion Laser

The third-order polarization follows in the usual way from Eq. (28),

$$\mathcal{P}_{n}^{(3)}(t) = -n^{(0)} \mathscr{E}_{ab} \frac{2}{L} \int_{0}^{L} dZ U_{n}(Z) \int d^{3}PF_{ab}^{(3)}(P,Q,t) + c.c. \qquad (104)$$

From Eq. (56) the appropriate form for  $F_{ab}^{(3)}$  is found to be

$$F_{ab}^{(3)}(P,Q,t) = (iE_{o}\cdot\delta_{ab}^{o}/\hbar)\int_{0}^{\infty} d\tau e^{-i\Omega_{ab}\tau}A_{n}(t-\tau)U_{n}(Q-P\tau/m) \left[F_{ab}^{(2)}(P,Q-P\tau/m,t-\tau)\right]$$
(105)  
$$-F_{bb}^{(2)}(P,Q-P\tau/m,t-\tau)\right] + e\int_{0}^{\infty} d\tau e^{-i\Omega_{ab}\tau}E^{(2)}(Q-P\tau/m,t-\tau) \cdot \frac{\partial}{\partial P}F_{ab}^{(1)}(P,Q-P\tau/m,t-\tau) .$$

The spatial integration required in the evaluation of the polarization density is

$$\frac{2}{L} \int_{0}^{L} dZ U_{n}(Z) U_{n}(Z - V\tau) e^{ik_{y}Z} = \frac{1}{2} \left( 2 \partial_{z0} \cos k_{n} V\tau - \partial_{z,2n} e^{ik_{y}V\tau} - \partial_{z,-2n} e^{-ik_{z}V\tau} \right)$$
(106)

It is found that the dominant correction to the polarization comes from the  $\mathcal{L} = 0$  term in  $F_{aa} \stackrel{(2)}{-} F_{bb} \stackrel{(2)}{-}$ . All other terms yield contributions that are higher order in  $(k_n v_0)^{-1}$ . Hence we set

$$\Delta \mathcal{P}^{(3)} = \mathcal{P}_{n}^{(3)} - \mathcal{P}_{n}^{(30)} = - \mathscr{C}_{ab}(iE_{o} \cdot \mathscr{C}_{ob}/\hbar) \int_{0}^{\infty} d\tau e^{-i\Omega_{Gb}\tau} A_{n}(t-\tau) \int d^{3}P \int \frac{ds}{2\pi i} e^{s(t-\tau)}$$

$$\times e^{-ik_{f} \cdot P\tau/m} \frac{mn_{\ell}^{(20)}(s)\omega_{p}^{2}}{k_{\ell}^{2}(s^{2}+\omega_{p}^{2})} \left(\frac{1}{s+\gamma_{a}+ik_{f} \cdot P/m} - \frac{1}{s+\gamma_{b}+ik_{f} \cdot P/m}\right) \delta_{\ell,o} \cos k_{n} \cdot P\tau/m ,$$
(107)

where  $\omega_p^2 = 4 \pi n^{(0)} e^2 / m$ . The  $\mathcal{L} = 0$  form of the dielectric constant, namely

$$\epsilon_{o}(s) = 1 + s^{2} / \omega_{p}^{2} , \qquad (108)$$

-23-

has already been introduced in Eq. (107) since it is insensitive to the limiting process required in evaluating this expression at  $k_{\perp} = 0$ .

Performing the s-inversion first, the correction to the third-order polarization may be written

$$\Delta \mathcal{P}^{(3)} = W_0 + W_1 + c.c. \qquad (109)$$

where

$$W_{a} = -\delta_{ab}^{o} \left(\frac{iE_{o} \cdot \delta_{ab}}{\hbar}\right) \frac{m\omega_{p}}{2i} - \frac{\omega_{pa}^{2}}{k_{g}^{2}} \int_{0}^{\infty} d\tau e^{-i\Omega_{ab}\tau} A_{n}(\tau-\tau) \int d^{3} P e^{-ik_{g} \cdot P \tau/m} \cos k_{n} \cdot P \tau/m$$

$$\times \frac{ik_{g} \cdot \frac{\partial \Phi^{(0)}}{\partial P}}{\gamma_{a} + i\omega_{p} + ik_{g} \cdot P/m} \int_{0}^{\infty} d\tau' N_{g}^{(20)}(\tau-\tau-\tau') \left[e^{i\omega_{p}\tau'} - e^{-(\gamma_{a} + i\omega_{p})\tau'}\right]$$
(110)

+ (similar term with  $\omega_p \rightarrow -\omega_p$ ),

and  $\omega_{p_{C}}^{2} = 4 \pi n_{\alpha}^{(0)} e^{2}/m$ .

The second order density variation to be used in Eq. (110) is readily obtained from Eqs. (50) - (52). The result is

$$N_{o}^{(20)}(t) = -\frac{1}{2} N (i E_{o} \cdot \delta_{ob}^{*} / \hbar)^{2} \int_{0}^{\infty} d\tau d\tau e^{-\Omega_{o}^{2} \tau^{2}} (e^{-\gamma_{0} \tau} - e^{-\gamma_{b} \tau})$$

$$\times (e^{-i\Omega_{ob} \tau'} + e^{-i\Omega_{ba} \tau'}) A_{n} (t - \tau) A_{n} (t - \tau - \tau')$$
(111)

Integration over momentum in Eq. (110) requires the evaluation of the integral

$$I_{\alpha}(t_{1}, t_{2}) = \int d^{3}P \frac{ik_{\ell} \cdot \frac{\partial \Phi^{(0)}}{\partial P} e^{-ik_{\ell} \cdot P t_{2}/m} \cos k_{n} \cdot P t_{1}/m}{\gamma_{\alpha} + i\omega_{p} + ik_{\ell} \cdot P/m} \qquad (112)$$

-24-

Expanding the denominator and only retaining terms  $O(k_{\ell}^{2})$  yields

$$I_{\alpha}(t_{1},t_{2}) = -k_{\alpha}^{2}e^{-(\Omega_{0}t_{1})^{2}}\left[1-2(\Omega_{0}t_{1})^{2}\right]\left[1+t_{2}(\gamma_{\alpha}+i\omega_{p})\right]/(\gamma_{\alpha}+i\omega_{p})^{2} .$$
(113)

When Eqs. (110) and (111) are combined and the three time factors  $A_n(t)$  are written in exponential form, it is again found that only two terms give rise to resonant contributions in subsequent time integrations. Hence Eq. (110) may be written

$$W_{\alpha} = \frac{1}{32} N \omega_{p} \omega_{p\alpha}^{2} \mathscr{G}_{ab} (E_{o} \cdot \mathscr{G}_{ab} / h) e^{-i(\omega_{n}i + \phi)} \int_{0}^{\infty} d\tau e^{-\Omega_{o}^{2}\tau^{2} - i(\Omega_{ab} - \omega_{n})\tau} \left[ 1 - 2(\Omega_{o}\tau)^{2} \right]$$

$$\times (\gamma_{\alpha} + i\omega_{p})^{-2} \int_{0}^{\infty} d\tau' \left\{ e^{i\omega_{p}\tau'} \left[ 1 + \tau(\gamma_{\alpha} + i\omega_{p}) \right] - e^{-\gamma_{\alpha}\tau'} \left[ 1 + (\tau + \tau')(\gamma_{\alpha} + i\omega_{p}) \right] \right\}$$
(114)
$$\times \int_{0}^{\infty} d\tau'' d\tau''' e^{-\Omega_{o}^{2}\tau'''^{2}} (e^{-\gamma_{0}\tau''} - e^{-\gamma_{b}\tau''}) \left[ e^{-i(\Omega_{ab} - \omega_{n})\tau'''} + e^{-i(\Omega_{ba} + \omega_{n})\tau'''} \right]$$

+ (similar term with 
$$\omega_p - - \omega_p$$
)

After performing the simple integrations of  $\tau'$  and  $\tau''$ , the  $\tau'''$  integration is given in terms of the z function introduced in Eq. (71) and the  $\tau$  integration requires

$$\int_{0}^{\infty} d\zeta e^{-\zeta^{2} + i \epsilon \zeta} (1 - 2\zeta^{2}) = -\frac{i}{2} \epsilon + 0 (\epsilon^{2})$$

$$\int_{0}^{\infty} d\zeta e^{-\zeta^{2} + i \epsilon \zeta} \zeta (1 - 2\zeta^{2}) = -\frac{i}{2} + 0 (\epsilon)$$
(115)

One finally obtains

$$W_{a} = -\frac{i N \omega_{po}^{2} \delta_{ab}}{4 \gamma_{a} (n v_{o})^{3}} \left(\frac{E_{o} \delta_{ab}}{\hbar}\right)^{3} \left(\frac{1}{\gamma_{a}} - \frac{1}{\gamma_{b}}\right) I_{m} Z \left(\frac{\omega_{n} - \Omega_{ab}}{k_{n} v_{o}}\right) \left[1 + i \left(\frac{\omega_{n} - \Omega_{ab}}{\gamma_{a}}\right)\right] \quad . \quad (116)$$

The resulting change in the polarization density may now be obtained

-25-

from Eq. (109) The associated in quadrature component is found to be

$$\Delta S_{n}^{(3)} = \frac{\mathscr{B}_{ab}N}{4\gamma_{a}\gamma_{b}k_{n}v_{a}} \left(\frac{E_{o}\mathscr{B}_{ab}}{\hbar}\right)^{3} Im Z\left(\frac{\omega_{n}-\Omega_{ab}}{k_{n}v_{o}}\right) \left[\frac{1}{(\gamma_{a}k_{n}L_{o})^{2}} + \frac{1}{(\gamma_{b}k_{n}L_{h})^{2}}\right] , \quad (117)$$

where  $L_a^2 = kT/4\pi n_a^{(0)} e^2$ . Finally the equil

Finally, the equilibrium field amplitude is given by

$$\frac{\mathscr{G}_{ab}^{2} E_{n}^{2}}{8\hbar r_{a} r_{b}} = \frac{1 - \mathscr{N}_{e}^{-1} (\omega_{n} - \Omega_{ab})^{2} / (k_{n} v_{o})^{2}}{1 + r_{ab}^{2} \mathcal{I}(\omega_{n} - \omega_{ab}) + 2(r_{o} - r_{b})^{2} \left[ (r_{a} k_{n} L_{a})^{-2} + (r_{b} k_{n} L_{b})^{-2} \right]} .$$
(118)

Comparison of Eq. (118) with the corresponding result of Ref.(1)shows that the effect of long range forces is contained in the last term in the denominator of Eq. (118). This term is enhanced by increasing the wavelength of the laser radiation and by decreasing the Debye length associated with the particles in states a and b. Since these Debye lengths turn out to be on the order of  $10^{-3}$  cm while the optical wavelengths are  $10^{-5} - 10^{-4}$  cm, it is seen that the correction term would be quite small were it not for the fact that the transition rate  $\gamma_b$  of the lower level may be as much as  $10\gamma_0$ .

As a numerical example, consider  $1\mu$  radiation in a plasma having a temperature of  $2000^{\circ}$ K and  $n_{o}^{(0)} \sim n_{b}^{(0)} \sim 5 \times 10^{11} \text{ cm}^{-3}$ . Then  $L_{a}^{2} = L_{b}^{2} = 2 \times 10^{-7} \text{ cm}^{2}$ . Assuming that  $\gamma_{b} = 10 \gamma_{o}$ , the correction term is 0.2. The response of the laser at frequencies above  $\omega_{ob}$ both with and without this correction is shown in Fig. 1. The complete response curve is, of course, symmetric about the frequency  $\omega_{ob}$ . Although the shape of the response curve remains unchanged when long range forces are included, a superposition of the two curves in Fig. 1 shows that the effect of such forces is to fill in the center of the curve. Also, the requirement of positive curvature at the center of the response curve yields as a criterion for the existence of a dip,

$$\mathcal{N} > 1 + \frac{2\gamma_{ob}^{2}}{\Omega_{o}^{2}} \left(1 + \frac{c}{2}\right) \qquad (119)$$

where

$$\frac{c}{2} = (\gamma_{a} - \gamma_{b})^{2} \left[ (\gamma_{a} k_{n} L_{a})^{-2} + (\gamma_{b} k_{n} L_{b})^{-2} \right]$$
(120)

Finally, Eq. (118) shows that the correction term vanishes if  $\gamma_a = \gamma_b$ . This is clear on physical grounds since in the laser theory one finds that  $F_{aa}^{(2)} = -F_{bb}^{(2)}$  when  $\gamma_a$  and  $\gamma_b$  are equal. These two terms then have identical spatial dependences that are 180° out of phase with each other. Since the net force acting on a given two-level system is due to the inhomogeneous distribution of other charged particles, it is clear that for  $\gamma_a \neg \gamma_b$  the distribution of particles on state <u>a</u> plus those in state <u>b</u> gives rise to a completely homogeneous spatial distribution of such particles and no net force is felt by an individual twolevel system.

#### MEASURE LINE OF ARGON ION LASER LINE PROFILE

#### Introduction

In considering the output power of laser oscillators, it has been shown that the saturation characteristics of the optical medium depend upon the broadened line width of the medium and the frequency separation between modes. <sup>(4)</sup> A representation of this mechanism is most easily described as a hole being "burned" in the Doppler-broadened gain curve of a gas laser. The "hole burning" refers to the saturation of the atoms in the vicinity of the frequency of the saturating radiation. <sup>(5)</sup> At low power levels the distortions in the "hole width" are due to the broadening caused by collisio in a non-ionized plasma<sup>(6)</sup> and by the long range forces in an ionized plasma<sup>(7)</sup> (see the earlier section of this report).

#### Experimental Arrangements and Preliminary Results

For a cavity of very short length, the spacing of the resonances may exceed the Doppler-broadened line width and the laser will oscillate in a single mode. One expects oscillation at a single frequency to cause gain saturation of the medium only in the vicinity of this frequency. Two holes are "burned" in the gain curve due to the standing wave character of the optical fields in the cavity. If the length of the cavity is tuned through the center of the Doppler profile, a "dip" is observed which is due to the fact that fewer atoms are contributing to the power output of the laser. For the specific case of the ionized argon gas laser, the gain profile is so wide that a short single mode cavity could not be conveniently constructed and therefore the "Lamb Dip" could not be displayed by tuning the c/2L frequency through line center. If the output intensity of the argon laser was extremely stable (which it is not), it could be operated very close to threshold in one mode and the "Lamb Dip" could possibly be exhibited; however, no range of the excitation parameter would

-28-

ezist.

An alternate solution which has been successfully applied during the reporting period is to use a three mirror, variable reflectivity system at one end of the resonator.  $^{(8)}$  A /2 motion of this system will allow the two cavity system to pass through alignment and will provide a resonance for one of the modes of the original cavity. By linearly driving one of the mirrors of the variable reflectivity cavity the resonances may be scanned across the whole Doppler-broadened line and will exhibit the "Lamb Dip" as the particular oscillating modes are scanned in "ime. The "Lamb Dip" may now be plotted as a function of excitation and the area of the "hole" represents the relation power to be achieved by a single mode since the "hole" area represents the homogeneously broadened line width.

The experimental arrangement is shown pictorially in Fig. 2 and schematically in Fig. 3a. A d.c.  $Ar^+$  discharge is contained in a quartz capillary which is 60 cm in length and has a 2 mm ID. The excitation can be varied from 3 to 10 amperes. An intracavity prism is incorporated into the design to enable the various atomic lines to be selected. The three mirror, variable reflectivity system is located at one end of the cavity and has a multiwafer piezoelectric drive associated with mirror M, which provides the scan of the short cavity relative to the long cavity.

The system has been designed, constructed and made operative. Preliminary data, as shown in Fig. 3b, has illustrated the "Lamb Dip", and demonstrated the suitability of the technique for a detailed investigation of the shape of the hole burned in the Doppler gain curve. Detailed results of the experimental portions of this program and attempts to interpret the experiments in terms of this theory will be reported in a later report.

-29-

#### SIX-MONTH STATUS EVALUATION

Some aspects of the theory of a gas laser recently developed by W. E. Lamb, Jr. have been recast in a form which more fully displays the role played by the particle dynamics. The Wigner distribution function has been used to derive kinetic equations which govern the external center of mass motion of the two-level systems as well as their internal dynamics. The effect of long range forces has been discussed by treating the collision integral in a manner similar to that employed in plasma kinetic theory. A modification in the criterion for the existence of a dip in the output has been obtained. It has also been shown that effects due to long range forces are most noticeable at long optical wavelengths and when there is a large difference between the lifetimes of the two laser levels.

The experimental system for measuring the line profile of a d.c. excited argon laser has been designed, constructed, and made operative. Preliminary data on the "Lamb Dip" has been obtained as a function of pressure and excitation. Detailed results of the experimental portions of this program and attempts to interpret the experiments in terms of this theory will be reported in a later report.

### REFERENCES

- \* Work partially supported by Contract NO0014-66-C0344 as part of Project DEFENDER under the joint sponsorship of the Advanced Research Projects Agency, the Office of Naval Research, and the Department of Defense.
- 1. Lamb, W. E., Jr., Phys. Rev. <u>134</u>, A1429 (1964).
- Gyorffy, B. L. and W. E. Lamb, . ., (Proceedings of the International Conference on the Physics of Quantum Electronics, Puerto Rico, 1965, edited by P. L. Kelley, B. Lax, and P. E. Tannenwald (McGraw-Hill Book Co., Inc., New York, 1965)), p. 602, and to be published.
- 3. Irving, J. H. and R. W. Zwanzig, J. Chem. Phys. <u>19</u>, 1173 (1951).
- 4. Rigrod, W. W., J. Appl. Phys. <u>34</u>, 2602 (1963).
- 5. Bennett, W. R., Jr., Appl. Opt. Suppl. <u>1</u>, 24 (1962).
- 6. Smith, P. W., J. Appl. Phys. <u>37</u>, 2089 (1966).
- 7. Lamb, G. L., Jr., (to be published).
- 8. Smith, P. W., J. Quant. Elec. <u>QE-1</u>, 343 (1965).

-31-

## LIST OF FIGURES

Fig. l - Calculated Lamb Dip Line Profiles

- Fig. 2 Experimental Apparatus for Laser Line Profiles
- Fig. 3 Single Mode Operation of Argon Ion Laser





SINGLE MODE UPERATION OF ARGON ION LASER



<u>Unclassified</u>			•			
Security Classification						
DUCUMEN   CON (Security Camelficetion of title, body of abstract and indexin	NIKOL DATA - REI na annotation must be en	) tered when ti	he overall report is clessified)			
1. ORIGINATIN ; ACTIVITY (Corporate author)		20. REPORT SECURITY CLASSIFICATION				
United Aircraft Corporation		Unclas	ssified			
Research Laboratories		25 GROUP				
East Hartford, Connecticut						
3. REPORT TITLE	• _					
Semiannual Report on the Rese	earch Invest:	igation	n of			
Laser Line Profiles						
A DESCRIPTIVE NOTES (Type of report and inclusive dates)						
Semignnual Report for the Per	riod 1 August	1966	- 31 January 1967			
5. AUTHOR(S) (Leet name, first name, initial)						
		_				
DeMaria, Anthony J.; Lemb, Ge	eorge L., Jr.	; and	Stetser, David A.			
S. REPORT DATE	74. TOTAL NO. OF P.	AGES	75. NO. OF REFS			
March 2, 1967						
B. CONTRACT OR GRANT NO.	94. ORIGINATOR'S RE	PORT NUM	SER(S)			
N00014-66-C0344	<b>F-9204</b>	79-2				
& PROJECT NO.						
c.	this report)	40(3) (Any	other numbers that may be assigned			
<b>d</b> .						
10. AVAILABILITY/LIMITATION NOTICES						
11. SUPPLEMENTARY NOTES	12. SPONSORING MILL	TARY ACTI				
	Departmen	t of t	the Navy			
	Office of	Naval	Research			
13. ABSTRACT						
Some aspects of the the	orv of a gas	laser	recently			
developed by W E. Lamb. Jr. at	re recast in	a for	m which more			
fully displays the role played	by the part	icle d	vnamics. The			
Wigner distribution function is used to derive kinetic						
equations which govern the exte	ernal center	of ma	ss motion of			
the two-level system as well as their internal dynamics. The						
effect of long range forces is discussed by treating the						
collision integral in a manner similar to that employed in						
plasma kinetic theory. A modification in the criterion for						
the existence of a dip in the output is obtained. It is also						
shown that effects due to long range forces are most noticeable						
at long optical wavelengths and when there is a large difference						
between the lifetimes of the t	wo laser leve	ers.				
The experimental system	for measuri	ng the	line profile of			
a dc excited argon laser has been designed, constructed, and						
made operative. Preliminary data on the "Lamb Dip" of an argon						
ion laser has been obtained as a function of pressure and						
excitation. The results of the experimental portion of this						
program and attempts to interpret the experiments in terms of						
this theory will be reported in	n later repo	LLS.				
DD FORM 1/72		Unala	esified			

Security Classification

#### Unclassified

Security Classification

Laser Line Profiles Instructions	14 KEY WORDS	LIN	LINKA		_1NK		LINKC		
Laser Line Profiles		KEY WORDS	ROLE	₩T	ROLE	W T	ROLE	<b>W</b> 1	
Laser Line Profiles									
Laser Line Profiles									
Laser Line Profiles									
Laser Line Profiles									
INSTRUCTIONS		Leson Line Drafilae							
INSTRUCTIONS		Laser Line Provides							
INSTRUCTIONS									
INSTRUCTIONS									
INSTRUCTIONS									
INSTRUCTIONS									
INSTRUCTIONS									
INSTRUCTIONS									
INSTRUCTIONS				ļ			[		
INSTRUCTIONS				1					
INSTRUCTIONS									
INSTRUCTIONS									
		INSTRUCTIONS				يتكف مرجلا الأصياب			

1. ORIGINATING ACTIVITY: Enter the name and addrasa of the contractor, aubcontractor, grantee, Department of Defenae activity or other organization (corporate author) issuing the report.

2a. REPORT SECURITY CLASSIFICATION: Enter the overall security classification of the report. Indicate whethar "Reatricted Data" is included. Marking is to be in accordance with appropriate security regulations.

2b. GROUP: Automatic downgrading is specified in DoD Directive 5200.10 and Armed Forces Industrial Manual. Enter the group number. Also, when applicable, show that optional markings have been used for Group 3 and Group 4 as suthorized.

3. REFORT TITLE: Enter the complete report title in all capital letters. Titles in all cases should be unclassified. If a meaningful title cannot be selected without classification, show title classification in all capitals in parenthesis immediately following the title.

4. DESCRIPTIVE NOTES: If appropriate, entar the type of report, e.g., interim, progreas, aummary, annual, or final. Give the inclusive dates when a specific reporting period is covered.

5. AUTHOR(S): Enter the name(a) of author(s) as abown on or in the report. Enter last name, first name, middla Initial. If military, show rank and branch of aervice. The name of the principal author is an absolute minimum requirement.

6. REPORT DATE: Enter the date of the report as day, month, year; or month, year. If more than one date appear on the report, use date of publication.

7a. TOTAL NUMBER OF PAGES: The total page count should follow normal pagination procedures, i.e., enter the number of pages containing information.

7b. NUMBER OF REFERENCES Enter the total number of references cited in the report.

8. CONTRACT OR GRANT NUMBER: If appropriate, enter the applicable number of the contract or grant under which the report was written.

8b, 8c, & 8d. PROJECT NUMBER: Enter the appropriate military department identification, auch as project number, subproject number, aystem numbera, task number, etc.

9a. ORIGINATOR'S REPORT NUMBER(5): Enter the official report number by which the document will be identified and controlled by the originating octivity. This number must be unique to this report.

9b. OTHER REPORT NUMBER(S): If the seport has been assigned any other report numbers (either by the originator or by the sponsor), also enter this number(a).

10. AVAILABILITY/LIMITATION NOTICES: Enter any limitations on further disacmination of the report, other than those imposed by security classification, using standard statements such ar.

- (1) "Qualified requestors may obtain copies of this report from DDC."
- (2) "Foreign announcement and disagenination of this report by DDC is not authorized."
- (3) "U. S. Government agencies may obtain copies of this report directly from DDC Other qualified DDC use a shall request through
- (4) "U. S. military agencies may obtain copies of this report directly from DDC. Other qualified users shall request through
- (5) "All distribution of this report is controlled. Qualified DDC users shall request through

If the report has been furniahed to the Offica of Technical Servicea, Department of Commerca, for sale to the public, indicate this fact and enter the price, if known

11. SUPPLEMENTARY NOTES: Use for additional explanatory notes.

12. SPONSORING MILITARY ACTIVITY: Enter the name of the departmental project office or laboratory sponsoring  $(p \neq p)$  ing for) the research and devalopment. Includa addreas.

13. ABSTRACT: Enter an abatract giving s brief and factual summary of the document indicative of the report, even though it may also appear elaeschere in the body of the technical report. If additional space is required, a continuation sheet shall be attached.

It is highly drairable that the abatract of classified reports be unclassified. Each paragraph of the abatract shall end with an indication of the military accurity classification of the information in the paragraph, represented as (TS), (S), (C), or (U)

There is no limitation on the length of the abstract. However, the suggested length is from 150 to 225 words.

14. KEY WORDS: Key words are tachnically maaningful terms or short phrases that characterize a report and may be used as index entries for cataloging the report. Key words must be selected so that no security classification is required. Identifiers, such as equipment model designation, trade name, military project code name, geographic location, may be used as key words but will be followed by an indication of technical context. The assignment of links, rules, and weights is optional

Security Classification