

ARL 67-0044
MARCH 1967



Aerospace Research Laboratories

AD 653968


SIMULTANEOUS TESTS FOR THE EQUALITY OF COVARIANCE MATRICES AGAINST CERTAIN ALTERNATIVES

P. R. KRISHNAIAH
APPLIED MATHEMATICS RESEARCH LABORATORY


Project No. 7071

009
950

Distribution of this document is unlimited



OFFICE OF AEROSPACE RESEARCH
United States Air Force



ARCHIVE COPY

ERRATA

ARL 67-0044

MARCH 1967

Page 6, line 2: $|w - v|$ should read as $|w - v| > 1$

Page 8: Add "for $j = 1, 2, \dots, p$ " above last line

AEROSPACE RESEARCH LABORATORIES
OFFICE OF AEROSPACE RESEARCH
UNITED STATES AIR FORCE
WRIGHT-PATTERSON AIR FORCE BASE, OHIO

ARL 67-0044

**SIMULTANEOUS TESTS FOR THE EQUALITY OF
COVARIANCE MATRICES AGAINST
CERTAIN ALTERNATIVES**

**P. R. KRISHNAIAH
APPLIED MATHEMATICS RESEARCH LABORATORY**

MARCH 1967

Project 7071

Distribution of this document is unlimited

**AEROSPACE RESEARCH LABORATORIES
OFFICE OF AEROSPACE RESEARCH
UNITED STATES AIR FORCE
WRIGHT-PATTERSON AIR FORCE BASE, OHIO**

FOREWORD

This report was prepared for the Applied Mathematics Research Laboratory, Aerospace Research Laboratories by Dr. P. R. Krishnaiah under Project 7071, "Research in Applied Mathematics". It contains some procedures for testing the hypothesis of equality of covariance matrices against different alternatives when the underlying populations are multivariate normal.

The author wishes to thank Miss Eva Brandenburg for typing the manuscript carefully.

ABSTRACT

In this paper, we consider the problems of testing for the equality of covariance matrices against certain alternatives when the underlying populations are multivariate normal. The alternative hypotheses considered are (i) at least one covariance matrix is not equal to the covariance matrix of the next population, (ii) at least one covariance matrix is not equal to the covariance matrix of the standard population and (iii) at least one covariance matrix is not equal to the covariance matrix of another population.

Table of Contents

Section	Page
1. Introduction and Summary	1
2. Preliminaries and Statement of Problem	1
3. Test for H Against A_1	4
4. Test for H Against A_2 and A_3	8
5. General Remarks	9
References	10

1. Introduction and Summary

In many situations, it is of interest to test for the equality of variances or covariance matrices against certain alternatives. Hartley [6] considered the problem of testing for the equality of variances against the alternative that at least one variance is different from the other. Gnanadesikan [3] considered the problem of testing for the equality of variances against the alternative that at least one variance is not equal to the standard. Recently, Krishnan [11] considered testing for the equality of variances against the alternative that at least one variance is not equal to the next. In the above procedures, it was assumed that the underlying populations are univariate normal. In this paper, we consider multivariate generalizations of the above test procedures. The test procedures proposed in this paper are based upon expressing the total hypothesis as the intersection of some elementary hypotheses and testing these elementary hypotheses by using conditional distributions. In the two sample case, our procedures are similar (but not equivalent) to the procedure proposed by Roy [14]; the test statistics used by him in testing some of the elementary hypotheses are different from those used in this paper.

2. Preliminaries and Statement of Problems

Let $S_i = (s_{iqr})$ denote i th sample sums of squares and cross products (SP) matrix and let $n_i + 1$ denote i th sample size. Let Σ_{ij} denote top $j \times j$ left hand corner of $\Sigma_i = (\sigma_{iqr})$ and let S_{ij} denote the top $j \times j$ left hand corner of $S_i = (s_{iqr})$. Also, let $S_{ij}^{-1} = (s_{ijtu}^*)$ and let σ_{0j}^2 denote the common value of σ_{ij}^2 (defined below) when

$$\sigma_{1j}^2 = \dots = \sigma_{kj}^2.$$

In addition,

$$\underline{\beta}_{ij} = \begin{pmatrix} \beta_{ij1} \\ \vdots \\ \beta_{ijj} \end{pmatrix} = \Sigma_{ij}^{-1} \begin{pmatrix} \sigma_{i1,j+1} \\ \vdots \\ \sigma_{ij,j+1} \end{pmatrix}$$

$$\underline{b}_{ij} = \begin{pmatrix} b_{ij1} \\ \vdots \\ b_{ijj} \end{pmatrix} = S_{ij}^{-1} \begin{pmatrix} s_{i1,j+1} \\ \vdots \\ s_{ij,j+1} \end{pmatrix}$$

$$s_{i,j+1}^2 = \frac{|s_{i,j+1}|}{|s_{ij}|}, \quad \sigma_{i,j+1}^2 = \frac{|\Sigma_{i,j+1}|}{|\Sigma_{ij}|} \text{ for } j=1, 2, \dots, (p-1)$$

$$s_{i1}^2 = s_{i11}, \quad \sigma_{i1}^2 = \sigma_{i11}, \quad s_{.,j+1}^2 = \sum_i s_{i,j+1}^2.$$

$$D_{imtu} = (b_{itu} - b_{mtu})^2 / (s_{ituu}^* + s_{mtuu}^*), \quad N = \sum_{i=1}^k n_i$$

$$F_{imt} = \frac{s_{it}^2 (n_i - t + 1)}{s_{mt}^2 (n_i - t + 1)}, \quad F_{imtu} = \frac{(N - kt) D_{imtu}}{s_{.,t+1}^2}.$$

Also, let

$$H_{j1}: \sigma_{1j}^2 = \dots = \sigma_{kj}^2 \quad H_{j2}: \beta_{1j} = \dots = \beta_{kj}$$

$$A_{1j1} = \bigcup_{i=1}^{k-1} [\sigma_{ij}^2 \neq \sigma_{i+1,j}^2] \quad A_{1j2} = \bigcup_{i=1}^{k-1} [\beta_{ij} \neq \beta_{i+1,j}]$$

$$A_{2j1} = \bigcup_{i=1}^{k-1} [\sigma_{ij}^2 \neq \sigma_{kj}^2] \quad A_{2j2} = \bigcup_{i=1}^{k-1} [\beta_{ij} \neq \beta_{kj}]$$

$$A_{3j1} = \bigcup_{i \neq i'=1}^k [\sigma_{ij}^2 \neq \sigma_{i'j}^2] \quad A_{3j2} = \bigcup_{i \neq i'=1}^k [\beta_{ij} \neq \beta_{i'j}]$$

In this paper, we consider the problem of testing H against A_1 , A_2 and A_3 where
 $H: \Sigma_1 = \dots = \Sigma_k$, $A_1 = \bigcup_{j=1}^p A_{1j1} \bigcup_{j=1}^{p-1} A_{1j2}$, $A_2 = \bigcup_{j=1}^p A_{2j1} \bigcup_{j=1}^{p-1} A_{2j2}$ and
 $A_3 = \bigcup_{j=1}^p A_{3j1} \bigcup_{j=1}^{p-1} A_{3j2}$

The test procedures considered in this paper are based on the following method.
 We first test H_{11} against the alternative of interest. If H_{11} is rejected, we declare that H is rejected. If H_{11} is accepted, we proceed further and test H_{21} and H_{12} holding the first variate fixed. If $H_{12} \cap H_{21}$ is accepted, we proceed further and test H_{31} and H_{32} holding the second variate fixed. We continue this procedure until H is accepted or rejected. Here we note that $\bigcap_{j=1}^r H_{j1} \bigcap_{j=1}^{r-1} H_{j2}$ is equivalent to the hypothesis that

$$\Sigma_{1r} = \dots = \Sigma_{kr}$$

We need the following known results (see [14]) in the sequel:

When S_{ij} is fixed, the distribution of b_{ij} is independent of the distribution of $s_{i,j+1}^2$; the distribution of b_{ij} is j -variate normal with mean vector β_{ij} and covariance matrix

$\sigma_{i,j+1}^2 S_{ij}^{-1}$ and $s_{i,j+1}^2 / \sigma_{i,j+1}^2$ is distributed as χ^2 with $(n_i - j)$ degrees of freedom.

3. Test for H Against A_1

The following lemma is needed in the sequel.

Lemma 4.1

If x_1, x_2, \dots, x_k are distributed independently as central chi-square variates with m_1, m_2, \dots, m_k degrees of freedom, then

$$f(F_{12}, F_{23}, \dots, F_{k-1,k}) = \frac{(m_1/m_k) \left[\prod_{j=1}^{k-1} m_j \prod_{i=j}^{k-1} F_{i,i+1} / m_k \right]^{(m_k-2)/2}}{\prod_{j=1}^k \Gamma(m_j/2) \left[1 + \frac{1}{m_k} \sum_{j=1}^{k-1} m_j \prod_{i=j}^{k-1} F_{i,i+1} \right]^{\sum m_j/2}}$$

where $F_{ij} = \frac{x_i m_j}{x_j m_i}$

The proof of the above lemma is given in [11].

We will first consider the problem of testing H_{j1} against the alternative A_{1j1} when the first $(j-1)$ variates are held fixed (with the understanding that no variate is held fixed when H_{11} is tested). In this case, we accept H_{j1} if and only if

$$(4.1) \quad \lambda_{ij} \leq F_{i,i+1,j} \leq \mu_{ij}$$

where λ_{ij} and μ_{ij} are chosen such that

$$(4.2) \quad P[\lambda_{ij} \leq F_{i,i+1,j} \leq \mu_{ij}; i = 1, 2, \dots, k-1 | H_{j1}] = P_j$$

When H_{j1} is true, $s_{1j}^2/\sigma_{0j}^2, \dots, s_{kj}^2/\sigma_{0j}^2$ are independently distributed as chi-square

variates with $(n_1 - j + 1), \dots, (n_k - j + 1)$ degrees of freedom. So, using Lemma 4.1, we can write down the joint distribution of $F_{12}, F_{23}, \dots, F_{k-1,k}$ when H_{j1} is true. We will now discuss about a procedure for testing H_{j2} against A_{1j2} when H_{j1} is true and when the first j variates are held fixed.

When H_{j1} is true and the first j variates are held fixed, we accept H_{j2} if and only if

$$(4.3) \quad F_{i, i+1, ju} \leq c_{ij}, \quad \begin{matrix} u = 1, 2, \dots, j \\ i = 1, 2, \dots, k-1 \end{matrix}$$

where c_{ij} 's are chosen such that

$$(4.4) \quad P[F_{i, i+1, ju} \leq c_{ij}; u = 1, 2, \dots, j; i = 1, 2, \dots, k-1 | H_{j1} \cap H_{j2}] = P_j.$$

When H_{j1} is true, $s_{..j+1}^2 / \sigma_{0,j+1}^2$ is distributed as a chi-square variate with $(N - k - j)$ degrees of freedom and it is distributed independently of $D_{i, i+1, ju}$ for $i = 1, 2, \dots, k-1$ and $u = 1, 2, \dots, j$. Also, when $H_{j1} \cap H_{j2}$ is true, the joint distribution of

$$(D_{12j1}, \dots, D_{k-1, kj1}, D_{12j2}, \dots, D_{k-1, kj2}, \dots, D_{12jj}, \dots, D_{k-1, kjj})$$

is a central multivariate chi-square distribution with 1 degree of freedom and with Ω^j as the covariance matrix of the "accompanying" multivariate normal where

$$\Omega^j = \begin{bmatrix} \Omega_{11}^j & \Omega_{12}^j & \dots & \Omega_{1j}^j \\ \Omega_{21}^j & \Omega_{22}^j & \dots & \Omega_{2j}^j \\ \vdots & \vdots & \ddots & \vdots \\ \Omega_{j1}^j & \Omega_{j2}^j & \dots & \Omega_{jj}^j \end{bmatrix}$$

$$\Omega_{ii}^j = (\omega_{ii}^j)_{vw}^2 \sigma_{0,j+1}^2,$$

$$\omega_{ii}^j = \begin{cases} 0 & |w-v| > 1 \\ \frac{(s_{vji}^* + s_{v+1,ji}^*)}{\sqrt{(s_{vji}^* + s_{v+1,ji}^*)(s_{vji}^* + s_{v+1,ji}^*)}} & w = v \\ \frac{-s_{tji}^*}{\sqrt{(s_{vji}^* + s_{v+1,ji}^*)(s_{wji}^* + s_{w+1,ji}^*)}} & |w-v| = 1 \end{cases}$$

and $t = \max.(w, v)$; (for the definition of the "accompanying" multivariate normal, see [7]). So, the joint distribution of

$$(F_{12j1}, \dots, F_{k-1,kj1}, \dots, F_{12jj}, \dots, F_{k-1,kjj})$$

is a multivariate F distribution with $(1, N-kj)$ degrees of freedom and with Ω^j as the covariance matrix of the "accompanying" multivariate normal. For various details on the multivariate F distribution, the reader is referred to [9, 10].

Now, combining (4.1), (4.2), (4.3) and (4.4) we use the following procedure for testing H against A_1 .

Accept H against A_1 if and only if

$$(4.5) \quad \begin{cases} \lambda_{ij}^* \leq F_{i,i+1,j} \leq \mu_{ij}^* & i=1,2,\dots,k-1 \quad j=1,2,\dots,p \\ F_{i,i+1,ju} \leq c_{ij}^* & u=1,2,\dots,j \quad j=1,2,\dots,(p-1) \\ & i=1,2,\dots,k-1 \end{cases}$$

where λ_{ij}^* , μ_{ij}^* and c_{ij}^* are chosen such that the probability of (4.5) holding good,

when H is true, is $(1-\alpha)$. But this probability is equal to $\prod_{j=1}^p q_j \prod_{j=1}^{p-1} q'_j$ where

$$q_j = P[\lambda_{ij}^* \leq F_{i,i+1,j} \leq \mu_{ij}^*; i=1,2,\dots,k-1 \mid H_{1j}]$$

$$q'_j = P \left[F_{i,i+1,ju} \leq c_{ij}^*; u=1,2,\dots,j \quad j=1,2,\dots,(p-1) \mid H_{2j} \right] \\ i=1,2,\dots,(k-1)$$

The optimum choice of the critical values is not known. For practical purposes, we impose the following restrictions.

$$q_1 = \dots = q_p = q'_1 = \dots = q'_{p-1} = (1-\alpha)^{1/2p-1}$$

$$c_{ij}^* = c_j^*.$$

In addition, we impose the restriction that the test associated with testing H_{1j} is locally unbiased.

The $(1-\alpha)\%$ simultaneous confidence intervals associated with the above test procedure are given by

$$\frac{\lambda_{ij}^* s_{i+1,j}^2 (n_i - j + 1)}{s_{ij}^2 (n_{i+1} - j + 1)} \leq \frac{\sigma_{i+1,j}^2}{\sigma_{ij}^2} \leq \mu_{ij}^* \frac{(n_i - j + 1)}{(n_{i+1} - j + 1)} \frac{s_{i+1,j}^2}{s_{ij}^2}$$

$$i = 1, 2, \dots, k-1 \quad j = 1, 2, \dots, p$$

$$|b_{iju} - b_{i+1,ju} - \beta_{iju} + \beta_{i+1,ju}| \leq \frac{c_{ij}^* s_{ij+1}^2 (s_{iju}^* + s_{i+1,ju}^*)}{(N - kj)}$$

$$u = 1, 2, \dots, j \quad j = 1, \dots, (p-1)$$

$$i = 1, 2, \dots, k-1.$$

4. Tests for H Against A_2 and A_3 .

When H is tested against A_2 , we accept H if and only if

$$a_{ij} \leq F_{ikj} \leq b_{ij} \quad i = 1, 2, \dots, k-1 \quad j = 1, 2, \dots, p$$

$$F_{ikju} \leq c_{ij} \quad u = 1, 2, \dots, j \quad j = 1, 2, \dots, p \\ i = 1, 2, \dots, k-1$$

where a_{ij} , b_{ij} , c_{ij} are chosen such that

$$\prod_{j=1}^p Q_j \prod_{j=1}^{p-1} Q_j' = 1 - \alpha,$$

and

$$Q_j = P[a_{ij} \leq F_{ikj} \leq b_{ij}; i = 1, 2, \dots, k-1 \quad j = 1, 2, \dots, p | H_{1j}]$$

$$Q_j' = P[F_{ikju} \leq c_{ij}; u = 1, 2, \dots, j \quad i = 1, 2, \dots, k-1 \quad H_{2j}].$$

We can evaluate Q_1, \dots, Q_p by using the methods (or their modifications) discussed in [1, 4, 5, 8, 12] whereas Q_1', \dots, Q_{p-1}' can be evaluated by using the methods discussed in [9, 10]. The optimum choice (in terms of increasing power of the test) of the critical values is not known. But, for practical purposes, we can choose them by imposing restrictions similar to those in the previous section.

We will now propose a procedure to test H against A_3 when the sample sizes are equal to $(n+1)$. According to this procedure, we accept H if and only if

$$\begin{cases} 1/\lambda_j \leq F_{ii'j} \leq \lambda_j; i \neq i' = 1, 2, \dots, k \\ F_{ii'ju} \leq c_{ij}; i \neq i' = 1, 2, \dots, k \quad u = 1, 2, \dots, j \end{cases}$$

where

$$\prod_{j=1}^p R_j \prod_{j=1}^{p-1} R_j' = 1 - \alpha$$

$$R_j = P[1/\lambda_j \leq F_{iif} \leq \lambda_j; i \neq i' = 1, 2, \dots, k | H_{1j}]$$

$$R_j' = P[F_{iif'ju} \leq c_j; i \neq i' = 1, 2, \dots, k \quad u = 1, 2, \dots, j | H_{2j}].$$

Using the method discussed in [6], we can evaluate R_1, \dots, R_p . In order to evaluate R_1', \dots, R_{p-1}' we note that, when H_2 is true and j is fixed, the statistics $F_{iif'ju}$ are jointly distributed as a singular multivariate F distribution. So it is complicated to obtain exact values of R_j' . But, we can obtain approximate values of R_j' by using Bonferroni's inequalities [2; p. 100]. For practical purposes, we can choose the critical values such that

$$R_1 = \dots = R_p = R_1' = \dots = R_{p-1}' = (1-\alpha)^{1/2p-1}$$

The simultaneous confidence intervals associated with the above test procedures can be obtained easily.

5. General Remarks

Roy [14] proposed a procedure, based on conditional distributions, for testing the equality of two covariance matrices. But, the lengths of the confidence intervals associated with the procedures proposed in this paper are at least as short as the lengths of the corresponding confidence intervals associated with the procedure by Roy [14]. In the univariate case, the procedures proposed in this paper for testing H against A_1 , A_2 and A_3 are respectively equivalent to the procedures considered by Krishnaiah [11], Gnanadesikan [3] and Hartley [6].

REFERENCES

1. Armitage, J. V. and Krishnaiah, P. R. (1964). Tables for the studentized largest chi-square distribution and their applications. ARL 64-188, Aerospace Research Laboratories, Wright-Patterson Air Force Base, Ohio.
2. Feller, W. (1950). Introduction to Probability Theory and Its Applications. Wiley, New York.
3. Gnanadesikan, P. (1959). Equality of more than two variances and more than two dispersion matrices against certain alternatives. Ann. Math. Statist., 30, 177-184.
4. Gupta, S. S. and Sobel, M. (1962). On the smallest of several correlated F statistics. Biometrika, 49, 509-523.
5. Gupta, S. S. (1963). On a selection and ranking procedure for gamma populations. Ann. Inst. Statist. Math., 14, 199-216.
6. Hartley, H. O. (1950). The maximum F-ratio as a short cut test for heterogeneity of variances. Biometrika, 37, 308-312.
7. Krishnaiah, P. R. and Rao M. M. (1961). Remarks on a multivariate gamma distribution, American Mathematical Monthly, 68, 342-346.
8. Krishnaiah, P. R. and Armitage J. V. (1964). Distribution of the studentized smallest chi-square, with tables and applications. ARL 64-218, Aerospace Research Laboratories, Wright-Patterson Air Force Base, Ohio.
9. Krishnaiah, P. R. and Armitage, J. V. (1965). Probability integrals of the multivariate F distribution, with tables and applications. ARL 65-236, Aerospace Research Laboratories, Wright-Patterson Air Force Base, Ohio.
10. Krishnaiah, P. R. (1965). On the simultaneous ANOVA and MANOVA tests. Ann. Inst. Statist. Math., 17, 35-53.
11. Krishnaiah, P. R. (1965). Simultaneous tests for the equality of variances against certain alternatives. Austral. J. Statist., 7, 105-109.
12. Ramachandran, K. V. (1956). On the simultaneous analysis of variance test. Ann. Math. Statist., 27, 521-528.
13. Roy, S. N. (1953). On a heuristic method of test construction and its use in multivariate analysis. Ann. Math. Statist., 24, 220-238.
14. Roy, J. (1958). Step-down procedure in multivariate analysis. Ann. Math. Statist., 29, 1177-1187.

Unclassified
Security Classification

DOCUMENT CONTROL DATA - R&D		
(Security classification of title, body of abstract and indexing annotation must be entered when the overall report is classified)		
1. ORIGINATING ACTIVITY (Corporate author)		2a. REPORT SECURITY CLASSIFICATION
Aerospace Research Laboratories Wright-Patterson Air Force Base, Ohio		Unclassified
		2b. GROUP
3. REPORT TITLE		
Simultaneous Tests for the Equality of Covariance Matrices Against Certain Alternatives		
4. DESCRIPTIVE NOTES (Type of report and inclusive dates)		
Scientific. Final.		
5. AUTHOR(S) (Last name, first name, initial)		
Krishnaiah, P. R.		
6. REPORT DATE	7a. TOTAL NO. OF PAGES	7b. NO. OF REFS
March 1967	15	14
8a. RESEARCH ORIGIN IN-house Research		9a. ORIGINATOR'S REPORT NUMBER(S)
A. PROJECT NO. 7071		
c. 61445014		9b. OTHER REPORT NO(S) (Any other numbers that may be assigned this report)
d. 681307		ARL 67-0044
10. AVAILABILITY/LIMITATION NOTICES		
1. Distribution of this document is unlimited		
11. SUPPLEMENTARY NOTES		12. SPONSORING MILITARY ACTIVITY
		Aerospace Research Laboratories (ARM) Office of Aerospace Research, USAF Wright-Patterson AFB, Ohio
13. ABSTRACT		
<p>In this paper, we consider the problems of testing for the equality of covariance matrices against certain alternatives when the underlying populations are multivariate normal. The alternative hypotheses considered are (i) at least one covariance matrix is not equal to the covariance matrix of the next population, (ii) at least one covariance matrix is not equal to the covariance matrix of the standard population and (iii) at least one covariance matrix is not equal to the covariance matrix of another population.</p>		

DD FORM 1473
1 JAN 64

Unclassified
Security Classification

Unclassified

Security Classification

14. KEY WORDS	LINK A		LINK B		LINK C	
	ROLE	WT	ROLE	WT	ROLE	WT
Simultaneous tests Covariance matrices Multivariate normal						

INSTRUCTIONS

1. **ORIGINATING ACTIVITY:** Enter the name and address of the contractor, subcontractor, grantee, Department of Defense activity or other organization (corporate author) issuing the report.

2a. **REPORT SECURITY CLASSIFICATION:** Enter the overall security classification of the report. Indicate whether "Restricted Data" is included. Marking is to be in accordance with appropriate security regulations.

2b. **GROUP:** Automatic downgrading is specified in DoD Directive 5200.10 and Armed Forces Industrial Manual. Enter the group number. Also, when applicable, show that optional markings have been used for Group 3 and Group 4 as authorized.

3. **REPORT TITLE:** Enter the complete report title in all capital letters. Titles in all cases should be unclassified. If a meaningful title cannot be selected without classification, show title classification in all capitals in parenthesis immediately following the title.

4. **DESCRIPTIVE NOTES:** If appropriate, enter the type of report, e.g., interim, progress, summary, annual, or final. Give the inclusive dates when a specific reporting period is covered.

5. **AUTHOR(S):** Enter the name(s) of author(s) as shown on or in the report. Enter last name, first name, middle initial. If military, show rank and branch of service. The name of the principal author is an absolute minimum requirement.

6. **REPORT DATE:** Enter the date of the report as day, month, year, or month, year. If more than one date appears on the report, use date of publication.

7a. **TOTAL NUMBER OF PAGES:** The total page count should follow normal pagination procedures, i.e., enter the number of pages containing information.

7b. **NUMBER OF REFERENCES:** Enter the total number of references cited in the report.

8a. **CONTRACT OR GRANT NUMBER:** If appropriate, enter the applicable number of the contract or grant under which the report was written.

8b, 8c, & 8d. **PROJECT NUMBER:** Enter the appropriate military department identification, such as project number, subproject number, system numbers, task number, etc.

9a. **ORIGINATOR'S REPORT NUMBER(S):** Enter the official report number by which the document will be identified and controlled by the originating activity. This number must be unique to this report.

9b. **OTHER REPORT NUMBER(S):** If the report has been assigned any other report numbers (either by the originator or by the sponsor), also enter this number(s).

10. **AVAILABILITY/LIMITATION NOTICES:** Enter any limitations on further dissemination of the report, other than those

imposed by security classification, using standard statements such as:

- (1) "Qualified requesters may obtain copies of this report from DDC."
- (2) "Foreign announcement and dissemination of this report by DDC is not authorized."
- (3) "U. S. Government agencies may obtain copies of this report directly from DDC. Other qualified DDC users shall request through _____."
- (4) "U. S. military agencies may obtain copies of this report directly from DDC. Other qualified users shall request through _____."
- (5) "All distribution of this report is controlled. Qualified DDC users shall request through _____."

If the report has been furnished to the Office of Technical Services, Department of Commerce, for sale to the public, indicate this fact and enter the price, if known.

11. **SUPPLEMENTARY NOTES:** Use for additional explanatory notes.

12. **SPONSORING MILITARY ACTIVITY:** Enter the name of the departmental project office or laboratory sponsoring (paying for) the research and development. Include address.

13. **ABSTRACT:** Enter an abstract giving a brief and factual summary of the document indicative of the report, even though it may also appear elsewhere in the body of the technical report. If additional space is required, a continuation sheet shall be attached.

It is highly desirable that the abstract of classified reports be unclassified. Each paragraph of the abstract shall end with an indication of the military security classification of the information in the paragraph, represented as (TS), (S), (C), or (U).

There is no limitation on the length of the abstract. However, the suggested length is from 150 to 225 words.

14. **KEY WORDS:** Key words are technically meaningful terms or short phrases that characterize a report and may be used as index entries for cataloging the report. Key words must be selected so that no security classification is required. Identifiers, such as equipment model designation, trade name, military project code name, geographic location, may be used as key words but will be followed by an indication of technical content. The assignment of links, rules, and weights is optional.

Unclassified

Security Classification