

AD

US ARMY MEDICAL RESEARCH LABORATORY

FORT KNOX, KENTUCKY 40121

REPORT NO. 744

IMMUNIZATION STUDIES WITH NAJA NAJA
VENOM DETOXIFIED BY PHOTOOXIDATION

(Final Report)

by

Walter F. Kocholaty, Ph.D.

Edith B. Ledford, A. B.

Thomas A. Billings, B.S.

Joyce C. Goetz, B.A.

and

Billy D. Ashley, M. S.

30 June 1967

AD658973

Distribution of this document is unlimited.

UNITED STATES ARMY
MEDICAL RESEARCH AND DEVELOPMENT COMMAND



REC'D
OCT 2 1967
RECEIVED

C

AD

REPORT NO. 744

IMMUNIZATION STUDIES WITH NAJA NAJA
VENOM DETOXIFIED BY PHOTOOXIDATION

(Final Report)

by

Walter F. Kocholaty, Ph. D.
Edith B. Ledford, A. B.
Thomas A. Billings, B. S.
Joyce C. Goetz, B. A.
and
Billy D. Ashley, M. S.

Biochemistry Division
US ARMY MEDICAL RESEARCH LABORATORY
Fort Knox, Kentucky 40121

30 June 1967

Production of Polyvalent Antivenins
Work Unit No. 149
In-House Laboratory Independent Research
Task No. 00
In-House Laboratory Independent Research
DA Project No. 3A013001A91C

Distribution of this document is unlimited.

USAMRL Report No. 744
DA Project No. 3A013001A91C

ABSTRACT

IMMUNIZATION STUDIES WITH NAJA NAJA
VENOM DETOXIFIED BY PHOTOOXIDATION

OBJECTIVE

The object was to detoxify cobra venom by photooxidation in such a way that the immunogenic properties of this venom were preserved.

RESULTS

Photooxidation of cobra venom by visible light in the presence of methylene blue resulted in a relatively slow detoxification of the venom. The detoxified venom when injected in rabbits caused a very weak immunogenic response.

CONCLUSION

The phenomenon of photooxidation in the detoxification of venoms appears to be generally applicable; however, the conditions for the immunogenic response may vary from venom to venom.

IMMUNIZATION STUDIES WITH NAJA NAJA VENOM DETOXIFIED BY PHOTOOXIDATION

INTRODUCTION

Detoxification of snake venoms by photooxidation with visible light in the presence of methylene blue was demonstrated first on Crotalus atrox venom (1) and extended to other venoms (2). Three of the photooxidized venoms tested for their immunogenic properties gave rise in rabbits to antibodies which protected mice from the toxic effects of these venoms. In an attempt to apply similar techniques to Naja naja venom some differences and difficulties were encountered which necessitated more detailed investigations reported herewith.

EXPERIMENTAL PROCEDURE

The venom was collected at the snake colony of this laboratory under ice, quick-frozen and lyophilized. The pooled collections were ground and sieved as described previously (1,2) and stored at 4 °C in a desiccator in the dark. The protein content of the centrifuged and filtered (Millipore Filter of 0.45 µm pore size) venom solution was 82.5% (biuret). Phospholipase A activity was determined as previously described (3). No proteolytic activity against casein could be demonstrated and esterase activity against TAME (p-toluene sulfonyl-L-arginine methyl ester) was absent. Photooxidation was carried out at pH 4.0 and 8.5. Toxicity and qualitative precipitin tests were conducted as described previously (2). The LD₅₀ for this particular batch of venom was 7.5 γ I.V. and 8 γ I.P. (20-25 g mouse).

RESULTS

Effect of Photooxidation on Loss of Toxicity and Phospholipase A Activity. Table 1 summarizes the rapid loss of phospholipase A activity and concomitant loss in venom toxicity with increasing O₂ uptake and time of photooxidation.

Immunization of Rabbits and Mice with Photooxidized Venom of Naja naja. Venom of N. naja was photooxidized to an uptake of 20 µl O₂/mg protein using the reaction mixture and conditions described in Table 1 and stored in ice and in the dark. Prior to injections the photooxidized venom was diluted with physiological saline and mixed with an equal amount of Algivant (Colab Laboratories, Inc.). The weekly

immunization schedule for rabbits and mice was identical. Each animal received three injections a week (Monday, Wednesday, Friday), mice for 3 and rabbits for 4 weeks consecutively. The first two doses for mice (mature white Swiss, 30 to 40 g) were 32 γ of photooxidized venom and the rest 64 γ administered subcutaneously in a volume of 0.1 ml for a total of 512 γ .

Rabbits (white New Zealand) weighing 2-3 kg, received 1 mg venom protein in 1.0 ml for the first two injections and 2 mg in 1.6 ml for all subsequent injections for a total of 22 mg per animal. Injections were administered intramuscularly and alternated between the two hind legs. After a rest period of 1 week, mice were challenged with unaltered venom and rabbits were bled to death by heart puncture.

Neutralization and Protection Studies. When the mice which had been immunized with the photooxidized N. naja venom were challenged intravenously with increasing amounts of unaltered venom, all mice died within 24 hours even with the lowest dose (2 LD₅₀) employed.

Of the 10 rabbits (23-31, 33) immunized with photooxidized venom (Table 2), four did not show a positive (qualitative) percipitin test with the highest concentration of venom used, and one was positive only to the fourth dilution. The rest were weakly positive up to the sixth dilution.

The sera of the rabbits showing the highest (#23, 24, 25, 31, 33 = "A") and the lowest (#26-30 = "B") titer were combined and their γ -globulins isolated separately (4).

The neutralization and protection studies were carried out on mice using the γ -globulin obtained from rabbits of group A. As shown in Table 3, the globulins were practically inert in the neutralization as well as in the protection studies.

Since the results of the neutralization and protection studies were disappointing, five more rabbits were immunized with twice the amount of photooxidized venom for a total of 44 mg using the same adjuvant and an identical immunization and bleeding schedule as before. No lesions of significance were recognized in the tissues of all animals injected with N. naja venom. This information was furnished by Colonel M. A. Ross, VC, formerly Director, Pathology Division, USAMRL. One rabbit died during this period from causes not related to the immunization procedure.

All the rabbits (#32, 34, 37, 38 = "C") immunized with the higher amounts (44 mg total) of photooxidized venom responded with a positive though somewhat erratic precipitin test (Table 2). The γ -globulins were isolated as described before. The results shown in Table 3 are almost identical with the ones obtained in the previous experiment, where only one half as much photooxidized venom had been used in the immunization schedule.

Photooxidation of *Naja naja* Venom at Acidic pH and at Low Temperature. The comparatively low protection obtained with the rabbit immune globulins prompted a more detailed investigation into the various parameters of the photooxidation mechanism of the particular venom. Photooxidation may be carried out in a more selective manner utilizing an acidic reaction thus excluding histidine and tyrosine from photooxidative destruction (5, 6). Another specific variation may be introduced by working at temperatures below 37° (7).

Examination of the effect of pH on detoxification by photooxidation of this venom revealed the usual pattern with a pH optimum at about 8.5* and a steep drop toward the acid side, similar to that demonstrated with *Crotalus atrox* (1).

Studies of other parameters revealed that apparently the only factor related to the decrease in toxicity is the amount of O₂ uptake per mg venom protein irrespective of the concentration of the venom, the length of time of photooxidation, the pH of the reaction mixture (8.5 or 4.0), and the temperature (37° or 10° C). No significant volumes of CO₂ were developed during any of the variations in the photooxidation procedure. A summary of these experiments is illustrated in Figure 1.

DISCUSSION

The photooxidation of proteins by visible light in the presence of methylene blue permits gradual and subtle alterations of certain amino acid side-chains, which can be further modified by carrying out the reaction varying the pH and the temperature. The changes resulting in the molecule affected have provided an insight into functional groups of enzymes and a method for the detoxification of toxins and venoms(8). The latter has proven useful in providing modified venoms capable of

* At pH 9.0 and a concentration of 10 mg venom protein per ml a precipitate forms very rapidly in the initial stages of photooxidation.

eliciting antibodies in experimental animals with simultaneous abolishment of the necrogenic and hemolytic properties.

Why immunization with the photooxidized venom failed to produce an immunoglobulin capable of protecting mice is not understood at the present time. Attempts to vary the conditions for photooxidation (Fig. 1) revealed nothing of relevance to conduct photooxidation in a more promising way, or to indicate abnormalities in the chemical nature of this venom.

One may, however, speculate on two facets which stand out by contrast and, although not explainable at this time, may have a bearing on the failure to produce an immunoglobulin.

One, is the observation that the photooxidative detoxification of Naja naja venom proceeds at a considerably slower rate and with a high O_2 consumption compared with the other venoms investigated. From this one might suspect a unique arrangement of the four amino acids susceptible to photooxidation. The isolation of cobrotoxin, one of the toxic proteins from Formosan cobra venom, was reported recently by Yang (9). The amino acid composition of this compound has not yet been reported. Its molecular weight is 11,000. In this connection it is of interest to compare the rapid loss of phospholipase A activity with the very slow decrease in toxicity, a fact that would tend to indicate that this enzyme is, at least in its native form, a minor factor in the toxicity of this venom.

Two, is the observation of De Vries and his group (10) that the Vipera palestinae neurotoxin by itself is weakly antigenic, but when combined with carboxymethyl cellulose (CMC) the immunogenic properties are enhanced to such a degree that minute amounts of carboxymethyl cellulose-bound neurotoxin elicit a high antibody response. The Vipera palestinae neurotoxin recently isolated by these workers (11) has a low molecular weight (11,600) similar to that of the cobrotoxin of Yang.

De Vries and co-workers conclude that if a low molecular weight characterizes venom neurotoxins the increase in immunogenicity obtained by binding to carboxymethyl cellulose "may be a reflection of the increase in molecular weight of the antigen."

SUMMARY

Photooxidation of cobra venom by visible light in the presence of methylene blue resulted in a relatively slow detoxification of the venom.

The γ -globulin isolated from rabbits immunized with the detoxified venom gave poor protection in the animal experiments. Possible reasons for the weak response of the immune serum are discussed.

LITERATURE CITED

1. Kocholaty, W. Detoxification of Crotalus atrox venom by photooxidation in the presence of methylene blue. USAMRL Report No. 628, 1965 (DDC AD No. 466948); Toxicon, 3: 175, 1966.
2. Kocholaty, W. and B. D. Ansley. Detoxification of the venoms of the Russell's viper (Vipera russellii) and the water moccasin (Agkistrodon piscivorus) by photooxidation. USAMRL Report No. 629, 1965 (DDC AD No. 466949); Toxicon, 3: 187, 1966.
3. Kocholaty, W. A rapid and simple phospholipase A assay. USAMRL Report No. 652, 1965 (DDC AD No. 485846); Toxicon, 4: 1, 1966.
4. Campbell, D. H., J. S. Garvey, N. E. Cremer, and D. H. Sussdorf. Methods in Immunology. New York: W. A. Benjamin, Inc., 1965.
5. Sluyterman, L. A. AE. Photooxidation sensitized by proflavine of a number of protein constituents. Biochim. Biophys. Acta, 60: 557, 1962.
6. DasGupta, B. R. and D. A. Boroff. Increased selectivity of photooxidation for amino acid residues. Biochim. Biophys. Acta, 97: 157, 1965.
7. Weil, L. On the mechanism of the photooxidation of amino acids sensitized by methylene blue. Arch. Biochem. Biophys. 110: 57, 1965.
8. Spikes, J. D. and B. W. Glad. Photodynamic action. Photochem. & Photobiol. 3: 471, 1964.
9. Yang, C. C. Crystallization and properties of Cobrotoxin from Formosan cobra venom. J. Biol. Chem. 240: 1616, 1965.

10. Moroz, C., N. Goldblum, and A. De Vries. Preparation of Vipera palestinae antineurotoxin using carboxymethyl-cellulose-bound neurotoxin as antigen. *Nature*, 200: 697, 1963.
11. Moroz, C., A. De Vries, and M. Sela. Isolation and characterization of a neurotoxin from Vipera palestinae venom. *Biochim. Biophys. Acta*, 124: 136, 1966.

TABLE I
EFFECT OF PHOTOOXIDATION ON LOSS OF TOXICITY
AND PHOSPHOLIPASE A ACTIVITY

% loss in phospholipase A activity	μ l O ₂ uptake/ mg venom protein	Time of photooxidation (min)	LD ₅₀ tolerated (20 g mouse)
69	5	10	2
93	10	29	4
100	20	111	8
-	27	258	16

Reaction mixture: 5 mg venom protein and 0.1 mg methylene blue in 0.1 M TRIS buffer, pH 8.5; total volume 2 ml; t = 37°C. No measurable CO₂ production was observed during the time of photooxidation.

TABLE 2

RESULTS OF THE QUALITATIVE PRECIPITIN TESTS
OF SERA OF RABBITS IMMUNIZED WITH
PHOTOOXIDIZED VENOM OF NAJA NAJA

Total amount of photooxidized venom injected	Rabbit No.	Tube No. *								
		1	2	3	4	5	6	7	8	9
22 mg	23, 24, 25	+	+	+	+	+	+	-	-	-
	26	+	+	+	+	-	-	-	-	-
	27, 28, 29, 30	-	-	-	-	-	-	-	-	-
	31, 33	+	+	+	+	+	+	-	-	-
44 mg	32	-	-	+	+	+	-	-	-	-
	34	+	-	+	+	-	-	-	±	-
	37	+	-	+	-	+	-	-	-	-
	38	+	±	±	+	+	+	-	±	-

* First tube contained 10 mg venom protein in 0.5 ml "dilute saline-borate buffer," pH 8.4 (4). All subsequent dilutions were 1:5. To each tube, two drops of serum were added. After 60 minutes at 37° C, the tubes were placed in the cold and readings were taken 24 hours later. Controls contained either saline and serum or saline and venom (10 mg) in amounts stated above. Both controls were negative.

TABLE 3
NEUTRALIZATION OF, AND PROTECTION AGAINST, NAJA NAJA VENOM IN MICE BY
RABBIT IMMUNE GLOBULIN

LD ₅₀	Neutralization by Imm. Glob. "A"	Protection by Imm. Glob. "A"	Neutralization by Imm. Glob. "C"	Protection by Imm. Glob. "C"	Normal globulin control*	Venom control I. P.
1	0/3	3/3	0/3	0/4	1/4	5/8
2	0/3	3/3	0/4	0/4	4/4	8/8
3	3/3	3/3	1/4	2/4		
4	3/3	2/2	3/4	4/4		

Numbers are deaths over the total number of mice used. Readings were done 24 hours after injection.

Three or more mice received either 25 mg immune or normal rabbit globulin and the LD₅₀ stated above in the neutralization tests.

Twenty-five mg of rabbit immune globulin was injected (I. V.) in a volume of 0.25 ml followed in 30 minutes by injection (I. P.) of 1-4 LD₅₀ of Naja naja venom in the protection tests. Controls received venom only.

"A" Derived from rabbits (#23-25, 31, 33) immunized with a total of 22 mg photooxidized venom.

"C" Derived from rabbits (#32, 34, 37, 38) immunized with a total of 44 mg photooxidized venom.

* Rabbit γ-globulin, fraction II, purchased from Pentex Corporation.




Fig. 1. Conditions for photooxidation (pH temperature) are recorded on the O_2 uptake curves: mg refer to mg venom protein in 2 ml of reaction mixture. Numbers in squares indicate LD_{50} tolerated by 20 to 25 g mouse at the particular O_2 uptake.

DISTRIBUTION LIST

10 copies to:

US Army Medical Research and Development Command
Department of the Army, Washington, D. C. 20315

20 copies to:

Defense Documentation Center, Cameron Station,
Alexandria, Virginia 22314

1 copy to:

US Army Combat Development Command, Medical Service Agency,
Brooke Army Medical Center, Fort Sam Houston, Texas 78234

Walter Reed Army Institute of Research, WRAMC
Washington, D. C. 20012

US Army Medical Equipment Research and Development Laboratory,
Fort Totten, Flushing, Long Island, New York 11359

US Army Medical Biomechanical Research Laboratory, WRAMC,
Forest Glen Section, Washington, D. C. 20012

US Army Medical Research and Nutrition Laboratory, Fitzsimons
General Hospital, Denver, Colorado 80240

US Army Research Institute of Environmental Medicine, US Army
Natick Laboratories, Natick, Massachusetts 01762

US Army Aeromedical Research Unit, US Army Aviation Center,
Fort Rucker, Alabama 36362

The Historical Unit, US Army Medical Service, Forest Glen Section,
WRAMC, Washington, D. C. 20012

Technical Director of Research, Armed Forces Institute of
Pathology, Washington, D. C. 20012

Commanding Officer, US Army Environmental Health Agency,
Edgewood Arsenal, Maryland 21041

UNCLASSIFIED

Security Classification

DOCUMENT CONTROL DATA - R & D

(Security classification of title, body of abstract and indexing no. station must be entered when the overall report is classified)

1. ORIGINATING ACTIVITY (Corporate author) US Army Medical Research Laboratory Fort Knox, Kentucky 40121		2a. REPORT SECURITY CLASSIFICATION UNCLASSIFIED	
		2b. GROUP	
3. REPORT TITLE IMMUNIZATION STUDIES WITH <u>NAJA NAJA</u> VENOM DETOXIFIED BY PHOTO- OXIDATION			
4. DESCRIPTIVE NOTES (Type of report and inclusive dates) Final Report			
5. AUTHOR(S) (First name, middle initial, last name) Walter F. Kocholaty, Edith B. Ledford, Thomas A. Billings, Joyce C. Goetz, and Billy D. Ashley			
6. REPORT DATE 30 June 1967	7a. TOTAL NO. OF PAGES 10	7b. NO. OF REFS 11	
8a. CONTRACT OR GRANT NO.		8b. ORIGINATOR'S REPORT NUMBER(S)	
a. PROJECT NO. 3A013001A91C		744	
*Task No. 00		8c. OTHER REPORT NO(S) (Any other numbers that may be assigned this report)	
*Work Unit No. 149			
10. DISTRIBUTION STATEMENT Distribution of this document is unlimited.			
11. SUPPLEMENTARY NOTES		12. SPONSORING MILITARY ACTIVITY US Army Medical Research and Develop- ment Command, Washington, D. C. 20315	
13. ABSTRACT Photooxidation of cobra venom by visible light in the presence of methylene blue resulted in a relatively slow detoxification of the venom. The γ -globulin isolated from rabbits immunized with the detoxified venom gave poor protection in the animal experiments. Possible reasons for the weak response of the immune serum are discussed. (U)			

DD FORM 1473

REPLACES DD FORM 1473, 1 JAN 60, WHICH IS OBSOLETE FOR ARMY USE.

UNCLASSIFIED

Security Classification

UNCLASSIFIED

Security Classification

14. KEY WORDS	LINK A		LINK B		LINK C	
	ROLE	WT	ROLE	WT	ROLE	WT
Venoms Immune Serums Immunization Toxoid Toxicology						

AG 1869-I-Army-Knox-Oct 67-90

UNCLASSIFIED

Security Classification