


USNRDL-TR-87-III

24 June 1987

**FISSION PRODUCT DECAY CHAINS: SCHEMATICS WITH
BRANCHING FRACTIONS, HALF-LIVES, AND
LITERATURE REFERENCES**

by
G. R. Crocker



**U.S. NAVAL RADIOLOGICAL
DEFENSE LABORATORY**

SAN FRANCISCO • CALIFORNIA • 94135

Approved by the
CLEARINGHOUSE
for the Naval Radiological Defense Laboratory
Information Springfield, Va. 22151

This document has been approved
for public release and sale; its
distribution is unlimited.

40

PHYSICAL CHEMISTRY BRANCH
E. C. Freiling, Head

NUCLEAR TECHNOLOGY DIVISION
R. Cole, Head

ADMINISTRATIVE INFORMATION

The work reported is part of a project sponsored by the Division of Biology and Medicine of the Atomic Energy Commission under Contract AT(49-7)-1963, Mod. 6.

DDC AVAILABILITY NOTICE

This document has been approved for public release and sale; its distribution is unlimited.

ACCESSION NO.	
CPETI	KNIFE SECTION <input checked="" type="checkbox"/>
BOB	DIFF. SECTION <input type="checkbox"/>
(MATERIALS)	<input type="checkbox"/>
JUSTIFICATION	
BY	
DISTRIBUTION/AVAILABILITY CODE	
NOV.	APRIL and/or SPECIAL

Eugene P. Cooper

Eugene P. Cooper
Technical Director

D.C. Campbell

D.C. Campbell, CAPT USN
Commanding Officer and Director

ABSTRACT

Schematic diagrams of the decay chains for the fission products of uranium and plutonium are presented, with half-lives of the nuclides, branching fractions, and literature references. The updated data were obtained from a review of the literature through February 1967.

This document has been approved
for public release and sale; its
distribution is unlimited.

SUMMARY

The Problem

Precise descriptions of the genetic relationships in fission-product decay chains, as well as half-life values for the radionuclides comprising the chains, are required for the calculation of the radiation properties of fission-product mixtures.

Findings

Accurate knowledge of these properties is required in assessments of the consequences of reactor operations and accidents, nuclear weapon testing, and nuclear warfare. Updated individual radionuclide data were obtained by review of the literature through February 1967. The information is presented in schematic form, along with literature references.

INTRODUCTION

The neutron fission of uranium or plutonium produces several hundred radionuclides, mostly beta-emitters, belonging to about 90 mass chains. Calculation of the radiation properties of fission-product mixtures as a function of time requires precise descriptions of the decay chains, including the parent-daughter relationships of the nuclides, the branching fractions where the chains are branched, and the half-lives of all the nuclides.*

This kind of information has been generated and/or collected at this Laboratory and utilized in computer calculations of fission-product abundances and activities for several years. An early version of the decay chains was published in Ref. A. Many of the half-lives in that version were estimated and some chain-branching fractions were arbitrarily assigned. Also, a sizable number of metastable states among the fission products were not known to exist at that time. The appearance of Katcoff's (Ref. B) and Stehn's (Ref. C) chain data led to considerable improvement. In 1964, a new version (Ref. D) of the fission-product decay chains, based largely on the decay schemes through 1963, in the Nuclear Data Sheets (NDS), see Ref. F, was prepared for use in this Laboratory.

The collection of new data has continued and a large number of changes has accumulated since 1963. These were incorporated into the decay chains, of which the latest versions are presented here.

SOURCES

The list of references is divided into (1) general references, which are cited by the letters A through F; and (2) numbered references, which were sources of specific items of information. The decay chains and the data as presented evolved via References A, B, C, and the NDS data sheets, except where indicated by parenthesized citations of the sources in the specific reference list.

*The independent yields of the nuclides are required as well, but these are not treated here. Predicted independent yields for several fission types can be found in Ref. 71.

The data collected since 1963 which have been incorporated into this presentation were accumulated by regular and systematic perusal of periodicals and reports in the nuclear science field. The pertinent information tends to appear in a handful of major journals, as a cursory examination of the list of references will confirm, and it is believed that most of these were considered. A crosscheck with the 1964 and 1966 Recent References of NDS has provided welcome reassurance that no very sizable body of publication was overlooked. Another crosscheck was provided by the reference list (through 1963) in Herrmann's excellent historical account of fission-product radiochemistry, Ref. E.

Since most of the NDS decay schemes for the fission-product mass chains (on which Ref. D was based) have not been revised since 1961 or earlier, the chain data presented here incorporate a large number of more recently published values. Many of the changes are minor, but some are important. The reports on which the changes are based have not been subjected to the critical study and evaluation which the Nuclear Data Group will eventually provide. The chain data as presented here should therefore be regarded as an interim summary, subject to further emendation and amplification.

The intention was to include in the chains all the products, including isomers with half-lives greater than 0.5 seconds, known to occur in the neutron fission of uranium and plutonium. Mass chains lower than mass 72 or higher than mass 161 were disregarded because of their insignificant contributions. Many of the short-lived nuclides at the beginnings of the chains have never been observed in fission products, but their presence is predicted by the models of charge distribution. Some of the shielded nuclides shown at the ends of the chains may not be produced in detectable amounts. However, very small yields of ^{86}Rb and ^{134}Cs have been reported⁷⁰ and it seems possible that similar shielded end-members in other chains will eventually be identified in fission products.

Some important gaps in the information remain. The lack of half-life values for short-lived early chain members is conspicuous. Almost all the chains begin with a few members whose half-lives are only roughly estimated at best. Estimated half-lives are marked with an asterisk and are mostly taken from Ref. A. Experience has shown that they may be in error by several orders of magnitude. Branching fractions are frequently unknown or are uncertain, particularly in the more complicated chains. Estimated branching fractions, also asterisked, are usually an arbitrary 50-50 splitting, but in a few cases have some basis in reports of fragmentary or incomplete studies. It is often unknown which of two states of a nuclide is the ground state. For calculational and schematic purposes, distinguishing symbolism is desirable, nonetheless. Therefore arbitrary assignments of "metastable" and "ground" states were made in cases of uncertainty.


The relationship of several short-lived states to the fission process and to the chain-decay path (or paths) is unknown. On the other hand, most of the longer-lived chain members have been well characterized and their positions in the chains are fairly certainly known. However, some exceptions exist. It is not clear, for instance, whether the mass-98 chain decays mainly through a 51-minute or a 2-minute isomer; or, indeed, whether either of these half-lives is correct. Certain chain-decay features, once regarded as well established, have required important modifications. For instance, until recently the mass-117 chain was believed to decay partly through a 3-hour ^{117m}Cd and partly through a 50-minute ^{117g}Cd . Now both isomers of ^{117}Cd are known to have half-lives near 3 hours.^{20, 34, 44, 59, 60} (The further decay pathways of the mass-117 chain through two In isomers and two Sn isomers remain uncertain.) Similarly, the reported 1-hour state of ^{129}Sn recently was established as one of about 6 minutes.⁴⁰ It is quite possible that further important changes of this kind may occasionally be necessary.

NOTATION


The decay chains are represented in the conventional symbolism with arrows indicating parent-daughter relationships. Branching fractions are written along the arrows where chain branching occurs. Half-lives are written below the nuclides, with abbreviations s for seconds, M for minutes, H for hours, D for days and Y for years. The numbers in parentheses which accompany some of the data refer to the sources in the List of References. Half-lives and branching fractions marked with an asterisk are estimates (see preceding section).

The arrows ordinarily indicate negatron decay but in a few cases (^{122g}Sb , ^{128}I , ^{152m}Eu and ^{152g}Eu), where the direction of the arrow is reversed, either positron emission or electron capture, or a combination of the two, occurs. Decay by neutron emission to the mass chain of next lower number is indicated by dotted lines for ^{85}As , ^{87}Br , ^{88}Br , ^{89}Br , ^{90}Br , ^{137}I , ^{138}I and ^{139}I . Since neutron emission is immediately preceded by beta emission, these lines terminate at elements of higher atomic number. The unsubstantiated possible decay by neutron emission of one Sb isotope (^{135}Sb) is indicated by dotted lines and the symbol (-n?). Decay by alpha emission has not been indicated. The only fission-product mass chains which terminate in alpha-emitters are 144 (Nd, half-life of 10^{15} years) and 147 (Sm, half-life of 10^{11} years). The shielded nuclides ^{152}Eu and ^{146}Pm , which may be produced in very small quantities, decay partly to the alpha-emitters ^{152}Gd (10^{14} years) and ^{146}Sm (5×10^7 years), respectively. In some cases where isomerism occurs, it is not certain whether one of the isomers has a precursor or what its daughter nuclides are. In such cases dotted lines and question marks indicate the missing information.





SCHEMATIC DIAGRAMS OF FISSION PRODUCT DECAY
CHAINS: MASS NUMBERS 72 THROUGH 161





(a) Branching to $^{77\text{m}}\text{Se}$ and its $T_{1/2}$ taken from NDS.




(a) Decay of ^{78}Ge according to Ref. 84.











(a) ^{89}Sr branches $< 0.01\%$ to $^{89\text{m}}\text{Y}$.




(a) $^{90\text{m}}\text{Y}$ is not the daughter of ^{90}Sr , see Ref. 18.








(a) Branching fractions, suggested by the NDS scheme, are approximate.





(a) The 51-M Nb has been observed, Refs. 1 and 17, but is not the daughter of ${}^{98}\text{Zr}$. The 2-M isomer is postulated in Ref. 1.







(a) NNIS lists both a 3-M and a 11.5-M Nb state, but suspects they may be the same.











(a) Ref. 73 reports 0.056 % yield of 5.3-H ^{103}Mo in 14-Mev neutron fission of ^{238}U .












(a) Other versions, considerably different and equally plausible, could be constructed for the mass-117 chain branching from the available data; e.g. Refs. 33, 34, 59, 60, 64, and 65.




(a) Ref. 21 states that $4.7\text{-M } {}^{118\text{m}}\text{In}$ is not a daughter of ${}^{118}\text{Cd}$.





(a) Ref. 90 proposes a very different scheme for the Cd-In-Sn branching in this chain.





(a) In and Sn chain decay from Ref. 4.










(a) The In and Sn portion of the chain decay is according to Ref. 7.





(a) The NDS 1961 sheet gives 1.8 H for $T_{1/2}$ for ^{129}Sn . Hagebö et al., J. Inorg. Nucl. Chem. 24, 1301 (1962) reported a 1.0 H and an 8.8 M state. Dropesky and Orth, Ref. 6 (1962), found only a 6.2-M state. Chu and Marinaky, Ref. 40 (1966), showed that Hagebö's 1.0-H isomer was an artifact and at the same time reported the 2-M isomer. It is not stated whether the 2-M or the 6.2-M isomer is the ground state. The assignment here is arbitrary.









(a) ${}^{144}\text{Nd}$ is an alpha-emitter with half-life of about 2.4×10^{15} Y, according to Ref. 50.





(a) The long-lived ${}^{146}\text{Sm}$ decays by alpha emission to stable ${}^{142}\text{Nd}$.





(a) ${}^{147}\text{Sm}$ is an alpha-emitter with half-life of about 10^{11} Y.












(a) ${}^{148\text{m}}\text{Pm}$ $T_{1/2}$ and branching, from NDS.





(a) ${}^{152}\text{Gd}$ is an alpha-emitter with half-life of about 10^{14} Y.





GENERAL REFERENCES

- A. R. C. Bolles and N. E. Ballou, "Calculated Activities and Abundances of U^{235} Fission Products," U. S. Naval Radiological Defense Laboratory, USNRDL-456, 1956.
- B. S. Katcoff, "Fission Products from Neutron-Induced Fission," *Nucleonics* 18, 201 (1960).
- C. J. Stehn, "Table of Radioactive Nuclides," *Nucleonics* 18, 186 (1960).
- D. D. Kochendorfer, "Calculated Activities of U^{235} Fission Products for Very Short Nuclear Reactor Operation. Vol. I. Discussion, Diagrams, and Figures," U. S. Naval Radiological Defense Laboratory Technical Report USNRDL-TR-757, June 1964.
- E. G. Herrmann, "Low-Energy Fission of Heavy Nuclei," *Radiochimica Acta* 3, 169 (1964).
- F. Nuclear Data Sheets, National Academy of Sciences--National Research Council.

SPECIFIC REFERENCES

1. C. J. Orth and R. K. Smith, "Short-Lived Isotopes of Nb and Zr from Fission," J. Inorg. Nucl. Chem. 15, 4 (1960).
2. M. Karras and J. Kantele, "New Nuclear Species Sc^{50m} and Rh^{110} ," Phys. Lett. 6, 98 (1963).
3. H. Weiss (This Laboratory), private communication.
4. A. C. Wahl and D. R. Nethaway, "Nuclear Charge Distribution in Symmetric Fission-- Sn^{121} Independent Yield," Phys. Rev. 131, 830 (1963).
5. E. der Mateosian and M. L. Sehgal, "Note on the Decay of Sb^{122m} ," Phys. Rev. 129, 2195 (1963).
6. B. J. Dropesky and C. J. Orth, "A Summary of the Decay of Some Fission Product Tin and Antimony Isotopes," J. Inorg. Nucl. Chem. 24, 1301 (1964).
7. D. E. Troutner, A. C. Wahl and R. L. Ferguson, "Independent Fission Yield of Sb^{127} ," Phys. Rev. 134, B1027 (1964).
8. G. N. Rao, V. R. Potnis, and H. S. Hans, "The Energy Levels of I^{129} ," Nucl. Phys. 44, 443 (1963).
9. N. Ballou (This Laboratory), private communication.
10. C. E. Bemis et al., "Half-Life of ^{135}Sb and Evidence on the Half-Life of ^{134}Sb ," J. Inorg. Nucl. Chem. 26, 213 (1964).
11. W. B. Clarke and H. G. Thode, "Mass Spectrometric Measurements of the Half-Lives of Kr^{87} , Kr^{88} , Xe^{135} and Xe^{138} ," Can. J. Phys. 42, 213 (1964).
12. K. Takahashi et al., "New Activities Produced by Fast-Neutron Bombardments," J. Phys. Soc. (Japan) 16, 1664 (1961).
13. A. F. Stehney and E. P. Steinberg, " Br^{86} --A New Nuclide," Phys. Rev. 127, 563 (1962).
14. J. E. Sattizahn, J. D. Knight and M. Kahn, "Short-Lived Bromine and Selenium Nuclides from Fission," J. Inorg. Nucl. Chem. 12, 206 (1960).

15. I. F. Croall and H. H. Willis, "The Fission Products ^{81}Se and ^{83}Se ," J. Inorg. Nucl. Chem. 24, 221 (1962).
16. D. E. Troutner, R. L. Ferguson and G. D. O'Kelley, "Yields and Half-Lives in the Mass-99 Fission Product Chain," Phys. Rev. 130, 1466 (1963).
17. A. C. Wahl et al., "Nuclear Charge Distribution in Low-Energy Fission," Phys. Rev. 126, 1112 (1962).
18. L. Haskin and R. Vandenbosch, "New Yttrium Isomers of Masses 86 and 90," Phys. Rev. 123, 184 (1961); R. L. Heath et al., "3.14-Hr Isomeric Level in Y^{90} ," *ibid.* p. 903; W. Alford, D. R. Koehler and C. E. Mandeville, "Evidence for an Isomeric State of Y^{90} ," *ibid.* p. 1365; W. S. Lyon, J. S. Eldridge and L. C. Bate, "3.1-Hour Y^{90m} ," *ibid.* p. 1747.
19. K. F. Flynn, L. E. Glendenin and E. P. Steinberg, "Half-Life Determinations by Direct Decay," Nucl. Sci. and Eng. 22, 416 (1965).
20. C. W. Tang and G. E. Gordon, in Massachusetts Institute of Technology Laboratory for Nuclear Science Progress Report NYO-10063, May 1, 1963.
21. C. E. Gleit and C. D. Coryell, "Decay of 49-min Cd^{118} and 5.1-sec In^{118} ," Phys. Rev. 122, 229 (1961).
22. D. C. Hoffman, "Half-Lives of Some Rare Earth Nuclides," J. Inorg. Nucl. Chem. 25, 1196 (1963).
23. B. F. Rider, J. P. Peterson, Jr., and C. P. Ruiz, "The Half-Life of Nd-149," J. Inorg. Nucl. Chem. 28, 266 (1966).
24. M. A. Wahlgren and W. W. Meinke, "Radiations from Short-lived Rare Gas Fission Products," J. Inorg. Nucl. Chem. 24, 1527 (1962).
25. V. R. Pandharipande, R. M. Singru and R. P. Sharma, "Decay of Pd^{111} and Pd^{111m} ," Phys. Rev. 140, B1488 (1965).
26. J. F. Emery, "A New Isomer of Bromine: ^{82m}Br ," J. Inorg. Nucl. Chem. 27, 903 (1965).
27. William R. Pierson, "Decay of Rh^{105} ," Phys. Rev. 140, B1516 (1965).
28. R. L. Auble and W. H. Kelly, "A Study of the Excited States of ^{127}I Populated in the Decay of ^{127}Te and ^{127m}Te ," Nucl. Phys. 73, 25 (1965).

29. G. Andersson, G. Rudstam and G. Soerensen, "Decay Data on Some Xe, I and Te Isotopes," Ark. Fys. 28, 37 (1964).
30. J. B. van der Kooi and H. J. van den Bold, "The Decay of ^{77}Ge and $^{77}\text{Ge}^m$," Nucl. Phys. 70, 449 (1965).
31. P. Kauranen, "The Short-lived ^{127}Sn Isomer," J. Inorg. Nucl. Chem. 27, 1713 (1965).
32. W. B. Walters, C. E. Bemis, Jr., and G. E. Gordon, "Decay Scheme of 25-Minute Te^{131g} ," Phys. Rev. 140, B268 (1965).
33. R. V. Mancuso and R. G. Arns, "Directional Correlation of Gamma Rays in Sn^{117} ," Nucl. Phys. 68, 513 (1965).
34. R. V. Mancuso and R. G. Arns, "The Decay of Cd^{117} and Cd^{117m} and Directional Correlation of Gamma Rays in In^{117} ," Nucl. Phys. 68, 504 (1965).
35. J. F. Neeson and R. G. Arns, "Directional Correlation of Gamma Rays in Rh^{105} ," Nucl. Phys. 68, 401 (1965).
36. E. T. Williams and C. D. Coryell, "Decay of 54-sec Br^{86} and 55-sec Br^{87} ," Phys. Rev. 144, 945 (1966).
37. L. Tomlinson, "Gamma Rays and Delayed Neutrons from Antimony and Arsenic Isotopes," J. Inorg. Nucl. Chem. 28, 287 (1966).
38. W. R. Daniels and D. C. Hoffman, "Fission Yields of ^{156}Eu , ^{157}Eu , ^{158}Eu , and ^{159}Eu ," Phys. Rev. 145, 911 (1966).
39. G. Goldstein and J. A. Dean, "Half-Life and Specific Activity of ^{99}Tc ," J. Inorg. Nucl. Chem. 28, 285 (1966).
40. P. Chu and J. A. Marinsky, "Isomers of ^{129}Sn ," J. Inorg. Nucl. Chem. 28, 1339 (1966).
41. N. K. Aras and G. E. Gordon, "The Fractional Chain Yield of ^{126}Sb in Thermal-Neutron Fission of ^{235}U ," J. Inorg. Nucl. Chem. 28, 763 (1966).
42. G. A. Brinkman, A. H. W. Aten, Jr., and J. Th. Veenboer, "Natural Radioactivity of K-40, Rb-87 and Lu-176," Physica 31, 1305 (1965).
43. P. Meyers and K. E. G. Loebner, "Decay of the 4.7-m isomer of ^{118}In and of the 48-sec isomer of ^{120}In ," Physica 31, 1774 (1965).

44. J. Brzosko et al., "Isomer Cross-section Ratios for Some Reactions Induced in Sn and Te Isotopes," Nucl. Phys. 74, 438 (1965).
45. J. Kantele and M. Karras, "Even Sn Isotopes Populated in Decay of In Isomers," Phys. Rev. 135, B9 (1964).
46. V. Akamian, "Decay characteristics of Sn^{93} ," U. S. Naval Radiological Defense Laboratory Technical Report USNRDL-TR-733 (1964).
47. H. W. Boyd and J. H. Hamilton, "Low-Energy Transitions in the Decay of ^{132}I ," Nucl. Phys. 72, 604 (1965).
48. V. V. Ovechkin and N. N. Demidovich, "Study of the Gamma-Radiation from Xe^{137} and Xe^{138} ," Soviet Physics (JETP) 20, 1123 (1965).
49. Janet S. Merritt and J. G. V. Taylor, "Decay of Cesium-137 Determined by Absolute Counting Methods," Anal. Chem. 37, 351 (1965).
50. R. D. Macfarlane, J. O. Rasmussen and M. Rho, "Theoretical Calculations of Properties of Rare-Earth Alpha Emitters," Phys. Rev. 134, B1196 (1964).
51. P. Boivinet, E. Calvet and Y. Ozias, "Détermination microcalorimétrique des caractéristiques de l'yttrium 90 et de la perte d'énergie due au freinage des betas," Comp. Rend. 256, 1103 (1963).
52. R. L. Auble and W. H. Kelly, "Beta and Gamma Spectroscopic Studies of 9.7-d ^{125}Sn ," Nucl. Phys. 79, 577 (1966).
53. L. Funke, H. Graber, K.-H. Kaun, H. Sodan and L. Werner, "Neue Niveaus im Dy^{161} ," Nucl. Phys. 55, 401 (1964).
54. T. Iwashita, "Decay of Eu^{159} ," J. Phys. Soc. (Japan) 20, 2105 (1965).
55. W. R. Pierson, H. C. Griffin and C. D. Coryell, "Characterization of Ru^{107} , Ru^{108} , Rh^{107} and Rh^{108} ," Phys. Rev. 127, 1708 (1962).
56. K. Fritze and T. J. Kennett, "Decay Measurements of Ba^{141} and Ba^{142} ," Phys. Rev. 127, 1262 (1962).
57. N. R. Johnson, G. D. O'Kelley and E. Eichler, "Decay of Rb^{90} ," Phys. Rev. 135, B36 (1964).
58. G. D. O'Kelley, N. H. Lazar and E. Eichler, "Decay of Rb^{89} ," Phys. Rev. 102, 223 (1956).

59. E. Hagebø, "Isomeric Yield Ratios and Fractional Chain Yields for ^{117}In and ^{117}Cd ," J. Inorg. Nucl. Chem. 27, 927 (1965).
60. K. Baskova et al., "Investigation of the Radiation of ^{117}Cd ," Sov. J. Nucl. Phys. 2, 288 (1966).
61. A. Delucci and A. Greendale (This Laboratory), personal communication.
62. F. Wunderlich, Thesis, Institut fuer allgemeine Wissenschaften der technische Hochschule, Muenchen, (1966).
63. K. Fritze, T. J. Kennett and W. V. Prestwich, "Half-Lives of ^{94}Rb , ^{94}Sr , ^{94}Y , ^{95}Rb , ^{95}Sr and ^{95}Y ," Can. J. Chem. 39, 675 (1961).
64. C. Tang, "Decay Schemes of the Cadmium-117 Isomers," Thesis, Massachusetts Institute of Technology (1965).
65. T. Kivikas and B. Forkman, "Low Energy Photofission of U^{238} ," Nucl. Phys. 64, 420 (1965).
66. R. E. Snyder and G. B. Beard, "Decay of Nb^{94} and Nb^{94m} ," Phys. Rev. 147, 867 (1966).
67. F. R. Roberts, E. J. Wheelwright and H. H. van Tuyl, "Detection of ^{146}Pm among the Products of Uranium Fission," J. Inorg. Nucl. Chem. 25, 1298 (1963).
68. C. E. Gleit and C. D. Coryell, "Decay of Cd^{119} and In^{119} Isomers," Phys. Rev. 124, 1914 (1961).
69. K. Wolfsberg et al., "Identification of ^{142}Xe and Measurements of Its Cumulative Yield from Thermal-Neutron Fission of ^{235}U ," J. Inorg. Nucl. Chem. 12, 201 (1959).
70. J. A. McHugh, "The Independent Yields of ^{86}Rb , ^{132}Cs , ^{134}Cs and ^{136}Cs in the Thermal Neutron Fission of ^{235}U ," J. Inorg. Nucl. Chem. 28, 1787 (1966).
71. L. E. Weaver, P. O. Strom and P. A. Killeen, "Estimated Total Chain and Independent Fission Yields for Several Neutron-Induced Fission Processes," U. S. Naval Radiological Defense Laboratory Technical Report USNRDL-TR-633, 5 March 1963.
72. Von P. Kienle et al., "Nachweis kurzlebiger Molybdaen-Spaltprodukte und Zerfall ihrer Technetium-Toechter III. Zerfallsreihe: ^{103}Mo - ^{103}Tc - ^{103}Ru ," Radiochim. Acta 1, 84 (1963).

73. R. Ganapathy, H. Inochi and P. Kuroda, "New Isomer: 5.3-h ^{103}Mo ," Phys. Rev. 151, 960 (1966).
74. J. A. Cooper, R. C. Gatti, J. O. Rasmussen and S. G. Thompson, "Gamma-Ray Spectrum of Sb^{121} ," J. Inorg. Nucl. Chem. 26, 2041 (1964).
75. G. P. Tercho and J. A. Marinsky, "The Fission Nuclide, 1.6-min ^{104}Mo ," J. Inorg. Nucl. Chem. 26, 1129 (1964).
76. H. Warhanek and R. Vandenbosch, "Relative Cross-Sections for Formation of the Shielded Isomeric Pair $^{134\text{m}}\text{Cs}$ and ^{134}Cs in Medium Energy Fission," J. Inorg. Nucl. Chem. 26, 669 (1964).
77. P. O. Strom et al., "Nuclear-Charge Distributions of Several Fission-Product Mass Chains," Phys. Rev. 144, 984 (1966).
78. M. Ishihara, "Decay of Zinc 72," J. Phys. Soc. (Japan) 18, 1111 (1963).
79. H. V. Weiss and N. E. Ballou, "Rapid Separation of Indium from Tin by Sublimation," J. Inorg. Nucl. Chem. 27, 1917 (1965).
80. K. F. Flynn, L. E. Glendenin, A. L. Harkness and E. P. Steinberg, "Half-Lives of ^{90}Sr and ^{137}Cs ," J. Inorg. Nucl. Chem. 27, 21 (1965).
81. H. A. Grench, K. J. Coop, H. O. Menlove and F. J. Vaughn, "A Study of the Spin Dependence of the Nuclear Level Density by Means of the $^{89}\text{Y}(n,\gamma)^{90\text{g}},^{90\text{m}}\text{Y}$ Reactions," Nucl. Phys. A94, 157 (1967).
82. P. del Marmol and M. Nève de Mévergnies, "Investigation of Delayed Neutron Precursors of As, Sb and Ge," J. Inorg. Nucl. Chem. 29, 273 (1967).
83. F. O. Lawrence, W. R. Daniels and D. C. Hoffman, "Radiochemical Purification of Tin and the Half-Lives of ^{121}Sn , ^{123}Sn , ^{125}Sn and ^{125}Sb ," J. Inorg. Nucl. Chem. 28, 2477 (1966).
84. K. Fritze, "Decay of Ge^{78} ," Nucl. Phys. 64, 303 (1965).
85. J. Lerner, "Half-Life of ^{85}Kr ," J. Inorg. Nucl. Chem. 25, 749 (1963).
86. I. M. H. Pagden, R. Jakeways and F. C. Flack, "The decays of Pm^{143} , Pm^{144} and Pm^{146} ," Nuclear Physics 48, 555 (1963).
87. L. R. Bunney and E. M. Scadden, "Decay Characteristics of Pm^{146} ," U. S. Naval Radiological Defense Laboratory Technical Report USNRDL-TR-1109, February 1967.

88. N. P. Archer and G. L. Keech, "Direct Measurements of the Half-Lives of Some Short-Lived Xenon and Cesium Isotopes," *Can. J. Phys.* 44, 1823 (1966).
89. A. von Raeckmann and H. Feuerstein, "Ueber Spalt-Technetium I," *Radiochimica Acta* 4, 111 (1965).
90. W. Kiefer, "Kernladungsverteilung bei der symmetrischen Spaltung von Uran-233 and Uran-238," Dissertation, Johannes Gutenberg-Universitaet, Mainz, 1966.
91. H. V. Weiss, "Near Symmetric Fission--Identification of Cd^{121} ," *Phys. Rev.* 139, B304 (1965).
92. Chart of the Nuclides, Federal Minister for Scientific Research, Federal Republic of Germany. Gersbach und Sohn Verlag (Munich), second edition (1965).

UNCLASSIFIED

Security Classification

DOCUMENT CONTROL DATA - R & D		
<small>(Security classification of title, body of abstract and indexing annotation must be entered when the overall report is classified)</small>		
1. ORIGINATING ACTIVITY (Corporate author) U. S. Naval Radiological Defense Laboratory San Francisco, California 94135		2a. REPORT SECURITY CLASSIFICATION UNCLASSIFIED 2b. GROUP
3. REPORT TITLE FISSION PRODUCT DECAY CHAINS: SCHEMATICS WITH BRANCHING FRACTIONS, HALF-LIVES, AND LITERATURE REFERENCES		
4. DESCRIPTIVE NOTES (Type of report and inclusive dates)		
5. AUTHOR(S) (First name, middle initial, last name) Glenn R. Crocker		
6. REPORT DATE 17 October 1967	7a. TOTAL NO. OF PAGES 41	7b. NO. OF REFS
8a. CONTRACT OR GRANT NO. AEC, Contract AT(49-7)-1963, Mod. 6	9a. ORIGINATOR'S REPORT NUMBER(S) USNRDL-TR-67-111	
b. PROJECT NO.		
c.	9b. OTHER REPORT NO(S) (Any other numbers that may be assigned this report)	
d.		
10. DISTRIBUTION STATEMENT This document has been approved for public release and sale; its distribution is unlimited.		
11. SUPPLEMENTARY NOTES		12. SPONSORING MILITARY ACTIVITY Atomic Energy Commission Washington, D. C. 20545
13. ABSTRACT Schematic diagrams of the decay chains for the fission products of uranium and plutonium are presented, with half-lives of the nuclides, branching fractions, and literature references. The updated data were obtained from a review of the literature through February 1967.		

DD FORM 1473 (PAGE 1)

S/N 0101-807-6801

UNCLASSIFIED
Security Classification

UNCLASSIFIED

Security Classification

14. KEY WORDS	LINK A		LINK B		LINK C	
	ROLE	WT	ROLE	WT	ROLE	WT
Fission-products Decay chains Radionuclide Half-life Beta decay						