

AD 669138

MOLECULAR MIXING LENGTHS FOR TURBULENT WAKES

George W. Sutton

AVCO EVERETT RESEARCH LABORATORY

RESEARCH REPORT 286
APRIL 1968

prepared for
ADVANCED RESEARCH PROJECTS AGENCY
DEPARTMENT OF DEFENSE
ARPA Order No. 1092

and
SPACE AND MISSILE SYSTEMS ORGANIZATION
AIR FORCE SYSTEMS COMMAND
DEPUTY FOR RE-ENTRY SYSTEMS (SMY)
Norton Air Force Base, California 92409

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED.

DDC
RECEIVED
MAY 17 1968
RECORDED
C

CLEARINGHOUSE

18

UNCLASSIFIED

AD 669 138

MOLECULAR MIXING LENGTHS FOR TURBULENT WAKES

George W. Sutton

Avco Everett Research Laboratory
Everett, Massachusetts

April 1968

Processed for . .

DEFENSE DOCUMENTATION CENTER
DEFENSE SUPPLY AGENCY



U. S. DEPARTMENT OF COMMERCE / NATIONAL BUREAU OF STANDARDS / INSTITUTE FOR APPLIED TECHNOLOGY

ABSTRACT

A simplified analysis is performed of the turbulent mixing of a trace chemical species with the bulk chemical species. The general relationship between the mixing on a molecular scale and the concentration fluctuation intensity of the trace species is given. This is then applied to a turbulent wake, to determine the axial length that it takes for injected gas to become mixed. The results are sensitive to the assumed turbulent dissipation rate; giving values of the mixing length from the order of to two orders of magnitude larger than the wake radius.

MOLECULAR MIXING LENGTHS FOR TURBULENT WAKES*

by

George W. Sutton

April 1968

AVCO EVERETT RESEARCH LABORATORY
a division of
AVCO CORPORATION
Everett, Massachusetts

*This research was supported by the Advanced Research Projects Agency of the Department of Defense and Space and Missile Systems Organization, Air Force Systems Command and was monitored by Space and Missile Systems Organization, Air Force Systems Command under Contract F04701-68-C-0036.

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED.

FOREWORD

Distribution of this document is unlimited. This indicates document has been cleared for public release by competent authority.

"This research was supported by the Advanced Research Projects Agency of the Department of Defense and Space and Missile Systems Organization, Air Force Systems Command and was monitored by Space and Missile Systems Organization, Air Force Systems Command under Contract F04701-68-C-0036." The secondary report number assigned by AERL is Avco Everett Research Report 286, and the author is G. W. Sutton. The Air Force program monitor for this contract is Major Walter D. McComb, Jr., SMYSE.

Publication of this report does not constitute Air Force approval of the report's findings or conclusions; it is published only for the exchange and stimulation of ideas.

Major Walter D. McComb, Jr.
SMYSE

LIST OF SYMBOLS

- k_e = energy-containing wave number
 m = 1 (planar wakes), 2 (axisymmetric wakes)
 $\overline{\frac{1}{2}q'^2}$ = turbulent kinetic energy
 R_n = drag radius, $\sqrt{C_D A/\pi}$
 r = radius
 S_c = Schmidt number
 t = time
 u = axial wake velocity
 x = axial distance behind body

 ϵ = rate of dissipation of turbulent kinetic energy
 ϕ = turbulent mixing parameter
 ξ = x/R_n
 $\Delta\xi$ = non-dimensional mixing length

Subscripts and superscripts

- = cross-sectional average
 A, B = species
 f = turbulent front
' = fluctuation
 l = point of entrainment

BLANK PAGE

Introduction

In performing calculations of chemical reactions in turbulent wakes, it is necessary to account for the rate at which the gas ingested by the turbulent front becomes mixed on a molecular scale with the gas on the interior of the turbulent wake. One proposed method uses a "mixing lag length" L , which is deduced on the basis of isotropic turbulence theory;¹ another method uses a transport equation for the electron density fluctuations to determine the mixing rate.² In this note, it is shown that the latter equation can be used to determine the mixing length.

Analysis

The model for determining this length is a one-dimensional wake of uniform properties of species A which ingests a small amount of species B at a distance x_1 behind the body. The flow field exterior of the

wake is also assumed to consist only of species A.

A mixing parameter ϕ is defined as follows:

$$\phi = \overline{C_A C_B} / \bar{C}_A \bar{C}_B \quad (1)$$

where C_A and C_B are the mass concentrations of species A and B respectively, and the overbar refers to a time average at a particular station in body-fixed coordinates. If the gas is unmixed, then either $C_A = 1$, $C_B = 0$, or $C_A = 0$, $C_B = 1$; so that $\overline{C_A C_B} = 0$; that is $\phi = 0$. If species B is intimately mixed with A, then $C_B(t)/C_A(t) = \text{constant} = \bar{C}_B/\bar{C}_A$, but since then $C_A(t) = \bar{C}_A$, $\phi = 1$. Thus ϕ is a measurement of mixing on a molecular scale. Also, since there are only two species in this model

$$C_A(t) + C_B(t) = \bar{C}_A + \bar{C}_B = 1 \quad (2)$$

If we decompose each concentration into a mean value and a fluctuating component ()', from eq. (2)

$$C_A' + C_B' = 0 \quad (3)$$

Combination of eqs. (1, 2, 3) yields:

$$\phi = 1 - \overline{C_B'^2} / \bar{C}_A \bar{C}_B \quad (4)$$

If B is a trace species, $\bar{C}_A \approx 1$, and eq. (4) can be rewritten as:

$$\phi(x) = 1 - \frac{\overline{C_B'^2}(x)}{\overline{C_B'^2}(x_1)} \cdot \frac{\bar{C}_B(x_1)}{\bar{C}_B(x)} \cdot \frac{\overline{C_B'^2}(x_1)}{\bar{C}_B(x_1)} \quad (5)$$

where x is the distance downstream and x_1 is the point of ingestion of species B which is initially unmixed at the point of ingestion; hence $\phi(x_1) = 0$ so that the last factor in eq. (5) is equal to unity, e.g.

$$\overline{C_B^{-1/2}}(x_1) = \overline{C_B}(x_1) \quad (6)$$

The second factor in eq. (5) is determined only by dilution. Thus the problem is reduced to that of determining the decay of mean square fluctuations of species B, for which Lin's equation (14)² can be used, e.g.:

$$(d/d\xi)(r_f^m \bar{u} \overline{C_B^{-1/2}}) - \overline{C_B}^2 u_f d(r_f^m)/d\xi = -R_n \epsilon^{1/3} k_e^{2/3} r_f^m \overline{C_B^{-1/2}} \quad (7)$$

where $m = 1$ for plane wakes and 2 for axially symmetric wakes; $r_f(x)$ is the mean distance from the axis to the wake turbulent front; $\xi = x/R_n$; R_n is the wake drag or momentum defect radius, \bar{u} is the mean wake speed, ϵ is the rate of dissipation of turbulent kinetic energy and k_e is the turbulent energy-containing wave number. The dissipation rate given in eq. (7) can be deduced from eqs. (7, 11, 20, and 23) of ref. 3, for a Schmidt S_c number of unity. This form is also applicable for $S_c < 1$, if multiplied by $\frac{1}{2}(3 - S_c^2)$.

From eq. (6), $\overline{C_B}$ is of the order of $\overline{C_B^{-1/2}}$. Since $\overline{C_B} \ll 1$, $\overline{C_B}^2 \ll \overline{C_B^{-1/2}}$ and the second term in eq. (7) can be neglected. In addition, one may approximate u_f as \bar{u} . Then eq. (7) can be integrated to:

$$\ln \left[r_f^m \overline{C_B^{-1/2}} / r_f^m(x_1) \overline{C_B^{-1/2}}(x_1) \right] = -R_n \int_{x_1}^x \bar{u}^{-1} \epsilon^{1/3} k_e^{2/3} d\xi \quad (8)$$

Now, from conservation of species B, $r_f^m \bar{C}_B \approx r_f^m(x_1) \bar{C}_B(x_1)$, and use of eq. (6) in eq. (8) yields:

$$\psi(x) = 1 - \exp\left(-R_n \int_{s_1}^{\xi} \bar{u}^{-1} \epsilon^{1/3} k_e^{2/3} d\xi\right) \quad (9)$$

We will next apply eq. (9) to three cases: (1) self-preserving wake; (2) the turbulence parameters calculated by Lin²; and (3) the turbulence parameters deduced by Schapker⁴ from turbulent front measurements.

Self Preserving Wake

The rate of destruction of turbulent kinetic energy for a wake can be estimated from the generalized form of eq. (8.4) of ref. 5 for an incompressible wake:

$$\int_0^{\infty} \epsilon dr^m = -\frac{1}{2} \frac{d}{dx} \int_0^{\infty} \bar{u} \left[(\Delta \bar{u})^2 + \overline{q'^2} \right] dr^m \quad (10)$$

where $\Delta \bar{u}$ is the mean velocity defect and $\frac{1}{2} \overline{q'^2}$ is the turbulent kinetic energy. For a self preserving wake, $\overline{q'^2} \sim (\Delta \bar{u})^2$. Also, from conservation of mass,

$$\Delta \bar{u} \sim U_{\infty} (R_n/r_f)^m \quad (10a)$$

where U_{∞} is the free stream velocity. Thus,

$$\epsilon(x) \sim U_{\infty}^3 (R_n/r_f)^{3m} x^{-1} \quad (11)$$

Since the self preserving wake is also geometrically similar, the energy-containing wave number is inversely proportional to the wake radius,

given by

$$(r_f/R_n) \sim \xi^{\frac{1}{m+1}} \quad (12)$$

and the mean wake velocity is taken as approximately U_∞ . Use of eqs. (11, 12) in (9) yields:

$$\phi = 1 - (\xi_1/\xi)^c \quad (13)$$

where c is the constant representing the numerical constants of proportionality in eqs. (11, 12) and between k_e and $1/r_f$. The important conclusion from (13) is that the distance required to achieve a given amount of mixing depends linearly on ξ_1 , the distance at which the species was initially ingested: that is, the further downstream the ingestion, the longer the length to become well-mixed. From eq. (13) the length $\Delta\xi = \xi - \xi_1$, to achieve a certain degree of mixing is

$$\Delta\xi/\xi_1 = (1 - \phi)^{-1/c} - 1 \quad (14)$$

From eq. (14), $\Delta\xi \rightarrow \infty$ as $\phi \rightarrow 1$ but for finite values of ϕ and $c^{-1} \ln(1 - \phi) \ll 1$

$$\Delta\xi \approx -c^{-1} \xi_1 \ln(1 - \phi) \quad (15)$$

In general, the integrand of eq. (9) can be represented as:

$$R_n \bar{u}^{-1} \epsilon^{1/3} k_e^{2/3} = c \xi^{-n} \quad (16)$$

and eq. (9) can be integrated as follows:

$$\phi = 1 - \exp \left[-c (n-1)^{-1} (\xi_1^{1-n} - \xi^{1-n}) \right] \quad (17)$$

For either c large or ϕ small, eq. (17) becomes:

$$\Delta \xi \approx -c^{-1} (\xi_1)^n \ln (1 - \phi) \quad (18)$$

which is the generalization of eq. (15). In order for the mixing length $\Delta \xi$ to be a constant multiple of the distance from the axis to the turbulent front from eq. (12) $n = (m+1)^{-1}$; e.g. $1/2$ for plane wakes and $1/3$ for axially symmetric wakes. This condition is almost fulfilled for Schapker's data⁴ for $\xi < 6 \times 10^3$, where $n = 0.43$, as shown below.

Equation (18) can also be written as

$$\Delta \xi = -\ln (1 - \phi) \left[R_n \bar{u}^{-1} \epsilon^{1/3} k_e^{2/3} \right]_{\xi_1}^{-1} \quad (19)$$

which could have been obtained directly from eq. (9) by taking the integrand constant at its value at ξ_1 . If one defines a characteristic decay time τ as

$$\tau \approx \epsilon^{-1/3} k_e^{-2/3} \quad (20)$$

and a non-dimensional decay length $\Lambda = U_\infty \tau / R_n$, then eq. (19) becomes

$$\Delta \xi = -\Lambda \ln (1 - \phi) \quad (21)$$

which yields the expected exponential decay of ϕ . The use of a definite mixing lag is obviously somewhat arbitrary, since $\Delta \xi$ depends on the criteria for ϕ . To compare various theories, an arbitrary value of ϕ equal to $1/2$ has been chosen.

Hypersonic Turbulence Parameters

Lin^2 assumed $k_e \approx r_n^{-1}$ and also calculated ϵ . A good match to those results for $\xi > 10^2$ is $c = 0.465$, $n = 2/3$, so that for $\phi = \frac{1}{2}$

$$\Delta \xi \approx 1.49 \xi_1^{2/3} \quad (22)$$

which is shown in Fig. 1 in comparison to the wake radius. It is seen that the mixing is between one and two orders of magnitude larger than the mean radius of the turbulent front, which is primarily a consequence of the low values of ϵ . Schapker's results⁴ can be represented as follows

$$\begin{aligned} R_N \bar{u}^{-1} \epsilon^{1/3} k_e^{2/3} &= 86.5 \beta^{5/6} (T_f/T_\infty)^{0.4} R_{ed\infty}^{-0.25} \xi^{-0.43}; \xi \leq 2000 \\ &= 6.75 \times 10^3 (T_f/T_\infty)^{0.4} \beta^{5/6} R_{ed\infty}^{-0.25} \xi^{-1}; \xi > 2000 \end{aligned} \quad (23)$$

where $\beta = u_f/U_\infty \approx 1$ and T is temperature. Neglecting the factor T_f/T_∞ , the mixing length for $\phi = \frac{1}{2}$ is then given by:

$$\begin{aligned} \Delta \xi &= 1.15 \times 10^{-2} (\ln 2) R_{ed\infty}^{0.25} \xi_1^{0.43} \quad \xi_1 \leq 2 \times 10^3 \\ &= 1.48 \times 10^{-4} (\ln 2) R_{ed\infty}^{0.25} \xi_1 \quad \xi_1 > 2 \times 10^3 \end{aligned} \quad (24)$$




Fig. 1 Present mixing length calculations based on Lin^2 and Schapker's⁴ turbulence parameters for $Re_{d\infty} = 10^6$, compared with the wake radius and previous estimates.¹

and is also shown in Fig. 1. These values result in a short mixing length, approximately equal to the wake radius within about a factor of 2. These short mixing lengths are caused primarily by the large values of both ϵ and k_e deduced by Schapker.⁴

Proudian and Feldman's analysis¹ results in:

$$\Lambda = \xi_1 \left[1 - \exp - (\rho/\rho_\infty)^{2/3} n^{-1} \right] \quad (25)$$

where ρ is mass density and $n^{1/2}$ is the exponent of the defect velocity decay. For a self similar wake $n = 4/3$, and for $\rho = \rho_\infty$, $\phi = \frac{1}{2}$,

$$\Delta\xi = 0.37 \xi_1 \quad (26)$$

also shown in Fig. 1. Here it is seen that this mixing length is much larger than any of the others, with a considerably larger slope. The values recommended in ref. 1 are also shown in Fig. 1, adjusted for $\phi = \frac{1}{2}$, which has an even larger average slope. (For $\xi > 600$, a constant mixing time of 2 msec is recommended.)

Discussion

The mixing length, is deduced from various theoretical calculations and measurements are obviously in wide disagreement by as much as two orders of magnitude. Unfortunately, direct measurements are not available, but calculations have been performed of the effect of various mixing lengths on electron densities in wakes.⁶ The calculations shown in Fig. 10 of ref. 6 indicate that the results are insensitive to the mixing length for $\xi < 2500$, but becomes very sensitive beyond that.

Furthermore, Figs. 3 and 10 of ref. 6 which compares the calculations with experiments⁷ indicate that the mixing length is quite short, e. g. $\Delta \xi / r_f \leq 10$ for $x/R_N < 4000$. These values are less than those predicted by Lin by only a factor of about 3 at $x/R_N = 4000$. The small values predicted by Schapker may be implausible. Additional experimental evidence would be desirable to resolve these differences.

References

1. A. P. Proudian, and S. Feldman, "A New Model for Mixing and Fluctuations in a Turbulent Wake", AIAA J. 3: 602-609 (1965).
2. S. C. Lin, "A Bimodal Approximation for Reacting Turbulent Flows: II. Example of Quasi One-Dimensional Wake Flow", AIAA J. 4: No. 2, 210-216, February 1966.
3. S. Corrsin, "The Isotropic Turbulent Mixer: Part II. Arbitrary Schmidt Number", A. I. Ch. E. Journal, 10: 870-877 (1964).
4. R. L. Schapker, "Statistics of High Speed Turbulent Wake Boundaries", AIAA J. 4: 1979-1987 (1966).
5. A. A. Townsend, "The Mechanism of Entrainment in Free Turbulent Flows", J. Fluid Mech., Vol. 26, Part 4, pp. 689-715 (1966).
6. E. A. Sutton, "The Chemistry of Electrons in Pure Air Hypersonic Wakes", Avco Everett Research Report, 266, July 1967 (submitted for publication to the AIAA Journal).
7. W. M. Kornegay, "Electron Density Decay in Wakes", AIAA J. 3: 1819-1823 (1965).

DOCUMENT CONTROL DATA - R&D

(Security classification of title, body of abstract and indexing annotation must be entered when the overall report is classified)

1 ORIGINATING ACTIVITY (Corporate author) Avco Everett Research Laboratory 2385 Revere Beach Parkway Everett, Massachusetts		2a REPORT SECURITY CLASSIFICATION Unclassified	
		2b GROUP	
3 REPORT TITLE MOLECULAR MIXING LENGTHS FOR TURBULENT WAKES			
4 DESCRIPTIVE NOTES (Type of report and inclusive dates) Research Report 286			
5 AUTHOR(S) (Last name, first name, initial) Sutton, George W.			
6 REPORT DATE April 1968	7a TOTAL NO OF PAGES 11	7b NO OF REFS 7	
8a CONTRACT OR GRANT NO. F04701-68-C-0036	9a ORIGINATOR'S REPORT NUMBER(S) Research Report 286		
b PROJECT NO	9b OTHER REPORT NO(S) (Any other numbers that may be assigned this report) SAMSO-TR-68-129		
c			
d			
10 AVAILABILITY/LIMITATION NOTICES Distribution of this document is unlimited. This indicates document has been cleared for public release by competent authority.			
11 SUPPLEMENTARY NOTES		12 SPONSORING MILITARY ACTIVITY Advanced Research Projects Agency, Department of Defense, ARPA Order #1092 and SAMSO, AFSC, Deputy for Re-entry Systems (SMY), Norton Air Force Base, Cal. 92409.	
13 ABSTRACT A simplified analysis is performed of the turbulent mixing of a trace chemical species with the bulk chemical species. The general relationship between the mixing on a molecular scale and the concentration fluctuation intensity of the trace species is given. This is then applied to a turbulent wake, to determine the axial length that it takes for ingested gas to become mixed. The results are sensitive to the assumed turbulent dissipation rate; giving values of the mixing length from the order of the wake radius to two orders of magnitude larger.			

4 KEY WORDS	LINK A		LINK B		LINK C	
	ROLE	WT	ROLE	WT	ROLE	WT
1. Turbulent mixing 2. Wakes 3. Re-entry Turbulence 4. Turbulent wakes 5. Turbulence 6. Mixing lengths 7. Re-entry physics						

INSTRUCTIONS

1. **ORIGINATING ACTIVITY:** Enter the name and address of the contractor, subcontractor, grantee, Department of Defense activity or other organization (*corporate author*) issuing the report.

2a. **REPORT SECURITY CLASSIFICATION:** Enter the overall security classification of the report. Indicate whether "Restricted Data" is included. Marking is to be in accordance with appropriate security regulations.

2b. **GROUP:** Automatic downgrading is specified in DoD Directive 5290.10 and Armed Forces Industrial Manual. Enter the group number. Also, when applicable, show that optional markings have been used for Group 3 and Group 4 as authorized.

3. **REPORT TITLE:** Enter the complete report title in all capital letters. Titles in all cases should be unclassified. If a meaningful title cannot be selected without classification, show title classification in all capitals in parentheses immediately following the title.

4. **DESCRIPTIVE NOTES:** If appropriate, enter the type of report, e.g., interim, progress, summary, annual, or final. Give the inclusive dates when a specific reporting period is covered.

5. **AUTHOR(S):** Enter the name(s) of author(s) as shown on or in the report. Enter last name, first name, middle initial. If military, show rank and branch of service. The name of the principal author is an absolute minimum requirement.

6. **REPORT DATE:** Enter the date of the report as day, month, year, or month, year. If more than one date appears on the report, use date of publication.

7a. **TOTAL NUMBER OF PAGES:** The total page count should follow normal pagination procedures, i.e., enter the number of pages containing information.

7b. **NUMBER OF REFERENCES:** Enter the total number of references cited in the report.

8a. **CONTRACT OR GRANT NUMBER:** If appropriate, enter the applicable number of the contract or grant under which the report was written.

8b, 8c, & 8d. **PROJECT NUMBER:** Enter the appropriate military department identification, such as project number, subproject number, system numbers, task number, etc.

9a. **ORIGINATOR'S REPORT NUMBER(S):** Enter the official report number by which the document will be identified and controlled by the originating activity. This number must be unique to this report.

9b. **OTHER REPORT NUMBER(S):** If the report has been assigned any other report numbers (*either by the originator or by the sponsor*), also enter this number(s).

10. **AVAILABILITY/LIMITATION NOTICES:** Enter any limitations on further dissemination of the report, other than those

imposed by security classification, using standard statements such as:

- (1) "Qualified requesters may obtain copies of this report from DDC."
- (2) "Foreign announcement and dissemination of this report by DDC is not authorized."
- (3) "U. S. Government agencies may obtain copies of this report directly from DDC. Other qualified DDC users shall request through _____."
- (4) "U. S. military agencies may obtain copies of this report directly from DDC. Other qualified users shall request through _____."
- (5) "All distribution of this report is controlled. Qualified DDC users shall request through _____."

If the report has been furnished to the Office of Technical Services, Department of Commerce, for sale to the public, indicate this fact and enter the price, if known.

11. **SUPPLEMENTARY NOTES:** Use for additional explanatory notes.

12. **SPONSORING MILITARY ACTIVITY:** Enter the name of the departmental project office or laboratory sponsoring (*paying for*) the research and development. Include address.

13. **ABSTRACT:** Enter an abstract giving a brief and factual summary of the document indicative of the report, even though it may also appear elsewhere in the body of the technical report. If additional space is required, a continuation sheet shall be attached.

It is highly desirable that the abstract of classified reports be unclassified. Each paragraph of the abstract shall end with an indication of the military security classification of the information in the paragraph, represented as (TS), (S), (C) or (U).

There is no limitation on the length of the abstract. However, the suggested length is from 150 to 225 words.

14. **KEY WORDS:** Key words are technically meaningful terms or short phrases that characterize a report and may be used as index entries for cataloging the report. Key words must be selected so that no security classification is required. Identifiers, such as equipment model designation, trade name, military project code name, geographic location, may be used as key words but will be followed by an indication of technical context. The assignment of links, rules, and weights is optional.