

AD 673 503


ENERGETICS OF MUSCLE ACTIVITY IN RELATION TO THE
MOLECULAR PHYSIOLOGY OF THE CONTRACTION PROCESS

Wilfried F. H. M. Mommaerts

California University
Los Angeles, California

15 December 1966

DISCLAIMER NOTICE



THIS DOCUMENT IS BEST QUALITY AVAILABLE. THE COPY FURNISHED TO DTIC CONTAINED A SIGNIFICANT NUMBER OF PAGES WHICH DO NOT REPRODUCE LEGIBLY.

OSR 68-1635 FINAL PROGRESS REPORT

Granting Agency: Air Force Office of Scientific Research
Period: October 2, 1964 - December 15, 1966
Title of Research: Energetics of Muscle Activity in Relation to the Molecular
Physiology of the Contraction Process
Prin. Investigator: Wilfried F. H. M. Mommaerts
Institution: University of California
405 Hilgard Avenue
Los Angeles, California 90024

AUG 27 1968

UNCLASSIFIED
A

The heat measuring techniques progressed to the stage where several significant investigations on the heat production during isometric contractions were carried out.

Activation heat, which is the heat produced in response to a stimulus when no tension or shortening is allowed, was determined by three different methods:

1. The heat is measured at successively shorter muscle lengths until the developed tension is very near zero. An extrapolation to zero tension gives the activation heat.
2. The tension is reduced to zero by uncoupling with deuterium oxide. The heat measured when no tension is developed is the activation heat.
3. The increment in heat is measured in response to a second stimulus given at a time when a new complete activation cycle is initiated out while the mechanical effects of the first are still present.

All three methods give essentially the same value of about 40% of the total heat in an isometric twitch which is in good agreement with earlier results of Hill and his colleagues. Equally significant is the fact that methods 2 and 3 have shown the activation heat to be independent of length. These results indicate that activation does not depend on the amount of overlap between the actin and myosin rods.

The first measurements of heat production of isolated papillary muscles have been accomplished due in large part to the development of very short, high sensitivity piles whose geometry have been adapted for this tissue.

This research was supported by the
Biological Sciences Division, AFOSR,
SRLA

under ~~Contract/Grant~~ AF-AFOSR-146-65

Reproduced by the
CLEARINGHOUSE
for Federal Scientific & Technical
Information Springfield Va. 22151

1. This document has been approved for public
release and sale; its distribution is unlimited.

AD 673503

Resting heat rate has been found to be ten times that of skeletal muscle. At 20°C. and at resting length its value is 24.8 mcal/g. muscle x min. The relation between heat production and actually developed tension is similar to that of skeletal muscle. An extra amount of heat is produced when the muscle is allowed to shorten.

Preliminary results indicate that the rate of heat production in stimulated muscle is much slower than in skeletal muscles. This point is under further investigation.

Resulting Publications:

Gibbs, C. L., Mommaerts, W. F. H. M. and Ricchiuti. Energetics of Cardiac Contractions. *J. Physiol.* 191: 25-46 (1967).

Gibbs, C. L., Ricchiuti, N. V. and Mommaerts, W. F. H. M. Activation Heat in Frog Sartorius Muscle. *J. Gen. Physiol.* 49:3, 517-535 (1966).

Gibbs, C. L., Ricchiuti, N. V. and Brady, A. J. Heat Production of Rabbit Papillary Muscle. *The Physiologist* 8:3 (1965).

Gibbs, C. L. and Mommaerts, W. F. H. M. The Heat Produced and the Work Performed by Rabbit Papillary Muscle. *Fed. Proc.*, 25:2 (1966).

Ricchiuti, N. V. and Gibbs, C. L. Heat Production in a Cardiac Contraction. *Nature* 208:5013, 897-898 (1965).

Ricchiuti, N. V. and Mommaerts, W. F. H. M. Technique for Myothermic Measurements. *The Physiologist* 8:3 (1965).

Ricchiuti, N., Gibbs, C. and Mommaerts, W. F. H. M. Two Methods for the Estimation of the Activation Heat in Muscle. *Fed. Proc.* 24:2 (1965).

Mommaerts, Wilfried F. H. M. The Rheology of Muscle. *Lab. Practice*. February 1966.

Mommaerts, W. F. H. M. and Wallner, A. The Break-Down of Adenosine Triphosphate in the Contraction Cycle of the Frog Sartorius Muscle. *J. Physiol.* 193: 343-357 (1967).

Seraydarian, K. and Mommaerts, W. F. H. M. Density Gradient Separation of Sarcotubular Vesicles and Other Particulate Constituents of Rabbit Muscle. *J. Cell Biol.* 26:2, 641-656 (1965).

Security Classification

DOCUMENT CONTROL DATA - R & D

Security classification of title, body of abstract and indexing annotation must be entered when the overall report is classified

1. ORIGINATING ACTIVITY (Corporate author) University of California Brain Research Institution Los Angeles, California 90024		2a. REPORT SECURITY CLASSIFICATION UNCLASSIFIED	
		2b. GROUP	
3. REPORT TITLE ENERGETICS OF MUSCLE ACTIVITY IN RELATION TO THE MOLECULAR PHYSIOLOGY OF THE CONTRACTION PROCESS			
4. DESCRIPTIVE NOTES (Type of report and inclusive dates) Scientific Final			
5. AUTHOR'S (First name, middle initial, last name) Wilfried F. H. M. Mommaerts			
6. REPORT DATE 15 December 1966	7a. TOTAL NO. OF PAGES 2	7b. NO. OF REFS 10	
8a. CONTRACT OR GRANT NO. AF-AFOSR-146-65		9a. ORIGINATOR'S REPORT NUMBER(S)	
b. PROJECT NO. 9777-01 6144501F		9b. OTHER REPORT NO(S) (Any other numbers that may be assigned this report) OSR 68-1635	
c. 681312			
10. DISTRIBUTION STATEMENT 1. This document has been approved for public release and sale; its distribution is unlimited.			
11. SUPPLEMENTARY NOTES T. CH, OTHER		12. SPONSORING MILITARY ACTIVITY Air Force Office of Scientific Research 1400 Wilson Boulevard (SRLA) Arlington, Virginia 22209	
13. ABSTRACT The heat measuring techniques progressed to the stage where several significant investigations on the heat production during isometric contractions were carried out. Activation heat, which is the heat produced in response to a stimulus when no tension or shortening is allowed, was determined by three different methods: (1) The heat is measured at successively shorter muscle lengths until the developed tension is very near zero. An extrapolation to zero tension gives the activation heat. (2) The tension is reduced to zero by uncoupling with deuterium oxide. The heat measured when no tension is developed is the activation heat. (3) The increment in heat is measured in response to a second stimulus given at a time when a new complete activation cycle is initiated but while the mechanical effects of the first are still present. The first measurements of heat production of isolated papillary muscles have been accomplished due in large part to the development of very short, high sensitivity piles whose geometry have been adapted for this tissue. Resting heat rate has been found to be ten times that of skeletal muscle. At 20°C. and at resting length its value is 24.8 mcal/g. muscle x min. The relation between heat production and actually developed tension is similar to that of skeletal muscle. An extra amount of heat is produced when the muscle is allowed to shorten. Preliminary results indicate that the rate of heat production in stimulated muscle is much slower than in skeletal muscles. This point is under further investigation.			

DD FORM 1473

Security Classification