


AD 67 3968




Carnegie-Mellon University

PITTSBURGH, PENNSYLVANIA 15213

GRADUATE SCHOOL OF INDUSTRIAL ADMINISTRATION

WILLIAM LARIMER MELLON, FOUNDER



Reproduced by the
CLEARINGHOUSE
for Federal Scientific & Technical
Information Springfield Va 22151

SEP 6 1968

This document is available
for public use and is not
distributed for profit

ON SOME NONSTANDARD SEMI-INFINITE
PROGRAMMING PROBLEMS

BY

A. Charnes*
W. W. Cooper**
K. O. Kortanek***

March, 1968

* Northwestern University
** Carnegie-Mellon University
*** Cornell University

This report was prepared as part of the activities of the Management Sciences Research Group, Carnegie-Mellon University, (Under Contract NONR 760(24), NR 047-048 with the U. S. Office of Naval Research and as part of the activities of the Systems Research Group, Northwestern University (under Contract NONR-1228 (10), Project NR 047-021 and the U. S. Army Research Office, Durham, Contract No. DA-31-124 ARO-D-322 and the National Science Foundation Grant NSF Project GP 7550 at Northwestern University, and the National Science Foundation Grant GK-1581 at Cornell University. Reproduction in whole or in part is permitted for any purpose of the United States Government.

Management Sciences Research Group
Graduate School of Industrial Administration
Carnegie-Mellon University
Pittsburgh, Pennsylvania

1. Introduction

Recently new structures and models of arithmetic developed by A. Robinson [13], [14] and others have provided a logical foundation for a nonarchimedean approach to the Calculus. These structures of nonstandard arithmetic offer "alternatives to familiar infinitary constructions and passages to the limit" (Robinson [14], p. 841). The basis for these methods is embedded in Foundations of Logic and permits consideration of a vast class of number fields which are richer in many ways than the more familiar extended number systems, such as Hilbert's Field of quotients of finite polynomials in one indeterminate.¹ It has long been known, for example, that finite linear programming theory (see [2], [4], and [5]) is valid over any ordered field. Charnes theorem (associating linear independence with extreme points) and the Charnes-Cooper Opposite Sign Theorem ([4]) are valid over any arbitrarily ordered field. Later Charnes-Cooper-Kortanek [8] proved that these two theorems are also valid for generalized finite sequence spaces over any arbitrarily ordered field in the context of semi-infinite programming. Historically, developments of extended number systems in linear programming arose in the context of resolving degeneracy (Charnes [2]) and Goal-programming (see [5], Chapter VI, Appendix B). For recent examples involving economic interpretations see Charnes-Clower-Kortanek [3]. The addition of relative infinitesimals and relative infinites to the real number

¹ Acknowledgement is due to Professor A. Nerode, Department of Mathematics, Cornell University for bringing these observations to our attention.

field avoids the necessity of considering familiar infinitary constructions in these particular contexts. While much evidence today is being compiled on the importance of extended number fields by outstanding mathematicians, the only textbook discussing linear programming over arbitrarily ordered fields is that of Charnes-Cooper, [5], in particular, pp. 756-757.

The main purpose of this paper is to develop and prove a duality theorem for a special nonstandard semi-infinite programming problem over Hilbert's Field. Attention will center on semi-infinite programs having real coefficients but whose complete regularizations (see [4], [5], and [8]) involve powers of the Nonarchimedean element. A duality result first proved in [12] but never published will be given which states conditions under which the infimum of the primal problem equals the maximum of the dual problem, possibly involving finite powers of the relative infinites. Research is currently under way into the construction of other Nonarchimedean fields in an attempt to obtain strong duality results which resolve, in yet another way, the phenomenon of duality gaps (see [9]).

We first review the basic properties of ordered fields, state the definition of Hilbert's Field, and discuss its topological properties.

2. Ordered Fields¹

A commutative field F is called "ordered" if the property of positiveness (> 0) is defined for all its elements, and if it satisfies the following postulates

1. For every a in F just one of the relations $a = 0$, $a > 0$, $-a > 0$ is valid.

¹ The results on ordered fields are presented here as in Van der Waerden[15].

2. If $a > 0$ and $b > 0$, then $a + b > 0$ and $ab > 0$.

If the absolute value $|a|$ of an element a in F is defined as the non-negative one of the elements, $a, -a$, then the following usual rules hold:

$$|ab| = |a| |b|$$

$$|a + b| \leq |a| + |b|$$

Furthermore, $a^2 = (-a)^2 = |a|^2 \geq 0$ with equality only for $a = 0$.

Thus a sum of squares is always positive. For completeness, other properties of inequalities in addition to those above are listed below:¹

(1) $0 < 1/a \iff a > 0$

(2) $a/b < c/d \iff abd^2 < b^2 cd$

(3) $0 < a < b \iff 0 < 1/b < 1/a$

(4) $a < b < 0 \iff 0 > 1/a > 1/b$

and (5) $a_1^2 + a_2^2 + \dots + a_n^2 \geq 0$, where all elements here are in F .

Again it is emphasized that these rules hold in any ordered field F .

Observe that by (5), the characteristic of any ordered field is zero.

If a commutative ring R is ordered, then the quotient field F of R is ordered in a unique way so that the ordering is preserved.

Further the ordering of F is given by

$$a/b > 0 \iff ab > 0$$
²

¹ See Birkhoff and MacLane [1] p. 50.

² See Birkhoff and MacLane [1], p. 49 Theorem 12.

The ordering of a field is called Archimedean¹ if there exists a "natural number" $n > a$ for every field element a . For example, the ordering of the rational number field is Archimedean. If the ordering of a field is not Archimedean, there exist "infinitely large" elements, larger than any rational number and "infinitely small" elements which are smaller than any positive rational number but larger than zero.

3. A Non-Archimedean (Order Preserving) Field Extension of the Real Numbers: The Hilbert Field

Let R be the field of real numbers with the usual ordering and let U be an indeterminate. Define an ordering of the polynomial ring $R[U]$ as follows

$$p \in R[U], p > 0$$

if and only if the coefficient of the highest^{nonvanishing} power of U is positive.

This defines an ordering in $R[U]$ since the laws of trichotomy, addition, and multiplication hold, and the ordering preserves the ordering of R . Further the quotient field $R[U]$ becomes ordered uniquely from $R[U]$ by:

$$(0) p/q > 0 \text{ if and only if } pq > 0,$$

preserving the ordering of $R[U]$ and hence that of R .²

¹ See Van der Waerden [15], p. 210.

² An equivalent and very intuitive definition of this ordering is given by $p(t) \geq q(t)$ if and only if $q(t)/p(t) < 1$ as $t \rightarrow \infty$, where t ranges through real values.

Theorem $R(U)$ under the ordering (0) is non-Archimedean.

Proof For any natural number n , we have $n - U < 0$ since the coefficient of the highest power of U is -1 . Hence $n < U$; \therefore it is impossible for the Archimedean property to hold.

U is an element of $R(U)$ which is to be regarded as infinitely large, that is, it is larger than any real number. Similarly observe that $1/U$ is infinitely small. Note also that real numbers can be regarded as polynomials in U of 0^{th} power. The field $R(U)$ with this ordering is usually called the Hilbert field and it is to this field which a major portion of our attention will be drawn. It is the largest order preserving extension of the real numbers which we consider in this paper.

Again, observe that as far as order relations among elements of $R(U)$ are concerned, they are the same as for the real numbers. Algorithms which use the order operations of real numbers go over into the big field, as is well known for the simplex algorithm.

4. The Order Topology of $R(U)$ and the Cauchy Completion

One of the basic properties of the real numbers that is used in semi-infinite programming is completeness, i.e. any Cauchy sequence is convergent. In fact every compact subset of Euclidean n -space is complete. Roughly speaking compactness requires certain limit points to be there and completeness assures that they are in the space itself. Thus we would like the Hilbert field to be complete in a topology that agrees with the subspace of reals.

Now it is known that every ordered field has an ordered extension field which is unique and complete. Of the various constructions of this extension field, perhaps the most well known is Cantor's construction by fundamental sequences.¹ Recall that an infinite sequence $\{a_n\}$ in an ordered field K is fundamental if given $\epsilon > 0$, and $\epsilon \in K$, there exists an integer n such that $|a_p - a_q| < \epsilon$ for $p > n$ and $q > n$. Observe that the topology induced on K by the " $| \quad |$ ", which is used in this construction, is the same as the order topology², i.e. if the topology for K has a subbase consisting of all sets of the form $\{x: x < a\}$ or $\{x: a < x\}$ for some $a \in K$.

It follows that the usual topology for the reals is identical with the order topology. Thus in this topology, which is actually a uniform topology, K is uniformly isomorphic to a dense subspace of its completion K^* .

Theorem of the Least Upper Bound:³ If K has an Archimedean ordering, then every non-empty subset of K^* bounded from above has a least upper bound in K^* .

In Kelly's terminology K^* is order-complete. As expected if $K = \mathbb{R}(U)$, then $\mathbb{R}^*(U)$ is also non-Archimedean and therefore not order complete. Thus there are sets which are closed and bounded but not compact e.g.,

¹ See Van Der Waerden [15] pp. 212-218.

² See Kelly [11] pp. 58-59.

³ See Van Der Waerden [15] pp. 216-217.

$$A = \{a \in R^*(U) : 0 \leq a \leq U\}.$$

To display the nature of the difficulties that can be encountered in nonstandard programming over $R^*(U)$, consider first the sequence $\{n\}$, $n = 1, 2, \dots$, in $R^*(U)$. This sequence is not a Cauchy sequence since the minimum distance between members is 1, and therefore no element in $R^*(U)$ is a limit point of this sequence. Consider the system of linear inequalities in one variable over $R^*(U)$,

$$u \geq n \quad u = 1, 2, \dots \quad (\text{integers})$$

Even over $R(U)$, this system is feasible, e.g. $u_0 = \alpha U$ with $\alpha > 0$ and real is feasible. The question here however, is whether there can be defined in some sense an extended concept of "canonical closure"¹, so that the duality theory applies as it does if u is required to be real. Two points need emphasis.

- (a) Clearly as it stands no inequality over $R^*(U)$ is a limiting inequality of the system since the sequence $\{n\}$, $n = 1, 2, \dots$ has no limit point in $R^*(U)$.
- (b) If we consider the equivalent inequality system (or any one which is bounded in the real sense),

$\frac{1}{n} u \geq 1$, $u = 1, 2, \dots$, then the limiting inequality $0 \cdot u \geq 1$ should be adjoined for canonical closure. However, with this addition the constraint set over $R^*(U)$ drastically changes, i.e. becomes empty, violating the important property that real systems possess as given previously by the following theorem.

¹ See [9] for the definition of canonical closure for semi-infinite programming problems.

Theorem. A canonically closed (real) system has the same constraint set as the original system. (See [8] p. 217).

Therefore we conclude that if we wish to maintain the above "constraint set preserving" property of the process of canonical closure, we cannot close up systems over $R^*(U)$ in the same manner as in the real case. In this situation the extended dual theorem, which provides 4 mutually exclusive and collectively exhaustive cases for the dual problems,¹ does not hold as evidenced by our example.

(I)	(II)
$\min u$	$\max \sum n \lambda_n$
$u \geq n, n = 1, 2, \dots$	$\sum \lambda_n = 1$
	$\lambda_n \geq 0 \quad (\text{all } n).$

(I) is consistent as well as (II), while $\inf u$

$$u \geq n \quad n = 1, 2, \dots$$

does not exist. This exactly displays the lack of order completeness of the extended field $R^*(U)$.

5. A Special Nonstandard Semi-Infinite Problem

We consider the following dual semi-infinite programming problems, where I is an arbitrary index set, $\{P_i : i \in I\} \subset R_m$, $P_0 \in R_m$, and $c_i \in R_1$ for all $i \in I$.²

¹ See [7], [8], and [9].

² See [8] and [9] where the Charnes-Cooper device of adding bounding constraints and an extra variable was first applied to arbitrary semi-infinite programs to render them bounded and consistent.

(I_R)

$$\min u_0 M + u^T P_0$$

$$\text{subject to } u_0 + u^T P_i \geq c_i \quad \text{all } i \in I$$

$$u^T I_m \geq -Ue_m^T$$

$$-u^T I_m \geq -Ue_m^T$$

$$u_0 \geq 0$$

(II_R)

$$\max \sum_{i \in I} c_i \lambda_i - Ue_m^T (v^+ + v^-)$$

$$\text{subject to } \lambda_0 + \sum_{i \in I} \lambda_i = M$$

$$\sum_{i \in I} P_i \lambda_i + I_m (v^+ - v^-) = P_0$$

$$\text{where } \lambda_0, \lambda_j \geq 0, i \in I,$$

$$\text{and } v_j^+, v_j^- \geq 0.$$

Since in the theory of semi-infinite programming, the (λ_i) -vectors have only finitely many non-zero entries, the summation " Σ " is well defined. Here I_m is the identity matrix of order m and e_m^T is the m -vector whose entries are all "ones." Clearly I_R is consistent, for take $u^T = 0$ and $u_0 \geq \sup \{c_i\}$, since $\{c_i : i \in I\}$ is a compact set. As for (II_R) , take $\lambda_i = 0$ ($i \in I$) and $v^+ - v^- = P_0$ with $\lambda_0 = M$. We shall assume that the set of coefficients $\{(P_i, c_i) : i \in I\}$ is compact in R_{m+1} and that if the system $u^T P_i \geq c_i, i \in I$, is consistent, then it has an interior point. Then by the Extended Dual

Theorem ([8] and [9]) it follows that there exist vectors λ^* , v^* such that $\inf \{u_0 M + u^T P_0\} = \sum_{i \in I} c_i \lambda_i^* - U \sum_{j=1}^m v_j^*$, where $v_j = v_j^+ - v_j^-$. Further, the relevant set¹ (u_0, u^T) for (I_R) is non-empty and compact since it is the intersection of closed sets (half-spaces) and by regularization is non-empty and bounded. Thus by compactness the "inf" is actually assumed for some u_0^* , u^{*T} .

We propose to study problems (I_R) and (II_R) by relaxing U and M through real values and observing the behavior of the solutions. Thus, we shall take a sequence of positive numbers, $U_n \rightarrow \infty$ as $n \rightarrow \infty$ and set $M_n = U_n^2$. We claim that in the limit it is possible to attain a non-Archimedean dual equation, where $U \in R(U)$, of a special type gotten from regularizing the minimization problem in powers of U . The non-Archimedean dual equation really represents a summary equation of the types (or orders) of divergence to infinity that may exist between unbounded variables. In order to provide for this variation, it is necessary to introduce powers of the indeterminate U .

Before we state and prove the theorem we turn to some notions of order as developed by Hardy [1].

Assume that f and g are positive and continuous. We say the order of f is greater than that of g if $f/g \rightarrow \infty$ when $x \rightarrow \infty$ and we write $f \succ g$.² If $f/g \rightarrow 0$, we say that the order of f is less

¹ Only bounded u_0 are relevant.

² See Hardy [10] P. 1-5.

than that of g and write $f \prec g$. If f/g remains bounded for all values of x after a certain point, then we say that the order of f equals the order of g and write $f \asymp g$. If f/g tends to a definite limit, we write $f \approx g$. If f and g are monotonic as well as f/g , then the following trichotomy holds:

$$(TR) \text{ either } f \succ g \text{ or } f \prec g \text{ or } f \asymp g.$$

Similarly we may introduce order notions for negative functions, which tend to $-\infty$, and we use the same symbols, e.g. $-\log x \prec -x$ having the same meaning as $\log x \prec x$.

With this introduction we now state our theorem.

Theorem Assume that the semi-infinite system (I_R) has a compact set of coefficients in R_{m+1} . Then there exist non-Archimedean dual optimal solutions to (I_R) and (II_R) respectively involving only powers of the indeterminate U and $P_i \in R_m$, such that

$$u_0^* M + u_*^T P_0 = \sum_{i \in I} c_i \lambda_i^* - \sum_{j=1}^m U^{\alpha(j)} v_j^*$$

where $\alpha: [1, 2, \dots, m] \rightarrow [0, 1, \dots, m]$, $M = U^{m+1}$, and the base field is $R(U)$.

We may speak of the semi-infinite program $\min u^T P_0$, subject to $u^T P_i \geq c_i$ all $i \in I$ as being embedded in a non-Archimedean extension. Each real semi-infinite program may in general have a different non-Archimedean embedding. From the viewpoint of the process of forming regularizations, the theorem states that it is possible to do non-Archimedean semi-infinite programming of a very special type, namely,

a "regularized" problem that originates from a compact semi-infinite problem with vectors having only real entries. The idea of the proof is to attain a finite non-Archimedean problem having only a finite number of half-spaces as constraints which are either limits of the original ones, (P_i, c_i) or involve powers of the indeterminate U as a bound. The limiting case will then be a finite non-Archimedean problem to which we can apply known finite duality theory.

Proof: We may simplify notation at the outset by omitting the variable u_0 and assume the problem to be consistent with interior point over the reals. This is because only bounded u_0 are relevant, and u_0 is introduced only to provide consistency. Let $Q(U)$ denote the cube $u^T I_m \geq -U e^T_m, -u^T I_m \geq -U e^T_m$, with the understanding that U always denotes the indeterminate. Let U_n be a real positive sequence $U_n \rightarrow \infty$ as $n \rightarrow \infty$.

Consider the following sequence of minimization problems as $U_n \rightarrow \infty$:

$$\begin{aligned} & \min u^T P_0 \\ & \text{subject to } u^T P_i \geq c_i \quad (i \in I) \\ & u \in Q(U_n) \end{aligned}$$

Optimal solutions exist by compactness or Haar duality. Let $u^{(n)}$ be an optimal solution and let T_n be a maximal linearly independent set of tangent planes from the collection $\{(P_i, c_i): i \in I\}$, i.e., $P_i \in T_n$ means $u^{(n)T} P_i = c_i$. Furthermore, at each stage, the associated finite problem,

$$\begin{aligned} & \min u^T P_0 \\ \text{subject to} & \quad u^T P_i \geq c_i \quad \text{all } P_i \in T_n \\ & \quad u \in Q(U_n) \end{aligned}$$

has $u^{(n)}$ as an optimal solution. Since the collection of P-planes is compact, there exists a limiting set $\{P_{\infty(1)}, \dots, P_{\infty(k)}\}$ of (maximal) linearly independent vectors of the sets T_n as $n \rightarrow \infty$.

Now as $n \rightarrow \infty$, $u^{(n)}$ will exhibit coordinates which are bounded and coordinates which are not. Let $S \equiv$ coordinate positions which are bounded, $W \equiv$ coordinate positions which are not bounded. The bounded coordinates will have limit points, say $u_i^{(\infty)}$ for all $i \in S$. Thus, at the n^{th} stage of the procedure (regularization with $u \in Q(U_n)$), the optimal solution will have the form,

$$u^{(n)} = \begin{cases} u_j^{(n)} : j \in S \\ f_j^{(n)} : j \in W \end{cases}$$

where $u_j^{(n)}$ and $f_j^{(n)}$ are, of course, real numbers. However, the $u_j^{(n)}$'s approach limits, while the $f_j^{(n)}$'s do not. Observe that feasibility is expressed as follows:

$$\sum_{j \in S} u_j^{(n)} a_j^{(i)} + \sum_{j \in W} f_j^{(n)} a_j^{(i)} \geq c_i, \quad i \in I \quad \text{where } P_j = (a_j^{(i)}).$$

We need some lemmas about the respective orders of the f_j 's.

Lemma 2: The trichotomy (TR) introduced above holds for the f_j 's.

That is, for $i \neq j$ in W one of the following hold:

$$f_i > f_j \quad \text{or} \quad f_i < f_j \quad \text{or} \quad f_i \sim f_j.$$

Proof: At some large n^{th} stage, the approximate order of greatest decrease for minimization will be established, i.e., if the dominance of $f_i^{(n)}$ over $f_j^{(n)}$ is observed, it cannot reverse itself at some later stage since the T_n -planes are approaching a limit set, and the finite coordinates (if there are any) of the optimal $u^{(n)}$ are also approaching limits. If at some large stage n , $f_i^{(n)} > f_j^{(n)}$, then at further stages the relative direction of $f_i^{(n)}$ will be sought, only possibly to be blocked by some finite plane in the T_n -set. However, at further stages these planes approach limits and their changes and effects on such limitation are negligible. This also shows the continuity of the functions f_j .

Thus, we may partition the index set W into subsets $W^{(h)} \subset W$ such that each subset corresponds to functions of the same order, i.e.

$$W^{(1)} \cup W^{(2)} \cup \dots \cup W^{(k)} = W$$

and functions in $W^{(i)}$ are of smaller order than those in $W^{(j)}$ if $i < j$.

For each order class choose a (positive) representative $\phi^{(h)}$, $h \in W^{(h)}$ in terms of which other members of the class differ by multiplicative constants in the large, say v_j . Thus, in terms of the partitioning of W , feasibility may be expressed as follows at the n^{th} stage:

$$(*) \quad \sum_{j \in S} u_j^{(n)} a_j^{(i)} + \sum_{j \in W^{(1)}} (1 + v_j) \phi^{(1)} a_j^{(i)} + \dots$$

$$+ \sum_{j \in W^{(k)}} (1 + v_j) \phi^{(k)} a_j^{(i)} \geq c_i \quad \text{for } i \in I.$$

These inequalities may only be approximate, in which case the error $\rightarrow 0$ as $n \rightarrow \infty$. This is not to say feasibility is attained with real values.

Lemma 3: In the above partitioning of W as expressed in the (*) inequalities, we may replace the $\phi^{(h)}$ representatives by $\pm U^h$, where $U \in R(U)$, and attain feasibility.

Proof: Observe that in each class the coefficients $\sum_{j \in W^{(h)}} (1 + v_j)$ are essentially constant and express the order relations between functions within the class. Two functions of the same order may differ by a multiplicative constant "at ∞ ." We now proceed by induction. The induction assumption is as follows. For $\leq k - 1$ order classes in W it is possible to assign appropriate powers to attain feasibility. For the case $k = 1$, feasibility is expressed at the n^{th} stage by

$$\sum_{j \in S} u_j^{(n)} a_j^{(i)} + f_n a_w^{(i)} \geq c_i,$$

where f_n is unbounded. Since $\{c_i\}$ is compact as well as $\{a_j^{(i)} : j \in S\}$ and $u_j^{(n)} \rightarrow u_j^{(\infty)}$, it follows that the choice of U or $-U$ (depending on which direction f_n moves) maintains the system of inequalities, even with the limit variables $u_j^{(\infty)}$ replacing $u_j^{(n)}$, $j \in S$. For the case $k = 2$, we are confronted with a situation as follows:

$$f_1 a_1^{(i)} + f_2 a_2^{(i)} \geq c_i$$

all $i \in I$, where $f_1 \succ f_2$ and bounded variables have been omitted. We must show, that $U^2 a_1^{(i)} + U a_2^{(i)} \geq c_i$, $i \in I$.¹ Suppose not, i.e., $U^2 a_1^{(i_0)} + U a_2^{(i_0)} < c_{i_0}$, for some i_0 . First, $a_1^{(i_0)} = a_2^{(i_0)}$ leads to a contradiction, $0 \geq c_{i_0}$ and $0 < c_{i_0}$. Therefore either $a_1^{(i_0)} < 0$

¹ Here we have assumed for simplicity that the coefficients have absorbed possible negative signs so that only directions at $(+\infty)$ are relevant.

or $a_1^{(i_0)} = 0$ and $a_2^{(i_0)} < 0$, which again is impossible since $f_1 > f_2$ and coefficients of the system are compact. Thus, $U^2 a_1^{(i)} + U a_2^{(i)} \geq c_i$, $i \in I$. Now assume that for $\leq k - 1$ unbounded variables the required assignment is possible, and consider the case for k order classes. Again, at the n^{th} stage, feasibility is expressed by,

$$\sum_{j \in S} u_j^{(n)} a_j^{(i)} + \sum_{j \in W^{(1)}} f_j^{(1)} a_j^{(i)} + \dots + \sum_{j \in W^{(k)}} f_j^{(k)} a_j^{(i)} \geq c_i, \quad i \in I.$$

In terms of the representative $\phi^{(k)}$ of $W^{(k)}$ we cannot have

$\sum_{j \in W^{(k)}} (1 + v_j) a_j^{(i)} < 0$ in the limit because this class is dominant and would destroy feasibility in such a situation. Thus, the net effect,

$\sum_{j \in W^{(k)}} (1 + v_j) a_j^{(i)}$ is positive, and if strictly positive, the assignment of U^k (or $-U^k$) will maintain feasibility. Further, if the net effect is zero, the induction assumption now applies to the remaining $k - 1$ variables. Therefore the indicated substitutions of U powers as expressed in the lemma are valid.

Thus, for all n beyond some fixed point, feasibility is attained with the U -power substitutions. In the limit we get an optimal solution with coordinates $u_i^{(\infty)}$, $i \in S$ and U -power coordinates for $i \in W$, which is determined by the T_∞ set and the U -power bounding hyperplanes. Taken together these hyperplanes form a finite non-Archimedean problem with the same minimum as the semi-infinite one due to our construction above. Hence, finite non-Archimedean duality theory now applies to

provide dual feasibility and the dual equation of the form required of the theorem. Observe that inconsistency is handled by bringing in the additional variable u_0 with functional value larger than any of the powers of U assigned to unbounded variables, e.g. with coefficients U^{m+1} . This completes the proof of the Theorem.

We conclude with three examples of problems whose variables have different rates of growth to $+\infty$.

Example 1

$$\begin{aligned} & \min u_0 + u_1 \\ & \text{subject to } u_0 \geq 0 \\ & \quad u_0 + 1/k u_1 + 1/k^2 u_2 \geq 0, \text{ for } k = 2, 3, \dots \end{aligned}$$

This system has interior points and the coefficient set is compact. Therefore it is canonically closed. Further, if we regularize with U_n (real) large, we find the solution to be

$$u_0^* = 1/2^2 U_n, u_1^* = -U_n, \text{ and } u_2^* = U_n.$$

These all have the same order, and therefore there is only one order class for unbounded coordinate positions. In fact, adjoining linear powers of U as constraints yields the optimum $u_0^* = 1/2^2 U$, $u_1^* = -U$, and $u_2^* = U$. Furthermore, the regularized dual reads:

$$\begin{aligned} & \max \quad -U [v_1^+ + v_2^+ + v_1^- + v_2^-] \\ & \text{subject to } \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix} \lambda_0 + \sum \begin{pmatrix} 1 \\ 1/k \\ 1/k^2 \end{pmatrix} \lambda_k + \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix} v_1^+ + \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix} v_2^+ + \begin{pmatrix} 0 \\ -1 \\ 0 \end{pmatrix} v_1^- + \begin{pmatrix} 0 \\ 0 \\ -1 \end{pmatrix} v_2^- = \begin{pmatrix} 1 \\ 1 \\ 0 \end{pmatrix} \\ & \quad \text{and } \lambda\text{'s, } v\text{'s } \geq 0. \end{aligned}$$

The dual optimum reads: $\lambda_2^* = 1$, $v_1^{+*} = 1/2$, $v_2^{-*} = 1/2^2$ with dual functional, $-U (1/2 + 1/2^2) = -3/4 U$. Since the direct functional is $u_0^* + u_1^* = -3/4 U$, we attain the dual equation.

Example 2. min $-u_2$

$$u_1 \frac{1}{x} - u_2 \geq 1 - \log x, x \geq 2$$

$$u_1 \geq 1$$

$$0 u_1 - 0 u_2 \geq -1$$

$$-u_1 \geq -U^2$$

$$-u_2 \geq -U$$




Figure 3

Regularizing with U_n yields an optimum

$$u_1^* = U_n \text{ and } u_2^* = \log U_n.$$

In this case there are two order classes, and variable u_2 is regularized with respect to U and u_1 with respect to U^2 . Thus, the non-Archimedean optimum reads $u_1^* = U^2$, $u_2^* = U$. The dual problem has an optimum at $v_2^{-*} = 1$ and all other variables zero, with equation of dual functionals prevailing, i.e.

$$\max_x \sum (1 - \log x) \lambda_x + \lambda_1 - \lambda_2 - U^2 v_1^- - U v_2^-$$

$$\text{s.t. } \sum_x \begin{pmatrix} 1/x \\ -1 \end{pmatrix} \lambda_x + \begin{pmatrix} 1 \\ 0 \end{pmatrix} \lambda_1 + \begin{pmatrix} -1 \\ 0 \end{pmatrix} v_1^- + \begin{pmatrix} 0 \\ -1 \end{pmatrix} v_2^- = \begin{pmatrix} 0 \\ -1 \end{pmatrix}$$

$$\lambda_x, \lambda_1, \lambda_2, v_1^-, v_2^- \geq 0.$$

Example 3¹ Consider as the constraint set the points under the curve $y = \tan^{-1}(x)$, with $x \geq 0$ and above the x-axis.




Figure 1

We observe that the equation of the tangent line at the point $(x, \tan^{-1} x)$ is given by $(u_2 - \tan^{-1} x)/(u_1 - x) = 1/(1 + x^2)$ and therefore our constraint set is given by the following system of inequalities

(I)

$$u_1 (1 + x^2)^{-1} - u_2 \geq -\tan^{-1} x + x (1 + x^2)^{-1} = c_x$$

$$u_1 \geq 0$$

$$-u_2 \geq -(\pi/2)$$

Let the direct problem, I, be $\min(-u_2)$ subject to the above constraints.

In this example the set of coefficient points of

$$\{(1 + x^2)^{-1}, -1, -\tan^{-1} x + x (1 + x^2)^{-1} : x \geq 0\}$$

in addition to the points $(1, 0, 0)$ and $(0, -1, -\pi/2)$. Clearly this set is compact, since the limit point $(0, -1, -\pi/2)$ as $x \rightarrow \infty$ is added. Note that $\min -u_2 = -\max u_2 = -\pi/2$ is never attained because feasible points must lie on or under the curve.

¹ See [8] pp. 214-215.

For this example, the dual II is as follows.

If we let

$$P_x = \begin{pmatrix} (1+x^2)^{-1} \\ 1 \end{pmatrix}, P_\alpha = \begin{pmatrix} 1 \\ 0 \end{pmatrix}, P_\beta = \begin{pmatrix} 0 \\ -1 \end{pmatrix}$$

then we have

$$\max \sum c_x \lambda_x + 0 \cdot \lambda_\alpha + (-\pi/2) \lambda_\beta$$

subject to

$$\sum P_x \lambda_x + P_\alpha \lambda_\alpha + P_\beta \lambda_\beta = P_0 = \begin{pmatrix} 0 \\ -1 \end{pmatrix}, \quad \text{all } \lambda\text{'s} \geq 0.$$

We see that if $\lambda_\beta = 1$ and all other $\lambda\text{'s} = 0$, then the maximum $-\pi/2$ is attained. Observe, however, that if we adjoin the constraint $u_2 \geq \pi/2$, then problem (I) becomes inconsistent over the real field. The dual solution, however, prevails. Over the Hilbert field $R(U)$, as previously introduced, problem (I) is consistent, e.g., $u_1 = U$ and $u_2 = \pi/2$. We view this type of inconsistency or duality gap over the real field as one which may be removed by considering the semi-infinite problem in a nonstandard way, namely programming over $R(U)$.

References

- [1] Birkhoff, G., and S. MacLane, A Survey of Modern Algebra, The Macmillan Company, New York, 1953.
- [2] Charnes, A., "Optimality and Degeneracy in Linear Programming," Econometrica, Vol. 20, No. 2, pp. 160-170.
- [3] Charnes, A., R. W. Clower, and K. Kortanek, "Effective Control Through Coherent Decentralization with Preemptive Goals," forthcoming Econometrica.
- [4] Charnes, A. and W. W. Cooper, "The Strong Minkowski-Farkas-Weyl Theorem for Vector Spaces over Ordered Fields," Proc. Nat. Acad. Sci. U.S.A., 44, No. 9, Sept. 1958, pp. 914-916.
- [5] Charnes, A. and W. W. Cooper, Management Models and Industrial Applications of Linear Programming, Vols. I and II, New York, J. Wiley and Sons, 1961.
- [6] Charnes, A., W. W. Cooper and K. Kortanek, "Duality, Haar Programs, and Finite Sequence Spaces," Proc. Nat. Acad. Sci. USA, 48, No. 5, May 1962, pp. 783-786.
- [7] Charnes, A., W. W. Cooper and K. Kortanek, "A Duality Theory for Convex Programs with Convex Constraints," Bull. of the Amer. Math. Soc., Vol. 68, No. 6, Nov. 1962, pp. 605-608.
- [8] Charnes, A., W. W. Cooper and K. Kortanek, "Duality in Semi-infinite Programs and Some Works of Haar and Caratheodory," Management Science, 9, No. 2, Jan. 1963, pp. 209-228.
- [9] Charnes, A., W. W. Cooper and K. Kortanek, "On Representations of Semi-Infinite Programs Which Have No Duality Gaps," Management Science, Vol. 12, No. 1, Sept. 1965.
- [10] Hardy, G. H., Orders of Infinity, Cambridge Tracts in Mathematics and Mathematical Physics, No. 12, Cambridge University Press, 1954.
- [11] Kelley, J. L., General Topology, D. Van Nostrand Co., Inc. 1955.
- [12] Kortanek, K. O., "Duality, Semi-Infinite Programming, and Some Aspects of Control in Business and Economic Systems, Ph.D. Thesis, Northwestern University, Department of Engineering Sciences, Evanston, June, 1964.
- [13] Robinson, A., Non-Standard Analysis, Studies in Logic and the Foundations of Mathematics, North-Holland, Amsterdam, 1966.
- [14] Robinson, A., "Non-standard Arithmetic", Bulletin of the American Mathematical Society, Vol. 73, No. 6, Nov. 1967.
- [15] Van der Waerden, B. L., Modern Algebra, Vol. I, Frederick Ungar Publishing Co., New York, 1953.

- [16] Springer, T.A. (ed.) Proceedings of a Conference on Local Fields
a. Monna, A.F. Linear Topological Spaces over Non-Archimedean
Valued Fields
b. Remmert, R. Algebraische Aspekte in der nichtarchimedischen
Analysis.
Springer, New York, 1967.

Unclassified

Security Classification

DOCUMENT CONTROL DATA - R&D

(Security classification of title, body of abstract and indexing annotation must be entered when the overall report is classified)

1. ORIGINATING ACTIVITY (Corporate author) Graduate School of Industrial Administration Carnegie Mellon University		2a. REPORT SECURITY CLASSIFICATION Unclassified	
		2b. GROUP Not applicable	
3. REPORT TITLE ON SOME NONSTANDARD SEMI-INFINITE PROGRAMMING PROBLEMS			
4. DESCRIPTIVE NOTES (Type of report and inclusive dates) Technical Report, March, 1968			
5. AUTHOR(S) (Last name, first name, initial) A. Charnes, W. W. Cooper, K. O. Kortanek			
6. REPORT DATE March, 1968		7a. TOTAL NO. OF PAGES 23	7b. NO. OF REFS 16
8a. CONTRACT OR GRANT NO. NONR 760(24)		8a. ORIGINATOR'S REPORT NUMBER(S) Management Sciences Research Report No. 139	
8b. PROJECT NO. NR 047-048		8b. OTHER REPORT NO(S) (Any other numbers that may be assigned to the report) Systems Research No. SRM 201	
10. AVAILABILITY/LIMITATION NOTICES Distribution of this document is unlimited			
11. SUPPLEMENTARY NOTES Also Contract NONR-1228(10), Project NR 047-021; U.S. Army Research Office, Durham, Contract No. DA-31-124 AROD 322		12. SPONSORING MILITARY ACTIVITY Logistics and Mathematical Statistics Br. Office of Naval Research Washington, D. C. 20360	
13. ABSTRACT NSF Project GP 7550, Northwestern and NSF GK-1581, Cornell University. Aspects of the duality theory for semi-infinite programming are here extended to fields with properties of non-Archimedean order. Emphasis is on nonstandard semi-infinite programming problems in Hilbert's Field. The ideas of regularization are generalized to include powers of the relative infinites in terms of the indeterminates.			

14. KEY WORDS	LINK A		LINK B		LINK C	
	ROLE	WT	ROLE	WT	ROLE	WT
Linear Programming						
Regularization						
Semi-Infinite Programming						
Duality						

INSTRUCTIONS

1. **ORIGINATING ACTIVITY:** Enter the name and address of the contractor, subcontractor, grantee, Department of Defense activity or other organization (*corporate author*) issuing the report.
- 2a. **REPORT SECURITY CLASSIFICATION:** Enter the overall security classification of the report. Indicate whether "Restricted Data" is included. Marking is to be in accordance with appropriate security regulations.
- 2b. **GROUP:** Automatic downgrading is specified in DoD Directive 5200.10 and Armed Forces Industrial Manual. Enter the group number. Also, when applicable, show that optional markings have been used for Group 3 and Group 4 as authorized.
3. **REPORT TITLE:** Enter the complete report title in all capital letters. Titles in all cases should be unclassified. If a meaningful title cannot be selected without classification, show title classification in all capitals in parenthesis immediately following the title.
4. **DESCRIPTIVE NOTES:** If appropriate, enter the type of report, e.g., interim, progress, summary, annual, or final. Give the inclusive dates when a specific reporting period is covered.
5. **AUTHOR(S):** Enter the name(s) of author(s) as shown on or in the report. Enter last name, first name, middle initial. If military, show rank and branch of service. The name of the principal author is an absolute minimum requirement.
6. **REPORT DATE:** Enter the date of the report as day, month, year; or month, year. If more than one date appears on the report, use date of publication.
- 7a. **TOTAL NUMBER OF PAGES:** The total page count should follow normal pagination procedures, i.e., enter the number of pages containing information.
- 7b. **NUMBER OF REFERENCES:** Enter the total number of references cited in the report.
- 8a. **CONTRACT OR GRANT NUMBER:** If appropriate, enter the applicable number of the contract or grant under which the report was written.
- 8b, 8c, & 8d. **PROJECT NUMBER:** Enter the appropriate military department identification, such as project number, subproject number, system numbers, task number, etc.
- 9a. **ORIGINATOR'S REPORT NUMBER(S):** Enter the official report number by which the document will be identified and controlled by the originating activity. This number must be unique to this report.
- 9b. **OTHER REPORT NUMBER(S):** If the report has been assigned any other report numbers (*either by the originator or by the sponsor*), also enter this number(s).
10. **AVAILABILITY/LIMITATION NOTICES:** Enter any limitations on further dissemination of the report, other than those

imposed by security classification, using standard statements such as:

- (1) "Qualified requesters may obtain copies of this report from DDC."
- (2) "Foreign announcement and dissemination of this report by DDC is not authorized."
- (3) "U. S. Government agencies may obtain copies of this report directly from DDC. Other qualified DDC users shall request through _____."
- (4) "U. S. military agencies may obtain copies of this report directly from DDC. Other qualified users shall request through _____."
- (5) "All distribution of this report is controlled. Qualified DDC users shall request through _____."

If the report has been furnished to the Office of Technical Services, Department of Commerce, for sale to the public, indicate this fact and enter the price, if known.

11. **SUPPLEMENTARY NOTES:** Use for additional explanatory notes.
12. **SPONSORING MILITARY ACTIVITY:** Enter the name of the departmental project office or laboratory sponsoring (*paying for*) the research and development. Include address.
13. **ABSTRACT:** Enter an abstract giving a brief and factual summary of the document indicative of the report, even though it may also appear elsewhere in the body of the technical report. If additional space is required, a continuation sheet shall be attached.

It is highly desirable that the abstract of classified reports be unclassified. Each paragraph of the abstract shall end with an indication of the military security classification of the information in the paragraph, represented as (TS), (S), (C), or (U).

There is no limitation on the length of the abstract. However, the suggested length is from 150 to 225 words.

14. **KEY WORDS:** Key words are technically meaningful terms or short phrases that characterize a report and may be used as index entries for cataloging the report. Key words must be selected so that no security classification is required. Identifiers, such as equipment model designation, trade name, military project code name, geographic location, may be used as key words but will be followed by an indication of technical context. The assignment of links, roles, and weights is optional.