

NEW CONDITIONS FOR CENTRAL LIMIT THEOREMS

Percy A. Pierre*

1. Introduction

A general formulation of the central limit problem for sums of independent random variables is the following (see Loève [1], p. 291).

Let

$$S_n = \sum_k X_{nk}$$

where $k=1, \dots, k_n$, $k_n \rightarrow \infty$ as $n \rightarrow \infty$, and for each n $\{X_{nk}\}$ are independent random variables with probability distribution functions F_{nk} and $EX_{nk} = 0$. Let $\{F_n\}$ be the distribution functions of $\{S_n\}$ and let $\phi(x)$ be the distribution function of a Normal random variable with zero-mean and variance σ^2 . Under these conditions it is possible to show the following.

Theorem 1.1: Let $\max_k \text{Var } X_{nk} \rightarrow 0$ and $\sum_k \text{Var } X_{nk} \rightarrow \sigma^2 < \infty$ where σ^2 is a positive constant. The sums S_n are asymptotically Normal (i.e., $F_n(x) \rightarrow \phi(x)$) if and only if for every $\epsilon > 0$

$$(1.1) \quad g_n(\epsilon) = \sum_k \int_{|x| \geq \epsilon} x^2 dF_{nk} \rightarrow 0.$$

Except in special cases, the application of condition 1.1 is difficult because of the integrals involved. By assuming the existence of

*This research was done at the University of Michigan while the author was on a Post-Doctoral Research Fellowship in the Program of Information and Control Engineering.

Any views expressed in this paper are those of the author. They should not be interpreted as reflecting the views of The Rand Corporation or the official opinion or policy of any of its governmental or private research sponsors. Papers are reproduced by The Rand Corporation as a courtesy to members of its staff.

fourth-order moments, we are able to prove new necessary and sufficient conditions for both Normal and Poisson convergence which involve only moments. The proof of the theorem makes use of a characterization of the Normal distribution among infinitely divisible (I.D.) laws which was perhaps first recognized by Borges [3] and later independently by the author [4].

2. Normal Convergence

Theorem 2.1: Let $E|S_n|^{(4+\delta)}$ be uniformly bounded for some $\delta > 0$. Let $\max_k \text{Var } X_{nk} \rightarrow 0$, $\sum_k \text{Var } X_{nk} \rightarrow \sigma^2 < \infty$ where σ^2 is a positive constant. Then S_n is asymptotically Normal if and only if

$$(2.1) \quad ES_n^4 - 3\{ES_n^2\}^2 \rightarrow 0.$$

Proof: The asymptotic normality of S_n implies condition 2.1 by the moment convergence theorem (see Loeve [1], p. 184) and the fact that for a zero-mean Normal random variable S , $ES^4 - 3\{ES^2\}^2 = 0$.

To prove the converse it is sufficient to show that every convergent subsequence $\{F_{n'}\}$ of $\{F_n\}$ converges to $\phi(x)$ (see Feller [2], p. 261).

Let F be the limit of $F_{n'}$, then F is an infinitely divisible law with characteristic function $f(u)$ such that (see Loève [1], p. 293),

$$(2.2) \quad \log f(u) = \int (e^{iux} - 1 - iux) \frac{1}{x^2} dK(x)$$

where $K(x)$ is monotone increasing and of bounded variation, $K(-\infty) = 0$, and $K(\infty) = \sigma^2 < \infty$. The integrand is defined by continuity

at the origin. It is known that $F(x)$ is a Normal law if and only if $K(x)$ increases only at $x = 0$.

Since $E|S_n|^{(4+\delta)}$ are uniformly bounded, all moments of S_n , of order 4 or less converge to those of F . Since $ES_n^4 - 3\{ES_n^2\}^2 = 0$,

$$\begin{aligned}
 0 &= \int x^4 dF(x) - 3 \int x^2 dF(x)^2 \\
 &= \frac{\partial^4}{\partial u^4} \log f(u) \Big|_{u=0} \\
 (2.3) \quad &= \int x^2 dK(x).
 \end{aligned}$$

This last equation is obtained by differentiating the RHS of Eq. (2.2) under the integral sign. This is justified in the following way. That

$$(2.4) \quad - \frac{d^2}{du^2} \log f(u) = \int e^{iux} dK(x)$$

is shown by Loeve [1], p. 293. Thus the L.H.S. of Eq. (2.4) is a characteristic function. Since this characteristic function is twice differentiable, its second derivative is given by (Loève [1], p. 200)

$$\frac{d^4}{du^4} \log f(u) = \int x^2 e^{iux} dK(x)$$

and Eq. (2.3) follows. Thus $K(x)$ increases only at $x = 0$.

Finally, since $\sum_k \text{Var } X_{nk} = \sigma^2$ we have shown that $F_n \rightarrow \phi(x)$ and the proof is complete.

Condition 2.1 becomes even simpler when we note that

$$ES_n^4 - 3\{ES_n^2\}^2 = \sum_k \left[EX_{nk}^4 - 3\{EX_{nk}^2\}^2 \right].$$

The L.H.S. is usually called the fourth cumulant of S_n . This identity says that the fourth cumulant of a sum of independent random variables equals the sum of the fourth cumulants.

If the distributions F_n are known to be infinitely divisible (I.D.), then moments higher than 4 are not required.

Theorem 2.2: If F_n are I.D., then $F_n(x) \rightarrow \phi(x)$ if and only if $ES_n^2 \rightarrow \sigma^2$ and

$$ES_n^4 - 3\{ES_n^2\}^2 \rightarrow 0.$$

Proof: The characteristic functions $f_n(u)$ are given by

$$\log f_n(u) = \int (e^{iux} - 1 - iux) \frac{1}{x^2} dK_n(x).$$

For any $\epsilon > 0$

$$\int_{|x| > \epsilon} dK_n(x) \leq \int \frac{x^2}{\epsilon^2} dK_n(x) \rightarrow 0$$

as $n \rightarrow \infty$. Thus $K_n(x)$ converges to a step function at the origin of size σ^2 .

The converse is obtained from the moment convergence theorem.

3. Poisson Convergence

The key feature of the results above is that the fourth cumulant (i.e., $ES_n^4 - 3\{ES_n^2\}^2$) corresponds to $\int x^2 dK(x)$ and is a good test for a jump of $K(x)$ at the origin. However, this method can be used to test for jumps at other points also. For example, the equation

$$\int (x-1)^2 dK(x) = 0$$

implies that $K(x)$ can only have a jump at $x = 1$. This leads to the following results. First we note that

$$\begin{aligned} \int (x-1)^2 dK_n(x) &= ES_n^4 - 3\{ES_n^2\}^2 - 2ES_n^3 + ES_n^2 \\ &= \sum_k \left[EX_{nk}^4 - 3\{EX_{nk}^2\}^2 - 2EX_{nk}^3 + EX_{nk}^2 \right]. \end{aligned}$$

Theorem 3.1: Let $E|S_n|^{(4+\delta)}$ be uniformly bounded for some $\delta > 0$. Then S_n is asymptotically Poisson (i.e., $\log f_n(u) \rightarrow [\sigma^2(e^{iu}-1) - iu\sigma^2]$) if and only if

$$ES_n^4 - 3\{ES_n^2\}^2 - 2ES_n^3 + ES_n^2 \rightarrow 0.$$

In a similar way we obtain:

Theorem 3.2: If F_n are I.D., then F_n is asymptotically Poisson if and only if $ES_n^2 \rightarrow \sigma^2$ and

$$ES_n^4 - 3\{ES_n^2\}^2 - 2ES_n^3 + ES_n^2 \rightarrow 0.$$

REFERENCES

1. M. Loève, Probability Theory, Van Norstrand, 3rd Ed., 1963.
2. W. Feller, An Introduction to Probability Theory and Its Applications, Vol. II, John Wiley and Sons, New York, 1966.
3. R. Borges, "A Characterization of the Normal Distribution (A note on a paper by Kozin)," Z. Wahrscheinlichkeitstheorie, Vol. 5, No. 3, 1966.
4. P. Pierre, "Properties of Non-Gaussian, Continuous Parameter, Random Processes as Used in Detection Theory," Dissertation, The Johns Hopkins University, Baltimore, Md.