

ESD RECORD COPY

RETURN TO
SCIENTIFIC & TECHNICAL INFORMATION DIVISION
(ESTI), BUILDING 1211

ESD ACCESSION LIST

ESTI Call No. 62845

Copy No. 1 of 1 cys.

Prepared under Electronic Systems Division

Contract AF 19(628)-5167 by

Lincoln Laboratory

MASSACHUSETTS INSTITUTE OF TECHNOLOGY

LINCOLN LABORATORY LIBRARY

33rd REFERENCE BIBLIOGRAPHY

Properties of the Compounds
 RbMnF_3 , KMnF_3 and CsMnF_3

Joan S. Friebely

William J. Ince

8 August 1968



AD677285

LINCOLN LABORATORY LIBRARY

PROPERTIES OF THE COMPOUNDS

RbMnF_3 , KMnF_3 and CsMnF_3

JOAN S. FRIEBELY

WILLIAM J. INCE

33rd REFERENCE BIBLIOGRAPHY

8 AUGUST 1968

Lincoln Laboratory is a center for research operated by Massachusetts Institute of Technology, with the support of the U.S. Air Force under Contract AF 19(628)-5167.

This document has been approved for public release and sale; its distribution is unlimited.

MASSACHUSETTS INSTITUTE OF TECHNOLOGY

LINCOLN LABORATORY

LEXINGTON, MASSACHUSETTS

INTRODUCTION

This bibliography lists publications related to the double fluorides XMnF_3 , where X is K, Rb or Cs. These materials are antiferromagnetic with Néel temperatures of 88.3°K, 82.6°K and 53.5°K, respectively. Considerable interest has been shown in their properties since about 1960. They are characterized by having large exchange ($H_{\text{ex}} \approx 10^6$ Oe) and small crystalline anisotropy ($H_A < 10$ Oe). Antiferromagnetic resonance (AFMR) can be observed, therefore, at microwave frequencies without the need for very large magnetizing fields. Further, there is a strong hyperfine interaction that causes pronounced coupling between the nuclear magnetic resonance (NMR) of the Mn^{55} nuclei and the AFMR modes. Hence much of the published work to date has dealt with magnetic resonance.

References are listed alphabetically by author. The consecutive numbers beside each citation provide unique identification for the articles as they appear in the subject outline and index.

The sources consulted for the literature search include the abstracting services listed here, lists of references in related published papers, and the scientific journals currently being received by the Lincoln Laboratory Library. We also extend our appreciation to NASA and DDC for their request bibliography services, which were helpful to the comprehensiveness of the survey.

Accepted for the Air Force
Franklin C. Hudson
Chief, Lincoln Laboratory Office

SOURCES CONSULTED

Chemical Abstracts	1957 - July 1968
Dissertation Abstracts	1963 - 1967
Electronics and Communications Abstracts	1961 - August 1966
Physics Abstracts/Science	1957 - July 1968
Solid State Abstracts	1960 - 1965

SUBJECT OUTLINE AND INDEX

Antiferromagnetic Domains

Cole et al. (14); Farztdinov (29); Ince (60); Waring (133).

Antiferromagnetic Resonance (AFMR)

Cole (12); Cole et al. (13, 14); Eastman (22); Eastman et al. (23, 24, 25); Freiser et al. (36, 37); Heeger (44); Heeger et al. (46); Hinderks et al. (51); Ince (60); Kedzie et al. (65, 66); Lee et al. (68); Minkiewicz et al. (80); Ninio et al. (88); Seiden et al. (109); Teaney (124); Teaney et al. (125, 126, 127); Turov et al. (129).

Basic Theory

Blume et al. (8); Bullock (10); DeGennes (19); Farztdinov (29); Freeman et al. (35); Gondaira (39); Gondaira et al. (40); Gorodetsky et al. (41); Hubbard et al. (58); Joseph (64); Lvov et al. (69); Mahler (71); Mehra et al. (75); Ninio et al. (88); Owen et al. (91); Pearson (94); Pincus et al. (99, 100); Prohofsky (104); Rimmer (107); Simanek et al. (113); Suhl (122, 123); Turov et al. (130).

Critical Phenomena

Cooper et al. (17); Golding (38); Teaney et al. (128).

Elasticity-Magnetoelastic Interactions

Collins et al. (15); Denison et al. (20); Eastman (22); Eastman et al. (23); Golding (38); Melcher et al. (77); Pickart et al. (98); Turov et al. (130).

Electron-Nuclear Double Resonance (ENDOR)

Heeger et al. (47, 48, 49); Lee et al. (68); Pincus et al. (99); Portis et al. (103); Shaltiel et al. (110); Witt (138); Witt et al. (139, 140).

Electron Spin Resonance (ESR)

Hall et al. (42); Horai et al. (57); Zhogolev (142).

Hyperfine Interactions

Cooke et al. (16); Egashira et al. (26); Freeman et al. (35); Montgomery et al. (82); Nakamura (85); Shulman et al. (111); Suhl (123); Walker (131).

Impurities

Eastman et al. (25); Holloway et al. (54, 56); Johnson et al. (61); Matyushkin et al. (73); Misetich et al. (81); Perry et al. (95); Pincus et al. (100); Shionoya et al. (112).

Neutron Diffraction

Collins et al. (15); Cooper et al. (17); Johnson et al. (61); Nathans et al. (87); Pickart et al. (97, 98); Satya Murthy et al. (108); Windsor (135, 136); Windsor et al. (137).

Nuclear Magnetic Resonance (NMR)

DeGennes (19); Denison et al. (20); Egashira et al. (26); Freiser et al. (36); Heeger et al. (50); Jones et al. (63); Mahler et al. (72); Minkiewicz (78); Minkiewicz et al. (79, 80); Nakamura (85); Nakamura et al. (86); Ninio et al. (88); Payne et al. (92, 93); Petrov et al. (96); Portis (102); Shulman et al. (111); Teaney (124); Turov et al. (129); Walker (131); Walker et al. (132); Welsh (134).

Optical Studies

Antonov et al. (2); Aoyagi (3); Dietz et al. (21); Elliott et al. (27); Eremenko et al. (28); Ferguson et al. (30, 31, 32, 33); Fleury (34); Holloway et al. (53, 54, 55, 56); Imbusch et al. (59); Johnson et al. (62); Kharchenko et al. (67); Matyushkin et al. (73); Mehra et al. (75, 76); Misetich et al. (81); Perry et al. (95); Prohofsky (104); Prohofsky et al. (105); Shionoya et al. (112); Stevenson (115, 116, 117, 119, 120); Young et al. (141).

Physical Properties

Aleksandrov et al. (1); Beckman et al. (4, 5); Belyaev et al. (6, 7); Breed (9); Chang (11); Collins et al. (15); Cooke et al. (16); Cooper et al. (17); Deenades et al. (18); Gorodetsky (41); Hashimoto (43); Heeger et al. (45); Hirakawa et al. (52); Joseph (64); Kedzie et al. (66); Lee et al. (68); Machin et al. (70); McGuire (74); Montgomery et al. (82); Moruzzi et al. (83, 84); Ogawa (89); Okazaki et al. (90); Pearson (94); Pickart et al. (97); Prohovsky et al. (105); Rao et al. (106); Simanov et al. (114); Stevenson et al. (118); Suemune et al. (121); Teaney (124).

Relaxation

Blume et al. (8); Heeger et al. (48); Hinderks et al. (51); Mahler (71); Mahler et al. (72); Portis (101); Suhl (123).

Spin Waves

Cole (12); Cole et al. (13); Freiser et al. (36); Heeger (44); Heeger et al. (46); Hinderks et al. (51); Nathans et al. (87); Ninio et al. (88); Windsor et al. (137).

PROPERTIES OF THE COMPOUNDS RbMnF_3 , KMnF_3 AND CsMnF_3

1. Aleksandrov, K. S., Reshchikova, L. M. and Beznosikov, B. V.

Behaviour of the Elastic Constants of KMnF_3 Single Crystals near the Transition of Puckering

Phys Stat Sol, Vol 18, No 1, pp K17-K20 (1 Nov 1966).

The velocities of longitudinal and shear elastic waves along principal crystalline directions were measured by an ultrasonic pulse method at a frequency of 30 MHz.

2. Antonov, A. V., Belyaeva, A. I. and Eremenko, V. V.

Low-Temperature Anomaly in the Absorption Spectra of the Antiferromagnets RbMnF_3 and KMnF_3

Fiz Tverd Tela, Vol 8, No 11, pp 3397-3399 (Nov 1966); Engl. transl. in Soviet Phys-Solid State, Vol 8, No 11, pp 2718-2720 (May 1967).

A discontinuity was found in the absorption of the $C_1(^6A_{1g} \rightarrow ^9E_g)$ and C_1 bands ($\sim 3900 \text{ \AA}$) at temperatures below 30°K .

3. Aoyagi, K.

Observation of Magnon Sideband in the Absorption Spectrum of KMnF_3

J Phys Soc Japan, Vol 22, No 6, pp 1516-1517 (Jun 1967).

The absorption lines B ($17,892 \text{ cm}^{-1}$) and C ($17,811 \text{ cm}^{-1}$) are attributed to pure electronic transitions (main lines) corresponding to the transition $^6A_{1g} \rightarrow ^4T_{1g}$ of Mn^{2+} , while the A ($17,944 \text{ cm}^{-1}$) line is attributed to the magnon sideband of C.

4. Beckman, O. and Knox, K.

Magnetic Properties of KMnF_3 . I. Crystallographic Studies

Phys Rev, Vol 121, No 2, pp 376-380 (15 Jan 1961).

Measurements of the lattice parameters for KMnF_3 by means of an x-ray rotation camera designed for temperatures down to 15°K are reported.

5. Beckman, O., Olovsson, I. and Knox, K.

Structural Changes of KMnF_3 at Low Temperatures

Acta Cryst, Vol 13, Pt 6, p 506 (10 Jun 1960).

Using x-ray analysis, the crystal structure of KMnF_3 was examined when the material was cooled through the Néel temperature.

6. Belyaev, I. N. and Revina, O. Ya.

The MF- MnF_2 Systems (M = Li, Na, K, Rb, Cs)

Zhur Neorgan Khim, Vol 11, No 6, pp 1446-1450 (1966); Engl. transl. in Russian J Inorgan Chem, Vol 11, No 6, pp 772-774 (Jun 1966).

The compositions of binary systems of the alkali metal fluorides were studied by thermography.

7. Belyaev, I. N. and Revina, O. Ya.

Ternary Systems of the Manganese and Alkali Metal Fluorides

Zhur Neorgan Khim, Vol 11, No 8, pp 1952-1958 (1966);

Engl. transl. in Russian J Inorgan Chem, Vol 11, No 8, pp 1041-1044 (Aug 1966).

Systems studied include LiF-CsF-MnF_2 , NaF-CsF-MnF_2 , KF-CsF_2 , CsF-KF , $\text{NaMnF}_3\text{-CsMnF}_3$, and $\text{CsMnF}_3\text{-KMnF}_3$.

8. Blume, M. and Orbach, H.

Spin-Lattice Relaxation of S-State Ions: Mn^{2+} in a Cubic Environment
Phys Rev, Vol 127, No 5, pp 1587-1592 (1 Sep 1962)

The theory of spin-lattice relaxation is developed for S-state ions; in particular, for Mn^{2+} in a cubic environment. The wave functions for Mn^{2+} are generated to 1st order in the spin-orbit coupling parameter, the orbit-lattice interaction is formulated in terms of spherical harmonics, and the rate equations are derived for this system.

9. Breed, D. J.

Experimental Investigation of Two Two-Dimensional Antiferromagnets with Small Anisotropy
Physica, Vol 37, No 1, pp 35-46 (1967).

The magnetic properties have been studied of two antiferromagnetic Mn compounds, K_2MnF_4 and Rb_2MnF_4 .

10. Bullock, D. L.

Perturbation Treatment of the Antiferromagnetic Ground State
Phys Rev, Vol 137, No 6A, pp 1877-1885 (15 Mar 1965).

Ground-state parameter series expansions are generated for the Heisenberg model. Experimental results for $KMnF_3$ and MnF_2 are not contradicted by this perturbation treatment.

11. Chang, T.-S.

Dielectric Properties of $RbMnF_3$
J Appl Phys, Vol 39, No 7, pp 3511-3512 (Jun 1967).

The dielectric constant and loss tangent of $RbMnF_3$ were measured as functions of frequency in the temperature range 25°C to 300°C.

12. Cole, P. H.

Nonlinear Coupling between Antiferromagnetic Resonance Modes in $RbMnF_3$
Appl Phys Lett, Vol 10, No 10, pp 272-275 (15 May 1967).

Nonlinear coupling between AFMR modes spaced in frequency an octave apart has been demonstrated experimentally. Calculations of the power-dependent conversion relation for up-conversion, and the critical power for subharmonic oscillation are given.

13. Cole, P. H. and Courtney, W. E.

Uniform Mode Resonance and Spin-Wave Instability in $RbMnF_3$
Proc 12th Ann Conf Magnetism and Magnetic Materials, Washington, D. C., 1966, ed. by E. G. Spencer and J. S. Kouvel; J Appl Phys, Vol 38, No 3, pp 1278-1279 (1 Mar 1967).

Low-power resonance linewidth and high-power 2nd-order spin-wave instability measurements were performed at 4.2°K.

14. Cole, P. H. and Ince, W. J.

Equilibrium Spin Configuration and Resonance Behavior of $RbMnF_3$
Phys Rev, Vol 150, No 2, pp 377-383 (14 Oct 1966).

Observations of the resonance spectrum at X-band frequencies for the applied field range 0 to 12 kOe are presented.

15. Collins, M. F. and Nathans, R.
Some Measurements of Exchange Energies by Paramagnetic Neutron Inelastic Scattering
Proc 10th Conf Magnetism and Magnetic Materials, Minneapolis, Minn., 1964, ed. by I. S. Jacobs and E. G. Spencer; J Appl Phys, Vol 36, No 3, Pt 2, pp 1092-1093 (Mar 1965).

Neutron paramagnetic scattering techniques have been applied to the measurement of exchange constants for a series of perovskite salts, KMnF_3 , NaMnF_3 , LaCrO_3 , LaMnO_3 , and LaFeO_3 .
16. Cooke, A. H. and Edmonds, D. T.
Nuclear Magnetic Interaction in an Antiferromagnetic Crystal
Proc Phys Soc, Vol 71, Pt 3, pp 517-519 (Jun 1958).

The measurement of the specific heat anomaly at the Néel temperature gave a value for the hyperfine coupling constant in MnF_2 in good agreement with previous determinations by other methods.
17. Cooper, M. J. and Nathans, R.
Critical Magnetic Scattering from KMnF_3
Proc 11th Ann Conf Magnetism and Magnetic Materials, San Francisco, Calif., 1965, ed. by E. G. Spencer and J. S. Kouvel; J Appl Phys, Vol 37, No 3, pp 1041-1047 (1 Mar 1966).

Magnetic scattering experiments on KMnF_3 are described.
18. Deenadas, D., Keer, H. V., Gopala Rao, R. V. and Biswas, A. B.
Heat Capacity of Potassium Manganese Trifluoride
British J Appl Phys, Vol 17, No 11, pp 1401-1404 (Nov 1966).

The heat capacity of KMnF_3 has been measured in the temperature range 80°K to 300°K.
19. DeGennes, P. G.
Dynamic Effects in the Nuclear Resonance of Magnetic Materials (in French)
"Magetic and Electric Resonance and Relaxation," Proc 11th Colloq AMPERE (Atomes Mol. Études Radio Elec.), Eindhoven, 1962, ed. by J. Smidt, North-Holland Publ. Co., Amsterdam, pp 88-95 (1963).

The nuclear spin-wave spectrum produced by the Suh-Nakamura indirect interaction is discussed.
20. Denison, A. B., James, L. W., Currin, J. D., Tanttila, W. H. and Mahler, R. J.
Ultrasonically Induced Nuclear Spin Transitions in Antiferromagnetic KMnF_3
Phys Rev Lett, Vol 12, No 10, pp 244-245 (9 Mar 1964).

A technique for producing nuclear spin transitions of F^{19} nuclei in KMnF_3 single crystals via ultrasonic modulation is described.
21. Dietz, R. E., Johnson, L. F. and Guggenheim, H. J.
Fluorescence from Magnetic Crystals
"Physics of Quantum Electronics, Conference Proceedings," ed. by P. L. Kelley, B. Lax and P. E. Tannenwald, McGraw-Hill Book Co., New York, N. Y., pp 361-369 (1966).

Near-infrared fluorescence and absorption have been investigated for Ni in KMnF_3 .
22. Eastman, D. E.
Magneto-Elastic Coupling in RbMnF_3
Phys Rev, Vol 156, No 2, pp 645-654 (10 Apr 1967).

Magnetoelastic coupling effects have been studied by observing shifts in AFMR frequency and changes in AFMR line shape with the application of axial stress.

23. Eastman, D. E., Joenk, R. J. and Teaney, D. T.
Antiferromagnetic Resonance in RbMnF₃ under Axial Stress
 Phys Rev Lett, Vol 17, No 6, pp 300-302 (8 Aug 1966).
 Magnetoelastic constants of antiferromagnetic RbMnF₃ were determined at 4.2°K.
24. Eastman, D. E. and Shafer, M. W.
Antiferromagnetic Resonance in Cubic TlMnF₃
 J Appl Phys, Vol 38, No 3, pp 1274-1276 (1 Mar 1967).
 Antiferromagnetic resonance studies at 25 GHz showed TlMnF₃ to be an undistorted cubic antiferromagnet at 4.2°K with <111> easy axes.
25. Eastman, D. E., Shafer, M. W. and Figat, R. A.
Cobalt-Doped TlMnF₃, a Zero Anisotropy Cubic Antiferromagnet
 J Appl Phys, Vol 38, No 13, pp 5209-5211 (Dec 1967).
 Single crystals of cobalt-doped thallium manganese fluoride (TlMn_{1-x}Co_xF₃) have been studied by AFMR for cobalt concentrations of $0 \leq x \leq 0.001$. Comparison with AFMR of undoped TlMnF₃ and RbMnF₃ is made.
26. Egashira, K. and Hirakawa, K.
Observation of the Hyperfine Interaction Constants in KMnF₃
 J Phys Soc Japan, Vol 22, No 1, p 344 (Jan 1967).
 The hyperfine constant is obtained from NMR experiments on the F¹⁹ nucleus.
27. Elliott, R. J., Thorpe, M. F., Imbusch, G. F., Loudon, R. and Parkinson, J. B.
Magnon-Magnon and Exciton-Magnon Interaction Effects on Antiferromagnetic Spectra
 Phys Rev Lett, Vol 21, No 3, pp 147-150 (15 Jul 1968).
 Green's function methods have been used to calculate the effect of magnon-magnon and exciton-magnon interactions on the sidebands in RbMnF₃.
28. Eremenko, V. V., Popkov, Yu. A., Novikov, V. P. and Belyaeva, A. I.
Characteristics of Exciton-Magnon Interaction in Antiferromagnetic Crystals with a Perovskite Structure
 Zhur Eksper Teor Fiz, Vol 52, No 2, pp 454-462 (Feb 1967); Engl. transl. in Soviet Phys - JETP, Vol 25, No 2, pp 297-302 (Jul 1967).
 The optical absorption spectra of RbMnF₃ and KMnF₃ are studied in the ${}^6A_{1g}({}^6S) \rightarrow {}^4E_g({}^4G)$ transition range in the Mn²⁺ ion.
29. Farztdinov, M. M.
Domains and S-Domain Boundaries in Antiferromagnetic Compounds
 Fiz Metal Metalloved, Vol 19, No 3, pp 321-332 (1965); Engl. transl. in Phys Met Metallog, Vol 19, No 3, pp 1-12 (1965).
 The possible domains and S-domain boundaries were investigated in antiferromagnetic compounds of the NiO type, and also uniaxial (Cr₂O₃, MnF₂) and orthorhombic (CuCl₂ · 2H₂O).
30. Ferguson, J., Guggenheim, H. J. and Tanabe, Y.
Absorption of Light by Pairs of Exchange-Coupled Manganese and Nickel Ions in Cubic Perovskite Fluorides
 J Chem Phys, Vol 45, No 4, pp 1134-1142 (15 Aug 1966).
 The absorption of light by exchange-coupled pairs of Mn and Ni ions in KZnF₃ is reported. Corresponding effects of exchange interaction have also been observed in the spectra of KMn_xNi_{1-x}F₃ crystals.

31. Ferguson, J., Guggenheim, H. J. and Tanabe, Y.
Absorption of Light by Pairs of Like and Unlike Transition-Metal Ions
 Phys Rev Lett, Vol 14, No 18, pp 737-738 (3 May 1965).
 Absorption of light by pairs of like and unlike transition-metal ions is described. Absorption by pairs of unlike ions is possible, and in all cases studied, the intensity of absorption is greater than for like ion pairs.

32. Ferguson, J., Guggenheim, H. J. and Tanabe, Y.
The Effects of Exchange Interactions in the Spectra of Octahedral Manganese. II. Compounds
 J Phys Soc Japan, Vol 21, No 4, pp 692-704 (Apr 1966).
 Experimental measurements of the Mn-pair spectrum of $\text{KZnF}_3\text{:Mn}$ and the spectrum of KMnF_3 are reported.

33. Ferguson, J., Guggenheim, H. J. and Tanabe, Y.
Exchange Effects in the Electronic Absorption Spectrum of Mn (II) in Perovskite Fluorides
 Proc 10th Conf Magnetism and Magnetic Materials, Minneapolis, Minn., 1964, ed. by I. S. Jacobs and E. G. Spencer; J Appl Phys, Vol 36, No 3, Pt 2, pp 1046-1047 (Mar 1965).
 The anomalously intense bands in the spectrum of KMnF_3 are interpreted on the basis of pure electronic transitions involving more than one ion.

34. Fleury, P. A.
Evidence for Magnon-Magnon Interactions in RbMnF_3
 Phys Rev Lett, Vol 21, No 3, pp 151-153 (15 Jul 1968).
 Light scattering from two-magnon excitations in RbMnF_3 has been detected. The experiments confirm the importance of magnon-magnon interactions.

35. Freeman, A. J. and Watson, R. E.
Origin of the F^{19} Hyperfine Structure in Transition Element Fluorides
 Phys Rev Lett, Vol 6, No 7, pp 343-345 (1 Apr 1961).
 The unpairing of the 1s electrons gives an appreciable contribution to the isotropic part of the hyperfine structure. This is illustrated with use of KMnF_3 as an example.

36. Freiser, M. J., Joenk, R. J., Seiden, P. E. and Teaney, D. T.
Magnetic Resonance Studies of the Cubic Antiferromagnet RbMnF_3
 Proc Int Conf Magnetism, Nottingham, 1964, publ. by Institute of Physics and the Physical Society, London, pp 432-436 (1965).
 The properties of the cubic antiferromagnet RbMnF_3 were studied by AFMR and Mn^{55} NMR.

37. Freiser, M. J., Seiden, P. E. and Teaney, D. T.
Field-Independent Longitudinal Antiferromagnetic Resonance
 Phys Rev Lett, Vol 10, No 7, pp 293-294 (1 Apr 1963).
 The field-independent AFMR mode for the flopped state was studied as a function of frequency and temperature.

38. Golding, B.
Ultrasonic Propagation in RbMnF_3 near the Magnetic Critical Point
 Phys Rev Lett, Vol 20, No 1, pp 5-7 (1 Jan 1968).
 Measurements of the ultrasonic attenuation and velocity near the magnetic critical point have been performed at UHF frequencies.

39. Gondaira, K.

Covalency Effects in KMnF_3

J Phys Soc Japan, Vol 21, No 5, pp 933-944 (May 1966).

The covalency parameters between Mn 3d orbitals and the neighboring F 2s, 2p orbitals in KMnF_3 are calculated by means of the molecular orbital method.

40. Gondaira, K. and Tanabe, Y.

A Note on the Theory of Superexchange Interaction

J Phys Soc Japan, Vol 21, No 8, pp 1527-1548 (Aug 1966).

Expressions are derived for the coupling constant in the interaction between a paramagnetic ion pair and an electric field. KMnF_3 is used as an example.

41. Gorodetsky, G.

Semiempirical Expression for the Paramagnetic Susceptibility of LuFeO_3 and KMnF_3 by the Padé Approximant

Solid State Commun, Vol 6, No 3, pp 159-162 (Mar 1968).

The theoretical susceptibility, based on the use of a Padé approximation, gives better fit to the experimental data in the region $(T/T_C) < 2$ than the expression obtained by using a high-temperature series expansion.

42. Hall, T. P. P., Hayes, W. and Williams, F. I. B.

Paramagnetic Resonance of Manganese

Proc Phys Soc, Vol 78, Pt 5, pp 883-894 (Nov 1961).

The paramagnetic resonance spectrum of bivalent Mn was studied in the crystals CdF_2 , KMgF_3 , and CdTe . Comparison is drawn with the spectrum of KMnF_3 .

43. Hashimoto, T.

Magnetic Properties of the Solid Solutions between KMnF_3 , KCoF_3 , and KNiF_3

J Phys Soc Japan, Vol 18, No 8, pp 1140-1147 (Aug 1963).

The magnetic susceptibilities of KNiF_3 and the solid solutions between KMnF_3 , KCoF_3 , and KNiF_3 have been measured in the temperature range between 80°K and 850°K.

44. Heeger, A. J.

Spin-Wave Instability and Premature Saturation in Antiferromagnetic Resonance

Phys Rev, Vol 131, No 2, pp 608-616 (15 Jul 1963).

Experimental evidence for the existence of spin-wave instability in the canted antiferromagnet KMnF_3 is presented and discussed.

45. Heeger, A. J., Beckman, O. and Portis, A. M.

Magnetic Properties of KMnF_3 . II. Weak Ferromagnetism

Phys Rev, Vol 123, No 5, pp 1652-1660 (1 Sep 1961).

The static magnetic properties of single-crystal KMnF_3 were studied by magnetic torsion measurements.

46. Heeger, A. J. and Pincus, P.

Spin-Wave Instability and Premature Saturation in Antiferromagnetic Resonance

Phys Rev Lett, Vol 10, No 2, pp 53-55 (15 Jan 1963).

Premature saturation and spin-wave instability were observed in KMnF_3 .

47. Heeger, A. J., Portis, A. M., Teaney, D. T. and Witt, G.
Double Resonance and Nuclear Cooling in an Antiferromagnet
 Phys Rev Lett, Vol 7, No 8, pp 307-309 (15 Oct 1961).
 Electron-nuclear double resonance was observed in KMnF_3 .
 Measurements of the saturation decay yielded a value of
 50 msec for T_1 .
48. Heeger, A. J., Portis, A. M. and Witt, G.
Relaxation of Mn^{55} Nuclear Magnetization in Antiferromagnetic KMnF_3
 Bull Amer Phys Soc, Ser 2, Vol 7, No 1, p 54 (24 Jan 1962).
 Observations of the nuclear resonance of Mn^{55} in antiferromagnetic
 KMnF_3 by the double resonance technique are discussed.
49. Heeger, A. J., Portis, A. M. and Witt, G.
Nuclear Antiferromagnetic Double Resonance
 "Magnetic and Electric Resonance and Relaxation," Proc 11th Colloq AMPERE
 (Atomes Mol. Études Radio Elec.), Eindhoven, 1962, ed. by J. Smidt, North-
 Holland Publ. Co., Amsterdam, pp 694-698 (1963).
 This technique was applied to demonstrate the dependence of the
 Mn^{55} nuclear resonance frequency in KMnF_3 on nuclear magnetization.
50. Heeger, A. J. and Teaney, D. T.
 Mn^{55} Nuclear Magnetic Resonance in Antiferromagnetic RbMnF_3
 Proc 9th Conf Magnetism and Magnetic Materials, Atlantic City, N. J.,
 1963, ed. by J. A. Osborn; J Appl Phys, Vol 35, No 3, Pt 2, pp 846-847
 (Mar 1964).
 The observation and study of the Mn^{55} NMR in antiferromagnetic
 RbMnF_3 at low temperatures are reported.
51. Hinderks, L. W. and Richards, P. M.
Excitation of Nuclear and Electronic Spin Waves in RbMnF_3 by Parallel Pumping
 J Appl Phys, Vol 39, No 2, Pt 1, pp 824-825 (1 Feb 1968).
 Observation of the simultaneous excitation of nuclear and electronic
 spin waves by parallel pumping is reported. From threshold fields,
 which are about 1.2 Oe minimum, it is calculated that at 4.2°K,
 $\eta K^n \eta K^e = 6.1 \times 10^{10} \text{ sec}^{-2}$ for $K \rightarrow 0$, and $9.5 \times 10^{10} \text{ sec}^{-2}$ for
 $K = 1.05 \times 10^5 \text{ cm}^{-1}$.
52. Hirakawa, K., Hirakawa, K. and Hashimoto, T.
Magnetic Properties of Potassium Iron Group Fluorides KMF_3
 J Phys Soc Japan, Vol 15, No 11, pp 2063-2068 (Nov 1960).
 The susceptibility of KMnF_3 , in particular, was measured as a
 function of temperature.
53. Holloway, W. W., Jr. and Kestigian, M.
Effect of Magnetic Ordering on the Fluorescence of MnF_2
 Phys Rev Lett, Vol 13, No 7, pp 235-237 (17 Aug 1964).
 Measurements were made of the wavelength, the relative intensity,
 and the lifetime of fluorescence in RbMnF_3 , CsMnF_3 , etc.
54. Holloway, W. W., Jr. and Kestigian, M.
Temperature Dependence of the $^3A_2 \rightarrow ^3T_1$ Absorption Band Peak of the Ni^{2+}
 Ion in Nickel and Manganese Salts
 Phys Rev Lett, Vol 15, No 1, pp 17-19 (5 Jul 1965).
 The effect of magnetic interaction on the temperature dependence
 of the $^3A_2 \rightarrow ^3T_1$ absorption band peak of the Ni^{2+} ion in RbMnF_3 is
 discussed with reference to RbMnF_3 and other compounds.

55. Holloway, W. W., Jr., Kestigian, M., Newman, R. and Prohofsky, E. W.
Anomalous Shifts in the Fluorescence of MnF_2 and KMnF_3
 Phys Rev Lett, Vol 11, No 2, pp 82-84 (15 Jul 1963).
 Large temperature-dependent changes in the fluorescence of MnF_2 and KMnF_3 were observed.
56. Holloway, W. W., Jr., Prohofsky, E. W. and Kestigian, M.
Magnetic Ordering and the Fluorescence of Concentrated Mn Systems
 Phys Rev, Vol 139, No 3A, pp A954-A961 (2 Aug 1965).
 The wavelength, lifetime, and relative intensity of the fluorescences of crystal samples of MnF_2 and the alkali Mn trifluorides were measured as functions of temperature above 21°K.
57. Horai, K. and Saiki, K.
Electron Spin Resonance of $(\text{KMnF}_3)_x(\text{KMgF}_3)_{1-x}$
 J Phys Soc Japan, Vol 21, No 2, p 397 (Feb 1966).
 The transition temperature at which the ESR field starts to shift from the room-temperature value has been measured for concentration x between 0.1 and 1.0.
58. Hubbard, J., Rimmer, D. E. and Hopgood, F. R. A.
Weak Covalency in Transition Metal Salts
 Proc Phys Soc, Vol 88, Pt 1, No 559, pp 13-36 (May 1966).
 A theory of weak covalency effects in transition-metal salts based upon a configuration-interaction approach is discussed. The method is applied to KMnF_3 , etc.
59. Imbusch, G. F. and Guggenheim, H. J.
Fine Structure in the ${}^6\text{A}_1 \rightarrow {}^4\text{T}_1$ (${}^4\text{G}$) Absorption Transition in RbMnF_3
 Phys Lett, Vol 26A, No 12, pp 625-626 (6 May 1968).
 Two sharp magnetic dipole, no-phonon, no-magnon lines are observed in absorption in RbMnF_3 along with a suggested magnon sideband.
60. Ince, W. J.
Antiferromagnetic Resonance in RbMnF_3 below the Spin-Flop Transition
 J Appl Phys, Vol 37, No 3, pp 1132-1133 (1 Mar 1966).
 This paper reports an investigation of resonance for the low-field region 0 kOe to 5 kOe.
61. Johnson, L. F., Dietz, R. E. and Guggenheim, H. J.
Exchange Splitting of the Ground State of Ni^{2+} Ions in Antiferromagnetic MnF_2 , KMnF_3 , and RbMnF_3
 Phys Rev Lett, Vol 17, No 1, pp 13-15 (4 Jul 1966).
 The splittings of some of the sidebands agree with the zone-boundary magnon frequencies determined by neutron scattering.
62. Johnson, L. F., Dietz, R. E. and Guggenheim, H. J.
Optical Studies of Ni^{2+} Ions in MnF_2 and KMnF_3
 Bull Amer Phys Soc, Vol 10, Ser 2, No 8, p 1193 (1965).
 This study concerns principally the ${}^3\text{A}_2$ (ground-state) $\rightarrow {}^3\text{T}_2$ (first excited state) transition.
63. Jones, E. D. and Jefferts, K. B.
Zero-Field Manganese Nuclear Magnetic Resonance in Antiferromagnetic Manganese Fluoride
 Phys Rev, Vol 135, No 5A, pp A1277-A1280 (31 Aug 1964).
 The extrapolated Mn^{55} NMR frequency at 0°K is found to be $\nu_0^{55} = 671.4 \pm 0.2$ MHz.

64. Joseph, R. I.

Theory of High-Temperature Susceptibility of Heisenberg Ferromagnets Having Nearest-Neighbor Bilinear and Biquadratic Exchange Interactions
Phys Rev, Vol 138, No 5A, pp A1441-A1444 (31 May 1965).

The high-temperature susceptibility series is derived for the nearest-neighbor Heisenberg ferromagnet when nearest-neighbor biquadratic exchange is included. KMnF_3 is chosen as an example.

65. Kedzie, R. W., Shane, J. R. and Kestigian, M.

Spin-Flop AFMR in an Oriented CsMnCl_3 Single Crystal
Bull Amer Phys Soc, Vol 10, Ser 2, No 3, p 315 (1965).

Antiferromagnetic resonances have been observed at low temperatures at X- and K-band frequencies in an oriented single crystal of CsMnCl_3 ($T_N = 69^\circ\text{K}$).

66. Kedzie, R. W., Shane, J. R., Kestigian, M. and Croft, W. J.

Resonance Observation of Antiferromagnetic Ordering in RbMnCl_3 , CsMnCl_3 , and KMnCl_3
Proc 10th Conf Magnetism and Magnetic Materials, Minneapolis, Minn., 1964, ed. by I. S. Jacobs and E. G. Spencer; J Appl Phys, Vol 36, No 3, Pt 2, pp 1195-1196 (Mar 1965).

Paramagnetic and antiferromagnetic resonances were observed as functions of temperature in KMnCl_3 , CsMnCl_3 , and RbMnCl_3 at 9.5 GHz and 25 GHz.

67. Kharchenko, N. F. and Eremenko, V. V.

Magnetoresonance Faraday Effect in the Antiferromagnets MnF_2 and RbMnF_3
Fiz Tverd Tela, Vol 9, No 6, pp 1655-1659 (Jun 1967); Engl. transl. in Soviet Phys - Solid State, Vol 9, No 6, pp 1302-1305 (Dec 1967).

The Faraday effect in MnF_2 and RbMnF_3 was measured at several wavelengths in the visible part of the spectrum in a magnetic field up to 150 kOe, over the temperature range 20°K to 150°K .

68. Lee, K., Portis, A. M. and Witt, G. L.

Magnetic Properties of the Hexagonal Antiferromagnet CsMnF_3
Phys Rev, Vol 132, No 1, pp 144-163 (1 Oct 1963).

The magnetic properties of the hexagonal antiferromagnet CsMnF_3 were investigated by magnetic susceptibility, torsion, ENDOR and electron resonance.

69. Lvov, V. S. and Petrov, M. P.

The Distribution of Spin Density in Paramagnetic Perovskite Crystals
Phys Stat Sol, Vol 13, No 2, pp K65-K68 (1 Feb 1966).

A tight-binding calculation has been made. It explains the observed dependence of the spin density on the nonmagnetic nucleus (Na^+ , Rb^+ ...) on the type of paramagnetic ion (Mn^{2+} , Ni^{2+} ...).

70. Machin, D. J., Martin, R. L. and Nyholm, R. S.

The Preparation and Magnetic Properties of Some Complex Fluorides Having the Perovskite Structure
J Chem Soc, Pt 2, No 281, pp 1490-1500 (1963).

The magnetic properties of compounds of the type $\text{KM}^{\text{II}}\text{F}_3$ ($\text{M}^{\text{II}} = \text{Mn}, \text{Fe}, \text{Co}, \text{Ni}, \text{Cu}, \text{and Zn}$) have been studied in the temperature range 80°K to 300°K . The manganese compound shows evidence of antiferromagnetism, with a Néel point below 80°K .

71. Mahler, R. J.
Phonon Induced Nuclear Dipole Transitions
 "Nuclear Magnetic Resonance and Relaxation in Solids," Proc 13th Colloq
 AMPERE (Atomes Mol. Études Radio Elec.), Leuven, 1964, ed. by L. Van Gerven,
 North-Holland Publ. Co., Amsterdam, pp 202-209 (1964).
 This paper outlines the theory of the interaction between the
 nuclear spin system and the lattice. Strong interaction was
 detected experimentally with KMnF_3 at 4.2°K.
72. Mahler, R. J., Daniel, A. C. and Parrish, P. T.
Observation of Two Intrinsic Nuclear Relaxation Rates in Antiferromagnetic KMnF_3
 Phys Rev Lett, Vol 19, No 2, pp 85-87 (10 Jul 1967).
 Measurement of nuclear spin-lattice relaxation rates of the F^{19}
 nuclei at two nonequivalent sites in KMnF_3 were made with a
 spin-echo technique.
73. Matyushkin, E. V., Kukushkin, L. S. and Eremenko, V. V.
Peculiarities of the $\text{RbMnF}_3\text{:Nd}^{3+}$ Crystals Decay Kinetics due to the Migration
 of Excitation Energy
 Phys Stat Sol, Vol 22, No 1, pp 65-69 (1 Jul 1967).
 The decay kinetics of Nd^{3+} luminescence in $\text{RbMnF}_3\text{:Nd}^{3+}$
 crystals were investigated experimentally.
74. McGuire, T. R.
Magnetic Susceptibility of RbMnF_3
 Bull Amer Phys Soc, Ser 2, Vol 8, No 7, p 55 (23 Jan 1963).
 Results of the temperature and field dependence of the single-
 crystal susceptibility are presented.
75. Mehra, A. and Venkateswarlu, P.
Spin-Orbit Coupling Parameter for Mn^{2+} in RbMnF_3
 Phys Rev Lett, Vol 19, No 4, pp 145-146 (24 Jul 1967).
 Fine structure measurements of the ${}^6\text{A}_{1g}(\text{S}) \rightarrow {}^4\text{T}_{2g}(\text{D})$ band in the
 absorption spectrum at 77°K were used to calculate the spin-orbit
 coupling parameter for Mn^{2+} .
76. Mehra, A. and Venkateswarlu, P.
Absorption Spectrum of RbMnF_3
 J Chem Phys, Vol 47, No 7, pp 2334-2342 (1 Oct 1967).
 The absorption spectrum of RbMnF_3 has been studied at room
 temperature and at 77°K. The observed band positions are fitted
 with four parameters, B, C, Dq, and α , and the values of the
 parameters for the 77°K spectrum are 840 cm^{-1} , 3080 cm^{-1} ,
 780 cm^{-1} , and 76 cm^{-1} , respectively.
77. Melcher, R. L., Bolef, D. I. and Stevenson, R. W. H.
Magnetic Field-Dependent Elastic Effects in the Cubic Antiferromagnet RbMnF_3
 Solid State Commun, Vol 5, No 9, pp 735-738 (Sep 1967).
 Measurements of velocity and attenuation of 10-MHz longitudinal
 acoustic waves in antiferromagnetic RbMnF_3 are reported.
78. Minkiewicz, V. J.
Nuclear Magnetic Resonance in Antiferromagnets
 Ph. D. Dissertation, University of Calif., Berkeley (1965); University
 Microfilms Order No. 66-3659.
 The Mn^{55} nuclear magnetic resonance in KMnF_3 has been ob-
 served in the frequency range between 580 MHz and 620 MHz.

79. Minkiewicz, V. and Nakamura, A.
Direct Observation of Mn^{55} Nuclear Magnetic Resonance in Antiferromagnetic $CsMnF_3$
 Phys Rev, Vol 143, No 2, pp 361-365 (11 Mar 1966).
 The direct NMR of the Mn^{55} nuclei in the hexagonal antiferromagnet $CsMnF_3$ was observed in the frequency range 250 MHz to 650 MHz.
80. Minkiewicz, V. and Nakamura, A.
Magnetic Properties of $KMnF_3$; III. Nuclear and Electron Spin Resonance
 Phys Rev, Vol 143, No 2, pp 356-360 (11 Mar 1966).
 Direct NMR experiments on Mn^{55} nuclei in $KMnF_3$ are reported.
81. Misetich, A. and Dietz, R. E.
Effect of Zero-Point Spin Deviation on Energy Levels of Magnetic Impurities in Antiferromagnets
 Phys Rev Lett, Vol 17, No 7, pp 392-395 (15 Aug 1966).
 The infrared emission spectrum of Ni^{2+} as an impurity in MnF_2 , $KMnF_3$, and $RbMnF_3$ was studied.
82. Montgomery, H., Teaney, D. T. and Walsh, W. M., Jr.
Hyperfine Field and Ground-State Spin Alignment in Antiferromagnetic $KMnF_3$
 Phys Rev, Vol 128, No 1, pp 80-81 (1 Oct 1962).
 The specific heat of antiferromagnetic $KMnF_3$ at low temperatures was measured and the contribution of the Mn^{55} polarization determined.
83. Moruzzi, V. L. and Teaney, D. T.
Observation of the Specific-Heat Anomaly Associated with the Antiferromagnetic Transition in $RbMnF_3$
 Bull Amer Phys Soc, Ser 2, Vol 8, No 4, p 382 (1963).
 The specific heat of a powder sample of $RbMnF_3$ has been measured from 25°K to 170°K.
84. Moruzzi, V. L. and Teaney, D. T.
Specific Heats of Antiferromagnet $XMnF_3$ Compounds
 Bull Amer Phys Soc, Vol 9, Ser 2, No 3, p 225 (1964).
 The specific heats of five powder samples of $XMnF_3$ compounds (Na, K, NH_4 , Rb, Cs) have been measured in the range of 15°K to 280°K.
85. Nakamura, T.
Indirect Coupling of Nuclear Spins in an Antiferromagnet with Particular Reference to MnF_2 at Very Low Temperatures
 Prog Theor Phys, Vol 20, No 4, pp 542-552 (Oct 1958).
 The linewidth of the F^{19} NMR in MnF_2 at 1.4°K is interpreted in terms of indirect coupling of nuclear spins through hyperfine interaction with spin waves.
86. Nakamura, A., Minkiewicz, V. and Portis, A. M.
Direct Mn^{55} NMR Absorption in Antiferromagnetic $KMnF_3$
 Proc 9th Conf Magnetism and Magnetic Materials, Atlantic City, N. J., 1963, ed. by J. A. Osborn; J Appl Phys, Vol 35, No 3, Pt 2, pp 842-843 (Mar 1964).
 The direct observation of Mn^{55} NMR absorption in $KMnF_3$ is reported.

87. Nathans, R., Menzinger, F. and Pickart, S. J.
Inelastic Magnetic Scattering from RbMnF_3 in the Neighborhood of Its Néel Point
 J Appl Phys, Vol 39, No 2, Pt 2, pp 1237-1238 (1 Feb 1968).
 A study of the inelastic neutron scattering in the vicinity of the Néel point reveals the presence of a diffusive-type peak, accompanied by spin-wave modes that persist above the transition temperature.

88. Ninio, F. and Keffer, F.
Simultaneous Parallel Pumping of Nuclear and Electronic Spin Waves
 Phys Rev, Vol 165, No 2, pp 735-750 (10 Jan 1968).
 The possibility is investigated of joint excitation of nuclear and electronic magnons by parallel pumping in RbMnF_3 .

89. Ogawa, S.
Antiferromagnetism in KMnF_3
 J Phys Soc Japan, Vol 14, No 8, p 1115 (Aug 1959).
 The susceptibility of KMnF_3 was measured as a function of temperature.

90. Okazaki, A. and Suemune, Y.
Electron Distribution in KMnF_3 , KFeF_3 , KCoF_3 , and KNiF_3
 J Phys Soc Japan, Vol 16, No 7, p 1474 (Jul 1961).
 The electron distributions in KMnF_3 , KFeF_3 , KCoF_3 , and KNiF_3 , determined by x-ray analysis, are reported.

91. Owen, J. and Taylor, D. R.
Zero-Point Spin Deviation in Antiferromagnets
 Phys Rev Lett, Vol 16, No 25, pp 1164-1169 (20 Jun 1966).
 The discrepancies between theoretical and experimentally observed zero-point spin deviations in CsMnF_3 , KMnF_3 , etc. are discussed.

92. Payne, R. E. and Forman, R. A.
Nuclear Magnetic Resonance in RbMnF_3
 Bull Amer Phys Soc, Vol 10, Ser 2, No 3, p 315 (1965).
 A NMR study of Rb^{88} , Rb^{87} , and F^{19} in paramagnetic RbMnF_3 is reported. This is the first report of measured shifts of the rubidium nuclear resonances of an alkali nucleus in paramagnetic XMF_3 perovskite compounds.

93. Payne, R. E., Forman, R. A. and Kahn, A. H.
Nuclear Magnetic Resonance in RbMnF_3
 J Chem Phys, Vol 42, No 11, pp 3806-3808 (1 June 1965).
 A NMR study of Rb^{85} , Rb^{87} , and F^{19} in paramagnetic RbMnF_3 is reported.

94. Pearson, J. J.
Theory of the Magnetic Anisotropy in KMnF_3
 Phys Rev, Vol 121, No 3, pp 695-702 (1 Feb 1961).
 A theoretical calculation is given of the magnetic anisotropy in KMnF_3 at room temperature and in its distorted structures at lower temperatures.

95. Perry, C. H. and Young, E. F.
Infrared Studies of Some Perovskite Fluorides. I. Fundamental Lattice Vibrations
 J Appl Phys, Vol 38, No 12, pp 4616-4624 (Nov 1967).
 The reflectance spectra of the cubic fluoride perovskites KMgF_3 , KMnF_3 , KCoF_3 , KNiF_3 , RbMnF_3 , and mixed crystals of $\text{K}(\text{Mg}_{0.8}\text{Ni}_{0.2})\text{F}_3$ and $\text{K}(\text{Mg}_{0.5}\text{Ni}_{0.5})\text{F}_3$ have been measured from 4000 cm^{-1} to 40 cm^{-1} at 300°K and 80°K .
96. Petrov, M. P., Smolenskii, G. S. and Syrnikov, P. P.
Nuclear Magnetic Resonance in RbMnF_3
 Fiz Tverd Tela, Vol 7, No 12, pp 3689-3690 (Dec 1965); Engl. transl. in Soviet Phys - Solid State, Vol 7, No 12, pp 2984-2985 (Jun 1966).
 F^{19} and Rb^{87} NMR have been observed in RbMnF_3 .
97. Pickart, S. J., Alperin, H. A. and Nathans, R.
Magnetic Structure of Binary Fluorides Containing Mn^{2+}
 J de Physique, Vol 25, No 5, pp 565-566 (May 1964) (In English).
 Spin orderings of the compounds XMnF_3 , where X was Na, Rb, Cs, and NH_4 , were studied by neutron diffraction.
98. Pickart, S. J., Collins, M. F. and Windsor, C. G.
Spin-Wave Dispersion in KMnF_3
 J Appl Phys, Vol 37, No 3, pp 1054-1055 (1 Mar 1966).
 The spin-wave dispersion curve of antiferromagnetic KMnF_3 at 4.2°K was measured by means of neutron inelastic scattering.
99. Pincus, P., DeGennes, P. G., Hartmann-Boutron, F. and Winter, J. M.
Dynamic Effects of the Suhl-Nakamura Interaction in Magnetic Materials
 Proc 8th Symp Magnetism and Magnetic Materials, Pittsburgh, Penn., 1962, ed. by I. S. Jacobs; J Appl Phys, Vol 34, No 4, Pt 2, pp 1036-1037 (Apr 1963).
 A discussion is given of nuclear spin waves and an attempt is made to explain the nonlinear effects observed in the NMR of KMnF_3 .
100. Pincus, P., Jaccarino, V., Hone, D. and Ngwe, T.
Suhl-Nakamura Interaction and Inhomogeneous Broadening
 Phys Lett, Vol 27A, No 1, pp 54-55 (20 May 1968).
 The effect of microscopic inhomogeneities on the Suhl-Nakamura indirect interaction between nuclear spins is shown to alter the NMR line shapes in the direction observed experimentally.
101. Portis, A. M.
Nuclear Relaxation in Magnetic Materials
 Proc Intl Conf Magnetism and Crystallography, Kyoto, 1961, ed. by T. Nagamiya; J Phys Soc Japan, Vol 17, Suppl B-1, pp 81-87 (Mar 1962).
 Studies of relaxation of nuclei in ferromagnetic and antiferromagnetic materials are reviewed.
102. Portis, A. M.
Nuclear Resonance in Magnetic Materials
 Nuovo Cimento Suppl, Vol 4, No 3, pp 603-606 (1966).
 The use of nuclear resonance in the study of magnetic materials is discussed.
103. Portis, A. M., Witt, G. L. and Heeger, A. J.
Excitation of Nuclear Magnetic Resonance Modes in Antiferromagnetic KMnF_3
 Proc 8th Symp Magnetism and Magnetic Materials, Pittsburgh, Penn., 1962, ed. by I. S. Jacobs; J Appl Phys, Vol 34, No 4, Pt 2, pp 1052-1053 (Apr 1963).
 The Mn^{55} nuclear resonance absorption in KMnF_3 was monitored through the observation of the AFMR at 4.2°K and below.

104. Prohofskey, E. W.
Magnetic Virtual Local Mode Condensation and Fluorescence Anomalies
 Phys Rev Lett, Vol 14, No 9, pp 302-305 (1 Mar 1965).
 This letter proposes a mechanism to explain the very abrupt increase in the frequency, intensity, and lifetime of fluorescent emission, with decreasing temperature, in antiferromagnetic Mn salts.

105. Prohofskey, E. W., Holloway, W. W., Jr. and Kestigian, M.
Magnetic Ordering and Fluorescence in Manganese Salts
 Proc 10th Conf Magnetism and Magnetic Materials, Minneapolis, Minn., 1964, ed. by I. S. Jacobs and E. G. Spencer; J Appl Phys, Vol 36, No 3, Pt 2, pp 1041-1042 (1965).
 The effect of antiferromagnetic ordering on the fluorescence of several Mn salts was observed. At temperatures below the Néel temperature the fluorescence moves to higher frequencies.

106. Rao, R. V. G., Das, C. D., Keer, H. V. and Biswas, A. B.
The Heat Capacities of Potassium Manganese Trifluoride
 Proc Phys Soc, Vol 81, Pt 1, No 519, pp 191-192 (1 Jan 1963).
 The results of measurements made using an isothermal calorimeter of the heat capacity over the range of temperature 78°K to 300°K are shown.

107. Rimmer, D. E.
Covalency Effects in Magnetic Salts by a Configurations Interaction Method
 Proc Intl Conf Magnetism, Nottingham, 1964, publ. by Institute of Physics and the Physical Society, London, pp 337-341 (1965).
 The configurations-interaction method of treating covalency has been used to calculate the fluorine spin density and cubic-crystal field parameter for KNiF_3 and KMnF_3 .

108. Satya Murthy, N. S., Venkataraman, G., Usha Deniz, K., Dasannacharya, B. A. and Iyengar, P. K.
Neutron Scattering by Paramagnetics
 "Inelastic Scattering of Neutrons," Proc Symp Inelastic Scattering of Neutrons, Bombay, 1964, International Atomic Energy Agency, Vienna, Vol 1, pp 433-442 (1965).
 Neutron scattering characteristics of MnO and KMnF_3 are reported.

109. Seiden, P. E., Freiser, M. J. and Teaney, D. T.
Field-Independent Parallel Antiferromagnetic Resonance in RbMnF_3
 Bull Amer Phys Soc, Ser 2, Vol 8, No 3, pp 213-214 (1963).
 This paper reports the observation of a field-independent antiferromagnetic resonance mode.

110. Shaltiel, D. and Fink, H. J.
Nuclear Antiferromagnetic Double Resonance in MnCO_3
 Proc 9th Conf Magnetism and Magnetic Materials, Atlantic City, N. J., 1963, ed. by J. A. Osborn; J Appl Phys, Vol 35, No 3, Pt 2, pp 848-849 (Mar 1964).
 The AFMR resonance was investigated as a function of UHF pumping power. Results are compared to similar experiments by other workers on KMnF_3 and CsMnF_3 .

111. Shulman, R. G. and Knox, K.
Nuclear Magnetic Resonance in KMnF_3
 Phys Rev, Vol 119, No 1, pp 94-101 (1 Jul 1960).
 The NMR of fluorine in KMnF_3 has been studied, and hyperfine interactions between the fluorine nucleus and the magnetic electrons measured.

112. Shionoya, S. and Hirano, M.
Spin-Wave Sideline in the Luminescence Spectra of Eu^{3+} Ion in KMnF_3
 Phys Lett, Vol 26A, No 11, pp 533-534 (22 Apr 1968).
 Spin-wave sidebands were observed in the luminescence spectra of Eu^{3+} doped in antiferromagnetic KMnF_3 crystals.
113. Simanek, E. and Tachiki, M.
A Note on the Theory of Covalency and Superexchange in Transition Metal Salts
 Phys Lett, Vol 21, No 6, pp 625-626 (1 Jul 1966).
 Reference is made to KNiF_3 and KMnF_3 .
114. Simanov, Yu. P., Batsanova, L.R. and Kovba, L.M.
X-Ray Investigation of Double Fluorides of Bivalent Manganese
 Zhur Neorgan. Khim, Vol 2, No 10, pp 2410-2415 (1957); Engl. transl. in J Inorgan. Chem USSR, Vol 2.
 NaMnF_3 , KMnF_3 , NH_4MnF_3 , RbMnF_3 and CsMnF_3 are studied by the powder x-ray method.
115. Stevenson, R.
Ultraviolet Absorption Spectra of MnF_2 , KMnF_3 , RbMnF_3 , and CsMnF_3
 AD 610 124, 20 pp (1964).
 Absorption spectra in the 3000-Å to 4000-Å region are presented for MnF_2 , KMnF_3 , RbMnF_3 , and CsMnF_3 at room temperature, liquid-air temperature, and liquid-helium temperature.
116. Stevenson, R.
Absorption Spectra of MnF_2 , KMnF_3 , RbMnF_3 , and CsMnF_3
 Canadian J Phys, Vol 43, No 10, pp 1732-1743 (Oct 1965).
 Absorption spectra in the 3000-Å to 4000-Å region are presented for MnF_2 , KMnF_3 , RbMnF_3 , and CsMnF_3 at room temperature, liquid-air temperature, and liquid-helium temperature.
117. Stevenson, R.
Absorption Spectra of MnF_2 and CsMnF_3
 AD 641 917, 15 pp (1966).
 Absorption spectra in the 3950-Å region of MnF_2 and CsMnF_3 at 4.2°K are presented.
118. Stevenson, R.
Compressions of CoCl_2 , FeCl_2 , NiCl_2 , EuS , MnF_2 , RbMnF_3 , and NiF_2
 Canadian J Phys, Vol 44, No 1, pp 281-283 (Jan 1966).
 Compressibilities in the pressure range 0 atm to 10^4 atm have been measured at 100°K, 80°K, and 4.2°K.
119. Stevenson, R.
Fine Structure in the Absorption Spectra of KMnF_3 and RbMnF_3
 Phys Rev, Vol 152, No 2, pp 531-535 (9 Dec 1966).
 Experimental studies are presented of the absorption spectra in the 3900-Å region of KMnF_3 and RbMnF_3 at 4.2°K.
120. Stevenson, R.
Magnon Sidebands in CsMnF_3
 Canadian J Phys, Vol 44, No 12, pp 3269-3270 (Dec 1966).
 Magnon sidebands similar to those observed in MnF_2 have been observed in CsMnF_3 .

121. Suemune, Y. and Ikawa, H.
Thermal Conductivity of KMnF_3 , KCoF_3 , KNiF_3 , and KZnF_3 Single Crystals
J Phys Soc Japan, Vol 19, No 9, pp 1686-1690 (Sep 1964).
 The thermal conductivities of single crystals of KMnF_3 , KCoF_3 , KNiF_3 , and KZnF_3 were measured at 15°K to 300°K by the absolute longitudinal heat-flow method.
122. Suhl, H.
Effective Nuclear Spin Interactions in Ferromagnets
Phys Rev, Vol 109, No 2, p 606 (15 Jan 1958).
 This paper presents the first discussion of the so-called Suhl-Nakamura interaction.
123. Suhl, H.
Nuclear Spin Interactions in Ferromagnetics and Antiferromagnets
J Phys Radium, Vol 20, No 2-3, pp 333-335 (Feb-Mar 1959) (In English).
 The influence of the hyperfine coupling on relaxation times is discussed.
124. Teaney, D. T.
Anomalous Dynamical Properties of the "Ideal" Antiferromagnet RbMnF_3
Bull Amer Phys Soc, Ser 2, Vol 13, No 2, p 164 (1968).
 Deviations from the ideal Heisenberg model for an antiferromagnet are discussed.
125. Teaney, D. T. and Freiser, M. J.
Antiferromagnetic Resonance in Cubic RbMnF_3
Proc 8th Symp Magnetism and Magnetic Materials, Pittsburgh, Penn., 1962,
ed. by I. S. Jacobs; J Appl Phys, Vol 34, No 4, Pt 2, p 1036 (Apr 1963).
 Observation of antiferromagnetic resonance in cubic RbMnF_3 is reported.
126. Teaney, D. T., Blackburn, J. S. and Stevenson, R. W. H.
Antiferromagnetic Resonance of Three XMnF_3 Single Crystals
Bull Amer Phys Soc, Ser 2, Vol 7, No 3, p 201 (23 Mar 1962).
 Magnetic resonances in RbMnF_3 , NaMnF_3 , and CsMnF_3 were investigated as functions of temperature.
127. Teaney, D. T., Freiser, M. J. and Stevenson, R. W. H.
Discovery of a Simple Cubic Antiferromagnet: Antiferromagnetic Resonance in RbMnF_3
Phys Rev Lett, Vol 9, No 5, pp 212-214 (1 Sep 1962).
 Antiferromagnetic resonance in RbMnF_3 is reported; measurements were performed at a frequency of 23.285 GHz.
128. Teaney, D. T., Moruzzi, V. L. and Argyle, B. E.
Critical Point of the Cubic Antiferromagnetic RbMnF_3
Proc 11th Ann Conf Magnetism and Magnetic Materials, San Francisco, Calif., 1965,
ed. by E. G. Spencer and J. S. Kouvel; J Appl Phys, Vol 37, No 3,
pp 1122-1123 (1 Mar 1966).
 The critical region of RbMnF_3 was studied by use of specific heat, x-ray, and strain-gage measurements.
129. Turov, E. A. and Kuleev, V. G.
Coupled Oscillations of Electronic and Nuclear Spins in Antiferromagnets
Zhur Eksper Teor Fiz, Vol 49, No 1, pp 248-256 (Jul 1965); Engl. transl.
in Soviet Phys. - JETP, Vol 22, No 1, pp 176-181 (Jan 1966).
 Coupled electron-nuclear spin resonance modes in antiferromagnets are analyzed.

130. Turov, E. A. and Timofeev, A. I.
Nuclear Magnetoacoustic Resonance in Ferro- and Antiferromagnetics
Fiz Metal Metalloved, Vol 24, No 5, pp 769-781 (1967); Engl. transl. in
preparation.

A theoretical discussion is given of the interactions of the
nuclear spins with lattice oscillations, of effective fields, and
of ultrasound absorption as studied via nuclear magnetoacoustic
resonance.
131. Walker, M. B.
Exchange Interactions from Nuclear Magnetic Resonance Linewidths
Proc Phys Soc, Vol 87, Pt 1, No 555, pp 45-48 (1 Jan 1966).

The exchange interactions in RbMnF_3 , and KMnF_3 , and MnF_2 have
been deduced from F^{19} NMR linewidth measurements.
132. Walker, M. B. and Stevenson, R. W. H.
Nuclear Magnetic Resonance in RbMnF_3
Proc Phys Soc, Vol 87, Pt 1, No 555, pp 35-43 (1 Jan 1966).

The F^{19} , Rb^{85} , and Rb^{87} NMR's observed at room temperature
in single crystals of RbMnF_3 are described.
133. Waring, R. K., Jr.
Magnetic Domains
Chapter 12 in "Magnetism and Magnetic Materials Digest," Vol 2, ed. by
C. Warren Haas and H. S. Jarrett, pp 149-162, Academic Press, New York
(1966).

Investigations of static domain patterns are reviewed.
134. Welsh, L. B.
Properties of the Mn^{55} Nuclear-Magnetic-Resonance Modes in CsMnF_3
Phys Rev, Vol 156, No 2, pp 370-382 (10 Apr 1967).

A new NMR mode of the Mn^{55} nuclei in the hexagonal antiferro-
magnet CsMnF_3 has been observed directly in the frequency range
673 MHz to 676 MHz.
135. Windsor, C. G.
Neutron Scattering from Paramagnetic RbMnF_3
Proc Phys Soc, Vol 89, Pt 4, No 566, pp 825-831 (1 Dec 1966).

The energy distribution of neutrons scattered from polycrystal-
line RbMnF_3 at room temperature ($T/T_N = 2.8$) has been meas-
ured using time-of-flight analysis.
136. Windsor, C. G.
Spin Correlations in a Classical Heisenberg Paramagnet
Proc Phys Soc, Vol 91, Pt 2, pp 353-355 (Jun 1967).

The neutron scattering from a polycrystalline sample of RbMnF_3
is predicted and compared with experiment.
137. Windsor, C. G. and Stevenson, R. W. H.
Spin Waves in RbMnF_3
Proc Phys Soc, Vol 87, Pt 2, No 556, pp 501-504 (1 Feb 1966).

The spin-wave dispersion relations for the simple cubic anti-
ferromagnet RbMnF_3 have been determined at 4.2°K by inelastic
neutron scattering.

138. Witt, G. L.

Nuclear and Electronic Resonance Modes in Antiferromagnets

Ph. D. Dissertation, University of Calif., Berkeley (1964); University Microfilms Order No 64-13, 22.

Electron-nuclear double resonance has been observed in KMnF_3 and CsMnF_3 .

139. Witt, G. L. and Portis, A. M.

Excitation of Coupled Nuclear Antiferromagnetic Resonance Modes

Phys Rev, Vol 136, No 5A, pp A1316-A1320 (30 Nov 1964).

The observation of ENDOR in KMnF_3 and CsMnF_3 is reported.

140. Witt, G. L. and Portis, A. M.

Nuclear Magnetic Resonance Modes in Magnetic Materials. II. Experiment

Phys Rev, Vol 135, No 6A, pp A1616-A1618 (14 Sep 1964).

Electron-nuclear double resonance in KMnF_3 is reported.
Nuclear absorption was observed for a range of frequencies
570 MHz to 685 MHz.

141. Young, E. F. and Perry, C. H.

Infrared Studies of Some Perovskite Fluorides. II. Multiphonon Spectra

J Appl Phys, Vol 38, No 12, pp 4624-4628 (Nov 1967).

The transmission spectra of thin single-crystal samples of KMgF_3 , KMnF_3 , KCoF_3 , KNiF_3 , KZnF_3 and $\text{K}(\text{Mg}_{0.5}\text{Ni}_{0.5})\text{F}_3$ have been investigated from 2.5μ to 400μ at 300°K and 85°K .

142. Zhogolev, D. A.

Shift of the NMR Frequencies in Paramagnetic TlMnF_3

Fiz Tverd Tela, Vol 8, No 9, pp 2798-2800 (Sep 1966); Engl. transl. in Soviet Phys - Solid State, Vol 8, No 9, pp 2237-2238 (Mar 1967).

The chemical shifts of Tl^{203} , Tl^{205} , and F^{19} in TlMnF_3 have been calculated.

DOCUMENT CONTROL DATA - R&D		
(Security classification of title, body of abstract and indexing annotation must be entered when the overall report is classified)		
1. ORIGINATING ACTIVITY (Corporate author) Lincoln Laboratory, M.I.T.	2a. REPORT SECURITY CLASSIFICATION Unclassified	
	2b. GROUP None	
3. REPORT TITLE Properties of the Compounds RbMnF ₃ , KMnF ₃ and CsMnF ₃		
4. DESCRIPTIVE NOTES (Type of report and inclusive dates) Lincoln Laboratory Library 33rd Reference Bibliography		
5. AUTHOR(S) (Last name, first name, initial) Friebely, Joan S. and Ince, William J.		
6. REPORT DATE 8 August 1968	7a. TOTAL NO. OF PAGES 26	7b. NO. OF REFS 142
8a. CONTRACT OR GRANT NO. AF 19(628)-5167	9a. ORIGINATOR'S REPORT NUMBER(S) 33rd Reference Bibliography	
b. PROJECT NO.		
c.	9b. OTHER REPORT NO(S) (Any other numbers that may be assigned this report)	
d.	ESD-TR-68-240	
10. AVAILABILITY/LIMITATION NOTICES This document has been approved for public release and sale; its distribution is unlimited.		
11. SUPPLEMENTARY NOTES None	12. SPONSORING MILITARY ACTIVITY Air Force Systems Command, USAF	
13. ABSTRACT This bibliography lists and briefly annotates publications related to the double fluorides XMnF ₃ , where X is K, Rb or Cs. References are listed alphabetically by author, and there is a subject outline and index. The period covered by the search is 1957 to July 1968.		
14. KEY WORDS nuclear magnetic resonance antiferromagnets		
perovskites RbMnF ₃		
CsMnF ₃ KMnF ₃		

