


SAM-TR-69-1

AD 691490

A FURTHER SURVEY OF COMPOUNDS FOR RADIATION PROTECTION


VIVIAN PLZAK, M.S.
JOHN DOULL, Ph. D., M.D.



USAF School of Aerospace Medicine
Aerospace Medical Division (AFSC)
Brooks Air Force Base, Texas

February 1969

This document has been approved for public release and sale;
its distribution is unlimited.



Qualified requesters may obtain copies of this report from DDC. Orders will be expedited if placed through the librarian or other person designated to request documents from DDC.

When U. S. Government drawings, specifications, or other data are used for any purpose other than a definitely related Government procurement operation, the Government thereby incurs no responsibility nor any obligation whatsoever; and the fact that the Government may have formulated, furnished, or in any way supplied the said drawings, specifications, or other data is not to be regarded by implication or otherwise, as in any manner licensing the holder or any other person or corporation, or conveying any rights or permission to manufacture, use, or sell any patented invention that may in any way be related thereto.

A FURTHER SURVEY OF COMPOUNDS FOR RADIATION PROTECTION

VIVIAN PLZAK, M.S.

JOHN DOULL, Ph. D., M.D.


This document has been approved for public release and sale;
its distribution is unlimited.

FOREWORD

This report was prepared at the USAF Radiation Laboratory of the University of Chicago and represents a summary of unpublished data. It is a sequel to "A Survey of Compounds for Radiation Protection" published in April 1962 as SAM-TR-62-29 by the USAF School of Aerospace Medicine. This compilation includes work accomplished from January 1961 through April 1965 under task No. 775703 and contract No. 41(609)-2977. The contract monitor was Lieutenant Colonel George S. Melville, Jr., Radiobiology Division, USAF School of Aerospace Medicine. The report was received for publication on 1 November 1968.

The animals involved in this study were maintained in accordance with the "Guide for Laboratory Animal Facilities and Care" as published by the National Academy of Sciences-National Research Council.

This report has been reviewed and is approved.



GEORGE E. SCHAFER
Colonel, USAF, MC
Commander

ABSTRACT

This report summarizes the results obtained with 617 compounds tested for their radioprotective activity in adult male mice irradiated with a control-demonstrated LD₉₉, of 800 R (250 kvp) x-rays. A compound was considered to exhibit radioprotective activity if it (1) permitted any of the treated mice to survive for 30 days after the otherwise lethal whole-body x-irradiation or (2) increased the median survival time of treated animals by 5 days or more beyond the median survival time of the untreated control mice (9 ± 3 days).

Of the 617 compounds tested, 245 successfully passed one or both of the stated specifications. Additionally, data are offered to allow comparisons of chemically related groups for structure-activity relationships, and to indicate the types of structures which offer the greatest promise as a source of more effective and less toxic radioprotective agents.

CONTENTS

	<i>Page</i>
INTRODUCTION	1
EXPERIMENTAL METHODS	1
RESULTS	2
Alcohols and thiols	5
Ethers	9
Ketones and phenones	11
Acids and thioacids	13
Esters	17
Amides and thioamides	19
Carbamates and thiocarbamates	23
Ureas and thioureas	25
Amines	29
Guanidines and nitriles	33
Hydrazines	37
Thiophenes, oxacyclopentanes, azacyclopentanes, and azacyclohexanes	39
Oxazoles and oxazines	45
Thiazoles and thiazines	47
Diazoles and diazines	49
Triazoles and dioxanes	55
Miscellaneous	57
INDEX	59

A FURTHER SURVEY OF COMPOUNDS FOR RADIATION PROTECTION

I. INTRODUCTION

The purpose of the radiation screening program of the University of Chicago was to find chemical or biologic agents which exhibited prophylactic or therapeutic effectiveness against radiation injury. Testing the ability of such agents to prolong survival or to reduce lethality in x-irradiated mice was one phase of our efforts to find practical methods to modify radiation injury in animals. Other phases of the program included: (1) the investigation of the physiologic and biochemical mechanisms responsible for the toxic and protective effects of radioprotective agents in mice and other species; (2) the evaluation of the efficacy of such agents against chronic radiation exposure and against exposure to other types of ionizing radiation; and (3) the ability of various environmental and pharmacologic factors to modify either the toxic or the protective properties of single agents or of combinations of radioprotective agents.

A previous survey of approximately 1,200 compounds was published in April 1962. The present report summarizes the results that were obtained with approximately 600 compounds tested from January 1961 through April 1965. It has a twofold purpose: (1) to facilitate the comparison of chemically related groups for structure activity relationships, and (2) to indicate the types of structures which offer the greatest promise as a source of more effective and less toxic radioprotective agents. Initially, the compounds used in the testing program were selected with the idea of obtaining examples of as many different chemical types as possible. Since more information has

been accumulated on potentially radioprotective agents of a rather large number of chemical types, it is now possible to depend less on the survey approach of selection and more on trying to obtain compounds designed to provide more specific data on the structure-activity relationships within each chemical type.

II. EXPERIMENTAL METHODS

Adult male mice (6 to 8 weeks old) from Carworth Farms (CF_1) were used in the studies. The average weight was from 20 to 25 gm. The mice were housed in an air-conditioned room (75° to 80° F.) and were provided with Rockland mouse pellets and water ad libitum. Both control and treated animals were selected at random from a single shipment so that their age and physical condition would be comparable. All the mice were kept under observation for at least one week before they were used. Those that appeared to be unhealthy or failed to gain weight at a normal rate were removed for autopsy.

To evaluate the radioprotective activity of the compounds, the mice were divided into groups of 10 and injected intraperitoneally 10 to 15 minutes prior to the administration of a lethal (800 R) dose of whole-body x-irradiation. Each compound was tested for protective effects at two or more dosage levels, one of which was close to the maximum tolerated dosage level for that particular compound. Distilled water was used as the vehicle whenever possible and the concentration of the solution was adjusted so that no animal received more than 2% of its body weight with each injection.

When necessary, the pH was adjusted to approximately 7.0 with either 1N HCl or 1N NaOH. Compounds which were insoluble in water or in propylene glycol (PG) were dissolved in cottonseed oil or suspended in a 0.5% solution of carboxymethylcellulose. Control animals were injected with a comparable amount of the vehicle used to prepare the solution and were irradiated simultaneously with the treated groups. The loss of weight and the mortality in both groups were recorded daily for 30 days after the x-ray exposure or until all of the mice were dead.

The radiation exposures were given as single whole-body 250 kvp, 15 ma. x-ray exposures using either a G.E. Maximar or a Keleket x-ray therapy unit. The added filtration consisted of 0.25 mm. of copper and 1.0 mm. of aluminum, and the target-skin distance was 75 cm. The dose rate in air was determined prior to each radiation period, by use of Victoreen ionization thimbles (100 R); it was found to average between 40 and 43 R per minute. The mice were placed in individual plastic tubes (50-ml. centrifuge tubes provided with numerous air holes) which, in turn, were positioned radially on a rotating turntable to equalize the dose of x-ray. The environmental temperature was maintained at 75° F. during the irradiation period to minimize the effect of variations in the environmental stress in both the control and treated groups.

III. RESULTS

Preliminary toxicity tests were conducted for each compound before the radiation studies to ascertain the maximum tolerated dose which could be given the animals without causing drug lethality. For these studies, small groups of mice were injected intraperitoneally with increasing dosage levels of each compound and the mortality observed for a week. The approximate LD₅₀ (7 days) thus obtained was used as a basis for selecting the dosage levels to be used for the radiation studies.

Doses exceeding 1,000 mg. kg. were not used, since the limited size of the sample and the disadvantages of injecting an increased

amount of vehicle made the larger doses impractical. The approximate LD₅₀ in milligrams per kilogram for each of the compounds tested is included in the tables which comprise the remaining part of this report.

The x-ray dosage level employed (800 R) uniformly caused a mortality greater than 99% within two weeks after the exposure under our experimental conditions. Radiation deaths usually began on the fifth or sixth day after the x-ray exposure and the median survival time (ST₅₀) of untreated animals was 9 ± 3 days. A compound was considered to exhibit radioprotective activity if it increased the median survival time by 5 days or more, or if it permitted any of the treated animals to survive for 30 days after the otherwise lethal whole-body x-irradiation.

Since all of these studies were carried out by use of a single dose of x-irradiation (800 R), the results do not provide a precise quantitation of the radioprotective effect of each compound tested. However, the protective effect or dose reduction factor (DRF) resulting from the preirradiation administration of a compound can be estimated on the basis of the x-ray dosage level which would be required to produce a comparable mortality at 30 days after the irradiation exposure. The LD₅₀ (30-day) value for the adult CF₁ male mice used in these studies was 542 ± 18 R. The estimated DRF values are given in table I.

TABLE I
Estimated dose reduction factors based on 30-day mortality data

Percent survival*	DRF†
10 to 20	1.3
30 to 40	1.4
50 to 60	1.5
70 to 80	1.6
90 to 100	1.7

*Percent of treated animals surviving at 30 days after 800 R x-irradiation.

†Admired radiation dose divided by effective radiation dose.

In the tables which comprise the remaining part of the report the compounds are listed in order of increasing complexity within each of the chemical types. The name of each compound is listed with the vehicle used.¹ In the radiation studies the change in ST₅₀ indicates

the increase or decrease (in days) of the median survival time of the treated animals when compared to that of the simultaneously irradiated control (vehicle only) groups.

¹The symbol representing vehicle (e.g., HOH) appears in parentheses below the name of the compound.

ALCOHOLS AND THIOLS

Compound and vehicle used for toxicity and radiation tests	Toxicity	Radiation studies		
	Approx. LD ₅₀ (mg./kg.)	Dose (mg./kg.)	Change in ST ₅₀ (days)	Mortality at 30 days
2-Methylaminoethanol (HOH)	250	200 100	-4 0	10/10 10/10
1-Amino-2-propanol (HOH)	400	300 100	-10 -4	10/10 10/10
Acetone sodium bisulfite (HOH)	> 1000	1000 500	+2 -3	10/10 10/10
Glyoxal (30% in water) (HOH)	750	500 300	-5 +3	9/10 5/10
Propylene glycol (1,2-propanediol) (100%)	17500	3000 5000 6250 7500 8750 10000 12500	+2 0 +2 -3 -4 -3 -7	10/10 8/10 9/10 10/10 10/10 10/10 10/10
2-Isobutylaminoethanol (HOH)	250	200 100	-3 -2	10/10 10/10
2,3-Butanediol (HOH)	> 1000	1000 500	-1 -2	10/10 10/10
Batyl alcohol (PG)	750	500 250	-1 -2	10/10 10/10
3-(N-Methylisopropylamino)-3-methyl-2-butanol hydrochloride (HOH)	250	200 100 50	0 -5 -8	10/10 10/10 10/10
Calcium gluconate (HOH)		10	0	10/10
2-Amino-1-phenyl-1-ethanol (HOH)	250	200 100	-5 -9	10/10 10/10
2,6-Dimethylphenol (PG)	150	100 50	0 -1	10/10 10/10
2,4-Di-tert-butylphenol (PG)	25	10 5	-3 -2	10/10 10/10
N-Propyl-p-hydroxybenzoate (PG)	200	100 50	-1 0	10/10 10/10

Sodium-p-hydroxyphenylarsonate (HOH)	1000	500 200	-6 +2	10/10 10/10
p-Methoxyphenol (PG)	250	200 100	-4 0	9 10 10/10
2-Methoxy-4-n-pr p'-phenol (PG)	150	100 50	-9 +2	10 10 7/10
m-Chloroacetylcatechol (HOH)	75	50 25	-6 +6	10 10 10/10
Pyrogallol (HOH)	400	300 100	-3 +2	10/10 10/10
Desoxypyridoxine hydrochloride (HOH)	150	100 50	-5 0	10 10 10/10
o-Aminophenol (PG)	200	100 50	-1 0	10 10 10/10
m-Aminophenol (PG)	150	100 50	0 -1	10/10 10/10
p-Nitrophenol (PG)	75	50 25 10	-2 -2 -2	10 10 10/10 10/10
p-Aminophenol hydrochloride (HOH)	750	500 300	-2 -4	10 10 10/10
4-Amino-3-methylphenol (PG)	200	100 50	-1 -2	10/10 10 10
p-Phenylphenol (PG)	150	100 50	-2 0	10 10 10/10
p-Hydroxydiphenyl (PG)	150	100 50	0 0	10 10 10/10
Naphthoresorcinol (PG)	200	100 50	0 0	8/10 10/10
o,o'-Biphenol (PG)	150	100 50	-1 -3	10 10 10/10
2,2'-Dipropyl-p,p'-biphenol (PG)	40	25 10	0 +1	10/10 10/10
4,4'-Di-tert-butyl-o,o'-biphenol (PG)	50	40 25	-6 -5	10/10 10/10
2,2'-Diallyl-p,p'-biphenol (PG)	100	50 25	-4 -2	10/10 9/10
p,p'-Isopropylidene-diphenol (PG)	150	100 50	-10 -9	10/10 10/10

4,4'-Isopropylidene-bis(2-tert-butylphenol) (PG)	40	25 10	-5 -6	10/10 10/10
p,p'-Oxydiphenol (PG)	150	100 50	-5 -4	10/10 10/10
p-Phenylazophenol (PG)	75	50 25	-3 -3	10/10 10/10
3-Methyl-4-phenylazophenol (PG)	25	10 5	+1 -1	10/10 10/10
Hydroquinone (HOH)	150	100 50	+3 0	8/10 10/10
2-Quinolinol (PG)	150	100 50	>+18 >+22	8/10 4/10
8-Quinolinol (PG)	50	25 10	+9 +1	5/10 9/10
2',5'-Salicyloxylidide, 3-nitro (PG)	25	10 5	+1 -1	10/10 10/10
2',6'-Salicyloxylidide, 3-nitro (PG)	40	25 10	-2 +6	10/10 7/10
o-Salicylotoluidide, 3,5'-dinitro (PG)	100	100 50	+5 +3	10/10 9/10
α -Benzoinoxime (PG)	150	100 50	-5 0	10/10 10/10
α -Thioglycerol (HOH)	400	300 100	0 +1	10/10 10/10
Thioglycolic acid - 98% (HOH)	150	100 50	-5 -1	10/10 10/10
Thiodiglycolic acid -- 98% (HOH)	250	200 100	-6 0	10/10 10/10
Thiomalic acid (HOH)	500	300 100	-4 +1	10/10 10/10
Methylmercaptoacetate (HOH)	400	100 50	+5 -2	10/10 10/10
1,2-Bis-(β -mercaptopethylaminomethyl)- cyclobutane (HOH)	250	200 100	>+18 +1	4/10 7/10
6-Methylthio-m-cresol (PG)	125	100 50	>+19 +1	3/10 9/10
4-Methylthio-m-cresol (PG)	150	100 50	>+19 -1	3/10 8/10

p-Chlorobenzenethiol (PG)	75	50 25	-4 +1	10/10 10/10
p-Dimethylaminobenzenethiol (PG)	150	100 50	-3 -2	8/10 9/10
1-Mercaptoanthraquinone (PG)	750	500 300	> +21 +2	4/10 10/10
1-Hydroxy-4-thiophenylanthraquinone (HOH and PG)	500	100 50	-2 +2	10/10 10/10
1-Amino-2-thiophenyl-4-hydroxyanthraquinone (PG and sesame oil)	200 750	500 300	-2 +1	10/10 10/10
Phenylmercaptodiaminoanthrarufin (PG and sesame oil)	150 300	300 200	-1 +1	10/10 10/10
1-Amino-2-tert-butylphenylmercapto-4-hydroxyanthraquinone (PG)	250	200 100	+1 +1	10/10 10/10
Thioflavine S (HOH)	400	300 100	-2 +1	10/10 10/10

ETHERS

Compound and vehicle used for toxicity and radiation tests	Toxicity		Radiation studies	
	Approx. LD ₅₀ (mg./kg.)	Dose (mg./kg.)	Change in S _T ₅₀ (days)	Mortality at 30 days
1-Methoxy-2-amino-2-methylpropane (HOH)	400	300 100	-7 0	10/10 10/10
Bis(2-nitroisobutyl)-2-ethylhexyl phosphate (HOH)	250	200 100	-1 +3	10/10 10/10
N-(3-Butoxypropyl)taurine, monosodium salt (HOH)	1000	1000 500	+4 +2	7/10 8/9
Cyclobutane-1,2-bis(methylammonium phosphate (HOH)	1000	1000 500	-3 +2	10/10 10/10
Cyclobutane-1,2-bis(methylammonium) dimonochloroacetate (HOH)	200	100 50 25	-4 -2 +3	10/10 10/10 9/10
β -Hydroxyethylbenzyl ether (PG)	250	200 100	0 +1	7/10 10/10
β -Hydroxyethyl ether of orthocresol (PG)	250	200 100	-3 +1	9/10 9/10
β -Hydroxyethyl ether of butylphenol (PG)	100	50 25	-1 -3	9/10 10/10
β -Hydroxyethyl ether of m-xylenol (PG)	150	100 50	-5 -1	10/10 10/10
p-Dimethoxybenzene (PG)	400	300 100	-7 +1	10/10 8/10
p-Nonylphenyl- β -dimethylaminoethyl ether (PG)	150	100 50	-6 -1	10/10 9/10
1,6-Bis(p-aminoethylphenoxy)hexane dihydrochloride (HOH)	100	50 25	-4 -3	10/10 7/10
1,6-Bis(p-methylaminomethylphenoxy) hexane dihydrochloride (HOH)	150	100 50	-2 0	9/10 8/10
Phenylmercurilauryl thioether (PG)	500	300 200	-3 0	9/10 9/10
2-Methyl-2-nitro-1-propyltosylate (PG)	250	200 100	+1 +3	10/10 10/10

Bis-[p-(α -diethylaminopropoxy)phenyl] sulfone (HOH)	75	50 25	-1 +1	10/10 10/10
Bis-p-phenyl methane sulfone (PG)	200	100 50	+1 -3	6/10 10/10
β -Naphthylethyl ether (PG)	100	50 25	-2 0	10/10 10/10
β -Hydroxyethyl ether of β -naphthol (PG)	150	100 50	-5 -1	9/10 7/10
β -Hydroxyethyl ether of orthophenylphenol (PG)	75	50 25	+2 +1	7/10 8/10

KETONES AND PHENONES

Compound and vehicle used for toxicity and radiation tests	Toxicity		Radiation studies	
	Approx. LD ₅₀ (mg./kg.)	Dose (mg./kg.)	Change in ST ₅₀ (days)	Mortality at 30 days
Methylallyl acetone (HOH)	1000	500 300	+1 -4	10/10 10/10
Acetylacetone (HOH)	750	500 300	-2 -6	10/10 10/10
p-Benzoyquinone (HOH)	7.5	5 1	-2 -2	10/10 10/10
2-Methylcyclohexanone (PG)	200	100 50	+1 -5	10/10 10/10
Pulegone (PG)	150	100 50	-1 +2	10/10 7/10
Acetone oxime (PG)	250	200 100	0 +2	10/10 10/10
2-Butanone oxime (PG)	1000	1000 500 250	-4 > +18 +2	10/10 3/10 10/10
Di(methylcyclopropyl)ketoxime (HOH)	250	200 100	-2 0	10/10 9/10
Cyclohexanone oxime (PG)	250	200 100	-2 0	10/10 10/10
Isonitrosopropiophenone (Lot No. FI-9) (PG)	250	200 100	0 -2	10/10 10/10
6-Amino-5-hydroxyimino-1,3-dimethyl hydrouracil (PG)	250	200 100	+1 -1	10/10 10/10
Pyridine-2-aldoxime (PG)	200	100 50	+3 +1	10/10 8/10
Pyridine-3-aldoxime (HOH)	200	100 50	+3 -1	10/10 10/10
Pyridine-4-aldoxime (PG)	150	100 50	+4 +6	10/10 9/10
Isonitrosopropiophenone (Lot No. EL-152) (PG)	200	100 50	+1 +8	7/10 8/10
Pyridine-2-aldoxime o-benzyl ether (PG)	100	50 25	+3 +3	10/10 10/10

2,3,4a,5,8,8a-Hexahydro-1,4-naphthoquinone (PG)	200	100 50	> +22 0	4/10 10/10
Methylnaphthoquinone (PG)	15	10 5	+4 +2	7/10 10/10
2-Methyl-1,4-naphthoquinone (Menadione) (PG)	50	40 25	-5 -2	10/10 10/10
7-Chloro-4-(3-octylaminopropyl)amino- quinoiline,1-oxide (PG)	40	25 10	-2 +2	10/10 10/10
Sodium β-naphthoquinone-4-sulfonate (HOH)	625	300 200	-4 0	10/10 10/10
2-Oxo-3-isobutyl-9,10-dimethoxy-1,3,4,6,7,11- β-hexahydro-2H-benzoguinolizine (PG)	250	200 100	+3 +5	10/10 10/10
N-1-Anthraquinonyl-2,3-dichloropropion- amide (PG)	200	100 50	-2 +1	10/10 10/10
1-Thiophenylanthraquinone (Sesame oil)	500	500 300	-1 +2	10/10 10/10
2-Thiophenylanthraquinone (PG)	150	100 50	-3 +3	10/10 10/10
1-Amino-2-thiophenylanthraquinone (PG)	> 1000	500 300	+2 +4	10/10 10/10
1-Amino-4-thiophenylanthraquinone (PG)	> 1000	500 300	0 +3	10/10 10/10
Dimethylsulfoxide (DMSO) (100%)	7500	6000 2500	> +22 +4	1/10 8/10
Tetramethyldiaminothiobenzophenone (PG)	250	200 100	+3 -1	10/10 10/10
2-Nitro-4-acetyl-2'-aminodiphenyl sulfide (PG)	150	100 50	+4 -5	10/10 10/10
1-Amino-2-benzylmercapto-4-hydroxy- anthraquinone (PG)	150	100 50	-1 +4	10/10 10/10

ACIDS AND THIOACIDS

Compound and vehicle used for toxicity and radiation tests	Toxicity		Radiation studies	
	Approx. LD ₅₀ (mg./kg.)	Dose (mg./kg.)	Change in ST ₅₀ (days)	Mortality at 30 days
Cacodylic acid (HOH)	750	500 200	-3 +4	10/10 6/10
Cyanoacetic acid (HOH)	200	100 50	-5 -3	10/10 10/10
Ethylenediaminetetraacetic acid (EDTA) (PG)	250	200 100	-1 +2	10/10 10/10
Bis(diaminoethyl ether)-N,N,N'-N'-tetraacetic acid (PG)	250	200 100	-4 +1	10/10 9/10
Ethylene glycol-bis-(aminoethyl ether) tetraacetic acid (PG)	150	300 200	-10 -1	10/10 10/10
Levulinic acid (HOH)	450	300 100	-10 >+22	10/10 4/9
Chrysanthemummonocarboxylic acid (PG)	150	100 50	-5 -1	10/10 10/10
3-Methylenecyclobutanecarboxylic acid (HOH + NaHCO ₃)	750	500 300 100	-4 0 -1	10/10 10/10 10/10
2-Aminocyclobutanecarboxylic acid hydrochloride (HOH)	>1000	1000 500 200	-4 -3 0	10/10 10/10 10/10
trans-1,2-Cyclobutanedicarboxylic acid (HOH)	400	300 200	0 -1	9/10 10/10
3-Methyl-2-cyclobutene-1,2-dicarboxylic acid (HOH)	200	100 50	0 0	9/10 10/10
1,2-Diaminocyclohexanetetraacetic acid (PG)	150	100 50	-3 -2	10/10 9/10
Trisodium-2-(hydroxycyclohexyl)ethylenediamine triacetate (PG)	400	300 200	-4 +3	10/10 9/10
α -Imino-p-methoxybenzylmercaptoacetic acid (PG)	250	200 100	-7 0	9/10 10/10

Indole-3-acetic acid (PG)	150	100 50	+1 +3	10/10 10/10
N-(2,3-Xylyl)anthranilic acid: mefenamic acid (PG)	200	100 50	+1 -2	10/10 10/10
N-(3,5-Dichloro-p-tolyl)anthranilic acid (HOH + NaHCO ₃)	100	50 25	0 -1	7/10 8/10
N-(a,a,a-Trifluoro-m-tolyl)anthranilic acid; flufenamic acid (PG)	150	50 25	+3 +1	10/10 10/10
N-(3-Sulfamoyl-2,6-xylyl)anthranilic acid (HOH + NaHCO ₃)	400	300 200	+3 +2	7/8 10/10
Ethylenediamine-di(o-hydroxyphenyl)acetic acid (PG)	350	100 50	-1 -1	10/10 10/10
N-(8-Amino-1-antraquinonyl)anthranilic acid (HOH + NaHCO ₃)	750	500 300	-5 -8	9/10 8/10
3-(2-Aminoethylthio)propionic acid (HOH)	>1000	500 300	-5 -5	10/10 10/10
meso-2,3-Dimercaptosuccinic acid (HOH)	500	300 200	-7 -1	10/10 10/10
5,8,11,14-Eicosatetraenoic acid (PG)	75	50 25	-5 -1	10/10 10/10
2-Brom-D-lysergic acid diethylamide (HOH)	75	25 10	-2 0	10/10 10/10
2-Aminoethanethiosulfuric acid (HOH)	400	300 100	+3 +3	7/10 9/10
3-Chloro-1-propanesulfonic acid, monosodium salt (HOH)	750	500 200	+6 +2	6/10 9/10
2-Carboxyphenylthioglycolic acid (HOH)	250	200 100	-8 -3	9/10 9/10
4-Chloro-2-methylphenylthioglycolic acid (PG)	150	100 50	-11 -2	10/10 10/10
Alanine: 2-methyl- (HOH)	750	500 300	-1 -1	10/10 10/10
DL-β-(3,4-Dihydroxyphenyl)alanine (HOH)	500	300 200	-1 0	10/10 10/10

L-Cysteine hydrochloride (HOH)	1250	1000 500	-6 -5	10/10 10/10
N-Acetyl-n-cysteine (HOH)	400	300 200	-4 0	10/10 10/10
(Carboxymethyl)trimethylammonium-N- acetyl-L-cysteinate (HOH)	>1000	1000 500	0 0	10/10 10/10
Glutathione (HOH)		1000	+1	10/10
Glutathione ascorbate, 63.57%-36.43% (HOH)	>100	100 50	+2 +1	9/10 9/10
Glutathione ascorbate, 1:10 (HOH)	>100	100 50	+2 -1	10/10 10/10
Glutathione ascorbate with calcium phytate (CaMg inositol hexaphosphate) (HOH)	>100	100 50	+2 -2	10/10 10/10
Glutathione (glutamyl-cysteinyl-glycine, in molecular relation) (HOH)	>100	100 50	+2 0	9/10 10/10
N-Acetyl-DL-penicillamine (PG)	400	300 200	-4 +1	10/10 10/10

This Document Contains
Missing Page/s That Are
Unavailable In The
Original Document

OR ARE
Blank pg.
that have
been removed

**BEST
AVAILABLE COPY**

ESTERS

Compound and vehicle used for toxicity and radiation tests	Toxicity		Radiation studies	
	Approx. LD ₅₀ (mg./kg.)	Dose (mg./kg.)	Change in ST ₅₀ (days)	Mortality at 30 days
Methyl chloroformate (HOH)	40	25 10	0 -5	10/10 10/10
Acetic acid: 2-methoxy-1-methyl ethyl ester (HOH)	750	500 200	+1 -5	10/10 10/10
3-(Amidino)propanoic acid, ethyl ester, monohydrochloride (HOH)	250	200 100	+2 -2	10/10 10/10
Nitriloacetic acid trisodium salt monohydrate (HOH)	500	300 100	-3 +2	10/10 10/10
Dipotassium methanearsonate (HOH)	750	500 200	-4 +1	10/10 10/10
Monoammonium methanearsonate (HOH)	>1000	1000 500	-3 -3	10/10 10/10
Sodium cacodylate (HOH)	>1000	1000 500	-5 -4	10/10 10/10
Calcium methanearsonate (HOH)	500	300 100	-3 +1	10/10 10/10
Magnesium cacodylate (HOH)	750	500 200	-4 -3	10/10 10/10
Magnesium methanearsonate (HOH)	>1000	1000 500	0 -1	10/10 10/10
Dimerchrous methanearsonate (HOH)	75	50 10	-11 -7	10/10 10/10
Disodium methanearsonate (HOH)	>1000	1000 500	-5 -2	10/10 S/10
Copper methanearsonate (HOH + NaHCO ₃)	15	10 5	-6 +2	10/10 9/10
Dibutyl fumarate (PG)	250	200 100	+1 +2	10/10 10/10
Di-2-ethylhexyl fumarate (PG)	250	200 100	+1 +1	10/10 10/10
Dibutyl maleate (PG)	150	100 50	-1 +2	10/10 10/10

Di-2-ethylhexyl maleate (PG)	>1000	1000 500	+1 0	10/10 10/10
Ethyl-1-methyl propyl diethyl malonate (PG)	200	100 50	+1 -2	10/10 10 10
Dimethyl-1,2-cyclobutanedicarboxylate (HOH)	>1000	1500 1000 500	+2 -2 0	9/10 8 10 10/10
cis-1,2-Diisopropylcyclobutanedicarboxylate (PG)	750	500 300 100	-4 -4 +5	10 10 10/10 5/10
trans-1,2-Diisopropylcyclobutanedicarboxylate (PG)	250	200 100	-3 +2	10/10 10/10
Diisodecyl-1,2-cyclobutanedicarboxylate (PG)	750	500 300	0 0	10/10 9 10
Hexamethylenediammonium cyclobutane-1,2-dicarboxylate (HOH)	>1000	1000 500	0 +2	10/10 10/10
Cyclobutane-1,2-bis(methylammonium) adipate (HOH)	>1000	1000 500	-3 -1	10 10 10/10
Bis-(2,4,5-trichlorophenyl)-1,2-cyclobutane dicarboxylate (PG)	250	200 100	0 +2	10 10 10/10
Methyl-p-nitrobenzoate (PG)	200	100 50	0 -2	9/10 10/10
Ethyl-p-nitrobenzoate (PG)	250	200 100	0 +2	10 10 10/10
Ethyl-N-(p-nitrobenzenesulfonyl)urethane (PG)	200	100 50	-3 +16	9/10 5/10
Ethyl-N-o-toluenesulfonylurethane (PG)	250	200 100	+6 +1	8/10 10/10

AMIDES AND THIOAMIDES

Compound and vehicle used for toxicity and radiation tests	Toxicity Approx. LD ₅₀ (mg./kg.)	Radiation studies		
		Dose (mg./kg.)	Change in ST ₅₀ (days)	Mortality at 30 days
N,N-Dimethylacetamide (HOH)	>1000	1000 500	+1 0	10/10 10/10
Acetamide: 2-(2-hydroxyethoxy)- (PG)	100	50 25	+2 -6	10/10 10/10
α -Cyanoacetamide (HOH)	>1000	1000 500	-3 0	10/10 10/10
α -Methylmercapto- α , α -diethylacetamide (PG)	750	500 200	-5 +2	10/10 10/10
α -Ethylmercaptoisobutyramide (PG)	>1000	1000 500	+5 -3	6/10 10/10
α -Allylmercaptoisobutyramide (PG)	250	200 100	0 +1	10/10 10/10
α -Allylmercapto- α , α -diethylacetamide (PG)	400	250 100	-2 0	10/10 10/10
α -Isopropylmercaptoisobutyramide (PG)	750	500 200	-1 +3	10/10 10/10
N-Acryloyltaurine, monosodium salt (HOH)	>1000	1000 500	0 +2	10/10 10/10
N-Methyl- α -crotylmercaptoisobutyramide (PG)	750	500 300	-7 -1	10/10 10/10
α -Propargylmercapto- α , α -diethylacetamide (PG)	250	200 100	-1 -3	9/10 10/10
α -Butylmercaptoisobutyramide (PG)	250	200 100	-1 -1	10/10 10/10
2,2'-Thio-bis-acetamide (PG)	625	500 300	+2 +3	10/10 10/10
trans-1,2-Cyclobutanedicarboxamide (HOH)	>1000	1000 500	+1 +1	10/10 10/10
N,N'-di-tert-Butylcyclobutane-1,2-dicarbox- amide (PG)	250	200 100	-2 +2	10/10 10/10
N,N'-Bis(1,1,3,3-tetramethylbutyl)-1,2- cyclobutanedicarboxamide (PG)	250	200 100	0 -4	10/10 10/10

N,N'-(α -Naphthyl)-1,2-cyclobutanedicarboxamide (PG)	275	250 200 100	-6 +2 -4	10/10 6/10 10/10
N,N'-(β -Naphthyl)-1,2-cyclobutanedicarboxamide (PG)	>1000	500 300	+2 -5	6/10 10/10
α -Hydroxy-N,N-diethylbenzamide (PG)	200	100 50	-2 -3	10/10 10/10
p-Toluenesulfanamide (PG)	250	200 100	-2 +3	10/10 10/10
Carbutamide (PG)	250	200 100	-1 +3	10/10 8/10
4',3-Dinitrosalicylanilide (PG)	25	10 5	-2 +3	10/10 10/10
4'-Chloro-3-nitrosalicylanilide (PG)	15	10 5	0 +2	9/10 10/10
4'-Chloro-5-nitrosalicylanilide (PG)	40	25 10	0 +4	10/10 9/10
3'-Chloro-3-nitrosalicylanilide (PG)	8	5 1	+3 0	8/10 10/10
3'-Chloro-5-nitrosalicylanilide (PG)	15	10 5	+1 +3	9/10 10/10
4'-Fluoro-3-nitrosalicylanilide (PG)	15	10 5	-7 +6	10/10 10/10
4'-Bromo-3-nitrosalicylanilide (PG)	15	10 5	0 +5	10/10 10/10
3'-Iodo-3-nitrosalicylanilide (PG)	15	10 5	+3 0	10/10 10/10
4'-Iodo-3-nitrosalicylanilide (PG)	15	10 5	-3 +4	10/10 10/10
3',4'-Dichloro-3-nitrosalicylanilide (PG)	15	10 5	+3 +8	10/10 8/10
2'-Chloro-5'-trifluoromethyl-3-nitrosalicylanilide (PG)	8	5 1	0 +2	10/10 10/10
N-p-Toluenesulfonyl, isoquinuclidylformamide (PG)	150	100 50	-4 -7	8/10 10/10

N,N-Bis(2-cyanoethyl)-N-4-hydroxy-1-anthraquinonylsulfonilamide (PG)	400	300 100	+2 -3	8/10 10/10
Bis-(2-amino-4-sulfonamidophenyl) disulfide (PG)	150	100 50	-4 -1	10/10 10/10
N,N'-Bis(2-methoxyethyl)dithioxamide (PG)	400	250 100	0 +3	10/10 9/10
N,N'-Di-tert-butyl dithioxamide (PG)	40	25 10	-2 +2	10/10 10/10
N,N'-Dioctadecyl dithioxamide (PG)	150	100 50	0 +7	10/10 6/10
Bis-didodecyl dithioxamide (PG)	250	200 100	-4 0	10/10 10/10
Thiopyrazinamide (PG)	150	100 50	0 -2	10/10 10/10
p-Aminothiobenzamide (PG)	75	50 25	0 -3	10/10 10/10
N,N-Dimethyl- α,α -diphenylthioacetamide (PG)	250	200 100	-5 -1	10/10 10/10
N,N'-Methylene-bis-(thioisonicotinamide; hydrate (PG)	250	200 100	-6 -1	10/10 10/10

This Document Contains
Missing Page/s That Are
Unavailable In The
Original Document

OR ARE
Blank pg.
that have
been removed

**BEST
AVAILABLE COPY**

CARBAMATES AND THIOCARBAMATES

Compound and vehicle used for toxicity and radiation tests	Toxicity		Radiation studies	
	Approx. LD ₅₀ (mg./kg.)	Dose (mg./kg.)	Change in ST ₅₀ (days)	Mortality at 30 days
Dimethylcarbamyl chloride (HOH)	>1000	1000 500	-2 +2	10/10 10/10
Diethylcarbamyl chloride (HOH)	750	500 300	-1 +1	10/10 10/10
Ethyl-N,N-di-N-butylcarbamate (PG)	150	100 50	-8 -7	10/10 10/10
Ethyl-N,N-butylcarbamate (PG)	250	200 100	-4 0	10/10 8/16
Carbamic acid: 2-aminoethyl ester (HOH)	>1000	1000 500	+3 -1	10/10 10/10
N,N'-Dicarbamylformamidine (PG)	250	200 100	+1 -2	8/10 10/10
2-Nitro-2-methyl-1,3-propanediolcarbamate (HOH)	>1000	1000 500	-3 -3	10/10 10/10
1,2-C ₃ -obutanedimethylcarbamate (PG)	>1000	1000 500	0 +3	9/10 10/10
1-Methylammonium-2-carbamylmethyl cyclobutane (HOH)	750	500 300	-3 +2	9/10 10/10
γ-Phenylpropylcarbamate (PG)	150	100 50	-4 0	10/10 10/10
Isopropyl-N-phenylethylcarbamate (PG)	150	100 50	-4 -4	10/10 10/10
Ethyl-N-(m-nitrophenylsulfonyl)carbamate (PG)	100	50 25	-10 -4	10/10 10/10
Ethyl-N-(p-chlorophenyl)carbamate (PG)	150	100 25	-6 0	10/10 6/10
2-Nitroisobutyl-N-phenylcarbamate (PG)	750	500 200	-2 -5	10/10 10/10
2-Nitro-2-methyl-1,3-propanediol-bis(N- phenylcarbamate) (PG)	150	100 50	-2 -3	9/10 10/10

n-Amyl-N-butyisulfonylcarbamate (sodium salt) (HOH)	750	500 200	-5 -2	10/10 10/10
Ethyl-N-(n-octylsulfonyl)carbamate (PG)	150	100 50	+6 -2	7/10 10/10
1-Methylammonium-2-dithiocarbamyl methylcyclobutane (HOH)	1000	300 100	+1 0	10/10 10/10
Thiocarbamic acid, p-hydroxyphenyl ester (PG)	150	100 50	+3 -1	7/10 10/10

UREAS AND THIOUREAS

Compound and vehicle used for toxicity and radiation tests	Toxicity		Radiation studies	
	Approx. LD ₅₀ (mg./kg.)	Dose (mg./kg.)	Change in ST ₅₀ (days)	Mortality at 30 days
Acetylurea (HOH)	>1000	1000 500	0 0	10/10 10/10
2-Aryl-2-thiopseudourea, monohydrobromide (HOH)	250	200 100	--1 +2	10/10 10/10
2-Methyl-2-thiopseudourea, monohydrobromide (HOH)	750	500 300	>+17 >+19	3/10 2/10
2-Methyl-2-thiopseudourea sulfate (HOH)	400	300 300 100	>+22 >+22 +7	3/10 1/9 6/10
2,2'-Trimethylene-bis(2-thiopseudourea), dihydrobromide (HOH)	250	200 100	0 +1	6/10 10/10
Tetramethylthiuram monosulfide (PG)	150	100 50	+2 >+15	10/10 3/10
β -Diisobutylaminoethylisothiourea dihydrochloride (HOH)	75	50 25	0 -2	10/10 10/10
2,2'-Pentamethylene-bis(1-acetyl-2-thiopseudourea), dihydrochloride (HOH)	250	200 100	-1 -3	10/10 10/10
2-Decyl-2-thiopseudourea, monohydrochloride (HOH)	75	25 10	0 +4	9/10 6/10
2-(Cyanomethyl)-2-thiopseudourea, monohydrochloride (HOH)	750	500 300	+7 0	5/10 8/10
2-(Amidinothio)butyric acid, monohydrobromide (HOH)	750	500 300	-10 -2	10/10 9/10
4-(Amidinothio)butyric acid, monohydrobromide (HOH)	100	50 25	-8 0	10/10 10/10
2-(2-Hydroxyethyl)-2-thiopseudourea, monohydrobromide (HOH)	400	300 200	-3 +1	10/10 7/10

Allylhydroxyethylthiourea (HOH)	>1000	1000 500	0 -1	9/10 10/10
2-(3-Hydroxypropyl)-2-thiopseudourea, monohydrobromide (HOH)	500	300 100 50	+2 -4 +2	4/5 10/10 10/10
2-(2-Methoxyethyl)-2-thiopseudourea, monohydrobromide (HOH)	400	300 200	-3 +7	7/10 7/10
1,1'-Dithioformamide dihydrochloride (HOH)	200	100 50	-4 +3	10/10 10/10
2,2'-Ethylene-bis(2-thiopseudourea), dihydrobromide (HOH)	75	50 25	-2 -1	10/10 10/10
2,2'-(2-Butenylene)bis(2-thiopseudourea), dihydrochloride (HOH)	75	50 25	+1 +3	10/10 10/10
2,2'-(2-Butynylene)bis(2-thiopseudourea), dihydrochloride (HOH)	100	50 25	> +22 +3	2/8 10/10
2,2'-Pentamethylene-bis(2-thiopseudourea), dihydrobromide (HOH)	150	100 50	+1 +2	10/10 10/10
2,2'-Oxydiethylene-bis(2-thiopseudourea), dihydrochloride (HOH)	400	200 100	0 +3	10/10 10/10
1,1'-Ethylene-bis[3-2-(amidinethio)ethyl urea], dihydrochloride (PG)	100	50 25	+2 -2	10/10 10/10
1,3-Bis-2-(amidinothio)ethylurea, dihydrochloride (HOH)	225	100 50	-1 +2	6/10 10/10
2,2'-Tetramethylene-bis(2-thiopseudourea), dihydrobromide (HOH)	250	200 100 50	-5 +3 +3	10/10 10/10 10/10
Tetraethylthiuram disulfide (PG)	75	50 25	-1 > +15	10/10 5/10
Cyclopentylisothiourea hydrobromide (HOH)	250	200 100 50	+4 0 -3	4/4 10/10 10/10
1-Phenyl-2-thiourea (HOH)	40	10 5	-6 +1	10/10 10/10

1-o-Tolyl-2-thiourea (PG)	150	100 50	+3 -2	8/10 10/10
2-(2-Phenoxyethyl)-2-thiopseudouracil, monohydrochloride (HOH)	40	25 10	-2 +1	10/10 10/10
p-Acetylaminoacetophenone thiosemicarbazone (PG)	50	25 10	-2 -1	10/10 7/10
2-(p-Nitrobenzyl)-2-thiopseudouracil, monohydrochloride (HOH)	75	50 25	+1 -4	10/10 10/10
p-Nitrobenzylisothiourea hydrobromide (HOH)	150	100 50	-2 -4	9/10 10/10
p-Nitrophenylsulfonylurea (PG)	750	500 300	+2 -2	8/10 10/10
N-Sulfanilylurea (HOH)	>1000	1000 500	+1 -1	10/10 9/10
N-(p-Acetylaminobenzenesulfonyl)ethylurethane (PG)	750	500 300	-3 -2	10/10 10/10
p,p'-Biphenylene-bis-1,1'-(2-thiourea) (PG)	75	50 25	-6 -6	10/10 10/10
Di-p-methoxyphenylthiourea (PG)	750	500 300	-4 -4	10/10 10/10
Di-p-chlorophenylthiourea (PG)	150	100 50	0 -3	10/10 10/10
1-Isothiourea-2-dibenzylaminoethane dihydrochloride (HOH)	100	50 25	-6 -1	10/10 9/10
1-(p-Phenylazo)phenyl-2-thiourea (PG)	150	100 50	+1 -2	7/10 9/10
Bis-[p,p-(isothioureidomethyl)phenyl] sulfone dihydrochloride (HOH)	75	50 25	-5 -5	10/10 10/10

This Document Contains
Missing Page/s That Are
Unavailable In The
Original Document

OR ARE
Blank pg.
that have
been removed

**BEST
AVAILABLE COPY**

AMINES

Compound and vehicle used for toxicity and radiation tests	Toxicity		Radiation studies	
	Approx. LD ₅₀ (mg./kg.)	Dose (mg./kg.)	Change in ST ₅₀ (days)	Mortality at 30 days
Methylamine hydrochloride (HOH)	> 1000	1000 500	-2 -2	10/10 10/10
N-(2-Nitro-2-methylbutyl)dimethylamine (PG)	200	100 50	+1 +3	10/10 10/10
N-(2-Nitroisobutyl)-N-methylglucamine (HOH)	> 1000	1000 500	-2 0	10/10 10/10
Benzyloctylmethylamine (PG)	100	50 25	+3 -1	10/10 8/10
Ethylamine: 2-bromohydrobromide (HOH)	> 1000	200 100	-2 +2	10/10 10/10
Ethylamine: 2-methoxy (65%-70% in HOH) (HOH)	400	200 100	-1 -4	10/10 10/10
2-Bromoethylamine hydrobromide (HOH)	400	200 100	-6 0	10/10 10/10
2,2'-Dithio-bis-(ethylamine) dihydrochloride (HOH)	750	300 200	> +16 0	0/10 7/10
Di(β -p-methoxyphenylethyl)amine hydrochloride (PG)	500	300 200	+3 -1	8/10 10/10
3-(N-Methylisopropylamino)-3-methyl-2-butanone hydrochloride (HOH)	150	100 50	-1 0	10/10 10/10
tert-Butyl-tert-amylamine hydrochloride (HOH)	100	50 25	-4 -2	10/10 10/10
3-tert-Butylamino-3-methyl-1-butene hydrochloride (HOH)	75	50 25	0 -5	10/10 10/10
3-tert-Butylamino-3-methyl-1-butyne hydrochloride (HOH)	75	50 25	-4 -2	10/10 10/10
3-Methyl-N-(2-o-propylphenoxyethyl)butylamine, monohydrochloride (HOH)	> 1000	1000 500	-10 -3	9/10 10/10
3,5-Dimethyl-3-amino-1-hexyne hydrochloride (HOH)	400	100 50	-5 -	10/10 10/10

N-(2-Nitroisobutyl)-2-ethylhexylamine (PG)	250	200 100	-2 +3	10/10 10/10
p-Isopropylbenzylhexylamine sulfate (PG)	150	100 50	-10 +1	10/10 10/10
N-Heptyltaurine, monosodium salt (HOH)	>1000	1000 500	+9 +3	5/10 10/10
p-Isopropylbenzylheptylamine sulfate (PG)	75	50 25	-7 -4	10/10 10/10
Octylamine (PG)	100	50 25	-1 +6	10/10 9/10
Benzioctylamine (PG)	100	50 25	-7 +1	10/10 9/10
p-Isopropylbenzyloctylamine (PG)	75	25 10	-8 +1	10/10 9/10
Dodecylamine (PG)	50	25 10	-3 -6	10/10 10/10
N-Dodecyltaurine, monosodium salt (HOH)	400	300 200	0 +2	10/10 10/10
Hexadecylamine (PG)	200	100 50	-1 +1	10/10 10/10
Octadecylamine (PG)	250	200 100	-3 0	10/10 10/10
N,N'-Dimethyl-1,2-butanediamine (PG)	150	100 50	-4 +1	10/10 10/10
N,N'-Bis(β -mercaptopethyl)hexamethylene diamine (PG)	350	200 100	-5 0	10/10 8/10
Armeen-CD-oleylamine (PG)	25	10 5	+2 +3	10/10 10/10
Armeen-TD-hydrogenated tallow amine (PG)	200	100 50	-1 +3	10/10 10/10
trans-1,2-Bis(aminooethyl)cyclobutane (HOH)	50	40 25	0 -3	9/10 10/10
1,3-Diamino-2,2,4, α -tetramethylcyclobutane (HOH)	450	200 100	-1 +1	10/10 10/10
Cyclobutane-1,2-bis(methylammonium) distearate (PG)	>1000	1000 750 500 500	-1 >+19 0 +2	6/10 3/10 5/10 9/10

Tetramethylammonium chloride (HOH)	25	7.5 5.0	+2 +2	9/10 10/10
Tetramethylammonium iodide (HOH)	40	10 5	0 +2	10/10 10/10
Aniline hydrochloride (HOH)	750	500 200	-4 -2	10/10 10/10
o-Toluidine (PG)	150	100 50	-1 -2	10/10 10/10
m-Toluidine (PG)	150	100 50	-1 -1	10/10 10/10
p-Toluidine (PG)	50	25 10	-2 +1	10/10 10/10
N-Acetyl-m-methylthioaniline (PG)	750	500 300	-2 -5	10/10 10/10
p-Nitroaniline (PG)	250	200 100	-3 +5	10/10 10/10
o-Phenylenediamine dihydrochloride (HOH)	250	200 100	-3 -1	10/10 10/10
m-Phenylenediamine dihydrochloride (HOH)	150	100 50	-4 0	9/10 10/10
2-Chlorobenzylethylenediamine (PG)	150	100 50	-2 -4	10/10 9/10
N,N-Dimethyl-p-phenylenediamine dihydrochloride (HOH)	40	25 10	+2 0	10/10 10/10
Diphenylbenzidine (sesame oil)	>1000	500 300	-2 +1	10/10 10/10
o,o-Dithio-bi-aniline (PG)	75	50 25	-6 -1	8/10 9/10
Benzyl-2,4-dichlorobenzylamine hydrochloride (PG)	250	200 100	-5 +2	10/10 6/10
Benzyl-3,4-dichlorobenzylamine hydrochloride (PG)	150	100 50	-3 -1	10/10 10/10
N-(2-Dimethylaminoethoxy)-N-(3,4,5-trimethoxybenzoyl)benzylamine hydrochloride (Tigan) (HOH)	400	300 100	+2 0	7/10 9/10
Dimethyl-a-naphthylamine (PG)	75	50 25	-3 0	10/10 10/10

N-(1-Naphthyl)ethylenediamine dihydrochloride (HOH)	150	100 50	+1 -2	10/10 10/10
2-Nitro-4-acetyl-2'-formylaminodiphenyl sulfide (PG)	250	200 100	-2 -7	10/10 10/10
N-(γ -Dimethylaminopropyl)-2-methyl thiadiphenylamine hydrochloride (HOH)	50	25 10	+2 -2	10/10 10/10
Hydroxylamine hydrochloride (HOH)	100	50 25	-4 -1	10/10 10/10
o-Benzylhydroxylamine hydrochloride (HOH)	250	200 100	-2 +1	10/10 9/10

GUANIDINES AND NITRILES

Compound and vehicle used for toxicity and radiation tests	Toxicity	Radiation studies		
	Approx LD ₅₀ (mg./kg.)	Dose (mg./kg.)	Change in ST ₅₀ (days)	Mortality at 30 days
Cyanoguanidine (HOH)	> 1000	1000 500	0 0	10/10 10/10
β -Phenethylbiguanide hydrochloride (HOH)	150	100 50 25	+2 -1 +1	4/4 10/10 10/10
Dipyridylguanidine trihydrochloride (HOH)	750	500 300	-2 -7	10/10 10/10
Methyl cyanoacetate (HOH)	750	500 300	+5 -2	10/10 10/10
Dimethylcyanamide (HOH)	40	25 10	-4 -2	10/10 10/10
Propionitrile (HOH)	50	25 10	-4 +1	9/10 10/10
Acrylonitrile (HOH)	75	75 50 25	+1 +18 -5	10/10 2/10 10/10
Cyanoacetamide (HOH)	750	500 300	0 -4	10/10 10/10
Adiponitrile (HOH)	40	25 10	+1 -4	10/10 10/10
Ethyl cyanoacetate (HOH)	750	500 300	+2 -4	10/10 10/10
Methacrylonitrile (PG)	100	50 75	+3 > +19	5/10 1/10
Hydroxyacrylonitrile (HOH)	> 1000	1000 500	-2 0	10/10 10/10
α,α' -Azodiisobutyronitrile (PG)	25	10 5	-2 -3	10/10 10/10
trans-1,2-Dicyanocyclobutane (HOH)	275	250 100 50 50 25	-7 -2 +10 0 +1	10/10 10/10 6/10 10/10 10/10
1,2-Dicyanocyclobutene-1 (HOH)	150	100 50	0 +1	9/10 10/10

3-Methylenecyclobutane carbonitrile (HOH)	250	200 100	0 0	10/10 10/10
Chlorinated-1,2-dicyanocyclobutane (Lot No. ST-15) (PG)	75	50 25	-4 -1	9/10 9/10
trans-2-Cyanocyclobutanecarboxamide (PG)	250	250 200 200 150 150 100 50	+3 +1 0 +15 +6 +3 +4	10/10 5/10 7/10 5/10 10/10 6/10 10/10
2,2-Dimethyl-3-methylenecyclobutane carbonitrile (HOH)	200	100 50	+2 0	10/10 10/10
Chlorinated-1,2-dicyanocyclobutane (Lot No. ST-24) (PG)	400	200 100 50	-4 -3 -4	10/10 10/10 10/10
Tetra- β -cyanoethyl-1,2-bis(aminomethyl) cyclobutane (HOH)	400	100 50	0 0	10/10 9/10
o-Chlorobenzonitrile (PG)	150	100 50	-4 +3	10/10 7/10
p-Chlorobenzonitrile (PG)	150	100 50	-2 0	10/10 10/10
Phenylacetonitrile (PG)	40	10 5	-6 -4	10/10 10/10
α -Dimethylamino-phenylacetonitrile (PG)	40	25 10	+3 -1	8/10 9/10
4-Hydroxybenzonitrile (HOH)	200	100 50	-6 -5	10/10 10/10
m-Trifluoromethylphenylacetonitrile (PG)	100	50 25	+1 +3	8/10 7/10
3,4-Dihydro-2-cyano-2H-pyran (HOH)	250	200 100	-1 +2	10/10 10/10
Thiocyanic acid, α -3, α -5-bis(dimethylamino)- 4-hydroxy-3,5-xylyl ester (PG)	75	50 25	+5 +2	7/10 9/10
Diphenylacetonitrile (PG)	200	100 50	-3 +2	10/10 10/10
4,4'-Biphenyldicarbonitrile (PG)	75	50 25	-5 -7	9/10 10/10

β -Cyanoethylmercaptan (HOH)	100	75 50	+1 +3	10/10 10/10
Sodium β -sulfopropionitrile (HOH)	>1000	500 300	-1 -2	10/10 10/10
p-Cetoxyphenylisothiocyanate (PG)	150	100 50	-5 +1	10/10 10/10
Cyclohexylcyanoethylethanolamine (PG)	200	100 25	-4 -1	10/10 10/10
p-(n)Dodecoxyphenylisothiocyanate (PG)	150	50 25	-2 -1	10/10 10/10
2-Thio-m-thiazone-2,4-dione (HOH)	750	400 200	-1 -4	10/10 10/10

This Document Contains
Missing Page/s That Are
Unavailable In The
Original Document

OR ARE
Blank pg.
that have
Been Removed

**BEST
AVAILABLE COPY**

HYDRAZINES

Compound and vehicle used for toxicity and radiation tests	Toxicity	Radiation studies		
		Approx. LD ₅₀ (mg./kg.)	Dose (mg./kg.)	Change in ST ₅₀ (days)
Hydrazine sulfate (HOH)	250	100 50		-2 -2
(3-Octylaminopropyl)hydrazine (PG)	15	10 5 1		+3 +2 -5
29 (2,2-Dimethylhydrazino)-3,6,9,12,15,18, 21,24,27-nonathianonacosane-1-thiol (PG)	400	400 300 250 200 50		-3 > +17 -4 -6 -4
Methylcyclopropanecarbonylhydrazine (HOH)	40	25 19		0 -3
1,2-Cyclobutanedicarboxylic acid dihydrazide (HOH)	>1000	500 300		-1 0
1,2-Cyclobutanedicarboxylic acid mono(2',2'-dimethyl)hydrazide (HOH)	>1000	1000 500		-1 +4
Acetphenylhydrazine (PG)	150	100 50		-1 -1
N-Benzyl-N'-isopropylhydrazine hydrochloride (HOH)	250	200 100 20		-2 0 -4
3,4,5-Trimethylacetophenonethiosemicarbazone (PG)	75	50 25		-10 -8
3,3'-Thiodipropionic acid, bis(D-galactosyl-methylenehydrazide) (PG)	750	500 300		0 -1
1-Isonicotinyl-2-dimethylhydrazine (PG)	400	300 100 30		+1 +1 -2
1-Ethyl-2-isonicotinylhydrazine (HOH)	150	100 50 10		+1 0 -5
1-Isonicotinyl-2-ethylidenehydrazine (PG)	150	100 50 10		+1 0 -2

[α -(Methylmercapto)-4-pyridylmethylenhydrazine (PG)]	200	100 50 10	-2 -2 -4	10/10 10/10 6/10
α -(Benzylmercapto)-4-pyridylmethylenhydrazine (PG)	3	1.0 0.5 0.1	+1 +2 -5	10/10 10/10 10/10
1-Isonicotinyl-2-glucosylhydrazine (PG)	400	300 100 30	-2 0 -4	10/10 10/10 10/10
N,N'-Ethyleneidene-bis(isonicotinylhydrazine) (Safflower oil)	>1000	1000 500	0 -3	10/10 10/10

THIOPHENES, OXACYCLOPENTANES, AZACYCLOPENTANES, AND AZACYCLOHEXANES

Compound and vehicle used for toxicity and radiation tests	Toxicity	Radiation studies		
	Approx. LD ₅₀ (mg./kg.)	Dose (mg./kg.)	Change in ST ₅₀ (days)	Mortality at 30 days
2-Acetothiophene (PG)	40	10 200	-2 -1	10/10 10/10
4'-Hydroxy-5-nitro-2-furanacrylanilide (PG)	400	300 200	+1 -1	7/10 10/10
4'-(2-Chloroethylthio)-5-nitro-2-furan-acrylanilide (PG)	200	100 50	-3 -1	10/10 10/10
p-[3-(5-Nitro-2-furyl)acrylamido]benzyl-thioacetic acid, methyl ester (PG)	750	500 300	-3 -1	5/10 10/10
N-Butylenepyrrolidine (HOH)	50	25 10	-2 +3	10/10 9/10
Monothiosuccinimide (PG)	150	100 50	+3 +10	6/10 6/10
N-Methylmonothiosuccinimide (PG)	250	200 100	-2 +3	7/10 10/10
2-[2-(1-Fyrrolidinyl)ethyl]-2-thiopseudourea, dihydrochloride (HOH)	150	100 50	-2 -1	10/10 10/10
N-Phenylmonothiosuccinimide (PG)	250	200 100	+22 +1	4/10 10/10
N-(p-Methoxyphenyl)monothiosuccinimide (PG)	250	200 100	-1 +4	5/10 6/10
Pyrrolidinopropiophenone hydrochloride (HOH)	75	50 25	-3 -2	10/10 10/10
1-(1-Pyrrolidinocyclohexyl)methylthio-sulfuric acid, zwitterion (I) (HOH)	150	100 50	+2 0	5/6 10/10
1,2-Cyclobutanedicarboxamide (HOH)	>1000	1500 1000 1000 500	0 +4 -1 +1	10/10 5/10 10/10 6/10
1-Aza-2-imino-7-oxacycloheptane (HOH)	>1000	1500 1000 1000 500	+2 +5 -1 -2	10/10 6/10 10/10 10/10

1-Aza-2,7-diiminocycloheptane (HOH)	250	200 100	0 +2	10/10 10/10
N-(α -Naphthyl)-1,2-cyclobutanedi-carboximide (PG)	750	500 300 100	-4 -3 +2	10/10 10/10 8/10
N-(β -Naphthyl)-1,2-cyclobutane dicarboximide (PG)	400	300 100	+2 +3	10/10 10/10
5-Acetyl-3-(2-aminoethyl)indole hydrochloride (HOH)	200	100 25	>+19 >+19	1/9 4/10
3-(2-Aminopropyl)-5-indolyl maleate salt (HOH)	625	500 300	+4 -1	3/8 6/10
DL-3-(2-Amino-1-methylethyl)-5-propionylindole hydrochloride (HOH)	200	100 50	+1 +2	13/17 8/10
DL-3-(Aminopropylene)indolyl-5-phenyl-ketone hydrochloride (HOH)	100	50 25	+3 -2	8/10 9/9
3-Amino-9-ethylcarbazole (PG)	150	100 50	-10 +1	10/10 5/10
3,6-Diamino-9-ethylcarbazole (PG)	150	100 50	-2 0	10/10 9/10
3-Methyl-5-carboxytetrahydro-4-caroline (PG)	150	100 50	0 +1	10/10 10/10
Ajmaline (PG)	75	50 25	+2 +1	9/10 10/10
Aspidospermine (PG)	40	25 10	-1 -2	10/10 8/10
Spiroxidine hydrochloride hemihydrate (HOH)	500	300 100	-3 -2	7/7 10/10
18-Benzylidene-17-hydroxy-yohimbane (PG)	250	200 100	>+19 -2	1/10 9/10
Monothioglutarimide (PG)	100	50 25	>+17 +3	3/9 10/10
2-Iminopiperidine hydrochloride (HOH)	150	50 25	+1 -5	8/10 10/10
N-(2-Nitroisobutyl)piperidine (PG)	200	100 50	-1 +3	10/10 10/10

1-Piperidine ethanethiol, monohydrochloride (HOH)	75	50 25	0 -1	10/10 10/10
1-Methyl-p-aminoethylthioazocyclohexane (HOH)	250	200 100	-4 0	10/10 10/10
1-[(2-Mercaptoethylthio)methyl]-4-methyl piperazine, polymer (PG)	200	100 50	-5 -1	9/10 10/10
4-Piperidinobenzaldehyde thiosemicarbazone (PG)	75	25 10	+1 -5	4/10 10/10
1-Phenyl-1-cyclopentyl-3-piperidino-1- propanol hydrochloride (HOH)	250	100 50	-3 -2	10/10 10/10
Isocinchomeric acid (pyridine 2,5- dicarboxylic acid) (HOH + NaHCO ₃)	>1000	1000 500	-6 0	10/10 10/10
2-Mercaptopyridine (PG)	250	200 100	-5 -5	10/10 10/10
Zinc derivative of 3-pyridinemethiol (PG)	150	100 50	-2 +4	10/10 8/10
Pyridoxal hydrochloride (HOH)	400	300 200	-1 -2	9/10 9/10
Thioisonicotinamide (PG)	250	200 100	+1 -1	10/10 10/10
Thioisonicotinamide-1-oxide (HOH)	750	500 300	0 0	10/10 10/10
Isonicotinaldehyde thiosemicarbazone (PG)	150	100 50	-2 -1	10/10 10/10
Nicotinaldehyde thiosemicarbazone hydrochloride (HOH)	150	100 50	-3 -3	10/10 10/10
Methyl-2-pyridyl ketone thiosemicarbazone (PG)	5	2 1	+2 -1	10/10 10/10
Methyl-3-pyridyl ketone thiosemicarbazone (PG)	25	10 5	-1 0	10/10 10/10
Nicotinylthiosemicbazide hydrochloride (HOH)	750	500 300	+1 0	10/10 10/10
Trimethylamine isonicotinamide (PG)	450	300 200	+2 +5	10/10 10/10

Isonicotinamido guanyl mercaptoacetic acid (HOH)	750	500 300	+1 0	10/10 10/10
Benzyl-4-benzylpyridinium chloride (HOH)	75	50 25	-3 0	10/10 10/10
Quinoline (synthetic) (PG)	200	100 50	+3 +9	7/11 6 10
Isoquinoline (PG)	150	100 50	-4 -1	10/10 10/10
6-Quinoliniccarboxylic acid (HOH + NaHCO ₃)	>1000	1000 500	-2 +2	10/10 8 9
3-Hydroxycinchoninic acid (PG)	250	200 100	-1 0	9/10 10 10
3-Hydrazinoquinoline dihydrochloride (HOH)	250	200 100	>+18 >+22	5 10 3 9
6-Methoxy-8-aminoquinoine (PG)	25	10 5	+2 -4	10 10 10/10
2-Hydroxy-3-n-butyl-9,10-dimethoxy-1, 2,3,4,5,7-hex-hydro-benzo(a)quinolizine hydrochloride (HOH)	150	100 50	-3 -1	10 10 10/10
6-Heptyl-4-quinolinol.1-oxide (PG)	450	300 200	-3 +1	10 10 10 10
2-Methyl-1,4-naphthoquinone-1-thiocsemi- carbazone (PG)	75	50 25	-2 -2	10 10 10/10
7-Chloro-4-[<i>(z</i> -diethyl)aminoethylamino] quinoline-1-oxide (PC)	150	100 50	+1 0	10/10 9 10
7-Chloro-4-[3-(1-pyrrolidinyl)propylamino] quinoline-1-oxide (PG)	150	100 50	0 +2	10 10 10 10
Cinchophen (PG)	200	100 50	-3 -5	10 10 10/10
1,7-Bis(N-tetrahydroisoquinolyl)heptane dihydrochloride (HOH)	250	50 25	-2 -3	10 10 9 10
1,10-Bis(N-tetrahydroisoquinolyl)decane dihydrochloride (PG)	150	50 25	+1 -2	10 10 10 10

2,3,3a,5,6,11,12,12a-Octahydro-8 hydroxy-1H-isoquinol(2,1a)(1)pyridinium bromide (HOH)	75	25 10	-2 -1	10/10 7/7
7-[2-Hydroxy-3-(N-2-hydroxyethyl-N-methyamino)-propyl]-1,3-dimethylxanthinepyridine-3-carboxylate (Complamex) (HOH)	>1000	1000 500 250 50	0 +3 -2 -4	10/10 10/10 10/10 10/10

OXAZOLES AND OXAZINES

Compound and vehicle used for toxicity and radiation tests	Toxicity		Radiation studies	
	Approx. LD ₅₀ (mg./kg.)	Dose (mg./kg.)	Change in ST ₅₀ (days)	Mortality at 30 days
3-(2-Hydroxyethyl)-2-oxazolidinone (HOH)	750	500 300	-1 -1	10/16 10/10
2-[2-(4-Morpholinyl)ether]-2-thiopseudourea, dihydrochloride (HOH)	40	25 10	0 -3	10/10 10/10
N-[2-Amidinothio)ethyl]-4-morpholine- carboxamide, monohydrochloride (HOH)	750	500 300	+4 +2	6/10 10/10
4-{[(2-Diethylaminoethyl)thio]methyl}-4- methylmorpholinium-p-toluenesulfonate metho-p-toluenesulfonate (HOH)	100	50 25	-2 0	10/10 9/10
4-[2-(Diethylaminomethylthio)ethyl]-4- methylmorpholinium-p-toluenesulfonate metho-p-toluenesulfonate (HOH)	50	25 10	-4 0	10/10 10/10
2-Imino-4-methyl-5-phenyloxazoline (PG)	10	5 1	+3 -4	10/10 9/10
4-Morpholinobenzaldehyde thiosemicarbazone (PG)	75	50 25	+5 -5	9/10 10/10
n-Heptyl ester of 4-(γ -morpholine propoxy) benzoic acid hydrochloride (HOH)	500	300 200	-7 -3	10/10 10/10

THIAZOLES AND THIAZINES

Compound and vehicle used for toxicity and radiation tests	Toxicity Approx. LD ₅₀ (mg./kg.)	Radiation studies		
		Dose (mg./kg.)	Change in ST ₅₀ (days)	Mortality at 30 days
2-Amino-2-thiazoline (PG)	150	100 50	-1 -5	10/10 10/10
2,4-Thiazolidinedione (PG)	750	300 200	-1 +2	10/10 10/10
2,4-Dimethylthiazole (PG)	250	200 100	0 0	9/10 10/10
Thiazolidine-4-carboxylic acid (HOH)	150	100 50	+7 0	10/10 7/10
Penicillamine (HOH)	350	300 100	+1 +1	8/10 10/10
6-Aminopenicillanic acid (PG)	>1000	500 300	+1 -1	10/10 10/10
5-(p-Chlorobenzylidene)-2-thiothiazolidine-2,4-dione (PG)	75	50 25	+6 -8	9/10 10/10
2-Thioureido-4-(p-nitrophenyl)thiazole (PG)	150	100 50	0 -8	7/10 10/10
5-(p-Dimethylaminobenzoylidene)rhodanine (PG)	150	100 50	+2 -3	9/10 9/10
5-Benzylidene-2-phenylimino-4-thiazolidinone (PG)	75	50 25	+1 -2	8/10 10/10
5-(p-Acetamidobenzylidene)-3-(p-acetamido-benzoylideneamino)rhodanine (PG)	100	50 25	+1 -4	6/10 8/10
6-Ethoxy-2-mercaptopbenzothiazole (PG)	250	200 100	-2 +2	10/10 8/10
5-Chloro-2-mercaptopbenzothiazole (PG)	150	100 50	+2 -5	10/10 10/10
2-(Dimethylaminomethylthio)benzothiazole (HOH)	150	100 50	+1 -6	10/10 10/10
Ethyl-3-methyl-4-oxo-5-piperidino-2-a-thiazolidine acetate (PG)	400	300 200	-1 0	9/10 9/10
2-Amino-4,5-diphenylthiazole (PG)	200	200 100	-7 -4	10/10 10/10

2,5-Bis(4-pyridyl)-1,3,4-thiadiazole (PG)	100	100 50	-4 0	10 10 10 10
5-Benzylidenerhodanine (PG)	75	50 25	-2 -2	10 10 10 10
3-Amino-5-benzylidenerhodanine (PG)	100	50 25	+3 +2	10/10 10/10
5-Benzylidene-3-benzylideneaminorhodanine (PG)	250	200 100	+1 0	8 10 9/10
5-Benzylidene-3-salicylideneaminorhodanine (PG)	75	50 40 25 15	-1 -3 > +22 0	10/10 10/10 4/10 10/10
2-(p-Methoxybenzylidenehydrazino)-4-oxo- 5-thiazolidineacetic acid (HOH + NaHCO ₃)	>1000	500 300	+3 +2	10/10 10/10
2-[(p-Acetamidobenzylidene)hydrazino]- 4-oxo-5-thiazolidineacetic acid (HOH + NaHCO ₃)	100	50 25	-1 +2	10/10 10/10
5,5'-Methylidynedirhodanine with triethyl- amine (PG)	150	100 50	-3 -5	10/10 10/10
2,2-Dimethyl-3-thiomorpholone (PG)	400	250 100	-3 +2	10 10 8/10
2,2-Diethyl-3-thiamorpholone (PG)	250	200 100	-1 -2	8/10 10 10
2-Methyl-1,5-benzothiazepin-4-(5H)-one (PG)	400	300 200	+2 +3	10/10 10/10
4-Methyl-1,5-benzothiazepin-2-(3H)-one (PG)	400	300 200	-1 +1	6/9 10/10
2,3-Dihydro-1,5-benzothiazepin-4-(5H)-one (PG)	750	500 360	0 +3	9/10 10/10
2,3-Dihydro-7-methyl-1,5-benzothiazepin- 4-(5H)-one,1,1-dioxide (PG)	400	300 200	+7 +2	7/10 10/10
2,3,4,5-Tetrahydro-1,5-benzothiazepin, monohydrochloride (HOH)	400	300 200	+19 +3	4 8 10 10
Thionine (HOH)	400	300 200	-3 -3	10/10 10/10
Methylene blue chloride (HOH)	150	100 50	-5 -2	10/10 10/10
(2-Chlorophenothiazino)propylbiguanide hydrochloride (HOH)	100	50 25	-4 -8	10/10 10/10

DIAZOLES AND DIAZINES

Compound and vehicle used for toxicity and radiation tests	Toxicity Approx. LD ₅₀ (mg./kg.)	Radiation studies		
		Dose (mg./kg.)	Change in ST ₅₀ (days)	Mortality at 30 days
1-Isopropyl-2-(2-furyl)-4,4-dimethyl-imidazolidine (PG)	250	200 100	-3 0	10/10 10/10
1,2-Diphenyl-4,4-dimethylimidazolidine (PG)	150	100 50	-1 +1	10/10 10/10
1-(1,1-Dimethyl-2-hydroxyethyl)-4,4-dimethyl-2-imidazolidinethione (PG)	250	200 100	0 0	10/10 10/10
1-Isopropyl-4,4-dimethyl-2-imidazolidine-thione (PG)	150	100 50	0 +5	8/10 10/10
1-Butyl-4,4-dimethyl-2-imidazolidinethione (PG)	150	100 50	-5 +1	10/10 10/10
1-(1,1,3,3-Tetramethylbutyl)-4,4-dimethyl-2-imidazolidinethione (PG)	100	50 25	-5 +1	10/10 10/10
1-Octadecyl-4,4-dimethyl-2-imidazolidine-thione (PG)	250	200 100	-1 -2	10/10 10/10
1-Phenyl-4,4-diethyl-2-imidazolidinethione (PG)	200	100 50	0 +1	10/10 10/10
2-[(p-Nitrobenzyl)thio]-2-imidazoline, monohydrobromide (HOH)	50	25 10	-1 +1	10/10 10/10
2-[(m-Chlorobenzyl)thio]-2-imidazoline, monohydrobromide (HOH)	75	25 10	+1 +1	10/10 10/10
2-{[(Tetrachloro-p-phenylene)methylene]thio}-di-2-imidazoline, dihydrochloride (HOH)	150	100 50	-1 0	6/7 10/10
2-[2-(Diethylamino)ethyl]-5-methyl-2H-(1)benzothiopyran-4,3,2-imidazole (PG)	200	100 50	-5 -2	10/10 10/10
6,7-Dihydro-3-methyl-1-5H-imidazo(2,1- β)thiazolium chloride (HOH)	75	50 25	-2 -1	10/10 10/10

2-Amino-1,3,4-thiadiazole, monohydrochloride (HOH)	250 100	200 100	-3 +2	10 10 10/10
5-Amino-1,3,4-thiadiazole-2-thiol (HOH)	350	200 100	+1 -2	8/10 10/10
2-Amino-5-mercaptop-1,3,4-thiadiazole (PG)	250 100	200 100	+2 -2	5/10 10/10
2,5-Dimercapto-1,3,4-thiadiazole (HOH)	750	200 100	-5 -3	10 10 10/10
1,3,4-Thiadiazole-2,5-dithiol (PG)	150	100 50	-2 +3	10/10 10/10
2,2'-Dithio-bis'5-amino-1,3,4-thiadiazole) (PG)	150	100 50	-2 -3	10/10 9/10
2-Amino-5-(p-aminophenylmercapto)-1,3,4-thiadiazole (PG)	250	200 100	-1 +3	8/10 10/10
5-Anilino-1,3,4-thiadiazole-2-thiol (PG)	250	200 100	-1 0	8/10 10 10
2-(4-Pyridyl)-5-phenyl-1,3,4-thiadiazole (PG)	150	100 50	-2 0	10/10 10 10
2-Methyl-1,3,4-thiadiazole-5-phenylhydrazine (PG)	75	50 25	-2 -2	10/10 10 10
2-Phenyl-1,3,4-thiadiazole-5-sulfonamide (PG)	250	200 100	-1 0	10 10 9/10
3-Allyl-2-thiohydantoin (HOH)	250	200 100	6 +1	10/10 10/10
3-(2-Aminoethyl)-2-thiohydantoin (HOH)	400	300 200	-1 -3	10/10 10 10
3-Phenyl-2-thionydantoin (PG)	75	25 10	+1 -5	9/10 10/10
3-Methyl-5-phenyl-5-(2-thienyl)hydantoin (PG)	150	100 50	-2 +3	10/10 10/10
5-Cyclohexylethyl-5-(2-thienyl)hydantoin (PG)	150	100 50	-5 0	10/10 8/10
3-Phenyl-5-(p-hydroxybenzyl)-2-thiohydantoin (PG)	150	100 50	-6 -7	10 10 10/10
5-Cyclohexylbutyl-5-(2-thienyl)hydantoin (PG)	200	100 50	-5 -4	10,10 10/10

1-Isonicotinyl-3,5-dimethylpyrazole (PG)	250	200 100	-2 -1	10/10 10/10
3,5-Dioxo-1,2-diphenyl-4-n-butylpyrazolidene (Butazolidin) (PG)	150	100 50	-5 0	9/10 9/10
3,5-Dioxo-1-phenyl-2-p-hydroxy-phenyl- 4-n-butylpyrazolidene (Oxyphenbutazone) (PG)	150	100 50	-2 0	10/10 9/10
N-[(2,3-Dimethyl-5-oxo-1-phenyl)-3- pyrazolin-4-yl]-anthranilic acid (HOH + NaHCO ₃)	100	50 25	+3 +2	8/10 10/10
1,2-Diphenyl-4-methylthioethyl-3,5- pyrazolidinedione (PG)	200	100 50	-1 > +15	10/10 9/10
1,2-Diphenyl-4-(2'-phenylsulfinethyl)- 3,5-pyrazolidinedione (Sulfinpyrazone) (PG)	150	100 50	0 -2	8/10 9/10
Quinoxaline (PG)	250	200 100	> +18 > +22	3/10 1/10
2,3-Quinoxalinediol (PG)	200	100 50	-7 -2	7/10 7/10
2,3-Quinoxalinedithiol (PG)	100	50 25	-5 +3	10/10 9/10
2,3-Di-(p,p'-dimethoxystyryl)quinoxaline (PG)	250	100 50	-4 -3	10/10 10/10
4-Quinazolinol, 3-oxide (PG)	250	200 100	+2 -1	10/10 10/10
4-Quinazolinethiol (PG)	750	500 300	> +17 > +19	4/10 4/10
6-Chloro-4-quinazolinethiol (PG)	>1000	1000 500	-2 +2	10/10 10/10
3-Propyl-4-(3H)-quinazolinethione (PG)	200	100 50	0 -1	10/10 10/10
3,4-Dihydro-4-phenyl-2(1H)-quinazoline- thione (PG)	75	50 25	0 +3	10/10 10/10
3-Benzyl-2,3-dihydro-2,2-dimethyl-4(1H)- quinazolinethione (PG)	450	300 200	-2 +4	10/10 10/10

3-(p-Acetylphenyl)-6-chloro-2-thio-2,4-(1H,3H)-quinazolininedione (PG)	250	200 100	-3 +2	10/10 10/10
7-Chloro-4-[(3-diethylaminopropyl)amino] quinoline-1-oxide (PG)	150	100 50	+1 +3	9/10 10/10
2-Dimethylaminopyrazine hydrochloride (HOH)	200	100 50	-3 -2	10/10 9/9
6-Amino-3-pyridazinethiol (PG)	400	300 100	+1 +2	10/10 8/10
Dithiothymine (PG)	200	100 50	0 +1	9/10 10/10
6-Aminouracil (sodium salt) (HOH)	>1000	500 300	-2 +1	10/10 10/10
2-Ethylthio-4-amino-6-oxypyrimidine (PG)	200	100 50	+2 0	6/10 7/10
5-Bromuracil (HOH + NaHCO₃)	>1000	1000 500	-6 -2	10/10 10/10
5-Carboxy-2-thiouracil (PG)	250	200 100	0 +1	10/10 5/10
4'-(1,4-Dihydro-2-mercaptop-4,4,6-trimethyl-1-pyrimidinyl)acetophenone (PG)	75	50 25	-2 0	7/10 10/10
1,4-Dihydro-1-(o-mercaptophenyl)-4,4,6-trimethyl-2-pyrimidinethiol (PG)	400	250 100	-3 +2	10/10 8/10
1-Methylpiperazine (PG)	150	100 50	+1 -2	10/10 10/10
1-m-Methylbenzylpiperazine (PG)	100	50 25	+1 -1	10/10 10/10
N-Benzyl N'-benzoylpiperazine (HOH)	400	100 50	-5 -1	10/10 10/10
1-(3,4-Dichlorophenyl)-4-methoxypropyl piperazine sulfate (HOH)	250	200 100	-7 -2	10/10 10/10
1,4-Piperazinediacetic acid, dihydrate (PG)	>1000	500 300	-4 -3	10/10 10/10
2-Pyridylethylpiperazine (PG)	150	100 25	-6 0	10/10 10/10

<i>a</i> -Ethyl-3-methyl-6-oxo-1(6H)-pyridazine acetic acid hydrazide (PG)	625	500 300	>+19 +3	4/10 5/10
1-[(Ethylthio)methyl]-1,4-dimethylpiperazinium di-p-toluenesulfonate, polymer (HOH)	7.5	5 1	0 -1	10/10 10/10
Thiocyanic acid, 4-hydroxy- <i>a</i> -3-(4-methyl-1-piperazinyl)-3,5-xylyl ester (PG)	75	50 25	+1 +1	10/10 10/10
1-(2-Methoxyphenyl)-4-thiofuran (PG)	400	300 100	-4 -2	10/10 10/10
1,4-Dimethyl-1,4-bis[2-(morpholinomethylthio)ethyl]-piperazinium di-p-toluenesulfonate bis(metho-p-toluene) sulfonate (HOH)	40	25 10	+1 +3	10/10 10/10
1,1,4-Trimethyl-4 { [2-(4-methyl-1-piperazinyl) ethylthio]methyl } piperazinium di-p-toluenesulfonate bis-(metho-p-toluene) sulfonate (HOH)	250	100 50	0 -3	9/10 10/10
6-Chloro- <i>o</i> -methyl-2 <i>H</i> -1,2,4-benzothiadiazin-7-sulfohydroxamic acid, 1,1-dioxide hemihydrate (PG)	400	300 100	-5 -7	10/10 10/10
7-Chloro-2-(cyclopropylamino)-5-phenyl-3 <i>H</i> -1,4-benzodiazepine-4-oxide (PG)	150	100 50	-1 -1	12/15 7/10

This Document Contains
Missing Page/s That Are
Unavailable In The
Original Document

OR ARE
Blank pg?
that have
been removed

**BEST
AVAILABLE COPY**

TRIAZOLES AND DIOXANES

Compound and vehicle used for toxicity and radiation tests	Toxicity		Radiation studies	
	Approx. LD ₅₀ (mg./kg.)	Dose (mg./kg.)	Change in ST ₅₀ (days)	Mortality at 30 days
S-Trithiane (Corn oil)	250	200 100	-2 -5	10/10 10/10
5-Amino-1H-1,2,4-triazole-3-carboxylic acid, sodium salt (HOH)	>1000	1000 500	0 -1	10/10 10/10
3-Mercapto-S-hydroxy-1,2,4-triazino-6- carboxylic acid (PG)	250	200 100	-3 -1	10/10 10/10
2,6-Diaminopurine sulfate (HOH + NaHCO ₃)	250	200 100	-1 0	10/10 10/10
2-Thio-6-oxypurine (HOH + NaHCO ₃)	625	500 300	-4 0	10/10 10/10
2,8-Dithio-6-oxypurine (HOH + NaHCO ₃)	400	300 100	-4 0	10/10 10/10
p-(2-Amino-6-hydroxy-8-purinylazo)- benzenesulfonic acid (HOH)	>1000	1000 500	+2 -6	10/10 10/10
2,6-Bis-(diethylolamine)-4,8-dipiperidino- pyrimido-(5,4-d) (PG)	150	100 50	-1 -1	10/10 10/10
1-Phenyl-5-mercaptotetrazole (PG)	250	200 50	-11 -1	10/10 10/10
Hypoxanthine (HOH + NaHCO ₃)	750	500 300	-1 -2	10/10 10/10
8-Azoguanine (HOH + NaHCO ₃)	>1000	1000 500	-8 -2	10/10 10/10
1-Methyl-2-mercapto-4,7-dihydroxy-1, 3,5,6-tetrazaindene (PG)	750	500 300	0 -7	8/10 10/10
1-Phenyl-2-mercapto-4,7-dihydroxy- 1,3,5,6-tetrazaindene (PG)	250	200 100	+1 -2	8/10 9/10
7-Theophylline acetamide (1,2,3,6-Tetrahydro- 1,3-dimethyl-2,6-dioxy-purine-7-acetamide) (PG)	750	500 300	-1 -2	9/10 10/10

Ethyl-7-theophylline acetate (Ethyl-1,2,3,6-tetrahydro-1,3-dimethyl-2,6-dioxopurine-7-acetate) (PG)	>1000	1000 500	+2 +1	7/10 7/10
8-(2-Dimethylaminoethoxy)theophylline (1,3-Dimethyl-8-(2-dimethylaminoethoxy) xanthine) (PG)	750	500 300	-3 >+19	8/10 4/10
7,7'-Trimethylene-bis { 8-[bis(2-hydroxyethyl)amino]theophylline } (PG)	1000	1000 500	>+19 -3	3/10 9/10
1,2-Cyclobutanedicarboguanamine (PG)	75	50 25	0 -2	10/10 10/10
1,2-Cyclobutanedicarboguanamine phosphate (HOH)	400	300 100	0 -1	10/10 10/10
1-Methyl-5-mercaptop-1,2,3,4-tetrazole (PG)	400	300 200	-6 -11	10/10 10/10
2-Phenyl-1,5-benzothiazepin-4(5H)-one (PG)	400	300 200	-2 +7	7/10 5/10
2-(Benzylmethylamino)-7-chloro-5-phenyl-3H-1,4-benzodiazepine-4-oxide (PG)	300	200 100	+3 -2	7/10 10/10
5-Methyl-7-phenyl-3H-1,2-dithiepin-3-thione (PG)	250	200 100	+3 +1	7/10 8/9
7-Chloro-1-methyl-5-phenyl-1H-1,4-benzodiazepine-2(3H)-thione (PG)	250	200 50	0 -1	6/10 10/10
5-Nitro-5-methyl-1,3-dioxane (PG)	150	100 50	+2 +3	10/10 10/10
2-Methyl-3-ethanol benzodioxane (HOH)	400	300 100	-3 +1	10/10 10/10

MISCELLANEOUS

Compound and vehicle used for toxicity and radiation tests	Toxicity	Radiation studies		
	Approx. LD ₅₀ (mg./kg.)	Dose (mg./kg.)	Change in ST ₅₀ (days)	Mortality at 30 days
Calcium chloride (HOH)	>200	200 100	+1 -2	10/10 10/10
Sodium chlorate (HOH)	>1000	1000 500	-1 +1	10/10 10/10
Sodium borate, tetra (HOH)	>1000	1000 500	-1 +1	10/10 10/10
Sodium nitrite (HOH)		125 100	+6 0	9/10 8/10
2-Nitropropane (PG)	75	50 25	-3 +2	10/10 10/10
2,3-Dibromopropane (PG)	75	50 25	-4 -3	10/10 10/10
2-Bromopentane (PG)	150	100 50	-7 -3	10/10 10/10
1,3-Dibromopropane (HOH)	750	500 200	-1 0	10/10 10/10
Naphthalene (PG)	150	100 50	-2 -4	10/10 10/10
α -(Methoxymethyl)naphthalene (PG)	200	100 50	-6 -3	9/10 10/10
Calcium flavonate (HOH)	625	500 300	-4 +3	10/10 10/10
Vitamin B ₁₂ with intrinsic factor concentrate (HOH)	>1000	1000 500	-4 0	10/10 10/10
Corn oil	>50000	50000	0	10/10
Olive oil	>50000	50000	-1	10/10
Petrolatum (liquid)	>50000	50000	0	10/10
Safflower oil	>50000	1 cc.	+2	10/10
Sesame oil	>50000	50000	-1	9/10

This Document Contains
Missing Page/s That Are
Unavailable In The
Original Document

OR ARE
Blank pg.
that have
been removed

**BEST
AVAILABLE COPY**

INDEX

	<i>Page</i>
Acetamide: 2-(2-hydroxyethoxy)	19
5-(p-Acetamidobenzylidene)-3-(p-acetamidobenzoylideneamino)rhodanine	47
2-[(p-Acetamidobenzylidene)hydrazino]-4-oxo-5-thiazolidineacetic acid	48
Acetic acid: 2-methoxy-l-methyl ethyl ester	17
Acetyl oxime	11
Acetone sodium bisulfite	5
2-Aceto thiophene	39
Acetphenylhydrazine	37
Acetylacetone	11
p-Acetylaminoacetophenone thiosemicarbazone	27
N-(p-Acetylaminobenzenesulfonyl)ethylurethane	27
5-Acetyl-3-(2-aminoethyl)indole hydrochloride	40
N-Acetyl-n-cysteine	15
N-Acetyl-m-methylthioaniline	31
N-Acetyl-DL-penicillamine	15
3-(p-Acetylphenyl)-6-chloro-2-thio-2,4-(1H,3H)-quinazolinedione	52
Acetylurea	25
Acrylonitrile	33
N-Acryloyltaurine, monosodium salt	19
Adiponitrile	33
Ajmaline	40
Alanine: 2-methyl-	14
Allylhydroxyethylthiourea	26
α -Allylmercapto- α,α -diethylacetamide	19
α -Allylmercaptoisobutyramide	19
3-Allyl-2-thiohydantoin	50

	<i>Page</i>
2-Allyl-2-thiopseudouracil, monohydrobromide	26
2-(Amidinothio)butyric acid, monohydrobromide	25
4-(Amidinothio)butyric acid, monohydrobromide	25
N-[2-(Amidinothio)ethyl]-4-morpholincarboxamide, monohydrochloride	45
3-(Amidinothio)propionic acid, ethyl ester, monohydrochloride	17
2-Amino-5-(p-aminophenylmercapto)-1,3,4-thiadiazole	50
N-(8-Amino-1-anthraquinonyl)anthranilic acid	14
3-Amino-5-benzylidenerhodanine	48
1-Amino-2-benzylmercapto-4-hydroxyanthraquinone	12
1-Amino-2-tert-butylphenylmercapto-4-hydroxyanthraquinone	8
2-Aminocyclobutanecarboxylic acid hydrochloride	13
2-Amino-4,5-diphenylthiazole	47
2-Aminoethanethiosulfuric acid	14
3-Amino-9-ethylcarbazole	40
3-(2-Aminoethyl)-2-thiohydantoin	50
3-(2-Aminoethylthio)propionic acid	14
6-Amino-5-hydroxymino-1,3-dimethyl hydouracil	11
p-(2-Amino-6-hydroxy-8-parinylazo)benzenesulfonic acid	55
2-Amino-5-mercaptop-1,3,4-thiadiazole	50
DL-3-(2-Amino-1-methylethyl)-5-propionylindole hydrochloride	40
4-Amino-3-methylphenol	6
6-Aminopenicillanic acid	47
o-Aminophenol	6
m-Aminophenol	6
p-Aminophenol hydrochloride	6
2-Amino-1-phenyl-1-ethanol	5
1-Amino-2-propanol	5

	<i>Page</i>
DL-3-(Aminopropylene)indolyl-5-phenylketone hydrochloride	40
3-(2-Aminopropyl)-5-indolyl maleate salt	40
6-Amino-3-pyridazinethiol	52
5-Amino-1,3,4-thiadiazole-2-thiol	50
2-Amino-1,3,4-thiadiazole, monohydrochloride	50
2-Amino-2-thiazoline	47
p-Aminothiobenzamide	21
1-Amino-2-thiophenylanthraquinone	12
1-Amino-4-thiophenylanthraquinone	12
1-Amino-2-thiophenyl-4-hydroxyanthraquinone	8
5-Amino-1H-1,2,4-triazole-3-carboxylic acid, sodium salt	55
6-Aminouracil (sodium salt)	52
n-Amyl-N-butylsulfonylcarbamate (sodium salt)	24
Aniline hydrochloride	31
5-Anilino-1,3,4-thiadiazole-2-thiol	50
N-1-Anthraquinonyl-2,3-dichloropropionamide	12
Armeen-TD-hydrogenated tallow amine	30
Armeen-CD-oleylamine	30
Aspidospermine	40
1-Aza-2,7-diiminocycloheptane	40
8-Azaguanine	55
1-Aza-2-imino-7-oxacycloheptane	39
α,α'-Azodiisobutyronitrile	33
Batyl alcohol	5
α-Benzoinoxime	7
p-Benzoquinone	11

	<i>Page</i>
N-Benzyl N'-benzoylpiperazine	52
Benzyl-4-benzylpyridinium chloride	42
Benzyl-2,4-dichlorobenzylamine hydrochloride	31
Benzyl-3,4-dichlorobenzylamine hydrochloride	31
3-Benzyl-2,3-dihydro-2,2-dimethyl-4(1H)-quinazolinethione	51
o-Benzylhydroxylamine hydrochloride	32
5-Benzylidene-3-benzylideneaminorhodanine	48
18-Benzylidene-17-hydroxy-yohimbane	40
5-Benzylidene-2-phenylimino-4-thiazolidinone	47
5-Benzylidenerhodanine	48
5-Benzylidene-3-salicylideneaminorhodanine	48
N-Benzyl-N'-isopropylhydrazine hydrochloride	37
α-(Benzylmercapto)-4-pyridylmethylenehydrazine	38
2-(Benzylmethylamino)-7-chloro-5-phenyl-3H-1,4-benzodiazepine-4-oxide	56
Benzyloctylamine	30
Benzyloctylmethylaniline	29
o,o'-Biphenol	6
4,4'-Biphenyldicarbonitrile	34
p,p'-Biphenylene-bis-1,1'-(2-thiourea)	27
1,3-Bis-(amidinothio)ethylurea, dihydrochloride	26
trans-1,2-Bis(aminoethyl)cyclobutane	30
1,6-Bis(p-aminoethylphenoxy)hexane dihydrochloride	21
Bis-(2-amino-4-sulfonamidophenyl) disulfide	9
N,N-Bis(2-cyanoethyl)-N-4-hydroxy-1-anthraquinonylsulfonilamide	21
Bis(Diaminoethyl ether)-N,N,N'-tetraacetic acid	13
Bis-didodecyldithioxamide	21
2,6-Bis-(diethylolamine)-4,8-dipiperidinopyrimido-(5,4-d)	55

	<i>Page</i>
Bis-[p-(α -diethylaminopropoxy)phenyl]sulfone	10
Bis-[p,p-(isothioureidomethyl)phenyl]sulfone dihydrochloride	27
1,2-Bis-(β -mercaptoethylaminomethyl)cyclobutane	7
N,N'-Bis(β -mercaptoethyl)hexamethylenediamine	30
1,6-Bis(p-methylaminomethylphenoxy)hexane dihydrochloride	9
N,N'-Bis(2-methoxyethyl)dithiooxamide	21
Bis(2-nitroisobutyl)-2-ethylhexyl phosphate	9
Bis-p-phenyl methane sulfone	10
2,5-Bis(4-pyridyl)-1,3,4-thiadiazole	48
1,10-Bis(N-tetrahydroisoquinolyl)decane dihydrochloride	42
1,7-Bis(N-tetrahydroisoquinolyl)heptane dihydrochloride	42
N,N'-Bis(1,1,3,3-tetramethylbutyl)-1,2-cyclobutanedicarboxamide	19
Bis-(2,4,5-trichlorophenyl)-1,2-cyclobutanedicarboxylate	18
2-Brom-D-lysergic acid diethylamide	14
2-Bromoethylamine hydrobromide	29
4'-Bromo-3-nitrosalicylanilide	20
2-Bromopentane	57
5-Bromuracil	52
2,3-Butanediol	5
2-Butanone oxime	11
2,2'-(2-Butenylene)bis(2-thiopseudourea), dihydrochloride	26
N-(3-Eutoxypropyl)taurine, monosodium salt	9
3-tert-Butylamino-3-methyl-1-butene hydrochloride	29
3-tert-Butylamino-3-methyl-1-butyne hydrochloride	29
N,N-di-tert-Butylcyclobutane-1,2-dicarboxamide	19
1-Butyl-4,4-dimethyl-2-imidazolidinethione	49
α -Butylmercaptoisobutyramide	19

	<i>Page</i>
N-Butylenepyrrolidine	29
tert-Butyl-tert-amylamine hydrochloride	29
2,2'-(2-Butynylene)bis(2-thiopseudocorea), dihydrochloride	26
Cacodylic acid	13
Calcium chloride	57
Calcium flavonate	57
Calcium gluconate	5
Calcium methane arsonate	17
Carbamic acid: 2-aminoethyl ester	23
5-Carbethoxy-2-thiouracil	52
(Carboxymethyl)trimethylammonium-N-acetyl-L-cysteinate	15
2-Carboxyphenylthioglycolic acid	14
Carbutamide	20
p-Cetoxyphenylisothiocyanate	35
Chlorinated-1,2-dicyanocyclobutane	34
m-Chloroacetylcatechol	6
p-Chlorobenzenethiol	8
o-Chlorobenzonitrile	34
p-Chlorobenzonitrile	34
2-Chlorobenzylethylenediamine	31
5-(p-Chlorobenzylidene)-2-thiothiazolidine-2,4-dione	47
2-[(m-Chlorobenzyl)thio]-2-imidazoline, monohydrobromide	49
7-Chloro-2-(cyclopropylamino)-5-phenyl-3H-1,4-benzodiazepine-4-oxide	53
7-Chloro-4-[(2-diethyl)aminoethylamino]quinoline-1-oxide	42
7-Chloro-4-[(3-diethylaminopropyl)amino]quinoline-1-oxide	52
4'-(2-Chloroethylthio)-5-nitro-2-furanacrylanilide	39

	<i>Page</i>
5-Chloro-2-mercaptopbenzothiazole	47
6-Chloro- α -methyl-2H-1,2,4-benzothiadiazin-7-sulfohydroxamic acid, 1,1-dioxide hemihydrate	53
7-Chloro-1-methyl-5-phenyl-1H-1,4-benzodiazepine-2(3H)-thione	56
4-Chloro-2-methylphenylthioglycolic acid	14
3'-Chloro-3-nitrosalicylanilide	20
3'-Chloro-5-nitrosalicylanilide	20
4'-Chloro-3-nitrosalicylanilide	20
4'-Chloro-5-nitrosalicylanilide	20
7-Chloro-4-(3-octylaminopropyl)aminoquinoline,1-oxide	12
(2-Chlorophenothiazino)propylbiguanide hydrochloride	48
3-Chloro-1-propanesulfonic acid, monosodium salt.....	14
7-Chloro-4-[3-(1-pyrrolidinyl)propylamino]quinoline, 1-oxide	42
6-Chloro-4-quinazolinethiol	51
2'-Chloro-5'-trifluoromethyl-3-nitrosalicylanilide	20
Chrysanthemummonocarboxylic acid	13
Cinchophen	42
Copper methanearsonate	17
Corn oil	57
α -C ₃ anoacetamide	19
Cyanoacetic acid	13
Cyanoacetamide	33
trans-2-Cyanocyclobutanecarboxamide	34
β -Cyanoethylmercaptan	35
Cyanoguanidine	33
2-(Cyanomethyl)-2-thiopseudourrea, monohydrochloride	25
Cyclobutane-1,2-bis(methylammonium) adipate	18
Cyclobutane-1,2-bis(methylammonium) dimonochloroacetate	9

	<i>Page</i>
Cyclobutane-1,2-bis(methylammonium) distearate	30
Cyclobutane-1,2-bis(methylammonium) phosphate	9
1,2-Cyclobutanedicarboguanamine	56
1,2-Cyclobutanedicarboguanamine phosphate	56
trans-1,2-Cyclobutanedicarboxamide	19
1,2-Cyclobutanedicarboxamide	39
trans-1,2-Cyclobutanedicarboxylic acid	13
1,2-Cyclobutanedicarboxylic acid dihydrazide	37
1,2-Cyclobutanedicarboxylic acid mono (2',2'-dimethyl)hydrazide	37
1,2-Cyclobutanedimethylcarbamate	23
Cyclohexanone oxime	11
5-Cyclohexylbutyl-5-(2-thienyl)hydantoin	50
Cyclohexylcyanoethyllethanolamine	35
5-Cyclohexylethyl-5-(2-thienyl)hydantoin	50
Cyclopentylisothiourea hydrobromide	26
L-Cysteine hydrochloride	15
2-Decyl-2-thiopseudourea, monohydrochloride	25
Desoxyribozyme hydrochloride	6
2,2'-Diallyl-p,p'-biphenol	6
1,2-Diaminocyclohexanetetraacetic acid	13
3,6-Diamino-9-ethylcarbazole	40
2,6-Diaminopurine sulfate	55
1,3-Diamino-2,2,4,4-tetramethylcyclobutane	30
1,3-Dibromopropane	57
2,3-Dibromopropane	57
Dibutyl fumarate	17
Dibutyl maleate	17

	<i>Page</i>
N,N'-Dicarbamylformamidine	23
3',4'-Dichloro-3-nitrosalicylanilide	20
1-(3,4-Dichlorophenyl)-4-methoxypropylpiperazine sulfate	52
Di-p-chlorophenylthiourea	27
N-(3,5-Dichloro-p-tolyl)anthranilic acid	14
trans-1,2-Dicyanocyclobutane	33
1,2-Dicyanocyclobutene-1	33
2,3-Di-(p,p'-dimethoxystyryl)quinoxaline	51
2-[2-(Diethylamino)ethyl]-5-methyl-2H-(1)benzothiopyrano-4,3,2-imidazole	49
4-{[(2-Diethylaminoethyl)thio]methyl}-4-methylmorpholinium-p-toluenesulfonate metho-p-toluenesulfonate	45
4-[2'-Diethylaminomethylthio)ethyl]-4-methylmorpholinium-p-toluene-sulfonate metho-p-toluenesulfonate	45
Diethylcarbamyl chloride	23
Di-2-ethylhexyl fumarate	17
Di-2-ethylhexyl maleate	18
2,2-Diethyl-3-thiamorpholone	48
2,3-Dihydro-1,5-benzothiazepin-4-(5H)-one	48
3,4-Dihydro-2-cyano-2H-pyran	34
1,4-Dihydro-1-(o-mercaptophenyl)-4,4,6-trimethyl-2-pyrimidinethiol	52
4'-(1,4-Dihydro-2-mercaptop-4,4,6-trimethyl-1-pyrimidinyl)acetophenone	52
2,3-Dihydro-7-methyl-1,5-benzothiazepin-4-(5H)-one,1,1-dioxide	48
6,7-Dihydro-3-methyl-1-5H-imidazo(2,1- β)thiazolium chloride	49
3,4-Dihydro-4-phenyl-2(1H)-quinazolinethione	51
DL- β -(3,4-Dihydroxyphenyl)alanine	14
β -Diisobutylaminoethylisothiourea dihydrochloride	25
Diisodecyl-1,2-cyclobutanedicarboxylate	18
cis-1,2-Diisopropylcyclobutanedicarboxylate	18

	<i>Page</i>
trans-1,2-Diisopropylcyclobutanedicarboxylate	18
meso-2,3-Dimercaptosuccinic acid	14
2,5-Dimercapto-1,3,4-thiadiazole	50
Dimercurous methanearsonate	17
p-Dimethoxybenzene	9
Di(β-p-methoxyphenylethyl)amine hydrochloride	29
Di-p-methoxyphenylthiourea	27
N,N-Dimethylacetamide	19
p-Dimethylaminobenzenethiol	8
5-(p-Dimethylaminobenzoylidene)rhodanine	47
8-(2-Dimethylaminoethoxy)theophylline	56
N-(2-Dimethylaminoethoxy)-N-(3,4,5-trimethoxybenzoyl)benzylamine hydrochloride	31
3,5-Dimethyl-3-amino-1-hexyne hydrochloride	29
2-(Dimethylaminomethylthio)benzothiazole	47
α-Dimethylaminophenylacetonitrile	34
N-(γ-Dimethylaminopropyl)-2-methylthiodiphenylamine hydrochloride	32
2-Dimethylaminopyrazine hydrochloride	52
1,4-Dimethyl-1,4-bis[2-(morpholinomethylthio)ethyl]-piperazinium di-p-toluenesulfonate bis(metho-p-toluene) sulfonate	53
N',N'-Dimethyl-1,2-butanediamine	30
Dime...y...carbamyl chloride	23
Dimethylcyanamide	33
Dimethyl-1,2-cyclobutanedicarboxylate	18
Di(methylcyclopropyl)ketoxime	11
N,N-Dimethyl-α,α-diphenylthioacetamide	21
29-(2,2-Dimethylhydrazino)-3,6,9,12,15,18,21,24,27-nonathianonacosane-1-thiol	37
1-(1,1-Dimethyl-2-hydroxyethyl)(-4,4-dimethyl-2-imidazolidinethione	49

	<i>Page</i>
2,2-Dimethyl-3-methylenecyclobutane carbonitrile	34
Dimethyl- α -naphthylamine	31
N-[(2,3-Dimethyl-5-oxo-1-phenyl)-3-pyrazolin-4-yl]-anthranilic acid	51
2,6-Dimethylphenol	5
N,N-Dimethyl-p-phenylenediamine dihydrochloride	81
Dimethylsulfoxide	12
2,4-Dimethylthiazole	47
2,2-Dimethyl-3-thiomorpholone	48
4',3-Dinitrosalicylanilide	20
N,N'-Dioctadecylidithioxamide	21
3,5-Dioxo-1,2-diphenyl-4-n-butylpyrazolidene	51
3,5-Dioxo-1-phenyl-2-p-hydroxy-phenyl-4-n-butylpyrazolidene	51
Diphenylacetonitrile	34
Diphenylbenzidine	31
1,2-Diphenyl-4,4-dimethylimidazolidine	49
1,2-Diphenyl-4-methylthioethyl-3,5-pyrazolidinedione	51
1,2-Diphenyl-4-(2'-phenylsulfinethyl)-3,5-pyrazolidinedione	51
Dipotassium methanearsonate	17
2,2'-Dipropyl-p,p'-biphenol	6
Dipyridylguanidine trihydrochloride	83
Disodium methanearsonate	17
4,4'-Di-tert-butyl-o,o'-biphenol	6
N,N'-Di-tert-butylidithioxamide	21
2,4-Di-tert-butylphenol	5
2,2'-Dithio-bis(5-amino-1,3,4-thiadiazole)	50
o,o'-Dithio-bis-aniline	31
2,2'-Dithio-bis-(ethylamine) dihydrochloride	29

	<i>Page</i>
1,1'-Dithioformamidine dihydrochloride	26
2,8-Dithio-6-oxypurine	55
Dithiothymine	52
p-(n)Dodecoxyphenylisothiocyanate	35
Dodecylamine	30
N-Dodecyltaurine, monosodium salt	30
5,8,11,14-Eicosatetraenoic acid	14
6-Ethoxy-2-mercaptopbenzothiazole	47
Ethylamine: 2-bromohydrbromide	29
Ethylamine: 2-methoxy-	29
Ethyl-N,N-butylcarbamate	23
Ethyl-N,N-di-N-butylcarbamate	23
Ethyl-N-(p-chlorophenyl)carbamate	23
Ethyl cyanoacetate	33
1,1'-Ethylene-bis[3-2-(amidinethio)ethylurea], dihydrochloride	2t
2,2'-Ethylene-bis(2-thiopseudourea), dihydribromide	26
Ethylenediamine-di(o-hydroxyphenyl)acetic acid	14
Ethylenediaminetetraacetic acid	13
Ethylene glycol-bis-(aminoethyl ether)tetraacetic acid	13
N,N'-Ethyleneidene-bis(isonicotinylhydrazine)	38
1-Ethyl-2-isonicotinylhydrazine	37
α-Ethylmercaptoisobutyramide	19
Ethyl-3-methyl-4-oxo-5-piperidino-2,α-thiazolidine acetate	47
-Ethyl-3-methyl-6-oxo-1(6H)-pyridazine acetic acid hydrazide	53
/l-1-methyl propyl diethyl malonate	18
ϵ-N-(p-nitrobenzenesulfonyl)urethane	18
μ- p- itrobenzoate	18

	<i>Page</i>
Ethyl-N-(m-nitrophenylsulfonyl)carbamate	23
Ethyl-N-(n-octylsulfonyl)carbamate	24
Ethyl-7-theophylline acetate	56
2-Ethylthio-4-amino-6-oxypyrimidine	52
1-[(Ethylthio)methyl]-1,4-dimethylpiperazinium di-p-toluenesulfonate	53
Ethyl-N-o-toluenesulfonylurethane	18
4'-Fluoro-3-nitrosalicylanilide	20
Glutathione	15
Glutathione ascorbate	15
Glutathione ascorbate, 1:10	15
Glutathione ascorbate with calcium phytate	15
Glutathione (glutamyl-cysteinyl-glycine)	15
Glyoxal	5
n-Heptyl ester of 4-(γ -morpholine propoxy)benzoic acid hydrochloride	45
6-Heptyl-4-quinolinol, 1-oxide	42
N-Heptyltaurine, monosodium salt	30
Hexadecylamine	30
2,3,4a,5,8,8a-Hexahydro-1,4-naphthoquinone	12
Hexamethylenediammonium cyclobutane-1,2-dicarboxylate	18
Hydrazine sulfate	37
3-Hydrazinoquinoline dihydrochloride	42
Hydroquinone	7
Hydroxyacrylonitrile	33
4-Hydroxybenzonitrile	34
2-Hydroxy-3-n-butyl-9,10-dimethoxy-1,2,3,4,5,7-hexahydro- benzo(a)quinolizine hydrochloride	42
3-Hydroxycinchoninic acid	42
o-Hydroxy-N,N-diethylbenzamide	20

	<i>Page</i>
p-Hydroxydiphenyl	6
β -Hydroxyethylbenzyl ether	9
β -Hydroxyethyl ether of butylphenol	9
β -Hydroxyethyl ether of β -naphthol	10
β -Hydroxyethyl ether of orthocresol	9
β -Hydroxyethyl ether of orthophenylphenol	10
β -Hydroxyethyl ether of m-xlenol	9
3-(2-Hydroxyethyl)-2-oxazolidinone	45
2-(2-Hydroxyethyl)-2-thiopseudourea, monohydrobromide	25
7-[2-Hydroxy-3-(N-2-hydroxyethyl-N-methylamino)-propyl]-1,3-dimethylxanthinepyridine-3-carboxylate	43
Hydroxylamine hydrochloride	32
4'-Hydroxy-5-nitro-2-furanacrylanilide	39
2-(3-Hydroxypropyl)-2-thiopseudourea, monohydrobromide	26
1-Hydroxy-4-thiophenylanthraquinone	8
Hypoxanthine	55
α -Imino-p-methoxybenzylmercaptoacetic acid	13
2-Imino-4-methyl-5-phenyloxazoline	45
2-Iminopiperidine hydrochloride	40
Indole-3-acetic acid	14
3'-Iodo-3-nitrosalicylanilide	20
4'-Iodo-3-nitrosalicylanilide	20
2-Isobutylaminoethanol	5
Isocinchomeronic acid	41
Isonicotinaldehyde thiosemicarbazone	41
Isonicotinamido guanylmercaptoacetic acid hydrochloride	42
1-Isonicotinyl-2-dimethylhydrazine	37
1-Isonicotinyl-3,5-dimethylpyrazole	51

	<i>Page</i>
1-Isonicotinyl-2-ethylidenehydrazine	37
1-Isonicotinyl-2-glucosylhydrazine	38
Isonitrosopropiophenone	11
p-Isopropylbenzylheptylamine sulfate	30
p-Isopropylbenzylhexylamine sulfate	30
p-Isopropylbenzyloctylamine	30
1-Isopropyl-4,4-dimethyl-2-imidazolidinethione	49
1-Isopropyl-2-(2-furyl)-4,4-dimethylimidazolidine	49
p,p'-Isopropylidene-diphenol	6
4,4'-Isopropylidene-bis(2-tert-butylphenol)	7
α-Isopropylmercaptoisobutyramide	19
Isopropyl-N-phenylethylcarbamate	23
Isoquinoline	42
1-Iothiourea-2-dibenzylaminocethane dihydrochloride	27
Levulinic acid	13
Magnesium cacodylate	17
Magnesium methanearsonate	17
1-Mercaptoanthraquinone	8
1-[(2-Mercaptoethylthio)methyl]-4-methyl piperazine, polymer	41
3-Mercapto-S-hydroxy-1,2,4-triazino-6-carboxylic acid	55
2-Mercaptopyridine	41
Methacrylonitrile	33
1-Methoxy-2-amino-2-methylpropane	9
6-Methoxy-8-aminoquinoline	42
2-(p-Methoxybenzylidenehydrazino)-4-oxo-5-thiazolidineacetic acid	48
2-(2-Methoxyethyl)-2-thiopoulosurea, monohydrobromide	26

	<i>Page</i>
<i>a</i> -(Methoxymethyl)naphthalene	57
<i>p</i> -Methoxyphenol	6
N-(<i>p</i> -Methoxyphenyl)monothiosuccinimide	39
1-(2-Methoxyphenyl)-4- <i>t</i> riofuran	53
2-Methoxy-4- <i>n</i> -propylphenol	6
Diethylallyl acetone	11
Methylamine hydrochloride	29
β -Methylaminoethanol	5
1-Ethyl- <i>p</i> -aminoethylthioazocyclohexane	41
1-Methylammonium-2-carbamylmethylcyclobutane	23
1-Methylammonium-2-dithiocarbamylmethylcyclobutane	24
2-Methyl-1,5-benzothiazepin-4-(5H)-one	48
4-Methyl-1,5-benzothiazepin-2-(3H)-one	48
1-m-Methylbenzylpiperazine	52
3-Methyl-5-carboxytetrahydro-4-caroline	40
Methyl chloroformate	17
N-Methyl- <i>a</i> -crotylmercaptoisobutyramide	19
Methyl cyanoacetate	33
3-Methyl-2-cyclobutene-1,2-dicarboxylic acid	13
2-Methylcyclohexanone	11
Methylcyclopropanecarbonylhydrazine	37
N,N'-Methylene-bis-(thioisonicotinamide)hydrate	21
Methylene blue chloride	48
3-Methylenecyclobutane carbonitrile	34
3-Methylenecyclobutanecarboxylic acid	13
2-Methyl-3-ethanol benzodioxane	56
5,5'-Methylidynedirhodanine with triethylamine	48

	<i>Page</i>
3-(N-Methylisopropylamino)-3-methyl-2-butanol hydrochloride	5
3-(N-Methylisopropylamino)-3-methyl-2-butanone hydrochloride	29
Methylmercaptoacetate	7
α -Methylmercapto- α,α -diethylacetamide	19
1-Methyl-2-mercapto-4,7-dihydroxy-1,3,5,6-tetrazaindene	55
[α -(Methylmercapto)-4-pyridylmethylene] hydrazine	38
1-Methyl-5-mercaptop-1,2,3,4-tetrazole	56
N-Methylmonothiosuccinimide	39
Methylnaphthoquinone	12
2-Methyl-1,4-naphthoquinone	12
2-Methyl-1,4-naphthoquinone-1-thiosemicarbazone	42
Methyl-p-nitrobenzoate	18
2-Methyl-2-nitro-1-propyltosylate	9
3-Methyl-4-phenylazophenol	7
5-Methyl-7-phenyl-3H-1,2-dithiepin-3-thione	56
3-Methyl-5-phenyl-5-(2-thienyl)hydantoin	50
1-Methylpiperazine	52
3-Methyl-N-(2-o-propylphenoxyethyl)butylamine, monohydrochloride	29
Methyl-2-pyridyl ketone thiosemicarbazone	41
Methyl-3-pyridyl ketone thiosemicarbazone	41
2-Methyl-1,3,4-thiadiazole-5-phenylhydrazine	50
4-Methylthio-m-cresol	7
6-Methylthio-m-cresol	7
2-Methyl-2-thiopseudourea, monohydrobromide	25
2-Methyl-2-thiopseudourea sulfate	25
Monoammonium methanearsonate	17
Monothioglutarimide	40

	<i>Page</i>
Monothiosuccinamide	39
4-Morpholinobenzaldehyde thiosemicarbazone	45
2-[2-(4-Morpholinyl)ether]-2-thiopseudourea, dihydrochloride	45
Naphthalene	57
Naphthoresorcinol	6
N-(α-Naphthyl)-1,2-cyclobutanedicarboximide	40
N-(β-Naphthyl)-1,2-cyclobutanedicarboximide	40
N,N'-(α-Naphthyl)-1,2-cyclobutanedicarboxamide	20
N,N'-(β-Naphthyl)-1,2-cyclobutanedicarboxamide	20
N-(1-Naphthyl)ethylenediamine dihydrochloride	32
β-Naphthylethyl ether	10
Nicotinaldehyde thiosemicarbazone hydrochloride	41
Nicotinylthiosemicarbazide hydrochloride	41
Nitriloacetic acid trisodium salt monohydrate	17
2-Nitro-4-acetyl-2'-aminodiphenyl sulfide	12
2-Nitro-4-acetyl-2'-formylaminodiphenyl sulfide	32
p-Nitroaniline	31
p-Nitrobenzylisothiourea hydrobromide	27
2-[(p-Nitrobenzyl)thio]-2-imidazoline, monohydrochloride	49
2-(p-Nitrobenzyl)-2-thiopseudourea, monohydrochloride	27
p-[3-(5-Nitro-2-furyl)acrylamido]benzylthioacetic acid, methyl ester	39
N-(2-Nitroisobutyl)-2-ethylhexylamine	30
N-(2-Nitroisobutyl)-N-methylglucamine	29
2-Nitroisobutyl-N-phenylcarbamate	23
N-(2-Nitroisobutyl)piperidine	40
N-(2-Nitro-2-methylbutyl)dimethylamine	29
5-Nitro-5-methyl-1,3-dioxane	56

	<i>Page</i>
2-Nitro-2-methyl-1,3-propanediol-bis(N-phenylcarbamate)	23
2-Nitro-2-methyl-1,3-propanedioldicarbamate	23
p-Nitrophenol	6
p-Nitrophenylsulfonylurea	27
2-Nitropropane	57
p-Nonylphenyl- β -dimethylaminoethyl ether	9
Octadecylamine	30
1-Octadecyl-4,4-dimethyl-2-imidazolidinethione	49
2,3,3a,5,6,11,12,12a-Octahydro-8-hydroxy-1H-isoquino(2,1a)(1)pyridinium bromide	43
Octylamine	30
(3-Octylaminopropyl)hydrazine	37
Olive oil	57
2-Oxo-3-isobutyl-9,10-dimethoxy-1,3,4,6,7,11- β -hexahydro-2H- benzoquinolizine	12
2,2'-Oxydiethylene-bis(2-thiopseudourea), dihydrochloride	26
p,p'-Oxydiphenol	7
Penicillamine	47
2,2'-Pentamethylene-bis(1-acetyl-2-thiopseudourea), dihydrochloride	25
2,2'-Pentamethylene-bis(2-thiopseudourea), dihydribromide	26
Petrolatum (liquid)	57
β -Phenethylbiguanide hydrochloride	33
2-(2-Phenoxyethyl)-2-thiopseudourea, monohydrochloride	27
Phenylacetonitrile	34
p-Phenylazophenol	7
1-(p-Phenylazo)phenyl-2-thiourea	27
2-Phenyl-1,5-benzothiazepin-4(5H)-one	56
1-Phenyl-1-cyclopentyl-3-piperidino-1-propanol hydrochloride	41

	<i>Page</i>
1-Phenyl-4,4-dimethyl-2-imidazolidinethione	49
m-Phenylenediamine dihydrochloride	31
o-Phenylenediamine dihydrochloride	31
3-Phenyl-5-(p-hydroxybenzyl)-2-thiohydantoin	50
Phenylmercaptodiaminoanthrarufin	8
1-Phenyl-2-mercapto-4,7-dihydroxy-1,3,6,6-tetrazaindene	55
1-Phenyl-5-mercaptotetrazole	55
Phenylmercurilauryl thioether	9
N-Phenylmonothiosuccinimide	39
p-Phenylphenol	6
γ-Phenylpropylcarbamate	23
2-Phenyl-1,3,4-thiadiazole-5-sulfonamide	50
3-Phenyl-2-thiohydantoin	50
1-Phenyl-2-thiourea	26
1,4-Piperazinediacetic acid, dihydrate	52
1-Piperidine ethanethiol, monohydrochloride	41
4-Piperidinobenzaldehyde thiosemicarbazone	41
α-Propargylmercapto-α,α-diethylacetamide	19
Propionitrile	33
Propylene glycol (1,2-propanediol)	5
N-Propyl-p-hydroxybenzoate	6
3-Propyl-4-(3H)-quinazolinethione	51
Pulegone	11
Pyridine-2-aldoxime	11
Pyridine-3-aldoxime	11
Pyridine-4-aldoxime	11

	<i>Page</i>
Pyridine-2-aldoxime <i>o</i> -benzyl ether	11
Pyridoxal hydrochloride	41
2-(4-Pyridyl)-5-phenyl-1,3,4-thiadiazole	50
2-Pyridylethylpiperazine	52
Pyrogallol	6
1-(1-Pyrrolidinocyclohexyl)methylthiosulfuric acid, zwitterion (I)	39
Pyrrolidinopropiophenone hydrochloride	39
2-[2-(1-Pyrrolidinyl)ethyl]-2-thiopseudourea, dihydrochloride	39
4-Quinazolinethiol	51
4-Quinazolinol, 3-oxide	51
Quinoline (synthetic)	42
6-Quinolincarboxylic acid	42
2-Quinolinol	7
8-Quinolinol	7
Quinoxaline	51
2,3-Quinoxalinediol	51
2,3-Quinoxalinedithiol	51
Safflower oil	57
<i>o</i> -Salicylotoluidide, 3,5'-dinitro	7
2',5'-Salicyloxylidide, 3-nitro	7
2',6'-Salicyloxylidide, 3-nitro	7
Sesame oil	57
Sodium borate, tetra	57
Sodium cacodylate	17
Sodium chlorate	57
Sodium-p-hydroxyphenylarsonate	6
Sodium β -naphthoquinone-4-sulfonate	12

	<i>Page</i>
Sodium nitrite	57
Sodium β -sulfopropionitrile	35
Spiroxyine hydrochloride hemihydrate	40
N-(3-Sulfamoyl-2,6-xylyl)anthranilic acid	14
N-Sulfanilylurea	27
2,2-{[(Tetrachloro-p-phenylene)methylene]thio}-di-2-imidazoline, dihydrochloride	49
Tetra- β -cyanoethyl-1,2-bis(aminomethyl)cyclobutane	34
Tetraethylthiuram disulfide	26
2,3,4,5-Tetrahydro-1,5-benzothiazepin, monohydrochloride	48
Tetramethylammonium chloride	31
Tetramethylammonium iodide	31
1-(1,1,3,3-Tetramethylbutyl)-4,4-dimethyl-2-imidazolidinethione	49
Tetramethyldiaminothiobenzophenone	12
2,2'-Tetramethylene-bis(2-thiopseudourea), dihydrobromide	26
Tetramethylthiuram monosulfide	25
7-Theophylline acetamide (1,2,3,6-tetrahydro-1,3-dimethyl-2,6-diaxo- purine-7-acetamide)	55
1,3,4-Thiadiazole-2,5-dithiol	50
Thiazolidine-4-carboxylic acid	47
2,4-Thiazolidinedione	47
2,2'-Thio-bis-acetamide	19
Thiocarbamic acid, p-hydroxyphenyl ester	24
Thiocyanic acid, α -3, α -5-bis(dimethylamino)-4-hydroxy-3,5-xylyl ester	34
Thiocyanic acid, 4-hydroxy- α -3-(4-methyl-1-piperazinyl)-3,5-xylyl ester	53
Thiodiglycolic acid	7
3,3'-Thiodipropionic acid, bis(D-galactosylmethylenehydrazide)	37
Thioflavine S	8

	<i>Page</i>
α -Thioglycerol	7
Thioglycolic acid	7
Thioisonicotinamide	41
Thioisonicotinamide-1-oxide	41
Thiomalic acid	7
Thionine	48
2-Thio-6-oxypurine	55
1-Thiophenylanthraquinone	12
2-Thiophenylanthraquinone	12
Thiopyrazinamide	21
2-Thio-m-thiazone-2,4-dione	35
2-Thioureido-4-(p-nitrophenyl)thiazole	47
p-Toluenesulfanamide	20
N-p-Toluenesulfonyl, isoquinuclidylformamide	20
m-Toluidine	31
o-Toluidine	31
p-Toluidine	31
1-o-Tolyl-2-thiourea	27
m-Trifluoromethylphenylacetonitrile	34
N-(α , α , α -Trifluoro-m-tolyl)anthranilic acid; flufenamic acid	14
3,4,5-Trimethylacetophenonethiosemicarbazone	37
Trimethylamine isonicotinamide	41
2,2'-Trimethylene-bis(2-thiopseudourea), dihydrobromide	25
7,7'-Trimethylene-bis { 8-[bis(2-hydroxyethyl)amino]theophylline }	56
1,1,4-Trimethyl-4 { [2-(4-methyl-1-piperazinyl)ethylthio]methyl } piperazinium di-p-toluenesulfonate bis-(metho-p-toluene) sulfonate	53
Trisodium-2-(hydroxycyclohexyl)ethylenediamine triacetate	12

	<i>Page</i>
S-Trithiane	55
Vitamin B ₁₂	57
N-(2,3-Xylyl)anthranilic acid: mefenamic acid	14
Zinc derivative of 3-pyridinethiol	41

Unclassified

Security Classification

DOCUMENT CONTROL DATA - R & D

(Security classification of title, body of abstract and indexing annotation must be entered when the overall report is classified)

1. ORIGINATING ACTIVITY (Corporate author) USAF Radiation Laboratory University of Chicago Chicago, Ill.		2a. REPORT SECURITY CLASSIFICATION Unclassified
2b. GROUP		
3. REPORT TITLE A FURTHER SURVEY OF COMPOUNDS FOR RADIATION PROTECTION		
4. DESCRIPTIVE NOTES (Type of report and inclusive dates) Final report Jan. 1961 - Apr. 1965		
5. AUTHOR(S) (First name, middle initial, last name) Vivian Plzak John Doull		
6. REPORT DATE February 1969	7a. TOTAL NO. OF PAGES 82	7b. NO. OF REFS —
8a. CONTRACT OR GRANT NO 41(609)-2977	9a. ORIGINATOR'S REPORT NUMBER(S) SAM-TR-69-1	
b. PROJECT NO	9b. OTHER REPORT NO(S) (Any other numbers that may be assigned this report)	
c. Task No. 775703	d.	
10. DISTRIBUTION STATEMENT This document has been approved for public release and sale; its distribution is unlimited.		
11. SUPPLEMENTARY NOTES	12. SPONSORING MILITARY ACTIVITY USAF School of Aerospace Medicine Aerospace Medical Division (AFSC) Brooks Air Force Base, Texas	
13. ABSTRACT This report summarizes the results obtained with 617 compounds tested for their radioprotective activity in adult male mice irradiated with a control-demonstrated LD ₉₉₊ of 800 R (250 kvp) x-rays. A compound was considered to exhibit radioprotective activity if it (1) permitted any of the treated mice to survive for 30 days after the otherwise lethal whole-body x-irradiation or (2) increased the median survival time of treated animals by 5 days or more beyond the median survival time of the untreated control mice (9 ± 3 days). Of the 617 compounds tested, 245 successfully passed one or both of the stated specifications. Additionally, data are offered to allow comparisons of chemically related groups for structure-activity relationships, and to indicate the types of structures which offer the greatest promise as a source of more effective and less toxic radioprotective agents.		

DD FORM 1 NOV 65 1473

Unclassified

Security Classification

Unclassified

Security Classification

14. KEY WORDS	LINK A		LINK B		LINK C	
	ROLE	WT	ROLE	WT	ROLE	WT
Radiobiology Radioprotective agents Toxicity of radioprotective agents Radioprotective activity Structure-activity relationships						

Unclassified

Security Classification