This document has been approved for public teleans and salet its distribution is unlimited.

Detailed Study of Contaminant Production in a Space Cabin Simulator at 258 mm. Mercury

J. P. CONKLE, J. D. ADAMS, W. E. MABSON, P. WOLF, H. J. ZEFT and B. E. WELCH

USAF School of Aerospace Medicine, Brooks Air Force Base, Texas 78235

CONKLE, J. P., J. D. ADAMS, W. E. MABSON, P. WOLF, H. J. FT and B. E. WELCH. Detailed study of contaminant production in a space cabin simulator at 258 mm. mercury. Aerospace Med. 41(9):1031-1038. 1970. Additional information has been accumulated concerning the

Additional information has been accumulated concerning the contaminants associated with habitation by man of a closed confined space. An experiment designed to determine man's contribution to trace contaminants was conducted jointly by the United States Air Force and the National Aeronautics and Space Administration. The experiment was divided into three phases: an unmanned period, a manned period, and a manned period coupled with an activated carbon scrubber. Direct analyses of the scaled environment were not adequate for this comprehensive survey; however, cryogenic fractionation and conceptration provided samples with sufficient concentration of contaminants for analysis by means of gas chromatography, infrared spectroscopy, and mass spectroscopy. Of the 142 compounds identified and quantifies during the experiment, only 45 were found during the manned phases.

. .

THE DEVELOPMENT AND USE of analytical procedures to detect trace atmospheric constituents, coupled with experiments specifically designed for the appraisal of man as a production or removal system, or both, will serve to define further the problems associated with man's subsistence in closed-loop conlined spaces. These endeavors are of value for ascertaining physiologic as well as instrumental requirements placed upon life support systems to maintain human existence without incurring debilitating conditions that might jeopardize mission accomplishment.

Extensive reporting of material off-gassing products has been combined with computer retrieval systems to make available data on the contributions that could be made from materials to the atmosphere.^{2,10} Detailed reports have been published which elucidate the matrix

050405027

Reprinted from Aerospace Medicine, Vol. 41, No. 9, September 1970

generated by man and his environment in submarines, spacecraft, and ground-based simulators. Likewise, an earlier experiment specifically designed for trace contaminant study provided information concerning man's contribution to the atmosphere.¹

REPRODUCED FROM BEST AVAILABLE COPY

28

1970

The purpose of this paper is to present the results of a joint study conducted by the United States Air Force and the National Aeronautics and Space Administration designed to define within the limits of the analytical procedures^{5,9,12} those contaminants resulting from human habitation of a sealed environment in an oxygen atmosphere at 258 mm. Hg total pressure.

METHODS

An experiment was conducted in a test cell¹ maintained at a total pressure of 258 mm. Hg. The study was divided into three phases: an 8-day unmanned background period, a 14-day manned period, and a 7day manned period with an activated carbon scrubber. The initial phase established the atmospheric matrix resulting from the test cell supply gases, and previous occupancy by man. The second phase provided information as to the change in the atmospheric matrix by the inclusion of man into the system. The final phase demonstrated the effect of an activated carbon scrubber on the man-influenced atmospheric matrix. The gross atmospheric composition during the three phases is presented in Table I.

Experimental Subjects—Four male Air Force volunteer subjects, ages 19-20 years, participated in the study: All of the subjects were in good physical condition, hav-

TABLE L	GROSS	ATMCSPHERIC	COMPOSITION
---------	-------	-------------	-------------

Major Atmospheric Components		Unmanned	Manned	Manned- phus- Charcoal
Oxygen	Maximum	98.8%	96.2%	92.0%
	Minimum	96.0%	90.0%	89.9%
	Mean	96.7%	92.7%	91.0%
CO:	Maximum	0.97%	1.62%	1.47%
	Minimum	0.0%	0.0%	0.15%
	Mesn	0.07%	06%	0.69%
Nitrogen	Maximum	2.8%	6.35%	6.90×4
	Minimum	0.35%	2.4%	4.35%
	Mean	1.96%	4.74%	6.47%
Total pressure	mean	256 mm. Hg	256.7 mm. Hg	237.8 mm, He

0

NATIONAL TECHNIC INFORMATION SERV Spermaneu, va. 22131

٦

6

The research reported in this paper was conducted by personnel of the Environmental Systems Division, USAF School of Aerospace Medicine, Aerospace Medical Division, AFSC, United States Air Force, Brooks AFB, Texas. This study was partially supported by NASA MIPR R-89. The voluntary informed consent of the subjects used in this research was obtained as required by AFR 169-8. Further reproduction is authorized to satisfy the needs of the U. S. Government.

Maj. W. E. Mabson, USAF, BSC, is in the AF Surgeon General's Office, Headquarters USAF; Washington, D. C. Capt. H. J. Zeft, USAF, MC, is in the Cardiovascular Labora-

Capt. H. J. Zeft, USAF, MC, is in the Cardiovascular Labora tory, Duke University Medical Center, Durham, N. C.

REPRODUC _D FROM BEST AVAILABLE COPY

CONTAMINANT PRODUCTION IN SPACE CABIN SIMULATOR-CONKLE, ET AL.

ing recently completed basic training. As in the previous study,¹ only individuals who produced methane in their flatus were selected. The flatus of the subjects was collected after the manner of Kirk^a and analyzed by gas chromatography. The physical characteristics of these individuals are presented in Table II. The subjects underwent complete physical examinations prior to entering the test cell and after completing the 21day experimental phase. The physical examination included ophthalmologic, ear-nose-throat, neurologic, and psychiatric evaluations. In addition, standard chest, abdominal, and paranasal sinus roentgenographs, electrocardiograms, and electroencephalograms were obtained. Throughout the entire 21-day manned phase and for four days pre- and post-experimentally, the subjects were maintained on a total liquid diet," ad libitum water intake, and a daily vitamin-mineral supplement** with an additional 26 mEq of potassium chloride in the enteric coated form. The 24-hour intake was adequate in all necessary dietary constituents.

The subjects were taught to obtain vital signs and to perform venipunctures on one another. Vital signs were measured daily and venipunctures were performed three times weekly. Blood, urine and feces samples were sent to the outside through a small passlock without altering the environmental conditions. A complete medical review of systems was performed by a physician at least once daily through an intercommunication system.

Test Cell Operation-The test cell was provisioned with food and other logistical support items before the unmanned background portion of the experiment was initiated. During the 21-day manned portion of the test, the fecal material was stored in plastic bags in a freezer before being removed from the test cell. Urine which was not retained for analysis was removed via a water trap from the test cell. The trap was well flushed with water following this procedure. The subjects did not shave and their hygienic activity was limited to dry brushing of the teeth and bathing with a damp cloth which was dried and stored in the test cell. Canisters containing approximately 14 kilograms of lithium hydroxide were used for carbon dioxide removal. These canisters were changed on an average every three days. The test cell had a double-walled shell, and a special control circuit was used to maintain the pressure of the annulus at the same pressure as the interior of the cell. This mode of operation provided a diffusion barrier be-

 Prepared by U. S. Army Natick Laboratories, Natick, Mass.
ABDEC with Minerals, Parke-Davis and Company, Detroit, Michigan.

TABLE II. PHYSICAL CHARACTERISTICS OF SUBJECTS

Subject	Age	Height	Weight
Number	(years)	(cma.)	(kg)
 90	19	184	76.7
91	20	170	68.0
92	19	188	79.4
93	20	176	77.1

1032

HERE MALL MATCHER MAL

ないてありまた

Ī.

記載

Aerospace Medicine • September, 1970

tween the ambient atmosphere and the test cell which prevented the loss of gas from the test cell and contamination from the ambient atmosphere. The portion of the test cell used during the study had a bound volume of 27.45 m.³

The background portion of the study was initiated by flushing the test cell with oxygen from a liquid source while the cell was being evacuated to 258 mm. Hg. This phase of the test was continued for 185.5 hours. The manned portion of the test was begun immediately following the background phase. The subjects entered the test cell through the airlock which was flushed with oxygen from a liquid source as it was reduced to the pressure of the test cell. The manned portion of the experiment continued for 503 hours. The final 172 hours of the study had a 454-gram bed of activated carbon (used without pretreatment from delivery containers) in operation with an average air flow through the bed of 0.07 m³/min.

Gas Analyses-A dual-flame gas chromatograph with two 3.05 meters by 3.2 millimeter columns packed with 4% Carbowax 4000 and 6% polyphenyl ether on 60-80 mesh Teflon was used to monitor the atmosphere of the test cell for contaminants in concentrations in excess of 10 p.p.m. Test-cell gas samples were introduced on a batch basis into the chromatograph from a sample loop which was continuously flushed with the test-cell gas. Methane concentrations were determined with a flame ionization gas chromatograph by analysis of batch samples obtained on a periodic basis. Carbon monoxide was evaluated using a Beckman IR-7 with a 10-meter multipath cell and a "Lira" 300 analyzer sensitized for carbon monoxide. Hydrogen was monitored with a Bendix 17-210 Time-of-Flight mass spectrometer. The techniques used were similar to those used in a previous study.1

Conventional techniques of analysis, such as gas chromatography, and infrared and mass spectrometry of unconcentrated samples, would have revealed only 5 to 10 of the 142 trace constituents reported; therefore, cryogenic trapping was utilized as the best and most practical method of obtaining concentrated samples for subsequent analysis. As in the previous study,¹ multistage cryogenic trapping systems were operated 20 to 22 hours per day. The samples were distributed to three analytical laboratories for detailed chemical analysis. The analyses were provided by the Von Karman Center of Aerojet-General Corporation,⁹ Arnold Engineering and Development Center,³ and Melpar, Inc.¹²

The average concentration of a compound in the chamber during a cryogenic trapping period was estimated from the total milligrams of the compound contained in the cryogenically obtained sample, the mass flow rate of the sample gas through the system, and the total trapping time. The total flow through the system in each trapping period was obtained by multiplying the average mass flow rate through the system by the time of the trapping period in minutes. The mass flow-meters used in the determination had an accuracy of one percent. The concentration in the test cell of the contaminant in mg/m³ was determined at 21.1°C. and 760 mm. Hg by the following equation:

REPRODUCED FROM **BEST AVAILABLE COPY**

VON KARMAN GENTER

11000

021

.17

100

VON KARMAN CENTER

.37

.14

083

ì

ELPAR

.43

11

WELPAR INC.

12

CONTAMINANT FRODUCTION IN SPACE CABIN SIMULATOR-CONKLE, ET AL.

A CONTRACTOR OF A CONTRACTOR

contaminant weight × 10¹ = milligrams/meter^a in milligrams

Where,

COMPOUND BY CHEMICAL CLA

COMPOUND BY HEMICAL CLASS PSEUDOCUMENE

The second second

milligrams

contaminant weight in an the amount of material in a sample as reported by an analysis group. = volume of gas passed

through the multistage cryogenic trapping system determined on a mass flow basis at 760 mm Hg and 21.1°C.

The observed concentration of a compound in mg/ m^a was corrected by a computer program for the

amount of each compound removed from the test cell and the addition of compounds from the oxygen supply.

Since the volume used in the calculations was measured with a mass flowmeter calibrated to read standard cc./minute at 760 mm. Hg rather than at 258 mm. Hg, the volume processed at the test-cell conditions was 2.95 (760/258) times greater. The concentrations in the inspired volume must be considered for all compounds (Figure 1) which are reported on a mass flowdetermined volume at 760 mm Hg and 21.1°C. Man's inspired volume at 258 mm. Hg is about the same as that at ground level;" therefore, in order to assess the physiological significance at 258 mm. Hg, the reported concentrations must be multiplied by 0.359.

Clinical Analyses-Numerous venous blood and urine studies were performed throughout the experimental phase and for at least one week before and after the ex-

				0 - Avis - 444, 753	ARNOLE	MELPAR	VON	OF STUDY GROUPS	ARNOLD	
OF PHILSE GROUPS	ARNOLD	MELPAR	VON KARMAN	COMPOUND BY	101	INC.	CENTER	COMPOUND BY	EA.	
COMPOUND BY		· 77	EZ DER	CHEMICAL CLASS		-	1917	ALDEHYDES		ŀ
KORGANIC COMPOUNDS NYDROGEN	[.		n	ise - PROPTL	22.0	-lin ha	18.	AROMATIC HYDROCARBONS BENZENE	.66	ŀ
NITROUS OXIDE		•	17	ise - BUTYL		0035	1 040	TOLUENE	1.44	f
DRGANIC ACIDS			1091	# - AMYL		1		STYRENE		ľ
PROPIONIC			11	FURFURYL		1000	073	XYLENE		#
YALERIC				CYCLOHEXANOL	20		•	ETNYL BENZERE	0175	ŀ
NLCOHOLS WETHYL	1	047		2 - MEXANOL	0045			CUMENE	<u>.</u>	ſ
ETNYL	- 114.	L	1	ALDENYDES ACETALDENYDE	147		7	WESITYLENE	070	ŀ
ALLYL				BUTYRALDEHYDE			817	*-PROPYL BENZENE		ľ
1	١.			<u> </u>	6			(:	İm
ANS STORES	ARNOLD	NELPAR INC.	VGN RARSAN GENTER	or this endures	ARHOLD	NELPAR INC.	VGN KARSMAN CENTER	OF STAR GROUPS	ARNOLD	
OMPOUND BY				COMPOUND BY				COMPOUND BY CHEMICAL CLASS		12
PSEUDOCUMENE		- 19 - 1		AROMATIC HYDRO CARBONS METHYL NAPHTHALENE			1000	BUTYL ACETATE		1
METHYL ETHYL BENZENE				DIMETHYL NAPHTHALERE			.012	ANTL ACETATE		
RAPHTRALENE			114	ANINES DINETNYL ANINE			24	ETNERS FURAN		
- BUTTL BENZEME				HIGHER AMINES			017	TETRAHYDROFURAN		
SOC-BUTYL BENZENE				ESTERS ETHYL FORMATE			00045	ETHYL ETHER	073	
tert - BUTYL BENZENE	."]			METHYL ACETATE	900034	041	**	METHYL FURAN		
9 - CYMENE	1.1.			ETHYL ACETATE		1	1000	DIMETHYL FURAN		
DIETHYL BENZENE		1		METHYL BUTYRATE				160 - PROPYL ETHER		-
TETRAHETHYL BENZENE			010	184 - PROPYL ACETATE			00073	BENZYL ETHER		
r)			B	7			12)	-

Fig. 1. Histogram of observed compounds. (For explanation see text.)

1033

REPRODUCED FROM EEST AVAILABLE COPY

CONTAMINANT PRODUCTION IN SPACE CABIN SIMULATOR-CONKLE, ET AL.

periment. Blood studies were performed three times weekly. Freshly voided morning urine specimens were analyzed for sugar, protein, acetone, specific gravity, and pH by standard laboratory methods. A careful microscopic examination was performed on the sediment. All stool samples were tested for occult blood with guaiac reagent. Complete blood counts, including hemoglobin and hematocrit determination, reticulocyte counts, white blood cell counts, and differential counts were performed by standard techniques. Serum and urine creatinines, fasting blood glucose, blood urea nitrogen, and serum alkaline phosphatase were determined by AutoAnalyzer techniques. Serum protein electrophoresis was performed with the Spinco Model R paper electrophoresis system and total serum proteins were measured by a modified biuret method.¹⁴ Serum glutamic-oxaloacetic transaminase (SCOT) and serum glutamic-pyruvic transaminase (SGPT) were ascer-

tained by the Sigma-Frankel procedure¹³ and serum bilirubin by the method of Malloy and Evelyn.⁶ Bromsulphalein retention was determined before and after the experiment utilizing standard dye injection techniques with a 45-minute venous blood measurement.

RESULTS

Atmospheric Gas Analyses—One hundred and fortytwo (142) different compounds (Figures 1 and 2) were reported by the analytical groups during the experiment. The data (Figure 1) are presented in mg/m³ in the cest cell at 21.1°C. and 760 mm. Hg corrected for the amount of a compound added to and removed from the test cell during the experiment. The greatest concentration, represented by the highest peak, appears at the top of each histogram with the base being equiva-

COMPOUND BY		CENTER	COMPOUND BY		CENTER	COMPOUND BY	122	ICE
CHEMICAL CLASS			CHEMICAL CLASS	1.05004		CHEMICAL GLASS		
AROMATIC AND CYCLIC MYDROCARDONS CHLOCORDONS	"	1912 	FREON 113	11	2 47	OF BETHANE CHLOROFORM	ه فسيا	000
TRIFLUOROBENZENE		10037	HALDGEN DERIVATIVES OF ETHYLENE VINYL CHLORIDE		1	FREON 12		20
TETRAFLUOROBENZENE		.0014	1,1-DICHLOROETHYLENE	10054 .00080	4	FREON 11		pors a
NEXAFLUGROBENZENE		.0017	TRICKLORGETHYLEIN	37	045	CARBON TETRACHLORIDE	K.	
NALOGEN DERIVATIVES OF ETHANE	. 4.1		PERCHLOROETHYLENE			INDOLE INDOLE		.012
1,2-DICHLOROETHANE		12	NALDGEN DERIVATIVES OF METHANE METHYL CHLORIDE	.032		KETONES ACETONE	130	1.1 1.4 A
NETHYL CHLOROFORM	1 14		METHYLENE CHLORIDE			METNYL ETNYL KETONE	"	1
TETRACHLOROETHANE	.081		FREON 22	.839		2,4 - PENTANEDIONE	860000	
FREON 114	.9017		FREDR 13		1.4	METNYL 100-SUTYL XETOHE	X	
COMPOUND BY		CENTER .	COMPOUND BY			CONPOUND BY		5728 272
KAPHTNERES	1000034		RAPHTHERES	100	54 090 j	STICKING CLASS	000534	<u>محملة المارم</u>
CYCLOPENTANE			HIDENE			2,2-DHETHYL BUTANE		1
HETHYL CYCLOPENTAME	055	1.25	DECALIN	inter a second	010	2,3-OMETHYL BUTANE		*
			PARAFFINS	I ·				
CYCLOHEXANE	10		WETHARE	23 0000		VEXAME	000091	
CYCLOHEXANE DIMETHYL CYCLOPENTANE	10	029	WETHANE	23 0000 a.a.b.a. 0072 0019		VEXANE 2-METHYL PENTANE	000031	
CYCLOHEXANE DIMETHYL CYCLOPENTANE METHYL CYCLONEXANE	017 9005	020 034 034	NETMARE ETMARE PROPANE	23 00m 0072 0019	10	HEXANE 2-METHYL PENTANE 3-METHYL PENTANE	0024	• • • • • •
CYCLOHEXANE DIMETHYL CYCLOPENTANE METHYL CYCLONEXANE DHMETHYL CYCLONEXANE) 0 017 00 00	070 034 034 0071 011	BETHARE ETHARE PROPARE BUTARE	23 0000 0072 0010	10	HEXANE 2-METHYL PENTANE 3-METHYL PENTANE 2,3-DMETHYL PENTANE	000051 0024 041 14	
CYCLOHEXANE DIMETHYL CYCLOPENTANE METHYL CYCLOHEXANE DHMETHYL CYCLOHEXANE 1,2 - DIMETHYL CYCLOHEXANE (cm)	017 0049 045	070 020 034 034 034	NETHARE ETHARE PROPARE BUTARE PERTARE	23 0000 	10 L	WEXAME 2-WETHYL PENTANE 3-WETHYL PENTANE 2,3-DIMETHYL PENTANE 2,4-DIMETHYL PENTAME	0000051 00224 041 14 	
СУССОНЕХАНЕ DIMETHYL CYCLOPENTANE METHYL CYCLOMEXANE DHMETHYL CYCLOMEXANE 1,2 - DIMETHYL CYCLOMEXAME (cro) 1,2 - DIMETHYL CYCLOMEXAME (roam)	007 0099 .0 009 .0 099 .0 099		NETNARE ETHARE PROPARE BUTARE PERTARE 150 - PERTARE	23 0000 0072 0079 	10 - 10	HEXAME 2-WETHYL PENTAME 3-WETHYL PENTAME 2,3-DIMETHYL PENTAME 2,4-DIMETHYL PENTAME MEPTAME	000091 0024 641 14 14 	00021 .081
СУССЮНЕХАНЕ DIMETHYL CYCLOPENTANE METHYL CYCLOMEXANE DIMETHYL CYCLOMEXANE DIMETHYL CYCLOMEXANE T,2-DIMETHYL CYCLOMEXANE (1100) ETHYL CYCLOMEXANE	0000 0000 0000 0000 0000 0000 0000 0000		МЕТНАЛЕ ЕТНАЛЕ РПОРАЛЕ ВИТАЛЕ РЕМТАЛЕ 150 - РЕПТАЛЕ DIMETHYL PPOPANE	23 0000 0072 0078 19 		HEXAME 2 - METHYL PENTANE 3 - WETHYL PENTANE 2,3 - DIMETHYL PENTANE 2,4 - DIMETHYL PENTANE HEPTAME 3 - METHYL NEXAME	0000051 0010 001 001 001 001 001	
СУСС. ОНЕХАНЕ DIMETHYL CYCLOPENTANE METHYL CYCLOMEXANE DIMETHYL CYCLOMEXANE DIMETHYL CYCLOMEXANE T,2 - DIMETHYL CYCLOMEXANE (111200) ETHYL CYCLOMEXANE	0045 044 043		ИЕТНАЛЕ ЕТНАЛЕ РПОРАНЕ ВИТАЛЕ РЕКТАЛЕ 150 - РЕКТАЛЕ DIMETHYL РРОРАНЕ		10 	HEXAME 2-METHYL PENTAME 3-METHYL PENTAME 2,3-DIMETHYL PENTAME 2,4-DHMETHYL PENTAME HEPTAME 3-METHYL NEXAME	0000051 0024 004 004 14 14 0023 14 0023 14 0023 14 0023 14 0023 14 0023	

REPRODUCED FROM BEST AVAILABLE COPY

CONTAMINANT PRODUCTION IN SPACE CABIN SIMULATOR-CONKLE, 5T AL.

lent to zero. Time is represented on the horizontal portion of the histogram with the unmanned phase ending at 185.5 hours, the manned phase at 516.5 hours, and the manned-plus-charcoal phase at 688.5 hours. The frequency of occurrence of each compound is approximate since an interpolation between data points was necessary in the computer program used for concentration corrections. Additionally, the corrected values in Figure 1 are approximately 10% greater than the observed values

Arnold Engineering and Development Center detected 73 compounds during the experiment of which all but carbon dioxide are shown in Figure 1. Six compounds were reported only during the unmanned phase, and 16 compounds during the manned or manned-pluscharcoal phases. The other 50 compounds were reported both in the unmanned phase and in one or both of the manned phases.

いたい かいたい おおからのためをえる

いいとかいでいています

A LINE CONTRACT

2 1214 200

Ļ

ž

OF ATHS GROUPS	ARNOLD	MELPAR INC.	YON KARMAN CENTER
COMPOUND BY CHEMICAL CLASS		122	22
PARAFFINS TRIMETHYL BUTANE			54
OCTANE			
no - OCTANE	1.117		
2,3,4 - TRIMETHYL PENTAME	1100.		012
n - NORANE			
TRIMETHYL HEXAME			00000
2,2,5-TRIMETHYL HEXARE	000083		
DECAME	. أسل	BOORT	
SULFIDES SHIETNYL, SULFIDE		20012	

M

Melpar, Inc. detected 48 compounds during the experiment of which all but carbon dioxide and ethyl benzene are shown in Figure 1. Ethyl benzene is not included here because its quantities, appearing in the manned-plus-charcoal phase only, were insufficient to be handled by the computer program. Two compounds were reported only during the unmanned phase, and 10 compounds during the manned or manned-plus-charcoal phases. The remaining 34 compounds were reported in both the unmanned phase, and in one or both of the manned phases.

Von Karman Center of Aerojet-General Corporation reported carbon dioxide and the 92 compounds presented in Figure 1. Two compounds were detected only during the unmanned phase, and 51 compounds during the manned or manned-plus-charcoal phases. The other 39 compounds were reported in both the unmanned phase and one or both of the manned phases.

Clinical Observations-The four experimental subjects tolerated the environmental phase without difficulty. Complaints were minimal and most were centered on the diet and relative boredom. Subject 93 experienced mild aches in the knee joints during the initial 12 hours, which disappeared with no further complications. His symptoms were interpreted as a transient, grade 1, decompression disturbance. Subjects 90 and 93 had minimal discomfort from pressure changes in the middle ear. The need for clearing their ears was manifested primarily during the sleeping hours. No further subjective symptoms were elicited during the experiment.

Subjects 90 and 93 exhibited abnormal urinary sediments on the fourth post-experimental day. The sediment from subject 90 showed numerous coarsely granular casts (2-4 per low-power field) and occasional hyaline casts. There was no proteinuria and only an oc-casional white blood cell. These abnormalities cleared completely over the next four days. The sediment from subject 93 exhibited numerous red blood cells (75-100 rbe per high power field) without casts, proteinuria, or

OF STUDY SHALLY SY ARMO	LD HELPAR KARMAN	OF MASS GROUP	ARNOLD MELPAR	VON KARMAN CENTER	OF PRIES GROUPS	ARNOLD	MELPAR INC.	KAS CEN
COMPOUND BY		COMPOUND BY			COMPOUND BY			123
CARBON DISULFIDE	037	OLEFINS 2-METHYL-2-BUTENE	.000034		2-OCTENE	1.0 .1		
SULFUR COMPOUNDS METHYL MERCAPTAN	00041	2-ETHYL -1-BUTENE	.016		2,4,4-TRIMETHYL-1-PENTENE	1.6	Į .	
PROPYL MERCAPTAN	7004	2-HEXERE			2,4,4-TRIMETHYL-2-PENTENE	024		
THIOPHENE	210	2-WETHYL-1-PENTENE			DIOLEFIN ISOPRENE			ľ
THOTOLENE		4-BETHYL-2-PERTENC	10040		CYCLOQLEFINS CYCLONEXENE	1)." 1		٥
THIOMENE	1000	HEXENE		00044	CYCLOPENTENE	0000 70		.004
METHYL ETHYL THIOPHENE	, 2034	R-HEPTENE	10E7	1.27	METHYL CYCLOHEXENE	.017		
OLEFINS ETHYLENE		5-HEPT HE	238					
PROPYLENE	20	1-OCTENE	.00024		·			
N			0		1	P		

Aerospace Medicine • September, 1970

CONTAMINANT PRODUCTION IN SPACE CABIN SIMULATOR-CONKLE, ET AL.

pyuria. Urinalysis repeated 2⁴ hours later was completely normal. Urinalyses on these subjects during the experimental phases were normal and numerous urine cultures were negative. It is noteworthy that approximately 16 hours prior to the abnormal post-experimental urinalyses, both subjects exercised moderately—their first exertion of any note since the confinement. All subjects displayed normal ability to concentrate their urine after only an 8-10 hour period of dehydration. Blood urea nitrogen, serum creatinine and endogenous creatinine clearances showed no deviations.

れたないである

Subject 92 exhibited mild hyperbilirubinemia with slight elevation of the indirect fraction before the experiment. This was associated with a reticulocytosis of three percent and was probably a manifestation of a mild hemolysis or possible Cilbert's syndrome. Otherwise, liver function studies were entirely normal.



Fig. 2. Carbon monoxide concentration corrected for gas lost from the test cell. The concentrations of carbon monoxide were corrected for the amount removed from the test cell and reported as mg/m³ at 258 mm. Hg and 21.1°C.



DISCUSSION

Variance in the analysis of similarly obtained samples may result in the misinterpretation that man does or does not produce certain compounds. Many explanations may be proposed for the inconsistency of data: (1) variabilities may occur in obtaining the samples with the cryogenic trapping system, (2) condensed compounds in the trapping cylinders may alter the thermal characteristics of the cylinders, (3) interaction between molecular species could increase or decrease the efficiency of the concentration of material, (4) each analytical group developed an independent procedure for analysis, (5) each analytical group had a number of unidentifiable compounds which may have been identified by another group, (6) compounds existed in very low concentration in the test cell or supply gases, and did not reach a detectable concentration until the manned phase of the experiment, (7) compounds reported in the last phase of the experiment (mannedplus-charcoal) may have been desorbed from the activated carbon, and (8) a compound identified only by gas chromatographic elution times may be two or more compounds having the same elution time. Furthermore, variations from the absolute values of contaminants were introduced by the operation of the test cell, controlling pressure, temperature, water vapor content, and in changing the lithium hydroxide sorption canisters. Large surface areas are available within the test cell for the sorption and desorption of compounds. Man also adds a variability factor. Man's efficiency as a contaminant removal system may change, and his contaminant production may be related to physical activity and biochemical needs.

Sixty-six (66) compounds were identified during the 258 mm. Hg experiment that were not reported in the previous study conducted at 760 mm. Hg.¹ These compounds could have been present in the earlier experiment but were not reported due to the state-of-the-art of the analytical techniques at that time. As a result of



Fig. 3. Hydrogen concentration corrected for gas lost from the test cell. The hydrogen concentrations were corrected for the amount of gas removed from the test cell and reported as mg/m^{-1} at 258 mm. Hg and 21.1°C.

1036 Aerospace Medicine • September, 1970



REPRODUCED FROM BEST AVAILABLE COPY

CONTAMINANT PRODUCTION IN SPACE CABIN SIMULATOR-CONKLE, ET AL.

the experience gained from the previous study, the analydcal groups improved their techniques for the analysis of cryogenically obtained samples. The experience gained as a result of this experiment will enable an even more detailed identification of the components of closed-system atmospheres.

Forty-five (45) of the 142 compounds were reported only during the manned and manned-plus-charcoal phases of this experiment. Of these compounds found during both manned phases, the following 10 were detected only after the carbon scrubber had been put into operation: valeric acid, n-propyl benzene, higher amines, benzyl ether, trifluorobenzene, tetrafluorobenzene, hexafluorobenzene, dimethyl propane, methyl mercaptan, and 2-methyl-2-butene. These compounds are subject to the same limitations imposed during the manned phase with the exception of the three fluorobenzenes; the latter compounds appeared in the last two days of the experiment which suggests that they may have been associated with the activated carbon.

The data, within the limits imposed by the experimental methods, suggest that the following compounds are associated with man: hydrogen, carbon monoxide," acetic acid, * propionic acid, * valeric acid, * methyl al-cohol, * ethyl alcohol, * n-propyl alcohol, furfuryl alcohol, acetaldehyde, butyraldehyde,* furfural, mesitylene,* methyl ethyl benzene, tert-butyl benzene, diethyl benzene, sec-butyl benzene, cumene, dimethyl amine, higher amines, ethyl formate, methyl n-butyrate,* isopropyl acetate, furan,* methyl furan, benzyl ether,* tetrahydrofuran, indole, acetone, cyclohexane, 1,2-dimethyl cyclohexane (cis), 1,2-dimethyl cyclohexane (trans), decalin,• methane,• butane, 2,3-dimethyl butane,• oc-tane, 2,4-dimethyl pentane, decane, 2-methyl pentane, trimethyl butane, 2,3,4-trimethyl pentane, 2,2,5-trimethyl hexane, n-nonane, dimethyl sulfide, methyl mercaptan, propyl mercaptan,* methyl ethyl thiophene, thiophene, thiotolene, thioxene, ethylene, propylene, 4-methyl-2pentene, n-heptene, 2-methyl-1-pentene, 2,4,4-trimethyl-1-pentene, hexene, 3-heptene, 1-octene, 2-octene, isoprene,* and cyclopentene.

Carbon monoxide, hydrogen, and methane increased rapidly with the inclusion of man into the test cell. The respective graphs (Figures 2, 3, 4) of the test cell concentrations, corrected for gas lost from the test cell (samples for analysis and lock operations), indicate an increase in production rate for methane and a slight decrease in the production rates of carbon monoxide and hydrogen during the manned-plus-charcoal phase of the experiment. The production rates (0°C. and 760 mm. Hg) of CO, CH₄ and H₂ during the manned phase were 0.54, 1.7, and 40.9 ml/man/hr, respectively. The values for hydrogen production are in accord with data reported by Kirk.⁴

٦.

Up to this point, the present data have been interpreted with respect to man as a source of compounds. Examples of man as a compound-removal system include the following: iso-p opyl alcohol, toluene, and methyl iso-butyl ketone.

No obvious ill effects attributable to trace contaminants were discovered in the four experimental subjects who lived for 21 days in the confined environment. Minimal bends and barotalgia have been reported in previous studies of 100% oxygen at 258 mm. Hg total pressure.⁴ Helvey et al.³ noted abnormal urinary sediments from individuals living in this environment for 14 days, with the abnormality persisting for as long as 3 months. Other environmental studies by Mautner and co-workers⁷ have revealed mitochondrial alterations in the proximal tubule of the monkey kidney as a result of the hyperoxic hypobaric atmosphere. In this study, the detectable but transient urinary abnormalities might possibly have been secondary to hyperoxia and less likely a result of toxicity from contaminants.

Conceivably, accumulation of trace amounts of organic compounds might have had an additive effec' and be harmful to certain body systems. Minimal deviations in liver function on single determination made before and after the experiment were observed in a previous study of contaminants at ambient conditions.¹ However, detailed studies on liver function in the present work revealed no deviations during or after the environmental phase. Likewise, absence of respiratory symptoms and clear lung fields by physical examinations and chest X-ray are evidence against pulmonary difficulties. Although carbon monoxide accumulation was present, no harmful effects from this compound were detected by careful medical observations, physical examinations, and post-experimental electroencephalograms.

CONCLUSIONS

Additional information has been accumulated concerning the contaminants associated with habitation by man of a closed confined space. However, the interaction of the many contaminant removal systems (such as man, lithium hydroxide, activated charcoal, water condensation, and the walls of the test cell) modified the concentration of the contaminants preventing meaningful production rate determinations. The large volume of the test cell buffers the results, indicating the need to conduct experiments with the developed techniques and analytical procedures with a single man in a minimum volume chamber containing very few support items.

ACKNOWLEDGMENT

The authors wish to express their appreciation to the cryogenic system operators, chamber operators, and subjects. Appreciation is also expressed to Harold N. Keiser of the Biometrics Division and Sgt. Charles A. Bond of the Environmental Systems Division for data reduction. The pressure control circuit used in the experiment was designed and built by Doyle White of the Biomedical Engineering Branch. Capt. Edward E. Lefebvre and Lt. J. T. Watson of the Environmental Systems Division provided assistance in CO determination and the activated carbon system, respectively.

Aerospace Medicine • September, 1970 1037

^{*}Compounds previously found to be suggestive of association with man in a closed system at 760 mm. Hg_1^{-1}

NY PRIMA IN THE CONTRACT OF THE PARTY OF

REFERES 285

- Consult of The MABON, J. D. ADAMS, H. J. ZEFT and B. E. 1975. 201: A detailed study of contaminants produced by man his space orbin simulator at 760 mm. Hg. SAM-TR-67-10; March 1967.
- 2. HARRIS, E. S.: Parts and materials data retrieval program relative to impterials selection in toxicology. AMRL-TR-66-120, December 1966.
- HELVEY, W. M., G. A. ALBRICHT, F. B. BENJAMIN, L. S. GALL, J. J. PETERS and H. RND: Effects of prolonged exposure to pure oxygen on human performance. Republic Aviation Corp. RAC 393-1 (ARD-807-701), November 1962.
- HERLOCHER, J. E., D. G. QUICLEY, V. S. BEHAR, E. G. SHAW and B. E. WELCH: Physiologic response to increased oxygen partial pressure: I. Clinical observations, Aerospace Med. 35:613, 1964.
- 5. KIRK, E.: The quantity and composition of human colonic flatus. Gastroenterology 12:782, 1940.
- 6. MALLOY, H. T., and K. A. EVELYN: The determination of bilirubin with the photoelectric colorimeter. J. Biol. Chem. 119:481, 1937.
- MAUTNER, W.: Electron microscopic investigations of oxygen effects on kidney tissue. Proceeding of the 2nd Annual Conference on Atmospheric Contamination in Con-

- fined Spaces, Wright-Patterson AFB, Ohio. AMRL-TR-659120, December 1966.
- MCCABE, J. R.: A trace contaminant analysis test on pir samples: Phase II. AEDC-TR-67-19, February 1967.
- 9. MOBERG, M. L.: Analyses of trace contaminants contained in samples from a closed environment at 258 mm. Hg. SAM-TR-67-6, February 1937.
- PUSTINGER, J. V., JR., F. N. HODGSON and W. D. Ross: Identification of volatile contaminants of space cabin materials. AMRL-TR-66-53, June 1966.
- ROBERTSON, W. G., J. J. HARGREAVES, J. E. HERLOCHER, et al.: Physiologic response to increased oxygen partial pressure: II. Respiratory studies. Aerospace Med. 35: 618-622, 1964.
- DESCHMERTZING, H., S. S. NELSON and H. G. EATON: Cryogenically trapped trace contaminants analyzed by ionizing gas chromatography. SAM-TR-37-68, August 1987.
- Sigma-Frankel Procedure. Sigma Technical Bull., No. 505, pp. 1-10, January 1981.
- WOLFSON, W. Q., C. COHN E. CALVERY and B. S. ICHIBA: Studies in serum proteins: V. A rapid procedure for the estimation of total protein, true albumin, total globulin, alpha globulin and gamma globulin in 1.0 ml. of serum. Amer. J. Clin. Path. 18:723, 1948.

REPRODUCED FROM BEST AVAILABLE COPY

ACCESSION	i far	
ersn	WHITE SECTION	11 6
000	BUFF SECTI	
UNAN	CES	
JUSTIFICAT	NON	
87	IROR/AVAILABILITY (200E3
MUL.	AVAIL and/or \$P	EGIAL
/	20	

1038 Aerospace Medicine • September, 1970

REPRODUCED FROM BEST AVAILABLE COPY

Security Classification	n i sheritikan n		
DOCUMENT CON	TROL DATA - R & D		* * * * * * * * * * * * * * * * * * *
(Security classification of title, body of abstract and indexing	ennotation must be entere	d when the ove	ell report la classified)
USAF School of Aerospace Medicine		Inclassif	on
Aerospace Medical Division (AFSC)	26.	GROUP	
Brooks Air Force Base, Texas	م ÷ الم مراجع ال		a transformation and the second se
REPORT TITLE			
Detailed study of contaminant production mercury	in a space cab	in simula	tor at 258 mm.
DESCRIPTIVE NOTES (Type of report and inclusive dates)		· · · · · · · · · · · · · · · · · · ·	
J. P. Conkle, J. D. Adams, Major W. Captain H. J. Zeft, and B. E. Welch	E. Mabson, Ca	ptain P.	Wolf,
REPORT DATE	78. TOTAL NO. OF PA	GTS 70.	NO. OF REFS
May 1970	8		14
	W. URIGINATOR'S RE	-UKI NUMBER	. . .
. PROJECT NO. 7930	SAM-TR-7	0-263	
.Task No. 793004	9b. OTHER REPORT N this report)	O(5) (Any other	numbers that may be seeigned
Work Unit No. 793004-002			
This document has been approved for pulunlimited.	blic release and	l sale; it	distribution is
. SUPPLEMENTARY NOTES	12. SPONSORING MILIT	ARY ACTIVIT	
Aerospace Med. (1(9):1031-1038 (1970)	USAF School	of Aeros	pace Medicine
	Aerospace M	eaicai Di	vision (AFSC)
ABSTRACT	1 DIOUKS AT D.	IEXas	· · · · · · · · · · · · · · · · · · ·
Additional information has been accu	mulated concern osed confined s	ning the c pace. An minants v	ontaminants experiment
associated with habitation by man of a cl designed to determine man's contribution jointly by the United States Air Force an Administration. The experiment was div period, a manned period, and a manned scrubber. Direct analyses of the sealed comprehensive survey; however, cryoger samples with sufficient concentration of chromatography, infrared spectroscopy, pounds identified and quantified during th the manned phases. (to trace contained the National A vided into three period coupled we environment we nic fractionation contaminants fo and mass spec e experiment, o	eronautic phases: with an ac ere not ac n and con r analysi troscopy only 45 w	as conducted, as and Space an unmanned stivated carbon lequate for this centration provided s by means of gas Of the 142 com- ere found during
associated with habitation by man of a cl designed to determine man's contribution jointly by the United States Air Force an Administration. The experiment was div period, a manned period, and a manned scrubber. Direct analyses of the sealed comprehensive survey; however, cryoger samples with sufficient concentration of chromatography, infrared spectroscopy, pounds identified and quantified during th the manned phases.	to trace contained the National A wided into three period coupled of environment we nic fractionation contaminants fo and mass spec e experiment, of	eronautic phases: with an ac ere not ac a and con r analysi troscopy only 45 w	as conducted, as and Space an unmanned tivated carbon lequate for this centration provided by means of gas Of the 142 com- ere found during
associated with habitation by man of a cl designed to determine man's contribution jointly by the United States Air Force an Administration. The experiment was div period, a manned period, and a manned scrubber. Direct analyses of the sealed comprehensive survey; however, cryoger samples with sufficient concentration of chromatography, infrared spectroscopy, pounds identified and quantified during th the manned phases.	to trace contained the National A wided into three period coupled we environment we nic fractionation contaminants fo and mass spec e experiment, o	eronautic phases: with an ac ere not ac n and con r analysi troscopy only 45 w	as conducted, as and Space an unmanned trivated carbon lequate for this centration provided by means of gas Of the 142 com- ere found during
associated with habitation by man of a cl designed to determine man's contribution jointly by the United States Air Force an Administration. The experiment was div period, a manned period, and a manned scrubber. Direct analyses of the sealed comprehensive survey; however, cryoger samples with sufficient concentration of chromatography, infrared spectroscopy, pounds identified and quantified during th the manned phases.	to trace contained the National A vided into three period coupled we environment we nic fractionation contaminants fo and mass spece e experiment, o	eronautic phases: with an ac ere not ac a and con r analysi troscopy only 45 w	as conducted, as and Space an unmanned tivated carbon lequate for this centration provided by means of gas Of the 142 com- ere found during
associated with habitation by man of a cl designed to determine man's contribution jointly by the United States Air Force an Administration. The experiment was div period, a manned period, and a manned scrubber. Direct analyses of the sealed comprehensive survey; however, cryoger samples with sufficient concentration of chromatography, infrared spectroscopy, pounds identified and quantified during th the manned phases.	to trace contained the National A vided into three period coupled we environment we nic fractionation contaminants fo and mass spec e experiment, o	eronautio phases: with an ac ere not ac a and con r analysi troscopy only 45 w	as conducted, an unmanned stivated carbon lequate for this centration provided s by means of gas Of the 142 com- ere found during
associated with habitation by man of a cl designed to determine man's contribution jointly by the United States Air Force an Administration. The experiment was div period, a manned period, and a manned scrubber. Direct analyses of the sealed comprehensive survey; however, cryoger samples with sufficient concentration of chromatography, infrared spectroscopy, pounds identified and quantified during th the manned phases.	to trace contained the National A wided into three period coupled of environment we nic fractionation contaminants fo and mass spec e experiment, of	eronautic phases: with an ac ere not ac a and con r analysi troscopy only 45 w	as conducted, as and Space an unmanned tivated carbon lequate for this centration provided by means of gas Of the 142 com- ere found during
associated with habitation by man of a cl designed to determine man's contribution jointly by the United States Air Force an Administration. The experiment was div period, a manned period, and a manned scrubber. Direct analyses of the sealed comprehensive survey; however, cryoger samples with sufficient concentration of chromatography, infrared spectroscopy, pounds identified and quantified during th the manned phases. (to trace contained the National A vided into three period coupled we not fractionation contaminants fo and mass spece e experiment, o	eronautic phases: with an ac ere not ac a and con r analysi troscopy only 45 w	as conducted, an unmanned tivated carbon lequate for this centration provided by means of gas Of the 142 com- ere found during
associated with habitation by man of a cl designed to determine man's contribution jointly by the United States Air Force an Administration. The experiment was div period, a manned period, and a manned scrubber. Direct analyses of the sealed comprehensive survey; however, cryoger samples with sufficient concentration of chromatography, infrared spectroscopy, pounds identified and quantified during th the manned phases.	to trace contained the National A vided into three period coupled we not fractionation contaminants fo and mass spece e experiment, o	eronautio phases: with an ac ere not ac a and con r analysi troscopy only 45 w	as conducted, as and Space an unmanned tivated carbon lequate for this centration provided s by means of gas Of the 142 com- ere found during
associated with habitation by man of a cl designed to determine man's contribution jointly by the United States Air Force an Administration. The experiment was div period, a manned period, and a manned scrubber. Direct analyses of the sealed comprehensive survey; however, cryoger samples with sufficient concentration of chromatography, infrared spectroscopy, pounds identified and quantified during th the manned phases. (to trace contained the National A vided into three period coupled we not fractionation contaminants fo and mass spece e experiment, o	eronautio phases: with an ac ere not ac a and con r analysi troscopy only 45 w	esified

.* X

4.						A LAN	
KEY WORDS	• • • · · ·	ROLE		BOLE		EIM BOLE	K C W7
			1]	1		
Aerospace medicine			1	Ì			
Closed environments	ч. н. н. н				· · · · ·		
Contaminants produced by man in closed			5	· ·	1	1.11.24	
environments			1				
Cryogenic fractionation		Ι	Í .	.			
Gas chrometography							
Infrared spectroscopy							
Mass spectroscopy							
space cabin environment			·	1.52			
eres a subscription of the second second	· • •	4.1	:	·			
		•.		e ~		· ·	
						- 1 e	
				1			
	1	[1	
					·		
· · · ·	1				2		
·							
•		[· ·]	
					1		
		. Viti i 🖡			÷.		
					·	5 - C - E	
· · · · ·		[[- 1	Ì	
		· .				<u>.</u> .	
2							
	ļ			·			
	2.4.5	.2	· · .	· ·			
			ł	. 1	· •	n l	••
		. [· · ·	-			
						1	
				· · ·	- 1		
				- [· · · · I .		
				. I.	s 1	. t. 🕴	
	ł	. J.		1			
· · · · · · · ·		.	.]		- e. Ja	~	
n n n n n n t		- e		· · .			

17

Unclassified Security Classification REFRODUCED FROM DEST AVAILABLE COPY