AD 740834

CELL DIVISION SYNCHRONIZATION

Final Report

by

David W. Rooney

April 15, 1972

The research was supported by the Office of Naval Research through Contract N00014-69-A-0206-0001, under Project Number NR 306-008

Reproduction in whole or in part is permitted for any purpose of the United States Government

> NATIONAL TECHNICAL INFORMATION SERVICE Springfiold, Va 22151

Approved for public release; distribution unlimited

Saint Louis University i Department of Biology St. Louis, Missouri 6310

h-13-2-

	CONTROL DATA - R & D lexing sumulation must be intered when the user off report is classified
A THE ACTIVES Content of the content of any market and market and market and the second secon	20. HE PORT SE CURITY CLESSIFICATION
Department of Biology	Unclassified
St. Louis University	26. CHOUP
St. Louis, Missouri 63074	
Cell Division Synchronizati	on
HPINE HOTES (7) pe of report and inclusive dates; Final report	
OR:S: (I 'Ist name, mid lie mittal, last name)	
David W. Rooney, Ph.D.	
ancil 15 1072	74. TOTAL NO OF PAGES 76. NO OF HEFS
April 15, 1972	3 6
N00014-69-3-0206-0001	
ГС7 NO	#2
	9b. OTHER REPORT NOISI (Any other numbers that may be a this report)
	nt is unlimited
Distribution of this docume	nt is unlimited.
Distribution of this docume	
Reported is progress i automatic equipment for syn by periodic hypoxia; probla carbohydrate sources for hu be subjected to periodic hypoxia; prodic hypoxia;	12 SPONSOHING WILITANY ACTIVITY

计结

L 1

To. U. S. Office of Naval Research

From:

David W. Rooney, Ph.D. Assistant Professor Saint Louis University, Department of Biology 1504 South Grand Boulevard, St. Louis, Missouri 63104

Date: March 15, 1972

Subject: Research under contract NOC14-69-A-0206-0001

FINAL REPORT

The specific objective at the onset of research under NO014-69-A-0206-0001 was to determine if periodic hypoxia is a suitable method for synchronizing division in cell cultures. The basic underlying purpose was to establish an additional useful biophysical technique for the study of the division cycle of eukaryotic (nucleated) cells.

A portion of the research has consisted in the design, testing, and refinement of automatic equipment to impose upon cell cultures any desired cyclic program of oxygen concentration. A preliminary description of this equipment may be found in a technical report (2) submitted to ONR. Subsequent refinements, such as nylon inserts to speed O_2 concentration changes have been made.

Another portion of the research has consisted of testing various sugars (glucose, fructose, galactose) at various concentrations in human (HeLa) cell suspension cultures. This testing was done to determine if rapid division (growth) rates of mammalian cells in suspension culture can be achieved with low glycolytic rates. Cnly when glycolysis is slow would oxygen deprivation be effective in limiting ATP production (since ATP is generated by both glycolysis and oxidative phosphorylation). Only when growth is rapid can biophysical perturbations (such as temporary 0, reductions) be expected to synchronize division. Extensivé testing of growth rate and pH charge (an index of glycolysis rate) with various sugars was carried cut by Miss Judith O'Brien who reported the results in h ster's Thesis (2). It was found that neither fractose nor guactose permitted rapid growth in suspension cultures.

Glucose permitted rapid growth -- but with rapid glycolysis.

In the course of these experiments, it became obvious that better facilities were needed to insure the absence of bacterial contamination in the human cell cultures. For this reason experiments with human cell suspensions have been temporarily halted. They will resume this summer (August, 1972) in a specially designed cell culture laboratory in the remodeled McDonnell Building at St. Louis University. Planned experiments will include steady-state culturing at <u>low</u> glucose concentration and imposition of periodic hypoxia.

Concurrent with the above described experimentation have been studies of periodic hypothermia as a method of synchronizing eukaryotic cell division. Because these studies are similar in objective, they are mentioned here. Briefly, I have found that periodic temperature reductions will affectively synchronize the division of an algal cell <u>Chlamydomonas</u>. The resulting publication (3) credits ONR support as do three aduitional publications (4, 5, 6) describing appropriate mathematical techniques for guantifying cell division synchrony in algal cultures.

REFERENCES

- O'Brien J., 1971, Effects of Sugar Substitutions on Growth and pH in Suspensions HELA-S3 Cell Cultures, M. S. (R) Thesis, St. Louis University.
- Rooney, D. W., 1970, Design Considerations in the Development of Automatic Equipment to Subject Suspension Cell Cultures to Repetitive Hypoxic Shock, Technical Report #1, submitted to ONR. AD. 707 c?c
- Rooney, D. W., Yen, B., and Mikita, E. J., 1971, Synchronization of <u>Chlamydomonas</u> Division with Intermittent. Hypothermia, Experimental Cell Research 65:94-98.
- Rooney, D. W., 1971, Simulated Algal Celi Cultures: Correlation of a Generalized Synchrony Index with Divisions Time Variation, Mathematical Biosciences 10:149-155.
- Rooney, D. W., 1971, Skewed Algal Division Patterns: Effects of Autospore Yield on Computed Synchrony Indices, Mathematical Biosciences 12:367-373.

 Rooney, D. W., 1972, A Quantitative Study of Factors Affecting Algal Division Synchrony Measurements, Mathematical Biosciences (in press).

and the month of the second

: