

AD 742204

ARL 72-0029
FEBRUARY 1972



Aerospace Research Laboratories


ON SOME NONCENTRAL DISTRIBUTIONS IN MULTIVARIATE ANALYSIS

P. R. KRISHNAIAH
A. K. CHATTOPADHYAY

APPLIED MATHEMATICS RESEARCH LABORATORY

PROJECT NO. 7071

Approved for public release; distribution unlimited.



Reproduced by
NATIONAL TECHNICAL
INFORMATION SERVICE
Springfield, Va. 22151

AIR FORCE SYSTEMS COMMAND
United States Air Force

23
R

NOTICES

When Government drawings, specifications, or other data are used for any purpose other than in connection with a definitely related Government procurement operation, the United States Government thereby incurs no responsibility nor any obligation whatsoever; and the fact that the Government may have formulated, furnished, or in any way supplied the said drawings, specifications, or other data, is not to be regarded by implication or otherwise as in any manner licensing the holder or any other person or corporation, or conveying any rights or permission to manufacture, use, or sell any patented invention that may in any way be related thereto.

Agencies of the Department of Defense, qualified contractors and other government agencies may obtain copies from the

**Defense Documentation Center
Cameron Station
Alexandria, Virginia 22314**

This document has been released to the

**CLEARINGHOUSE
U. S. Department of Commerce
Springfield, Virginia 22151**

for the sale to the public.

Copies of ARL Technical Documentary Reports should not be returned to Aerospace Research Laboratories unless return is required by security considerations, contractual obligations or notices on a specified document.

AIR FORCE: 86-72/500

A	DATE	DATE	DATE
	BY	BY	BY
DATE	DATE	DATE	DATE
BY	BY	BY	BY
DATE	DATE	DATE	DATE
BY	BY	BY	BY

Unclassified

Security Classification

DOCUMENT CONTROL DATA - R & D		
<i>(Security classification of title, body of abstract and indexing annotation must be entered when the overall report is classified)</i>		
1. ORIGINATING ACTIVITY (Corporate author) Aerospace Research Laboratories Wright-Patterson AFB, Ohio 45433		2a. REPORT SECURITY CLASSIFICATION Unclassified
		2b. GROUP M-2
3. REPORT TITLE On Some Noncentral Distributions in Multivariate Analysis		
4. DESCRIPTIVE NOTES (Type of report and inclusive dates) Scientific Final		
5. AUTHOR(S) (First name, middle initial, last name) P. R. Krishnaiah and A. K. Chattopadhyay		
6. REPORT DATE February 1972	7a. TOTAL NO. OF PAGES 20	7b. NO. OF REFS 9
8a. XXXXXXXXXXXX In-House Research 7071-00-12		8b. ORIGINATOR'S REPORT NUMBER(S)
8c. PROJECT NO. DoD Element 61102F DoD Subelement 681304		8d. OTHER REPORT NO(S) (Any other numbers that may be assigned this report) ARL 72-0029
9. DISTRIBUTION STATEMENT Approved for public release; distribution unlimited		
11. SUPPLEMENTARY NOTES TECH OTHER		12. SPONSORING MILITARY ACTIVITY Aerospace Research Laboratories (LB) Wright-Patterson AFB, Ohio 45433
13. ABSTRACT In this paper, the authors derived expressions for the marginal distributions of any few consecutive ordered roots, moments of the elementary symmetric functions of the ordered roots and the Laplace transformations of the traces of a class of random matrices in the noncentral cases. This class of random matrices includes the MANOVA, Canonical Correlation, and Wishart matrices. The expressions obtained here are in terms of the linear combinations of the products of double integrals; these double integrals can be evaluated without difficulty for the cases of the random matrices that occur commonly in multivariate statistical analysis. In deriving the results presented in this paper, the authors exploited the method of de Bruijn (J. Indian Math Soc. 19 133-152) for the evaluation of certain integral.		

DD FORM 1 NOV 68 1473

Unclassified

Security Classification

Unclassified

Security Classification

14. KEY WORDS	LINK A		LINK B		LINK C	
	ROLE	WT	ROLE	WT	ROLE	WT
Noncentral Distributions Multivariate Analysis Elementary Symmetric Functions Random Matrices Traces Ordered Roots						

Unclassified

Security Classification

ARL 72-0029

**ON SOME NONCENTRAL DISTRIBUTIONS
IN MULTIVARIATE ANALYSIS**

P. R. KRISHNAIAH

AND

A. K. CHATTOPADHYAY

APPLIED MATHEMATICS RESEARCH LABORATORY

FEBRUARY 1972

PROJECT 7071

Approved for public release; distribution unlimited.

AEROSPACE RESEARCH LABORATORIES
AIR FORCE SYSTEMS COMMAND
UNITED STATES AIR FORCE
WRIGHT-PATTERSON AIR FORCE BASE, OHIO

FOREWORD

This report was prepared for the Applied Mathematics Research Laboratory, Aerospace Research Laboratories, by P.R. Krishnaiah and A.K. Chattopadhyay under project 7071, "Research in Applied Mathematics". The work of A.K. Chattopadhyay is performed at the Aerospace Research Laboratories while in the capacity of Technology Incorporated Visiting Research Associate under contract F33615-71-C-1463, T.I. Project No. 4262B.

In this report, the authors consider the problems connected with certain non null distributions associated with the eigenvalues of a class of random matrices.

The authors wish to thank Dr. V.B. Waikar for some helpful discussions. Thanks are also due to Mrs. Georgene Graves for typing the manuscript carefully.

ABSTRACT

In this paper, the authors derived expressions for the marginal distributions of any few consecutive ordered roots, moments of the elementary symmetric functions of the ordered roots and the Laplace transformations of the traces of a class of random matrices in the noncentral cases. This class of random matrices includes the MANOVA, Canonical Correlation, and Wishart matrices. The expressions obtained here are in terms of the linear combinations of the products of double integrals; these double integrals can be evaluated without difficulty for the cases of the random matrices that occur commonly in multivariate statistical analysis. In deriving the results presented in this paper, the authors exploited the method of de Bruijn (J. Indian Math Soc. 19 133-152) for the evaluation of certain integral.

TABLE OF CONTENTS

SECTION		PAGE
I	INTRODUCTION	1
II	PRELIMINARIES	2
III	MOMENTS OF ELEMENTARY SYMMETRIC FUNCTIONS OF ROOTS	4
IV	DISTRIBUTION OF A FEW ORDERED ROOTS	7
V	PROBABILITY INTEGRALS OF ORDERED ROOTS	11
VI	NON NULL DISTRIBUTIONS OF TRACES	13
	REFERENCES	15

1. INTRODUCTION

Krishnaiah and his associates [4-6] derived exact expressions for the marginal distributions of any single or few ordered roots of a class of random matrices as well as the distributions of the traces of two random matrices. They derived the above results by exploiting the method of integration over alternate variables (e.g., see [7]). For a very brief summary of the literature on the distributions of the individual roots and traces of some random matrices the reader is referred to [4-6].

In general one can use any suitable function(s) (not necessarily symmetric) of the roots of the appropriate random matrices to test various hypotheses that arise on such problems as MANOVA, canonical correlation, tests for equality of covariance matrices, etc. Of course, the choice of these functions depends upon such factors as the optimum properties of the tests and the feasibility of the evaluation of the distributions. It will be of interest to examine the distributions of some of these functions which have at least intuitive appeal in testing some of the hypotheses.

In this note, we extend the results in [4-6] to the non-central cases by using the method of de Bruijn [1] for the evaluation of certain integral. Taking advantage of de Bruijn's method [1], we have also derived the moments of the elementary symmetric functions of roots of a class of random matrices in the non-central case. Here we note that Pillai and his associates (see [2,3,9] and the references there) derived the moments of elementary symmetric functions of roots explicitly in some special cases.

2. PRELIMINARIES

de Bruijn [1] proved the following useful result:^{*}

Lemma 2.1 (de Bruijn). Let (a, b) be any interval finite or infinite.

Then

$$\int_a^b \dots \int_a^b \phi(x_1, \dots, x_n) dx_1 \dots dx_n = \text{Pf}(A) \quad (2.1)$$

where

$$\phi(x_1, \dots, x_n) = |y_{ij}|, \quad y_{ij} = \phi_j(x_j).$$

$A = (a_{ij})$ and $\text{Pf}(A)$ denotes the Pfaffian of A . Here the elements a_{ij} are given by

$$a_{ij} = \int_a^b \int_a^b \phi_i(x) \phi_j(y) \text{Sgn}(y - x) dx dy \quad (2.2)$$

$i, j = 1, \dots, 2m$

if $n = 2m$; if $n = 2m + 1$, then they are given by (2.2) and by

$$a_{2m+2, 2m+2} = 0,$$

$$a_{i, 2m+2} = -a_{2m+2, i} = \int_a^b \phi_i(x) dx \quad i = 1, \dots, 2m+1.$$

We need the following in the sequel:

Lemma 2.2. Let $\eta(x_1, \dots, x_n)$ be a symmetric function of x_1, \dots, x_n and $\phi(x_1, \dots, x_n)$ be as defined before. Then

^{*} The integral in Eq. (2.2) of [6] is a special case of the integral in Eq. (2.1) of this paper.

$$\int_{a \leq x_1 \leq \dots \leq x_n \leq b} n(x_1, \dots, x_n) \phi(x_1, \dots, x_n) dx_1 \dots dx_n$$

$$= \int_a^b \dots \int_a^b E(x_1, \dots, x_n) n(x_1, \dots, x_n) \prod_{j=1}^n \{\phi_j(x_j) dx_j\} \quad (2.3)$$

where

$$E(x_1, \dots, x_n) = \prod_{1 \leq j < l \leq n} \text{Sgn}(x_j - x_l).$$

In addition, if

$$n(x_1, \dots, x_n) = \sum_{\kappa} C_{\kappa} x_1^{k_1} \dots x_n^{k_n}, \quad (2.3a)$$

where C_{κ} depends upon k_1, \dots, k_n and \sum_{κ} denotes the summation over the different elements of $\kappa = (k_1, \dots, k_n)$ subject to suitable restrictions, we have

$$\int_{a \leq x_1 \leq \dots \leq x_n \leq b} n(x_1, \dots, x_n) \phi(x_1, \dots, x_n) dx_1 \dots dx_n$$

$$= \sum_{\kappa} C_{\kappa} \text{Pf}(A_{\kappa}) \quad (2.4)$$

where $A_{\kappa} = (a_{ij}^{\kappa})$

In the above lemma, a_{ij}^{κ} are given by

$$a_{ij}^k = \int_a^b \int_a^b x^{k_i} \phi_i(x) y^{k_j} \phi_j(y) \operatorname{Sgn}(y - x) dx dy \quad (2.5)$$

$$i, j = 1, \dots, 2m$$

when $n = 2m$; if $n = 2m + 1$, the elements a_{ij}^k are given by (2.5) and

$$a_{2m+2, 2m+2}^k = 0$$

$$a_{i, 2m+2}^k = -a_{i, 2m+2}^k = \int_a^b x^{k_i} \phi_i(x) dx \quad i = 1, \dots, 2m + 1.$$

Lemma 2.2 can be proved by following the same lines as in de Bruijn [1].

3. MOMENTS OF ELEMENTARY SYMMETRIC FUNCTIONS OF ROOTS

Let $\lambda_1, \dots, \lambda_p$ be the latent roots of a random matrix and let us consider the situations when their joint density in the non-central case is of the form

$$f(\lambda_1, \dots, \lambda_p) = C \prod_{i=1}^p (\lambda_i) \prod_{i>j} (\lambda_i - \lambda_j) \sum_2 \sum_1 a(\kappa) \eta_\kappa(L) \quad (3.1)$$

$$a \leq \lambda_1 \leq \dots \leq \lambda_p \leq b$$

where $L = \operatorname{diag}(\lambda_1, \dots, \lambda_p)$, $\kappa = (k_1, \dots, k_p)$ is a partition of k subject to suitable restrictions, $\eta_\kappa(L)$ is a symmetric function and $a(\kappa)$ depends upon the population parameters and the elements of the partition κ .

Also \sum_1 denotes the summation over all partitions of k , whereas \sum_2 denotes the summation over k . Now, let

$$\eta_{\kappa}(L) = \sum_{\mathcal{Z}} b_{\kappa}^{\kappa} z_1^{r_1} \cdots z_p^{r_p} \quad (3.2)$$

where $\sum_{\mathcal{Z}}$ denotes the summation over r_1, \dots, r_p subject to some suitable restrictions and b_{κ}^{κ} depends on r_1, \dots, r_p and κ . If $\eta_{\kappa}(L)$ is the zonal polynomial and the elements of the partition $\kappa = (k_1, \dots, k_p)$ are subject to the restriction $k_1 \geq \dots \geq k_p \geq 0$, then $\eta_{\kappa}(L)$ is of the form (3.2).

The elementary symmetric function of order q is given by

$$\begin{aligned} z(z_1, \dots, z_p) &= \sum_{i_1 < \dots < i_q} z_{i_1} \cdots z_{i_q} \\ &= \sum_{\mathcal{A}} z_{i_1} \cdots z_{i_q} \quad \text{say} \end{aligned}$$

Now

$$\{z(z_1, \dots, z_p)\}^s = \sum_{\mathcal{S}} \binom{s}{s_1, \dots, s_{p^*}} z_1^{n_1} \cdots z_p^{n_p}, \quad (3.3)$$

where

$$p^* = \binom{p}{q} \text{ and } \sum_{\mathcal{S}}$$

denotes the summation over s_1, \dots, s_{p^*} such that $s_1 + \dots + s_{p^*} =$

$s, n_1 + \dots + n_p = qs$, and each n_i depends upon s_1, \dots, s_{p^*} . We know that

$$\prod_{i=1}^p \psi(z_i) \eta_{\kappa}(L) \{z(z_1, \dots, z_p)\}^s$$

is a symmetric function of z_1, \dots, z_p . So by (2.3) and (2.4) we get the sth moment of $z(z_1, \dots, z_p)$ as

$$\begin{aligned} \zeta_p [(z(z_1, \dots, z_p))^s] &= C \sum_2 \sum_1 a(\kappa) \int_a^b \dots \int_a^b E(z_1, \dots, z_p) \\ &\prod_{i=1}^p (\psi(z_i) z_i^{i-1}) n_\kappa(L) (z(z_1, \dots, z_p))^s dz_1 \dots dz_p \\ &= C \sum_2 \sum_1 \sum_3 \sum_5 a(\kappa) b_r^\kappa(s_1, \dots, s_{p^*})^s \\ &(A(r_1, \dots, r_p; n_1, \dots, n_p)) \end{aligned} \tag{3.4}$$

where

$$\begin{aligned} A(r_1, \dots, r_p; n_1, \dots, n_p) &= \int_a^b \dots \int_a^b E(x_1, \dots, x_p) \\ &\prod_{i=1}^p (\psi(x_i) x^{i+r_i+n_i-1} dx_i) \\ &= Pf(a_{ij}(r_1, \dots, r_p; n_1, \dots, n_p)) \end{aligned}$$

and

$$\begin{aligned} a_{ij}(r_1, \dots, r_p; n_1, \dots, n_p) &= \int_a^b \dots \int_a^b x^{i+r_i+n_i-1} y^{j+r_j+n_j-1} \\ &\psi(x) \psi(y) \operatorname{sgn}(y-x) dx dy \end{aligned}$$

when $p = 2m$; when $p = 2m+1$, the elements $a_{ij}(r_1, \dots, r_p; n_1, \dots, n_p)$ are as given above and

$$\begin{aligned}
& a_{2m+2, 2m+2}(r_1, \dots, r_p; n_1, \dots, n_p) = 0 \\
& a_{1, 2m+2}(r_1, \dots, r_p; n_1, \dots, n_p) \\
& = -a_{2m+2, 1}(r_1, \dots, r_p; n_1, \dots, n_p) \\
& = \int_a^b x^{r_i + n_i + 1} \psi(x) dx, \quad i = 1, \dots, 2m+1.
\end{aligned}$$

The summations $\sum_1, \sum_2, \sum_3, \sum_5$ are defined in (3.1), (3.2) and (3.3).

In the central case, let us assume that the joint density of x_1, \dots, x_p is of the form

$$f(x_1, \dots, x_p) = C \prod_{i=1}^p \psi(x_i) \prod_{i>j} (x_i - x_j).$$

In this case

$$E\{z(x_1, \dots, x_p)\}^S = C \sum_5 \binom{S}{s_1, \dots, s_p} \text{Pf}(A(0, \dots, 0; n_1, \dots, n_p))$$

where $A(0, \dots, 0; n_1, \dots, n_p)$ is given by (3.4).

4. DISTRIBUTION OF A FEW ORDERED ROOTS

We need the following in the sequel.

Remark 4.1. Let (a, b) be any interval finite or infinite. Also, let

$$I(k_1, \dots, k_n; a, b) = \int \dots \int_R \psi(x_1, \dots, x_n; k_1, \dots, k_n) dx_1 \dots dx_n$$

where

$$R: a \leq x_1 \leq \dots \leq x_n \leq b$$

and

$$\phi(x_1, \dots, x_n; k_1, \dots, k_n) = \begin{vmatrix} \phi_{k_1}(x_1) & \dots & \phi_{k_1}(x_n) \\ \vdots & & \vdots \\ \phi_{k_n}(x_1) & \dots & \phi_{k_n}(x_n) \end{vmatrix}$$

Then

$$I(k_1, \dots, k_n; a, b) = |D(k_1, \dots, k_n)|^{\frac{1}{2}} \quad (4.1)$$

where

$$D(k_1, \dots, k_n) = (d_{ij}(k_1, \dots, k_n)),$$

$$d_{ij}(k_1, \dots, k_n) = \int_a^b \int_a^b \phi_{k_i}(x) \phi_{k_j}(y) \operatorname{Sgn}(y-x) dx dy \quad (4.2)$$

$$i, j = 1, \dots, 2n$$

when $n = 2m$; if $n = 2m+1$, we have, in addition to (4.2),

$$d_{2m+2, 2m+2}(k_1, \dots, k_n) = 0$$

$$d_{1, 2m+2}(k_1, \dots, k_n) = -d_{2m+2, 1}(k_1, \dots, k_n) = \int_a^b \phi_{k_1}(x) dx$$

$$i = 1, \dots, 2m+1.$$

Formula (4.1) is implicit in lemma 2.1.

Let (a, b) be any interval finite or infinite and let

$$R^*: a \leq x_1 \leq \dots \leq x_r \leq x_{r+1} \leq x_{r+s} \leq x_{r+s+1} \leq \dots \leq x_p \leq b$$

Then

$$\int_{R^*} \dots \int \phi(x_1, \dots, x_p) n(x_1, \dots, x_p) dx_1 \dots dx_r dx_{r+s+1} \dots dx_p$$

$$= \sum_6 \sum_7 (-1)^{\frac{r(r+1)}{2} + \frac{s(s+1)}{2} + \sum_{i=1}^r k_i + \sum_{i=1}^s a_i}$$

$$V(x_{r+1}, \dots, x_{r+s}; a_1, \dots, a_s) \int_{R^{**}} \dots \int E(x_1, \dots, x_r) E(x_{r+s+1}, \dots, x_p) \quad (4.3)$$

$$V(x_1, \dots, x_r; k_1, \dots, k_r) V(x_{r+s+1}, \dots, x_p; \beta_{r+s+1}, \dots, \beta_p)$$

$$n(x_1, \dots, x_p) dx_1 \dots dx_r dx_{r+s+1} \dots dx_p; a \leq x_{r+1} \leq \dots \leq x_{r+s} \leq b,$$

and $(k_1 < \dots < k_r)$ is a subset of $(1, \dots, p)$, $(t_1 < \dots < t_{p-r})$ is its complementary set. Also $(a_1 < \dots < a_s)$ is a subset of $(t_1 < \dots < t_{p-r})$ and $(\beta_{r+s+1} < \dots < \beta_p)$ is the complementary subset of $(a_1 < \dots < a_s)$ with respect to $(t_1 < \dots < t_{p-r})$, \sum_6 denotes this summation over all possible $\binom{p}{r}$ choices of (k_1, \dots, k_r) and \sum_7 stand for the summation over $\binom{p-r}{s}$ possible choice of $(a_1 < \dots < a_s)$.

$$V(x_1, \dots, x_r; k_1, \dots, k_r) = \det(\phi_{k_i}(x_j))_{i,j=1, \dots, r}$$

and

$$R^{**}: (a \leq x_i \leq x_{r+1}, (i=1, \dots, r); x_{r+s} \leq x_j \leq b, (j = r+s+1, \dots, p)).$$

Formula (4.3) follows by repeated use of Laplace expansion of $\phi(x_1, \dots, x_p)$ starting with the first r columns and then with the first s columns of the remaining determinant. Now again if $n(x_1, \dots, x_p)$ is of the form $n(x_1, \dots, x_p) = \sum_{\mathcal{B}} C_{\mathcal{B}} x_1^{d_1} \dots x_p^{d_p}$ such that $n(x_1, \dots, x_p)$ is symmetric in x_1, \dots, x_p and $\sum_{\mathcal{B}}$ is similarly defined as in (2.3a)

$$\int \dots \int_{R^*} \phi(x_1, \dots, x_p) n(x_1, \dots, x_p) dx_1 \dots dx_r dx_{r+s+1} \dots dx_p$$

$$= \sum_{\mathcal{B}} \sum_{\mathcal{C}} \sum_{\mathcal{D}} (-1)^{\frac{r(r+1)}{2} + \frac{s(s+1)}{2} + \sum_{i=1}^r k_i + \sum_{i=1}^s a_i} C_{\mathcal{B}}$$

$$V(x_{r+1}, \dots, x_{r+s}; a_1, \dots, a_s) x_{r+1}^{d_{r+1}} \dots x_{r+s}^{d_{r+s}}$$

$$I(k_1, \dots, k_r, d_1, \dots, d_r; a, x_{r+1})$$

$$I(\beta_{r+s+1}, \dots, \beta_p, d_{r+s+1}, \dots, d_p; x_{r+s}, b) \tag{4.4}$$

Where $\sum_{\mathcal{B}}$ is given in (2.3a) and $I(k_1, \dots, k_r, d_1, \dots, d_r; a, b)$ is obtained from $I(k_1, \dots, k_r; a, b)$ replacing $\phi_{k_i}(x_j)$ with $x_j^{d_j} \phi_{k_i}(x_j)$.

Now let us take the joint density of the roots in the noncentral case as given by (3.1). Then applying formula (4.3) we get the density of roots x_{r+1}, \dots, x_{r+s} as

$$f(x_{r+1}, \dots, x_{r+s})$$

$$\begin{aligned}
&= C \sum_2 \sum_1 \sum_3 \sum_6 \sum_7 (-1)^{\frac{r(r+1)}{2} + \frac{s(s+1)}{2} + \sum_{i=1}^r k_i + \sum_{i=1}^s a_i} \\
& a(\kappa) b_{\lambda}^{\kappa} I(k_1, \dots, k_r, \lambda_1, \dots, \lambda_r; a, x_{r+1}) \\
& I(\beta_{r+s+1}, \dots, \beta_p, \lambda_{r+s+1}, \dots, \lambda_p; x_{r+s}, b) \\
& V(x_{r+1}, \dots, x_{r+s}; a_1, \dots, a_s) \\
& x_{r+1}^{d_{r+1}} \dots x_{r+s}^{d_{r+s}}, \quad a \leq x_{r+1} \leq \dots \leq x_{r+s} \leq b
\end{aligned}$$

where

$$\phi(x_1, \dots, x_p) = \det (x_j^{i-1})_{i,j=1, \dots, p} \quad (4.5)$$

$C, \sum_1, \sum_2, \sum_3, \sum_6, \sum_7$ are all defined earlier b_{λ}^{κ} and $\lambda_1, \dots, \lambda_p$ are defined in the same way as b_r, r_1, \dots, r_p in (3.2).

5. PROBABILITY INTEGRALS OF ORDERED ROOTS

As before, let us assume that the joint density of the roots $x_1 < \dots < x_p$ is given by (3.1). Then the density functions of the extreme roots can be obtained by following the same lines as in Krishnaiah and Chang [5] and using (2.3) and (2.4). Also the c.d.f. of the individual extreme root can be computed by applying (2.3) and (2.4). The probability integral associated with the joint density of any two ordered roots is easily obtained by following the same lines as in Krishnaiah and Waikar [6] and applying formulae (2.3) and (2.4). The expressions associated with the density functions

and c.d.f.s of the extreme roots and the probability integral associated with any two ordered roots are not given here for the sake of brevity. We will now derive the c.d.f.s associated with the intermediate roots.

It is known that

$$P[z_r \leq x] = P[z_{r+1} \leq x] + P[z_1 < \dots < z_r < x < z_{r+1} < z_p] \quad (5.1)$$

Now c.d.f.s of the extreme roots can be evaluated using (2.3) and (2.4). The second term in (5.1) can be calculated following similar lines as in Krishnaiah and Waikar [6] and using (4.3) and (4.4). It is given by

$$P[z_1 < \dots < z_r < x < z_{r+1} < \dots < z_p]$$

$$C \sum_2 \sum_1 \sum_3 \sum_6 (-1)^{\frac{1}{2}r(r+1) + \sum_{i=1}^r k_i} b_{\lambda}^{\kappa} a(\kappa)$$

$$I(k_1, \dots, k_r, \lambda_1, \dots, \lambda_r; a, x) I(t_1, \dots, t_{p-r}, t_{r+1}, \dots, \lambda_p; x, b)$$

where

$$C, \sum_1, \sum_2, \sum_3, \sum_6, (k_1 < \dots < k_r), (t_1 < \dots < t_{p-r})$$

and

$$\phi(z_1, \dots, z_r)$$

are as defined in (4.5).

6. NON NULL DISTRIBUTIONS OF TRACES

Here we use the results in [4] and treat both traces discussed there in a unified way. Let the joint density of the roots be given as (3.1).

Then

$$\begin{aligned} & E(e^{-t(\lambda_1 + \dots + \lambda_p)}) \\ &= C \sum_2 \sum_1 \sum_3 a(\kappa) b_r^k Pf(B_r) \end{aligned}$$

where

$$B_r = (b_{r,ij})_{i,j=1,\dots,p}$$

and

$$b_{r,ij} = \int_a^b \int_a^b x^{r_1+i-1} y^{r_j+j-1} \psi(x) \psi(y) e^{-tx} e^{-ty} \operatorname{sgn}(y-x) dx dy$$

$i, j=1, \dots, p$

If p is odd, the B_r matrix has to be augmented as given in Section 2.

Now to find the exact distribution we have to invert this Laplace transform. To this end we note that

$$Pf(B_r) = \sum \pm b_{r,i_1,i_2} \dots b_{r,i_{2a-1},i_{2a}}$$

Where the summation is over all possible choices of i_1, \dots, i_{2a} subject to this restriction $i_1 < i_2, \dots, i_{2a-1} < i_{2a}$ and the sign is +ve or -ve according as the permutation is even or odd. Thus inverting the Laplace transform we get

$$f(V) = C \sum_2 \sum_1 \sum_3 \sum_6 a(\kappa) b_r h_r (i_1, \dots, i_{2a}; V)$$

where $h_r(i_1, \dots, i_{2a})$ is the inverse Laplace transform of $\text{Pf}(B_r)$. For discussion on inverse transform of the above type for two important cases we refer to [4].

REFERENCES

- [1] deBruijn, N.G. (1955), "On Some Multiple Integrals Involving Determinants", J. Indian Math. Soc. 19, 133-152.
- [2] Khatri, C.G. and Pillai, K.C.S. (1968), "On the Non-Central Distributions of Two Tests Criteria in Multivariate Analysis of Variance", Ann. Math. Statist. 39, 215-225.
- [3] Khatri, C.G. and Pillai, K.C.S. (1968), "On the Moments of Elementary Symmetric Functions of the Roots of Two Matrices and Approximations to a Distribution", Ann. Math. Statist. 39, 1274-1281.
- [4] Krishnaiah, P.R. and Chang, T.C. (1970), "On the Exact Distributions of the Traces of $S_1 S^{-1}$ and $S_1(S_1 + S_2)^{-1}$ ", ARL 70 Aerospace Research Laboratories, Wright-Patterson AFB, Ohio.
- [5] Krishnaiah, P.R. and Chang, T.C. (1971), "On the Exact Distribution of Extreme Roots of the Wishart and MANOVA Matrices", J. Multivariate Analysis 1, 108-117.
- [6] Krishnaiah, P.R. and Walker, V.B. (1971), "Exact Joint Distributions of any Few Ordered Roots of a Class of Random Matrices", J. Multivariate Analysis 1, 308-315.
- [7] Mehta, M.L. (1967), Random Matrices and the Statistical Theory of Energy Levels, Academic Press, New York.
- [8] Pillai, K.C.S. (1968), "On the Moment Generating Function of Pillai's $V^{(s)}$ Criterion", Ann. Math. Statist. 39, 877-880.
- [9] Pillai, K.C.S. and Jouris, G.M. (1969), "On the Moments of Elementary Symmetric Functions of the Roots of Two Matrices", Ann. Inst. Statist. Math. 21, 309-320.