

A Preprint from the

AD 744678

OWENS VALLEY RADIO OBSERVATORY

California Institute of Technology

Pasadena, California

1972

:

3. A MODEL OF THE CRAB NEBULA DERIVED FROM DUAL-FREQUENCY RADIO MEASUREMENTS

by K. W. Weiler and G. A. Seielstad

DOCUMENT CO	NTROL DATA - R	& D	· · · · · · · · · · · · · · · · · · ·
Soundly clausilication of title, budy of abetract and index	ing annotation must be	entered when the	overall report in classified)
California Institute of Technology	7	28. REPORT SE	CUNITY CLASSIFICATION
Owone Valley Faile Chesrystory		UNCLA	SSIFIED
Decident California Gillo	,	25. GROUP	
rasadenan carronna sitos			•
A MODEL OF THE CRAB NEBILA DERIVER RADIO MEASUREMENTS.	FROM DUAL-	FREQUENCY	: .
OFSCRIPTIVE NGTES (Type of report and inclusive dates)			
AUT(10(13) (First rame, middle initial leases			
K. W. Weiler and G. A. Seielstad	:		:
: : :			
#3. 1972	. 74. TOTAL NO. O	F PAGES	78. NO. OF REFS
ju w produkter i station i statio station i station i station i station i station i station i st	28		26
LUNTRACT OR GRANT NO	Se. ORIGINATOR	S REPORT NUME	ER(\$)
Nonr N00014-67-A-0094-0019	* *	3, 1972 .	
· ·		• •	
	B. OTHER REPO	RT NOIS (Any of	her numbers that may be assigned
			• •
This document has been approved for	r nublic re	lease and	sale: its
distribution is unlimited.	r pustic le	!	· · · · · ·
distribution is unlimited.		!	
distribution is unlimited.	12. \$PON30#ING	MILITARY ACTIV	
distribution is unlimited.	12. SPONSOWING	MILITARY ACTIN	esearch
distribution is unlimited.	12. SPONSOPING Office o	f Naval R	esearch
distribution is unlimited.	Office o	f Naval R	esearch
distribution is unlimited. SUPPLEMENTANY NOTES ANATHACT The total intensity and linea	Office o arly polariz	f Naval R ed emissi	esearch on from the Crab
distribution is unlimited.	Office of a resol	f Naval R ed emissi	esearch on from the Crab approximately 1
distribution is unlimited.	Office o arly polariz	f Naval R ed emissi	esearch cn from the Crab approximately 1
distribution is unlimited.	Office o arly polariz to a resol	f Naval R ed emissi ution of hese data	esearch on from the Crab approximately 1 were calculated
distribution is unlimited.	Office of arly polariz to a resol Hz. From t	f Naval R ed emissi ution of hese data	esearch on from the Crab approximately 1 were calculated on angle, and
distribution is unlimited. 3UPPLEMENTALLY NOTES ANATHACT The total intensity and linea Nebula (Taurus A) were synthesized arc-minute at both 1420 and 2880 M the spectral index, rotation measu	Office o arly polariz to a resol Hz. From 't	f Naval R ed emissi ution of hese data	esearch on from the Crab approximately 1 were calculated on angle, and
distribution is unlimited. SUPPLEMENTATION NOTES ANATHACT The total intensity and linea Nebula (Taurus A) were synthesized arc-minute at both 1420 and 2880 M the spectral index, rotation measu depolarization ratio distributions	Office o Arly polariz to a resol MHz. From 't are, intrins 5. Then, co	f Naval R ed emissi ution of hese data ic positi mbined wi	esearch on from the Crab approximately 1 were calculated on angle, and th a source model
distribution is unlimited. 3UPPLEMENTATION NOTES ANALY ACT The total: intensity and linea Nebula (Taurus A) were synthesized arc-minute at both 1420 and 2880 M the spectral index, rotation measu depolarization ratio distributions the physical conditions within the	Office of office of arly polariz to a resol MHz. From t are, intrins then, co e supernova	f Naval R ed emissi ution of hese data ic positi mbined wi remnant w	esearch on from the Crab approximately 1 were calculated on angle, and th a source model ere established.
distribution is unlimited. 3UPPLEMENTATION NOTES ANALY NOTES ANALY ACT The total: intensity and linea Nebula (Taurus A) were synthesized arc-minute at both 1420 and 2880 M the spectral index, rotation measu depolarization ratio distributions the physical conditions within the	12. SPONSORING Office of arly polariz to a resol MHz. From t are, intrins 5. Then, co e supernova	f Naval R ed emissi ution of hese data ic positi mbined wi remnant w	esearch on from the Crab approximately 1 were calculated on angle, and th a source model ere established.
distribution is unlimited.	12 SPONSOMING Office of arly polariz to a resol MHz. From t are, intrins 5. Then, co e supernova oth the homo	f Naval R ed emissi ution of hese data ic positi mbined wi remnant w geneous a	esearch on from the Crab approximately 1 were calculated on angle, and th a source model ere established. nd random com-
distribution is unlimited.	12 SPONSOWING Office of arly polariz to a resol MHz. From t are, intrins 5. Then, co e supernova oth the homo	f Naval R ed emissi ution of hese data ic positi mbined wi remnant w geneous a and a me	esearch on from the Crab approximately 1 were calculated on angle, and th a source model ere established. nd random com- asurement of the
distribution is unlimited.	12 SPONSOWING Office of arly polariz to a resol MHz. From t are, intrins 5. Then, co e supernova oth the homo e determined	f Naval R ed emissi ution of hese data ic positi mbined wi remnant w geneous a and a me	esearch on from the Crab approximately 1 were calculated on angle, and th a source model ere established. nd random com- asurement of the
distribution is unlimited.	12. SPONSOWING Office of arly polariz to a resol MHz. From t are, intrins 5. Then, co e supernova oth the homo e determined ion obtained	f Naval R ed emissi ution of hese data ic positi mbined wi remnant w geneous a and a me	esearch on from the Crab approximately 1 were calculated on angle, and th a source model ere established. nd random com- asurement of the
distribution is unlimited. 30PPLEMENTATION NOTES ANALY ACT The total: intensity and linea Nebula (Taurus A) were synthesized arc-minute at both 1420 and 2880 M the spectral index, rotation measu depolarization ratio distributions the physical conditions within the The strength and orientation of bo ponents of the magnetic field were thermal electron plasma distributi	12 SPONSOFING Office of Arly polariz to a resol Miz. From t are, intrins 5. Then, co e supernova oth the homo e determined ion obtained	f Naval R ed emissi ution of hese data ic positi mbined wi remnant w geneous a and a me	esearch on from the Crab approximately 1 were calculated on angle, and th a source model ere established. nd random com- asurement of the
distribution is unlimited. 30PPLEMENTATION NOTES ANALY	Office of Office of arly polariz to a resol MHz. From t are, intrins to the nomo e supernova oth the homo e determined	f Naval R ed emissi ution of hese data ic positi mbined wi remnant w geneous a and a me	esearch cn from the Crab approximately 1 were calculated on angle, and th a source model ere established. nd random com- asurement of the
distribution is unlimited. 3UPPLEMENTATION NOTES ANALY	12 SPONSORING Office of arly polariz to a resol MHz. From t are, intrins to the nomo e supernova oth the homo e determined ton obtained	f Naval R ed emissi ution of hese data ic positi mbined wi remnant w geneous a and a me	esearch on from the Crab approximately 1 were calculated on angle, and th a source model ere established. nd random com- asurement of the
distribution is unlimited. 30PPLEMENTATIONOTES ANALY NOTES ANALY NO	12 SPONSORING Office of arly polariz to a resol Hz. From t are, intrins to the nomo e supernova oth the homo e determined ton obtained	f Naval R ed emissi ution of hese data ic positi mbined wi remnant w geneous a and a me	esearch on from the Crab approximately 1 were calculated on angle, and th a source model ere established. nd random com- asurement of the
distribution is unlimited.	12. SPONSORING Office of arly polariz to a resol MHz. From t are, intrins to the nomo e supernova oth the homo e determined ion obtained	f Naval R ed emissi ution of hese data ic positi mbined wi remnant w geneous a and a me	esearch on from the Crab approximately 1 were calculated on angle, and th a source model ere established. nd random com- asurement of the
distribution is unlimited.	12 SPONSOWING Office of arly polariz to a resol Hz. From t are, intrins 5. Then, co e supernova oth the homo e determined ton obtained	f Naval R ed emissi ution of hese data ic positi mbined wi remnant w geneous a and a me	esearch on from the Crab approximately 1 were calculated on angle, and th a source model ere established. nd random com- asurement of the
distribution is unlimited. SUPPLEMENTALLY NOTES ANALYMACT The total: intensity and linea Nebula (Taurus A) were synthesized arc-minute at both 1420 and 2880 M the spectral index, rotation measu depolarization ratio distributions the physical conditions within the The strength and orientation of bo ponents of the magnetic field were thermal electron plasma distributi	12 SPONSOWING Office of arly polariz to a resol Hz. From t are, intrins to the nomo e supernova oth the homo e determined ton obtained	f Naval R ed emissi ution of hese data ic positi mbined wi remnant w geneous a and a me	esearch on from the Crab approximately 1 were calculated on angle, and th a source model ere established. nd random com- asurement of the
distribution is unlimited. SUPPLEMENTALLY NOTES Anstruct The total intensity and linea Nebula (Taurus A) ware synthesized arc-minute at both 1420 and 2880 M the spectral index, rotation measu depolarization ratio distributions the physical conditions within the The strength and orientation of bo ponents of the magnetic field were thermal electron plasma distributions	12 SPONSOFING Office of arly polariz to a resol Hz. From t are, intrins 5. Then, co e supernova oth the homo e determined ion obtained	f Naval R ed emissi ution of hese data ic positi mbined wi remnant w geneous a and a me	esearch on from the Crab approximately 1 were calculated on angle, and th a source model ere established. nd random com- asurement of the
distribution is unlimited. SUPPLEMENTALLY NOTES ANSTRACT The total intensity and linea Nebula (Taurus A) ware synthesized arc-minute at both 1420 and 2880 M the spectral index, rotation measu depolarization ratio distributions the physical conditions within the The strength and orientation of bo ponents of the magnetic field were thermal electron plasma distributions	I'2 SPONSORING Office o Arly polariz to a resol Miz. From t are, intrins 5. Then, co e supernova oth the homo e determined ion obtained	f Naval R ed emissi ution of hese data ic positi mbined wi remnant w geneous a and a me	esearch on from the Crab approximately 1 were calculated on angle, and th a source model ere established. nd random com- asurement of the

Security Classification

212 3 M. P. 13.

مايمه ويستمينه بالمعادية والكمية والمسامية المسامية معتقاته والملاحظة المكما المعادية محالاته المحافية والمقاصات المحافية والمتحديد وفقا فالمتحديد وفقا فالمتحد والمحافية والمتحديد

UN 44 3 - 1 1

Elisteraria statistication

DO DE LADOS DE LA CARRENTE A RECENCIÓN DE LA COMPANYA DE LA

ABSTRACT

The total intensity and linearly polarized emission from the Crab Nebula (Taurus A) were synthesized to a resolution of approximately 1 arc-minute at both 1420 and 2880 MHz. From these data were calculated the spectral index, rotation measure, intrinsic position angle, and depolarization ratio distributions. Then, combined with a source model, the physical conditions within the supernova remnant were established. The strength and orientation of both the homogeneous and random components of the magnetic field were determined and a measurement of the thermal electron plasma distribution obtained. Address of Authors:

Dr. K. W. Weiler Radiosterrenwacht Westerbork Post Hooghalen, Netherlands

Dr. G. A. Seielstad Owens Valley Radio Observatory California Institute of Technology P.O. Box 387 Big Pine, California 93513 U.S.A.

Key Words: Supernova, Crab Nebula, Taurus A

-1-

I. INTRODUCTION

Earth rotation synthesis observations of the total and linearly polarized radiation from Taurus A (The Crab Nebula) were taken concurrently with those of the two supernova remnants 3Cl0 (Tycho's Supernova) and 3C58 (Weiler and Seielstad 1971, hereafter referred to as Paper I). The Owens Valley Radio Observatory's interferometer was used at frequencies of both 1420 and 2880 MHz at multiple spacings East-West and North-South and over all hour angles available to the telescopes $(-4^{h}$ to $+4^{h})$. The resulting coverage in the Fourier transform (u, v) plane is illustrated in Fig. 1.

The observing methods, dates of observation, and calibration procedure are fully described in Paper I. Briefly, the observations were done by alternating the receiving horns between three different configurations of orthogonal position angles to obtain measures of the Fourier components of the Stokes parameters I+Q, U-iV, and -Q-iV. Since we previously measured an upper limit of 0.2 per cent to the integrated degree of circular polarization of Taurus A at 1418 MHz (Seielstad and Weiler 1968)¹, we assumed that there was no circularly polarized radiation, i.e., V=0.

These three uncorrected sets of data were then inverted into the real plane. However, because of the large gaps evident in

¹G. L. Berge and G. A. Seielstad (unpublished) have measured an upper limit of 0.2 per cent at 1602 MHz also.

-2-

Fig. 1 in the information coverage of the Fourier plane, the inversion technique employed in Faper I was no longer appropriate. Instead, we used an inversion scheme developed by J. A. Högbom and described by Rogstad and Shostak (1971). Finally the instrumental corrections (< 0.3 per cent at 1420 MHz and < 0.4 per cent at 2880 MHz) were applied and the solutions combined to yield the distributions of the total radiation (Stokes parameter I), of the linearly polarized radiation ($\sqrt{Q^2 + U^2}$), and of the position angles of the maximum electric vectors ($k_2 \tan^{-1} (U/Q)$).

II. SOURCE DISTRIBUTIONS

The resulting synthesized maps are displayed in Figs. 2 and 3. All details of position and scale are identical among the several diagrams. The origin is

> $\alpha \quad (1950.0) = 05^{h} 31^{m} 31.20,$ $\delta \quad (1950.0) = +21^{\circ} 59^{\circ} 17.0^{\circ}$

aa waaraa waxaa waxaa waxay dhixadha ahaa ahaa ahaa ahaa ahaa ahaa waxadhi waxaa ka ahaa ahaa hiyadhi ahaa aha

Relative to the Sp of the central double stars, this position is 0.26° earlier and 22.2 north. The error analysis and the effects of ionospheric Faraday rotation are discussed in Paper I. The error estimates for the three fundamental maps (I, Q, U) are collected in Table 1.

A. Distributions of the Total Radiation

The brightness distributions of the total radiation at the two frequencies show the same smooth appearance already noted by several previous studies with comparable resolution (Branson

-3-

1967; Mayer and Hollinger 1968; Braun and Yen 1968; Johnston and Hobbs 1969; Wright 1970). Only the maps with resolution better by an order of magnitude (Hogg <u>et al</u>. 1969; Wilson 1971) begin to reveal further structural detail. It seems worthwhile repeating that the most remarkable feature of the brightness distribution is its central concentration. Milne (1971) has demonstrated that it is much more common for supernova remnants to show at least partial shell structure.

t is twentile as the three a bire to start schere bit is over we distance and a drate stration of the head the t

Wright (1970) has called attention to an extended low brightness region SW of the nebula. There is little evidence for such a feature in either the total or polarized radiation maps. Also, recent $\lambda 21$ cm maps made with the Westerbork Telescope fail to show such a region (R. M. Duin 1972, private communication).

B. Distributions of Linearly Polarized Radiation

Both distributions of linearly polarized radiation reveal similar structure, which is more detailed than that of the total radiation. There is a conspicuous gap in the NW which is near the most prominent "bay" in the optical distribution (Woltjer 1957). The rapid change of position angles in that region implies that the low radio polarization may result partly from beam-averaging vectors having a wide range of position angles. Higher resolution studies (Conway 1971) confirm this. We find it somewhat surprising that our polarization distributions do not exhibit similar gaps elsewhere, since other prominent

-4-

optical bays are known to exist. The center of the polarization distribution in both cases is displaced about 0:25 east and 0:3 south from the center of the total brightness distribution. Other observers (Mayer and Hollinger 1968; Downs and Thompson 1968; Johnston and Hobbs 1969; Wright 1970), confirm most of these features over a wide range of frequencies.

The degree of polarization reaches nearly 4 per cent at 1420 MHz and nearly 8 per cent at 2880 MHz. In both cases it is relatively constant over the main body of the nebula increasing slightly at 21 cm around the disturbed region in the NW. The position angles likewise show a constancy within the main body with a sharp rotation of angles occurring in the NW.

III. PARAMETER DISTRIBUTIONS

Since we have observations at two very different frequencies with the same resolution and with approximately the same Fourier plane coverage, processed in an identical manner, determinations of frequency-dependent parameter distributions are especially reliable. For these distributions we have chosen a coordinate system oriented along the major and minor axes of the source. A. Spectral Index Distribution

The spatial distribution of spectral indices for Taurus A (defined as $I_{\nu} v^{+\alpha}$) between 1420 and 2880 MHz is presented in Table 2. The spectrum is flattest near the center and steepens progressively along any radial direction outward. This steepening

-5-

of spectral index toward the periphery of the nebula was noted between 404 and 1420 MHz alsc (Davies <u>et al</u>. 1966). Furthermore, Davies <u>et al</u>. (1966) and Scargle (1969) have emphasized that the optical nebula is smaller than the radio, and that the X-ray nebula is smaller yet. These spatial variations in spectrum reflect corresponding differences in the electron-energy distributions at different points in the nebula. As expected, electrons with the highest energy are concentrated nearest the center, which is now universally recognized as the seat of acceleration. B. Rotation Measure and Intrinsic Position Angle Distributions.

The polarization parameters of the integrated radiation from Taurus A are accurately known at several frequencies. There is consequently no ambiguity in the determination of the integrated rotation measure. Burn (1966) finds -25.6 rad m⁻². To resolve any $\pm n\pi$ ambiguities in position-angle difference between 1420 and 2880 MHz, we require that our calculated values be near this integrated value and that variations in rotation measure not be discontinous over dimensions comparable to the beamwidth. Since a difference of π rad between 1420 and 2880 MHz changes the rotation measure by 93 rad m⁻²; we feel certain that the values presented in Table 3 are free from ambiguity.

The rotation-measure distribution varies extremely smoothly and over a surprisingly small range of values. A contribution to each of these rotation measures is made by the interstellar medium between the sun and the Crab Nebula. Manchester (1971)

-6-

has pointed out the similarities in rotation measure and dispersion measure of the two pulsars, 0531+21 and 0525+21, both believed associated with the Crab Nebula, and has argued convincingly that their rotation arises in the interstellar medium. We therefore adopt -4. rad m^{-2} as the interstellar rotation measure. By subtracting this value from those measured, we obtain the rotation measures produced internally in the Grab Nebul. These rights are presented in Table 4.

... rept near the eastern edge of the nebula, lines of constant rotation measure are parallel to the major axis. It seems unlikely that a gradient in the intervening interstellar medium would coincide with a principal axis of Taurus A^2 , so that the variation in Table 4 probably represents changes in its internal magneto-ionic medium.

We have also calculated the distribution of intrinsic position angles using the rotation measures in Table 3 and the position angles measured at 2880 MHz. These results are shown in Fig. 4. Except in the northwest corner, the intrinsic angles vary little across the source. Over a central region of angular diameter ~ 2', the total scatter in angles is less than 10°. Over the main body of the nebula the intrinsic position angles are ~ 15° greater than the position angle of the major axis.

²Since the major axis of the Crab Nebula is closely parallel to the galactic equator (Münch 1958), this argument may be weakened somewhat.

-7-

C. Depolarization Ratio Distribution

The depolarization ratio (P_{1420}/P_{2380}) is computed over the face of the Crab and presented in Table 5. The percentage polarization at the lower frequency is nearly everywhere very close to half that at the higher. There is no obvious systematic spatial trend in the ratios. Coupled with the previously noted systematic variation in rotation measures, this indicates that depolarization and rotation are not strongly correlated. Since correlation would be expected in a uniform Faraday medium, we conclude that the magneto-ionic medium internal to the Crab Nebula has non-uniformities on a scale much smaller than our beamwidth (i.e., <<0.5 parsec).

IV. DISCUSSION

With our resolution of ~l arcmin (corresponding to ~ 0.5 pc at a distance of 1.7 kpc), the overall impression of Taurus A is one of extraordinary large-scale uniformity in its structure, as revealed by the following evidence:

(a) The total brightness distribution is very smooth and simple (Figs. 2 and 3).

(b) The intrinsic polarization angles show little variation over most of the main body of the nebula (Fig. 4). This implies a uniform direction for H_{\perp} , the component of internal magnetic field normal to the line of sight.

-8-

(c) The rotation measures vary smoothly, slowly and only over a small range. (Table 3 or 4). This implies the product $n_e H_{||}$, where n_e is the density of thermal gas, is relatively constant.

(d) The depolarization ratios are nearly constant (Table 5), implying that the scale of irregularities and the energy density in turbulent magnetic fields changes little.

However, coexisting with this large-scale ordering of the Crab Nebula is a chaotic fine structure. This can be seen by:

(a) The prominent optical filaments (Woltjer 1958; Trimble 1968) reveal directly high density concentrations on a scale $\sim 10^{-2}$ pc.

(b) The optical continuum emission has a fine, fibrous structure with details at the limit of resolution (\leq 1 arcsec (corresponding to $< 10^{-2}$ pc; Scargle 1969).

(c) Fine structure is seen in high resolution radio studies(Wilson 1971).

(d) The continuum radiation from optical to long radio wavelengths is strongly depolarized. Burn (1966) has explained this in terms of a spread in Faraday depths over several orders of magnitude. The available decimetric wavelength polarization distributions reveal that portions of the polarized emission radiate through gaps in a highly depolarizing screen (Downs and Thompson 1968; Wright 1970; present paper). The highest resolution radio polarization study (Conway 1971) reveals that this depolarizing screen has a scale < 7" (or <0.06 pc). (e) The polarized radiation from the pulsar 0531+21, presumably located at the center of the nebula, apparently receives little or no Faraday rotation from the nebula (Manchester 1971). This is direct evidence for at least one hole in the magneto-plasma.

(f) The constant linear polarization of the optical continuum near the pulsar implies that the scale of the fine structure is less than 0.02 pc (Forman and Visvanathan 1971). n so say child K 1 Navácska i uce a nom ra nah. Ezerskiho uh si dezerlikozek sztrácké sztránk szterkezek sztrát

and a substant of the substant resolution of the state of the substant of a definition of the substance of the

A. The Uniform Magnetic Field

Our measurements clearly refer primarily to the overall large-scale magnetic field. Using synchrotron theory (Ginzburg and Syrovatskii 1965), we can estimate the distribution of magnetic-field strengths. These are presented in Table 6. We have assumed that the Crab Nebula is a uniformly filled prolate spheroid (prolate <u>vs</u>. oblate case argued in, e.g., Shklovsky 1968, pp. 311, 312) with semi-major and semi-minor axes, which lie in the plane of the sky, of dimensions 1.61 and 1.25 pc, respectively, at a distance of 1.7 kpc (Minkowski 1971). We have also assumed equipartition between the energies of the magnetic field and of the relativistic electrons (see Shklovsky 1968, pp. 3!6-318 for arguments for ignoring protons), and have used the spectral index distribution from Table 2 and the brightness distribution at 1420 MHz (Fig. 2). The field strengths in Table 6 represent the quantity $\sqrt{H_{O}^{-2}+H_{r}^{-2}}$, where H_{O} is a uniform field and H_{r} an isotropic random field, averaged along lines-of-sight through the nebula. This quantity decreases only slightly from ~ 2.5x10⁻⁴ G at the center to ~ 1.9x10⁻⁴ G near the edges.

We can also estimate the relative contributions of the random and uniform components of the field. The fractional polarization expected in a uniform magnetic field is $p(\alpha) = (3-3\alpha)/((5-3\alpha))$, where α is the spectral index. We find that $p(\alpha)$ varies from ~ 0.66 near the center of Taurus A to ~ 0.75 near the edges. However, even at the short wavelength of 1.55 cm the fractional f `larization p_i reaches a maximum of only ~ 0.16 near the center and falls to less than half this value near the edges (Mayer and Hollinge: 1968). The field obviously has a sizeable random component, therefore, which can be estimated from the relation (Burn 1966)

$$p_{i} = p(\alpha) \frac{H_{o}^{2}}{H_{o}^{2} + H_{r}^{2}}$$

Using the measurements at λ 1.55 cm as indicative of the intrinsic degree of polarization p_i and our p(α) distribution, we estimate that the ratio $(H_r/H_o)^2$ varies from ~ 3 near the center to ~ 9 near the edges.

Knowing the distributions of both $(H_r/H_0)^2$ and $\sqrt{H_0^2+H_r^2}$, we can estimate the distributior of the strengths of the uniform

magnetic field. Referred to the peak of the polarization distribution (~0!5 southeast of our coordinate origin), the uniform field H_{o} , is

- (a) \sim 1.2x10⁻⁴ G within the central 0:5;
- (b) ~ 0.9×10^{-4} G at 1:0-1:5;
- (c) ~ 0.6×10^{-4} G beyond 1!5.

The orientation of this uniform field can also be determined. The components normal to the line of sight are perpendicular to the intrinsic position angle vectors illustrated in Fig. 4. Since, except at the northwestern edge, nearly all of these vectors are within 10° of position angle 149°, the normal component, H_{o1} , is close to position angle 59° over the main body of the nebula, i.e., ~ 14° greater than the position angle of the minor axis. Further, from Table 4 we see that the rotation measure produced within the Crab Nebula is everywhere positive. Hence, the lineof-sight component of the uniform magnetic field, $H_{o||}$ is everywhere directed toward was. If we assume that $\prod_{o1}^{2} \approx \frac{H_{o}}{2} \sum_{o1}^{2} \approx \frac{1}{2}E_{o}$, then the strength of $H_{o||}$ is 0.7 times the values of H_{o} listed above.

B. The Thermal Plasma.

From the known rotation measures in Taurus A (Table 4) we obtain directly the product $\langle n_e \ ' \circ || L \rangle$, where L is the pathlength through the nebula parallel to the line of sight. Assuming again that the nebula is a uniformly filled prolate spheroid with major axis in the plane of the sky, we can easily calculate L. We then have the product $\langle n_e \ H_o || \rangle$ of thermal electron density n_e and line-of-sight uniform field $H_{0||}$. This product is displayed in Table 7. We can also estimate $\langle n_e | H_0 || \rangle$ from the depolarization produced by a uniform medium. The values calculated by this latter method are all larger. In other words, the measured depolarization exceeds that predicted from the rotation produced in a uniform slab. This reinforc.s the picture of coexisting uniform and finely structured media within the nebula: i.e., come radiation is depolarized by structure so fine that there is no net rotation.

We note that $\langle n_{e} | H_{o||} \rangle$ is approximately constant along lines parallel to the major axis. Since we previously estimated that H_{o} decreased from center to edge, the electron density appears to increase near the edges. We estimate n_{e} 0.25 cm⁻³ near the center of the nebula, increasing more or less smoothly in all radial directions to ~ 0.5 cm⁻³ at the periphery. Even larger values may be found at the northern boundary and at the extreme southwestern edge. Clearly our assumption of a uniformly dense medium was at best a crude approximation to reality. More likely, the thermal gas has a more shell-like distribution. This would support the ideas of Scargle (1969) and Rees (1971), who have argued that the pulsar activity in the Crab Nebula evacuates the thermal gas from the central region.

-13-

. SUMMARY

From our investigations of the Crab Nebula we find that: (1) The spectrum of the radiation is flattest at the center and progressively steepens in all outward directions.

(2) The magnetic field strength is $\sim 2.5 \times 10^{-4}$ G at the center and decreases to 75-80 per cent of this maximum value near the periphery.

(3) There is structure on a scale much less than our resolution (<<0.5 pc) co-mixed with a uniform large-scale medium.

(4) The strength of the small-scale random component of the magnetic field is ~ 1.8 times that of the uniform component near the center, changing to ≥ 3 times near the periphery.

(5) Throughout most of the nebula the projection of the uniform magnetic field onto the plane of the sky is near position angle 60°; there is also a component out of the plane of the sky toward us.

(6) The central region of the ...bula has a relatively less dense thermal plasma than the periphery.

-14-

ACKNOWLEDGEMENTS

and the second and the second of the second s

2

We wish to thank the members of the Owens Valley Staff and observers who have greatly assisted the collection and calibration of the data. We further wish to thank D. H. Rogstad for his assistance and discussion of inversion techniques and the members of the Kapteyn Laboratory, Groningen, Netherlands for their encouragement and the use of their facilities. We benefited greatly from discussions with M. C. H. Wright. Research at the Owens Valley Radio Observatory is supported by the Office of Naval Research under Contract N00014-67-A-0094-0019 and by the National Science Foundation under Grant GP30400-X. and a second and an and an analysis of the second second

-15-

REFERENCES

Branson, N.J.B.A. 1967, Nature 213, 1211.

Braun, L. D., and Yen, J. L. 1968, Ap. J. (Letters) 153, L123.

Burn, B. J. 1966, M.N.R.A.S. 133, 67.

Conway, R. G. 1971, Radio polarization of the Crab Nebula in

The Crab Nebula, Eds. R. D. Davies and F. G. Smith, D. Reidel Publishing Co., Dordrecht, Holland, p. 292.

<mark>kulukunda ana kanana sana da sana da suruku surukuna da surukunda sana sana sana suruku suruku suruku suruku sa</mark>

Davies, R. D., Gardner, F. F., Hazard, C., and Mackey, M. B. 1966, Australian J. Phys. 19, 409.

Downs, G. S., and Thompson, A. R. 1968, Ap. J. (Letters) 152, L65. Forman, W., and Visvanathan, N. 1971, <u>Nature 229</u>, 39.

Ginzburg, V. L., and Syrovatskii, S. I. 1965, Ann. Rev. Astron.

and Astrophys. 3, 297.

Hogg, D. E., Macdonald, G. H., Conway, R. G., and Wade, C. M.

1969, A. J. <u>74</u>, 1206.

Johnston, K. J., and Hobbs, R. W. 1969, Ap. J. 158, 145.

Manchester, R. N. 1971, Nature Phys. Sci. 231, 189.

Mayer, C. H., and Hollinger, J. P. 1968, Ap. J. 151, 53.

Milne, D. K. 1971, Padio emission from supernova remnants in

The Crab Nebula, Eds. R. D. Davies and F. G. Smith, D. Reidel Publishing Co., Dordrecht, Holland, p. 248.

Minkowski R. 1971, Comments on supernova remnants and ancient novae, ibid, p. 241.

Münch, G. 1958, Rev. Mod. Phys. 30, 1042.

Rees, M. J. 1971, Nature Phys. Sci. 230, 55.

Rogstad, D. H., and Shostak, G. S. 1971, Astron. and Astrophys.

13, 99.

Scargle, J. D. 1969, Ap. J. 156, 401.

Seielstad, G. A., and Weiler, K. W. 1968, Ap. J. 154, 817.

Shklovsky, I. S. 1968, Supernovae. John Wiley & Sons Ltd., London. Trimble, V. 1968, <u>A. J. 73</u>, 535.

Weiler, K. W., and Seielstad, G. A. 1971, <u>Ap. J. 163</u>, 455, Paper I. Wilson, A. S. 1971, High resolution maps of the Crab Nebula at

2700 MHz and 5000 MHz in <u>The Crab Nebula</u>, Eds. R. D. Davies and F. G. Smith, D. Reidel Publishing Co., Dordrecht, Holland, p. 68.

Woltjer, L. 1957, B.A.N. 13, 301.

1958, ibid. <u>14</u>, 39.

Wright, M. C. H. 1970, M.N.R.A.S. 150, 271.

Taurus A Error Estimates

	Error							
Stokes	1420 MHz -2	2880 MHz _2						
Parameter	(f.u. arcmin ²)	(f.u. arcmin ²)						
I	$4.5 \times 10^{\circ}$	5.7 x 10 ⁰						
Q	1.8×10^{-1}	1.1×10^{-1}						
U	6.0×10^{-2}	5.1×10^{-1}						

ł

South a state of the second of the second

فللطفط بالمعادلة والمتعادية والمتعالمة والمعالمة والمعارية والمتعادية

Samo

こうちょうシブラン

Leters straiter keiterstrauterieter der Let. "Auster eine Kritter erken aufenderte Verklichen Kerken des Verkrauteriete

Spectral Index Distribution $(I_v \sim v^{+\alpha})$

						17.000 00 1 0.1			
3:0				-0.75	-0.69				
2:5		-1.09	-0.69	-0.60	-0.53	-0.59	-0.99		
2:0		-0.72	-0.52	-0.50	-0.47	-0.58	-0.70		
1.5	-1.61	-0.69	-0.51	-0.47	-0.42	-0.41	-0.54	-0.91	
1:0	-0 . 99	-0.56	-0.45	-0.42	-0.40	-0.43	-0.54	-0.70	
n) 0:5	-0.83	-0.47	5 -0.34	5 -0.36	5 -0.33	6 -0.35	6 -0.48	3 -0.54	7 -0.65
(arcmi) 0:0	-0.81	-0.49	-0.35	-0.36	-0.36	-0.3	-0.4	• • • •	- 0 - ·
Axis 0.5	-0.65	-0.44	-0.37	-0.42	-0.46	-0.44	-0.51	-0.60	0 • 34
Major 1:0	-0.58	-0.44	-0.38	-0.42	-0.45	-0.44	-0.50	-0.61	·
1.5	-0.57	-0.50	-0.49	-0.54	-0.53	-0.48	-0.55	-0 67	
2:0	-0.69	-0.58	•0.55	-0.60	-0.58	-0.54	-0.64	-	
2:5	-1.00	-0.79	-0.6 .	-0.67	-0.66	-0.64		2 M	· ·
3.0		-1.02	-0.89	-0.96	-1.05			ш.	· .
	1.5	1.0	0.5	0:0	0 • 5	1,0	1,5	2:0	2.5
\bigvee			ເພາຍ)	kis (ar	linor A	Ŵ			-6

Rotation Measure Distribution (rad m^{-2})

							· <u>····</u> • •• ••	
2:5			28	-27		NK		
2:0		8 1	- 25				\rangle	
1.5		- 22	-22			ណ		
1:0		-21	- 23	-23	-21			
amin) 0:5		- 22	-23	-23	-21	-19	-19	
cis (arc 0:0		-24	- 23	-23	-21	-20	-18	-16
Aajor Ay 0.5	-25	-25	-24	-23	-21	-19	-16	
	-27	-26	-25	-23	-21	-19		
1.5	-28	- 28	-26	-24	-21	-18		
2:0			-30	-26				
	1.5	1,0	0.5	0	0.5	0 	1.5	5,0
			(uīm:		XA TORİ	W		

20

127.00

1000

ï

:

1 1

Internal Rotation Measure Distribution (rad m^{-2})



arterstel – Arterien Wester bereuternen Subjection here der päärmännin (1989) (1999)

いたいまたとうないというないとうとうとう

	•			• ;					
		+		1	22				
	:	:	. !,	i					
	2.5		:	0.55	0.62		ZĦ		
	2:0		- 0.41 	0.47	;	<i>,</i>		>	
-(08)	1.5		0.44	0.58			ш¥ ш		
420 ^{/P} 28	1:0		0.50	0.57	0.54	0•54			
ion (F _{.1}	in) 0.5.		0.48	0.48	0.46	6.47	0.54	0.63	
stribut	s (arcm 0.0		0.43	0.44	0.44	0.41	0.46	0.58	0.66
atio Di:	jor Axis	0.47	0.44	. 43	0.41 (. 36	0.37 (0.51 (-
tion Ra	Ma 0	.48 (.47 (.45 (.44 (.41 (.40 (0	
olarize	5. 1	.50.0	.52 0		- 51 C	• 21 • 21	.49 C		
Dep			•	.81 0	0.08.0	0	0		
;	5		•	0 1	0				
·		. <mark>1</mark> .			0 	0.5	1,0	1,5	2:0
		· ·		 (атстіп)	r Axis	oniM		- 1994 F	

ł

;

:

;,

ļ

í

Magnetic Field Strength Distribution (10⁻⁴ Gauss)

					02				
	·				25	<u></u>			
3.0				2.0	2.0				
2.5		2.7	2.1	2.0	2.0	2.0			
2.0		2.0	2.1	2.2	2.2	2.1	2.1		
1,5		2.1	2.2	۲. ۲	2.2	2.2	2.0	2.7	
1:0	2.7	2.1	2.3	2.3	2.3	2.2	2.1	2.1	
0,5	2.3	2.1	2.4	2.4	2.5	2.4	2.1	2.0	
0,0	2.3	2.2	2	2.4	2•5	2.4	2.2	2.0	2.5
0.5	2.0	2.1	2.3	2.4	2.4	2.3	2.2	2.0	:
1,0	1.9	2.1	2.2	2.3	2.3	2.2	2.1	1.9	:
1:5	1.8	2.0	2.1	2.2	2.2	2.2	2.0	1.9	
2:0	1.9	2.0	2.1	2,1	2.1	2.0	1.9	zĸ	,
2:5	3.2	2.4	2.2	2.1	2.1	1.9			> .
3:0			2.8	2.8	·: •:			ц у	
	1:5	1,0	0.5	0;0	0,5	1,0	 	2:0	5.5

Party (Statistics)

er.

Fi Ye

.---

Distribution of $\langle n_e H_o || \rangle$ (10⁻⁵ cm⁻³ Gauss)

<u> </u>]				24				
2.5			2.3	2.3		ZK		
2:0		2.1	2.2				\rangle	
1,5		2.4	2.4			д		
1,0		2.4	1.9	1.9	2.3			
cmin) 0.5		2.2	1.9	2.0	2.2	2.5	2.9	
tis (arc 0.0		1.9	1.9	2.0	2.1	2.5	3.0	4.3
4ajor A) 0.5	2.1	1.8	1.9	2.0	2.2	2.5	3.2	
1:0	2.1	1.9	1.9	2.0	2.3	2.7		
1.5	2.1	1.7	1.8	2.0	2.4	3.0		
2.0			1.6	2.0				
		1;0	0.5 5	0;0	0.55	1,0	1,5	5
V			(ແກ່ຫວ	ıs) six	A roniM	[

;

- - -

FIGURE CAPTIONS

Fig. 1. Portion of the Fourier (u, v)-plane covered at the declination of Taurus A.

(a) 1420 MHz; 0-1600 feet E/W and N/S in basic increments of 200 feet; 100 feet spacing included E/W; 1000 and 1200 foot spacings missing N/S. Crosses mark every hour of hour angle; scales in thousands (10^3) of wavelengths.

(b) 2880 MHz; 0-800 feet E/W in increments of 100 feet; 0-800 feet N/S in increments of 200 feet; 500 foot spacing missing E/W. Crosses mark every hour of hour angle; scales in thousands (10^3) of wavelengths.

Fig. 2. Distributions of total and linearly polarized radiation over Taurus A at 1420 MHz. North at top and east on left. Origin at $\alpha_{1950} = 05^{h}31^{m}31^{s}20$, $\delta_{1950} = 21^{\circ}59^{\circ}17$. and a start of the
(a) total radiation; half-power beamwidth shown in upper left-hand corner; $10=76.8 \times 10^{-26} \text{W m}^{-2} \text{Hz}^{-1} \text{ arcmin}^{-2}$.

With the state of the second state and the second state of the

(b) polarized radiation; 20 per cent contour darkened; $10=2.3 \times 10^{-26} \text{W m}^{-2} \text{ Hz}^{-1} \text{ arc min}^{-2}$.

(c) position angle of the polarization; orientation of the line gives position of the electric vector; degree of linear polarization indicated by line length; line length appropriate for 5 per cent polarization shown in lower left-hand corner. Fig. 3. Taurus A; 2880 MHz (see caption to Fig. 2).

(a) total radiation; $10=56.2 \times 10^{-26} W m^{-2} Hz^{-1} arcmin^{-2}$.

(b) polarized radiation; 20 per cent contour darkened; $10=4.3 \times 10^{-26} W m^{-2} Hz^{-1} arcmin^{-2}$.

(c) electric vectors; line length appropriate for 10 per cent polarization shown in lower left-hand corner.

Fig. 4. Intrinsic position angle of the electric vector for Taurus A. <u>Solid line</u>, delimiting 1420 MHz contour; <u>dashed line</u> delimiting 2880 MHz contour. Line length has no significance. tion is the construction of the state of the

UNE DENNESSERVER CONTRACTOR





the second s





2. Metransmichter of



Fig. 4

A street survey maker when a