



AD 745920

AD

Reports Control Symbol
OSD-1366


RESEARCH AND DEVELOPMENT TECHNICAL REPORT
ECCOM-5436

THE ACCURACY OF BALLISTIC DENSITY DEPARTURE TABLES 1934-1972

By

Marvin J. Lowenthal

April 1972



Approved for public release; distribution unlimited.

ECCOM

UNITED STATES ARMY ELECTRONICS COMMAND - FORT MONMOUTH, NEW JERSEY

Reproduced by
NATIONAL TECHNICAL
INFORMATION SERVICE
U S Department of Commerce
Springfield VA 22151

40

NOTICES

Disclaimers

The findings in this report are not to be construed as an official Department of the Army position, unless so designated by other authorized documents.

The citation of trade names and names of manufacturers in this report is not to be construed as official Government endorsement or approval of commercial products or services referenced herein.

Disposition

Destroy this report when it is no longer needed. Do not return it to the originator.

ACCESSION	
CFSTI	REF. SECTION <input checked="" type="checkbox"/>
DDC	DIFF. SECTION <input type="checkbox"/>
UNANNOUNCED	<input type="checkbox"/>
DISTRIBUTION
.....
E.V.
DISTRIBUTION/AVAILABILITY CODES	
DIST.	APPL. AND/OR SPECIAL
AI	

UNCLASSIFIED

Security Classification

DOCUMENT CONTROL DATA - R & D

(Security classification of title, body of abstract and indexing annotation must be entered when the overall report is classified)

1. ORIGINATING ACTIVITY (Corporate author) Atmospheric Sciences Laboratory White Sands Missile Range, New Mexico		2a. REPORT SECURITY CLASSIFICATION Unclassified	
		2b. GROUP	
3. REPORT TITLE THE ACCURACY OF BALLISTIC DENSITY DEPARTURE TABLES 1934-1972			
4. DESCRIPTIVE NOTES (Type of report and inclusive dates)			
5. AUTHOR(S) (First name, middle initial, last name) Marvin J. Lowenthal			
6. REPORT DATE April 1972		7a. TOTAL NO. OF PAGES 29	7b. NO. OF REFS 5
8a. CONTRACT OR GRANT NO.		9a. ORIGINATOR'S REPORT NUMBER(S) ECOM-5436	
b. PROJECT NO.		9b. OTHER REPORT NO(S) (Any other numbers that may be assigned this report)	
c. DA Project No. IT062111A126-05-57			
d.			
10. DISTRIBUTION STATEMENT Approved for public release; distribution unlimited.			
11. SUPPLEMENTARY NOTES		12. SPONSORING MILITARY ACTIVITY U.S. Army Electronics Command Fort Monmouth, New Jersey	
13. ABSTRACT The accuracy of ballistic density departure tables is examined, starting with the earliest available sets in 1934. The extension of the tables (originally developed for the US) to encompass the entire Northern Hemisphere is discussed and the shortcomings of the current climatological regional zones described. New tables, based on current data and used for a more limited geographical area, are shown to be accurate to one half of one percent, hence furnish excellent back-up information when a current sounding is not available for artillery firings. A procedure for minimizing ballistic density errors that accrue between observational periods is also presented.			

DD FORM 1473
1 NOV 65

REPLACES DD FORM 1473, 1 JAN 64, WHICH IS OBSOLETE FOR ARMY USE.

UNCLASSIFIED

Security Classification

UNCLASSIFIED

Security Classification

14. KEY WORDS	LINK A		LINK B		LINK C	
	ROLE	WT	ROLE	%	ROLE	WT
1. Ballistic Density 2. Air Density 3. Radiosondings 4. Station Elevation 5. Geographical Regions 6. Meteorological Messages						
<i>ii</i>						

UNCLASSIFIED

Security Classification

Reports Control Symbol
OSD-1366

Technical Report ECOM-5436

THE ACCURACY OF BALLISTIC DENSITY
DEPARTURE TABLES 1934-1972

By

Marvin J. Lowenthal

Atmospheric Sciences Laboratory
White Sands Missile Range, New Mexico

April 1972

DA Project No. IT062111A126-05-57

Approved for public release; distribution unlimited.

ija

U. S. Army Electronics Command

Fort Monmouth, New Jersey

ABSTRACT

The accuracy of ballistic density departure tables is examined, starting with the earliest available sets in 1934. The extension of the tables (originally developed for the US) to encompass the entire Northern Hemisphere is discussed and the shortcomings of the current climatological regional zones described.

New tables, based on current data and used for a more limited geographical area, are shown to be accurate to one half of one percent, hence furnish excellent back-up information when a current sounding is not available for artillery firings.

A procedure for minimizing ballistic density errors that accrue between observational periods is also presented.

CONTENTS

	Page
INTRODUCTION	1
CONSTRUCTION OF THE TABLES	16
CONCLUDING REMARKS	24
APPENDIX: COMPUTER DETERMINATION OF METRO MESSAGES FROM RADIO- SONDE DATA	25
Procedures	25
Method of Solution	25
Appendix I: Calculation of Zone Temperature	28

Preceding page blank

INTRODUCTION

One of the basic premises of all firing tables is the "Standard Atmosphere" - one whose properties are enumerated in the "1962 Standard Atmosphere" [1]. With such a density distribution and zero wind throughout, the shell should describe the calculated trajectory given in the firing tables. Those conditions are never encountered on earth, hence corrections for nonstandard conditions are always in order. Artillery Meteorological Sections, organic to the Army Artillery, provide meteorological information on winds and density to permit such corrections.

Accurate, representative, fresh observations are the best method of correction available today. There are, however, occasions when these measurements are not available, and alternate methods are necessary to provide the requisite information. In the case of atmospheric density such an option is available.

Air density at ground level can always be determined, the only equipment required being a good barometer and wet- and dry-bulb thermometers. If the upper air density can be inferred from the surface measurement, the ballistic density can be furnished for correction of artillery fire. The vital question is: Is the assumption justified that the surface density is a good predictor of ballistic density aloft?

To test the theory, some 3000 radiosondings from Vietnam and 2000 from the Korean region were examined. Figure 1 shows the mean of ballistic density aloft as a function of surface density over a wide range of surface values.* Values of ballistic density are expressed as a percent




FIG-1 VARIATION OF BALLISTIC DENSITY I

[1] US Standard Atmosphere, US Govt. Printing Office, Dec. 1962.

*The values on the abscissa of Figure 1 represent $\pm 0.2\%$ on each side of the central value.

of standard with the ordinate being the normal NATO zones (Table 1).

TABLE I
BALLISTIC ZONES, STANDARD ATMOSPHERE

<u>Line No.</u>	<u>Height at Top of Zone (Meters)</u>	<u>Temperature (°K)</u>	<u>Density (g/m³)</u>
0	Surface	288.2	1225.0
1	200	287.5	1213.3
2	500	286.9	1184.4
3	1000	283.3	1139.2
4	1500	280.0	1084.6
5	2000	276.8	1032.0
6	3000	271.9	957.0
7	4000	265.5	863.4
8	5000	259.0	777.0
9	6000	252.5	697.4
10	8000	242.7	590.0
11	10000	229.8	467.0
12	12000	216.8	364.8
13	14000	216.7	266.6
14	16000	216.7	194.8
15	18000	216.7	142.3

As can be seen, the ballistic densities are a function of the surface value. The various curves do not intersect, although the spread between the extremes continually decreases. The values all seem to trend toward a common value at some higher level, beyond the limit of our data. The curves do show that ballistic density is more independent of the surface value at the high line numbers than at low line numbers. Thus, a climatological value of ballistic density at high altitudes (line 10 or above) would have a very small error for all ranges of surface density. This is most fortunate since the range errors due to an incorrect assessment of ballistic density are greatest for high maximum ordinates.

In absolute values, the ballistic density curves exhibit a behavior similar to that of the Standard Atmosphere (Figure 2). Again, the convergence of all curves is seen with increasing altitude. It should be noted that the curves for the tropical region (Vietnam) converge toward a different value than those from the nontropical Korea. This difference indicates that one set of values is not satisfactory for the entire




FIG.2 VARIATION OF BALLISTIC DENSITY

globe, regional tables being required to minimize the error in the estimate of the ballistic density.

It has been shown, therefore, that a surface density measurement can be used to categorize ballistic densities aloft for a particular geographic region; that a climatological value of ballistic density becomes more accurate with increasing height (at least to the levels of interest for conventional tube artillery); and that all geographic localities cannot be accurately described by a single set of values.

Once the feasibility of the procedure is established, it is necessary to determine the accuracy of such a technique. It may well be that even though the mean curves of ballistic density do not intersect, the dispersion within each set is so great that the probability of an accurate estimate from the mean is quite small. This involves, of course, the study of the standard deviations of density for any given surface value. A portion of the computer analysis for both Vietnam and Korea is shown in Table II.

MET DATA ANALYSIS
CALCULATION OF DENSITY VS. SURFACE DENSITY ON SEASONAL BASIS - VIETNAM DATA -
ALTI 001 WGT 000 SEASONAL PERIOD SEP. 1 TO NOV. 31 YEARS 1964 TO 1967 TIME PERIOD 04-11

SEC.	LINE 1				LINE 2				LINE 3				LINE 4				LINE 5				SEC.	
	XRAN	STD.	NO.	XRAN	STD.	NO.	XRAN	STD.	NO.	XRAN	STD.	NO.	XRAN	STD.	NO.	XRAN	STD.	NO.	XRAN	STD.		NO.
80.0	94.0	0.00	1	94.0	0.00	1	94.0	0.00	1	94.0	0.00	1	94.0	0.00	1	94.0	0.00	1	94.0	0.00	1	89.4
80.1	94.0	0.00	1	94.0	0.00	1	94.0	0.00	1	94.0	0.00	1	94.0	0.00	1	94.0	0.00	1	94.0	0.00	1	89.0
80.2	94.0	0.00	1	94.0	0.00	1	94.0	0.00	1	94.0	0.00	1	94.0	0.00	1	94.0	0.00	1	94.0	0.00	1	89.1
80.3	94.0	0.00	1	94.0	0.00	1	94.0	0.00	1	94.0	0.00	1	94.0	0.00	1	94.0	0.00	1	94.0	0.00	1	89.2
80.4	94.0	0.00	1	94.0	0.00	1	94.0	0.00	1	94.0	0.00	1	94.0	0.00	1	94.0	0.00	1	94.0	0.00	1	89.3
80.5	94.0	0.00	1	94.0	0.00	1	94.0	0.00	1	94.0	0.00	1	94.0	0.00	1	94.0	0.00	1	94.0	0.00	1	89.4
80.6	94.0	0.00	1	94.0	0.00	1	94.0	0.00	1	94.0	0.00	1	94.0	0.00	1	94.0	0.00	1	94.0	0.00	1	89.5
80.7	94.0	0.00	1	94.0	0.00	1	94.0	0.00	1	94.0	0.00	1	94.0	0.00	1	94.0	0.00	1	94.0	0.00	1	89.6
80.8	94.0	0.00	1	94.0	0.00	1	94.0	0.00	1	94.0	0.00	1	94.0	0.00	1	94.0	0.00	1	94.0	0.00	1	89.7
80.9	94.0	0.00	1	94.0	0.00	1	94.0	0.00	1	94.0	0.00	1	94.0	0.00	1	94.0	0.00	1	94.0	0.00	1	89.8

Table II Vietnam

MET DATA ANALYSIS
CALCULATION OF DENSITY VS. SURFACE DENSITY ON SEASONAL BASIS - KOREA DATA - BALLISTIC ZUNFS
ALTI 001 WGT 000 SEASONAL PERIOD SEP. 1 TO NOV. 31 YEARS 1964 TO 1967 TIME PERIOD 20-07

SEC.	LINE 1				LINE 2				LINE 3				LINE 4				LINE 5				SEC.	
	XRAN	STD.	NO.	XRAN	STD.	NO.	XRAN	STD.	NO.	XRAN	STD.	NO.	XRAN	STD.	NO.	XRAN	STD.	NO.	XRAN	STD.		NO.
92.9	94.0	0.00	1	94.0	0.00	1	94.0	0.00	1	94.0	0.00	1	94.0	0.00	1	94.0	0.00	1	94.0	0.00	1	93.4
93.0	94.0	0.00	1	94.0	0.00	1	94.0	0.00	1	94.0	0.00	1	94.0	0.00	1	94.0	0.00	1	94.0	0.00	1	93.5
93.1	94.0	0.00	1	94.0	0.00	1	94.0	0.00	1	94.0	0.00	1	94.0	0.00	1	94.0	0.00	1	94.0	0.00	1	93.6
93.2	94.0	0.00	1	94.0	0.00	1	94.0	0.00	1	94.0	0.00	1	94.0	0.00	1	94.0	0.00	1	94.0	0.00	1	93.7
93.3	94.0	0.00	1	94.0	0.00	1	94.0	0.00	1	94.0	0.00	1	94.0	0.00	1	94.0	0.00	1	94.0	0.00	1	93.8
93.4	94.0	0.00	1	94.0	0.00	1	94.0	0.00	1	94.0	0.00	1	94.0	0.00	1	94.0	0.00	1	94.0	0.00	1	93.9
93.5	94.0	0.00	1	94.0	0.00	1	94.0	0.00	1	94.0	0.00	1	94.0	0.00	1	94.0	0.00	1	94.0	0.00	1	94.0

Table II Korea

The standard deviations (STD) are seen to be of the order of 0.5% or less, which suggests that ballistic density aloft can be estimated to about 1/2% from a measurement of surface density alone. Since density is calculated from measurements of pressure and temperature from the radiosonde, the accuracy of the density measurement is dependent upon the accuracy of the pressure and temperature sensors. By use of current specification criteria, the error of the density determination is found to be 0.3 - 0.4% at low line numbers and greater than 0.5% at high line numbers. An assessment of ballistic densities to about 1/2 to 1% by a surface measurement alone is an excellent back-up system that is always available where a radiosonde observation of upper air density cannot be made.

Historically, the use of density tables for the assessment of ballistic densities was the standard procedure before radiosondes were available. The earliest manual available for this report was TR-1236-1, "Meteorological Message for the Artillery," 1934. At that time it was stated:

"It is impractical to actually measure the temperature, pressure, and moisture content of the atmosphere at various heights, compute the ballistic densities, and get the computed data to the Artillery without the elapse of considerable time. As atmospheric conditions are continually changing, there must be as little delay as possible between the times that meteorological observations are made and the times that the completed reports are available to the Artillery. No attempt is made, therefore, to determine air densities from observations made at various heights above the ground. Instead, the air density is determined near the ground at the meteorological station and the air density is assumed to decrease at a definite rate in height above the ground." [2]

Three decades of technological advances have, of course, made it possible to obtain an atmospheric measurement of temperature and density and transmit it to the Fire-Direction Center. The other statements concerning atmospheric variability and the necessity for fresh metro information are as true today as when the original words were written.

The tables given in TR 1236-1 were meant to apply to the US only (page 24) and values of ballistic density were given for station elevations near sea level, and at 1000 and 2000 feet above mean sea level. For higher elevations, the data given in the table for stations located 2000 feet above sea level may be used [3].

[2] TR 1236-1, pp 23-24.

[3] Ibid., pp 82, 92.

The data, in part, are shown in Figure 3. The plot shows recognition




FIG.3 BALLISTIC AIR DENSITY FROM TR 1236-1 (1934)

of (1) the convergence of all values of ballistic density at higher levels, (2) the difference in behavior of ballistic densities with increasing station elevation.

The main sources of error in the tables are (1) use of the values at 2000 feet for stations at higher elevations, and (2) assumption that one set of values will suffice for the entire US. To correct these errors, a study was carried out by the Signal Corps General Development Laboratory, Fort Monmouth, New Jersey, to revise and extend the tables [4]. The results of this study appeared in Sep 1942 as Change 4 to Technical Manual TM 4-240 which superseded TR 1236-1 in Dec 1941.

As a result of that study, the US and its Western Hemisphere territories (Alaska, Hawaii, Canal Zone, Puerto Rico, etc.) were divided into six geographical regions**. The boundaries of the regions in the US were

[4] "Preparation and Evaluation of Revised Ballistic Density Tables," Brasefield, C., SCL Eng Report #760, SigC Gen Dev Lab, Ft. Monmouth, New Jersey.

**Region 7, the Pacific Northwest, is not mentioned in Ch 4 to TM 4-240, 17 Sep 1943. It first appears in TM 20-240 which superseded TM 4-240 in Nov 1944. No data are available on the introduction of this new region.

the same as shown in the current FM 6-16. Appendix I, Change 4 to TM 4-240 dated 17 Sep 1943, extended the valid geographical limits of the density regions to include the entire Northern Hemisphere. Figure 4 below, a reproduction of 10, shows the revisions.

TM 4-240
C 4

TECHNICAL MANUAL

APPENDIX I

METEOROLOGICAL TABLES

Table X-1n (added by C 1, 7 Sept. 1942), change title to read as follows: FOR ANTI-AIRCRAFT ARTILLERY AND OTHER HIGH-ANGLE FIRE; VALID DURING THE NIGHT IN REGION 1 (EASTERN U. S. A., BRITISH ISLES, COAST OF NORTH AFRICA, AND EUROPE, EXCEPTING ALPINE REGION, SCANDINAVIAN PENINSULA, AND RUSSIA NORTH OF LATITUDE 55° N.).

Table X-1a (added by C 1, 7 Sept. 1942), change title to read as follows: FOR ANTI-AIRCRAFT ARTILLERY AND OTHER HIGH-ANGLE FIRE; VALID DURING THE AFTERNOON IN REGION 1 (EASTERN U. S. A., BRITISH ISLES, COAST OF NORTH AFRICA, AND EUROPE, EXCEPTING ALPINE REGION, SCANDINAVIAN PENINSULA, AND RUSSIA NORTH OF LATITUDE 55° N.).

Table X-3n (added by C 1, 7 Sept. 1942), change title to read as follows: FOR ANTI-AIRCRAFT ARTILLERY AND OTHER HIGH-ANGLE FIRE; VALID DURING THE NIGHT IN REGION 3 (WESTERN U. S. A. AND ALPINE REGION OF SOUTHERN EUROPE).

Table X-3a (added by C 1, 7 Sept. 1942), change title to read as follows: FOR ANTI-AIRCRAFT ARTILLERY AND OTHER HIGH-ANGLE FIRE; VALID DURING THE AFTERNOON IN REGION 3 (WESTERN U. S. A. AND ALPINE REGION OF SOUTHERN EUROPE).

Table X-5 (added by C 1, 7 Sept. 1942), change title to read as follows: FOR ANTI-AIRCRAFT ARTILLERY AND OTHER HIGH-ANGLE FIRE; VALID IN REGION 5 (ALASKA, ICELAND, SCANDINAVIAN PENINSULA, AND RUSSIA NORTH OF LATITUDE 55° N.).

Table X-6n (added by C 1, 7 Sept. 1942), change title to read as follows: FOR ANTI-AIRCRAFT ARTILLERY AND OTHER HIGH-ANGLE FIRES; VALID DURING THE NIGHT IN REGION 6 (WEST INDIES, CANAL ZONE, HAWAII, AND SOUTHWEST PACIFIC AREA).


Table X-6a (added by C 1, 7 Sept. 1942), change title to read as follows: FOR ANTI-AIRCRAFT ARTILLERY AND OTHER HIGH-ANGLE FIRE; VALID DURING THE AFTERNOON IN REGION 6 (WEST INDIES, CANAL ZONE, HAWAII, AND SOUTHWEST PACIFIC AREA).

10

Figure 4. Meteorological Table (from TM 4-240)

C 4

An examination of the current Chart I (pp 5-6, FM 6-16, May 1961) Figure 5 shows an apparent inconsistency. Note the wording for Region 1 "Eastern USA, and Europe, excepting Alpine Regions, Scandinavian Peninsula, and Russia north of latitude 55°N." and that for Region 5, "Alaska, Iceland, Scandinavian Peninsula, and Russia north of latitude 55°N." Clearly the intent was to place Scandinavia and Northern Russia into Region 5, and exclude those areas from the rest of Europe which were included in Region 1. The numbers 1 and 5 in the Eastern Hemisphere are misplaced. Likewise the limits of Region 3, "Western USA and Alpine Region of Southern Europe" has been rather broadly interpreted - the eastern Mediterranean and Iraq are scarcely Alpine regions of Southern Europe. Corrections to the current regional density map will be proposed for inclusion into the next revision of FM 6-16, tentatively scheduled for the second quarter of FY-73.



Time is calculated from the meridian of Greenwich. The middle of the zero time zone passes through Greenwich with its east and west limits 7°30' on each side. Each 15 degree zone east and west of the initial zone represents one hour of time. The number of hours that must be added to or subtracted from local standard time to give Greenwich time is indicated for each zone.

Political boundaries in the various countries have caused modifications of the time zones. The vertical lines and clear sections are used to show to which zones these divisions belong. Where a half hour difference is legal, horizontal lines are used. Where no time system has yet been adopted, the area is represented by small dots. Where no legal time has been established, the larger dots are used. Variations from zone time are given in hours and minutes.


The seven climatic regions of the Northern Hemisphere are indicated and identified by the large black numbers 1 through 7. (See par. 164d, FM 6-15.)

Figure 5. Time zones and climatological regions of the Northern Hemisphere.

The departure tables themselves have been only slightly modified through the years, the greatest change occurring with the issuance of FM 6-16 in May 1961, owing to the change in units from yards to meters, and the adoption of a new standard surface density, 1225 g/m³ instead of the older value of 1203.4 g/m³.

Some of the ballistic density profiles from TR 1236-1 and the revision are shown in Figure 6. Surface densities 5% above the mean and 5% below the mean were chosen to emphasize the difference in high and low surface density conditions.***

***Prior to 1961 (FM 6-16) the mean surface density of a meteorological station at sea level was 103%. Currently, the standard mean surface density for a sea-level station is 100%.



The profiles show wide variations between regions, in the example, as high as 3%. The profile from TR 1236-1 falls between the extremes and is not coincident with any single region - nor is it the mean of all regional values. For 98% and 108%, the old profile is closest to that of Region 5, the Alaskan area. The figure clearly shows that a geographical area the size of the US cannot be adequately described with a single density profile. Thus, it is difficult to comprehend why almost the entire Eurasian Continent from the Baltic Sea all the way to the Sea of Japan was considered a single region, or why the Northern Siberian regions were adjudged similar to Eastern US rather than the Polar Regions of North America, or why the mountainous Western US is similar to the lands around the northern borders of the Mediterranean.

The revised tables do show, however, that the profiles are dependent upon the elevation of the station, since a given surface density may represent mean conditions, above-normal surface density, or abnormally low surface density depending on whether the station in question is at sea level or at an elevated location. The behavior of the profile of 100% surface density at a sea-level station, at 1000', and at 2000' above MSL is shown in Figure 7 for both old and revised tables, Region 5 being used since it appears closest to the values in TR 1236-1.




FIG7. VARIATION OF BALLISTIC DENSITY WITH STATION ELEVATION

The figure clearly indicates that ballistic density aloft decreases most rapidly in the cases of high surface densities and increases aloft when surface densities are low. Meteorologically, this is known as compensation, where lighter air overlays dense surface air, and below-normal densities at the surface gradually disappear aloft to be replaced with higher-than-normal densities. This emphasizes again the fact noted earlier, that at some altitude the atmospheric density is independent of surface density, and at this level horizontal density gradients are minimal.†

The distinction between atmospheric density and ballistic density should be emphasized here. The atmospheric density at any height is dependent on the pressure and temperature of the atmosphere at that elevation. The pressure at that height is the weight of the column of air above that level, irrespective of conditions below.

The ballistic density at any line "n" is the sum of the weighted densities from the surface up to line "n" in question, where the weighting

†This is the so-called isopycnic level of the atmosphere. In middle latitudes, it occurs near 8 km and was first discovered by A. Wagner in 1910 and more closely investigated by F. Linke in 1919.

function is described by

$$\sum_{j=1}^{j=n} a_j = 1$$

and listed in Table III.

TABLE III

DENSITY WEIGHTING FACTORS, IN % (from FM 6-16)

Height at Top of Zone		Zone #														
Line #	Meters	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15
1	200	100														
2	500	43	57													
3	1000	22	31	47												
4	1500	15	21	32	32											
5	2000	11	17	25	22	25										
6	3000	8	11	17	17	15	32									
7	4000	6	8	14	13	12	22	25								
8	5000	5	6	11	11	10	19	17	21							
9	6000	4	6	9	9	8	17	15	14	18						
10	8000	3	4	7	7	7	13	12	11	11	25					
11	10000	1	3	5	5	6	12	11	9	9	16	23				
12	12000	2	3	5	5	5	11	10	9	8	14	12	16			
13	14000	2	2	4	5	5	11	9	9	8	14	10	9	12		
14	16000	2	3	5	5	5	10	9	8	7	13	11	8	6	8	
15	18000	2	4	5	5	5	10	9	8	7	12	9	8	5	5	6

Hence, while the atmospheric density at the isopycnic level is nearly constant around the globe, the ballistic density at the same level will vary somewhat, although far less at that level than at lower altitudes, since the weighting factors are greatest at higher line numbers, as noted earlier.

It is undoubtedly true that each air mass has a distinctive density structure, since the source region determines the temperature and moisture content of the air mass. With cold frontal passages, the air mass changes and density changes at low levels (generally line 5 and below) are greater than normal for a short period of time. It would thus seem logical to categorize surface densities in terms of air mass, or relation to other meteorological factors (highs, lows, position in relation to a cold front, etc.). However, this technique would require trained meteorologists for each metro section, a luxury the current Army Table of Distribution and Allowances (TDA) cannot afford. Hence, the simplistic

climatological approach, that determines the values of ballistic density aloft from the surface density without regard to other meteorological conditions, was adopted.

The ultimate success of such tables - accurate assessment of ballistic density aloft - will be dependent on restriction of geographic extent of the regions to insure uniformity of conditions, adequate sample of data on which to base the tables, and diurnal or season breakdowns where required.

As an example, note the curves in Figures 8 and 9 that show the differences in density for night and day conditions for high and low line numbers.




FIG. 8 VIETNAM BALLISTIC DENSITIES




FIG.9 VARIATION IN BALLISTIC DENSITY (KOREA)

In both the tropical Vietnam and temperate Korean regions, there is a significant difference in upper air density for a given surface condition. Hence, separate tables are needed for nighttime and daytime, a fact realized in the first revision of the old tables in TR 1236-1. However, significant changes in day and night sounding are noted in all lines, not only in line 4 as promulgated in the revised tables ([4], pp 3-4).

Similarly, plotting the densities as a function of season reveals the necessity for constructing density tables on a seasonal rather than an annual basis. In Vietnam (Figure 10), the difference between summer and winter is significant at all levels above line 2, whereas a combination spring-fall table could be developed that would fit these transitional seasons.




FIG.10 SEASONAL VARIATION IN BALLISTIC DENSITY (VIETNAM)

Figures 11 and 12 show the seasonal density profiles for Korea. The contrast between summer and winter could not be depicted, since there were no corresponding surface densities (lowest surface density in winter 102%, highest in summer, 100%). There is, however, sufficient spread to make separate seasonal tables worthwhile.




FIG.11 SEASONAL VARIATION IN BALLISTIC DENSITY (KOREA)




FIG.12 SEASONAL VARIATION IN BALLISTIC DENSITY (KOREA)

The third factor, geographical extent of a region, is harder to define. For the Vietnam area, the density profiles from the most northerly station (near $16^{\circ}50'$ N) and the most southerly one (near $10^{\circ}20'$ N) were so nearly alike that it was obvious that all of South Vietnam could be included in a single region. Data were not available from North Vietnam to delineate the northern extent of the region. It is most probable that the tables can be extended to 20° N and possibly to 25° N. The westward extension to the Bay of Bengal is also probable but must await further checking with data from upper air stations in that area.

For the Korea Region, the ballistic densities in the southern portion (near Pusan) differ markedly from those in the north (around Pyongyang). The separation is most pronounced in winter (Figure 13) where variations of more than 1% are possible between the extreme limits of the Korean area.




FIG 13 VARIATION OF BALLISTIC DENSITY (KOREA-WINTER)

The one-sigma values shown as vertical lines between the northern and southern curves indicate that the differences are significant between lines 3 and 7. For that reason, three separate tables were constructed - one for the southern islands to 36°N; the second valid between 36°N and the DMZ; and the third to be used from the DMZ to 40°N.

Construction of the Tables

Once the differential criteria for the tables are established, the actual construction largely depends on the availability of the data. Where large numbers of Artillery Metro Sections are in operation as in South Vietnam, the problem is simplified. In fact, zone and ballistic densities are available on Forms DA 6-57 and DA 6-59 for processing by computer. Frequency distributions of upper air densities as a function of surface density are produced, and tables are constructed for surface density steps of 0.5%. Abrupt changes in slope of the profiles are avoided both from zone to zone for a given surface density and for a given zone as the surface density varies. Such smoothing rarely changes the tabular values by more than 0.2 - 0.3%, much less than the standard deviations of the values themselves.

Where sources other than Artillery Metro Sections are used, radiosonde data are available only at mandatory or significant levels. In those cases, the densities must be computed from the sounding (see Appendix). After this has been completed, the procedure follows that is stated above

for the Vietnam data. Recognizing that variations in station altitude cause the greatest difference in density profiles (see Figures 3 and 7), the original tables in TR 1236-1 specified density factors for sea level, 1000' above MSL, and 2000' above MSL. A similar procedure, incorporating metric rather than English units, was followed for the new Vietnam tables as sufficient data were available. Tables were constructed for stations whose altitudes lay between sea level and 200 meters^{††} Another set was valid for those stations in the range of elevation 200 to 450 meters, the next 450 - 650 meters, etc. This procedure has the advantage of allowing for discrimination between density profiles due to elevation while keeping the number of tables within reason. In Vietnam, five sets of tables covered all operational station elevations.

The question naturally arises: Can a density profile be constructed for an elevated station if a nearby sea-level station is available? A special series of Vietnam flights was made every three hours for a week at a pair of locations separated by approximately 20 kilometers, one 30 meters above sea level, the other 770 meters above sea level. These flights were ideal for determining if ballistic densities from the station near sea level could be used to construct a density profile for the elevated location.

A simplistic approach was attempted. The difference between the surface densities of the two stations was computed and the difference subtracted from the ballistic densities of the sea-level metro message. The resulting values were compared with the actual ballistic densities from the higher station. The results are shown in Table IV below.

TABLE IV
ERROR IN BALLISTIC DENSITY

Line No.	1	2	3	4	5	6	7	8	9	10
Density error in percent	.3	.4	.5	.5	.5	.6	.7	.7	.8	.9

It is readily seen that the error increases with height and becomes greater than 0.5% above line five. More complex schemes were tried. The most successful involved subtracting the sum of the difference in

^{††} It is believed that by presenting the density for each zone as a tabular value, rather than a departure from the mean (which involves algebraic addition before the ballistic density is obtained), arithmetic errors will be reduced.

surface densities and one tenth of the line number from each ballistic density at sea level to give the corresponding ballistic density at 770 meters. The results of the computations are shown in Table V.

TABLE V
ERROR IN BALLISTIC DENSITY

Line No.	1	2	3	4	5	6	7	8	9	10
Density error in percent	.3	.4	.5	.6	.4	.6	.6	.6	.6	.5

The errors for high line numbers have been reduced, but the results are still not encouraging.

Neither of the two techniques is as accurate as the results obtained using the new climatological density tables as shown in Table VI.

TABLE VI
DENSITY ERRORS USING CLIMATOLOGICAL TABLES

Line No.	1	2	3	4	5	6	7	8	9	10
Density error in percent	.2	.2	.3	.3	.4	.4	.4	.4	.4	.4

The mean error is seen to be less than 0.5% in all cases and does not continuously increase with height. It appears, therefore, that ballistic densities from a climatological table are superior to values obtained from extrapolating density profiles from a sea-level station.

It has been shown that climatological tables are feasible, seem to be reasonably accurate in one sample case investigated, and are easy to use. It remains to be seen whether they are accurate for a much larger sample and whether they are better or worse than the existing tables.

Since only the data from Artillery Metro Sections in Vietnam are available for direct comparison of tabular and actual ballistic densities, the analysis will be confined to those data. It should be noted that the absolute value of the error is the important factor in the application of the ballistic density to an actual firing. The resulting effects from a shell whose trajectory is 150 meters too short, for example, are

just as bad as those from a shell whose trajectory is 150 meters too long. The algebraic error is a measure of the amount of bias in the tables and indicates whether the tables could be improved by changing the individual values to eliminate the bias. Ideally the algebraic error should be zero, but, owing to the inherent inaccuracies in the determination of density, small biases of a few tenths of a percent appear. These, however, are insignificant.

Tables VII and VIII show sample distribution of errors for the new Vietnam tables. Shown are the errors for the summer daytime and the winter nighttime tables for stations below 200 meters elevation.

MONTHS- 6 TO 4
ELEVATION- 0 TO 200 METERS
AND HOURS TO 1900 HOURS

34 POINTS OUT OF RANGE

TABLE VII												
	1	2	3	4	5	6*	7	8	9	10	11	TOTAL
-15	0	0	3	1	0	0	0	0	0	1	0	5
-14	0	0	3	4	0	0	3	2	0	0	0	12
-13	0	3	8	2	1	5	2	1	2	0	0	24
-12	4	6	11	6	4	7	2	5	2	0	0	47
-11	5	7	15	9	6	3	6	5	5	3	4	64
-10	9	17	23	14	7	15	6	6	5	2	5	111
-9	6	20	17	25	13	28	14	5	24	10	25	187
-8	25	29	44	36	40	36	46	26	19	23	25	349
-7	33	48	53	54	31	52	45	44	48	27	45	480
-6	40	50	97	69	55	68	74	67	58	65	48	601
-5	52	92	109	123	75	116	103	93	81	84	56	1014
-4	72	86	129	125	118	137	136	116	115	124	94	1252
-3	113	146	141	174	110	140	142	125	130	152	107	1500
-2	164	163	178	191	190	151	179	145	152	145	99	1757
-1	214	174	190	187	183	170	169	139	148	145	94	1813
0	285	235	148	194	187	161	212	132	151	154	94	1813
1	291	232	157	169	181	165	104	144	159	136	76	1894
2	288	179	161	141	150	183	151	133	142	154	84	1766
3	191	174	130	117	130	133	123	119	113	99	85	1405
4	95	134	87	101	128	104	78	69	84	68	58	1026
5	45	62	78	67	94	84	77	60	68	40	50	725
6	25	45	56	54	71	68	55	61	40	32	33	546
7	14	33	45	33	60	48	56	52	33	18	26	318
8	4	13	27	29	37	41	28	34	22	24	25	284
9	5	10	17	19	32	19	18	24	23	26	11	204
10	5	8	11	16	19	11	21	16	22	20	23	174
11	2	3	10	14	18	13	7	17	10	13	12	119
12	2	6	3	3	6	4	5	10	8	20	16	83
13	0	3	2	4	5	5	4	11	5	4	13	60
14	0	1	3	4	4	2	1	8	1	5	8	40
15	0	0	1	0	1	2	1	2	2	3	6	18
1961	1961	1961	1961	1978	1977	1962	1966	1973	1610	1261	TOTALS	
0.2	0.3	0.4	0.3	0.3	0.4	0.3	0.4	0.3	0.4	0.4	AVERAGE ERROR(ARS)	
0.0	0.0	-0.1	-0.1	0.0	-0.0	-0.0	0.0	-0.0	0.0	0.0	AVERAGE ERROR(ALG)	

20081 DATA POINTS

* Values in this column represent density errors X 10

DISTRIBUTION OF ERRORS
NEW VIETNAM TABLES

TABLE VII

It can be seen that the mean absolute error is 0.5% or less for all lines in both seasons and the bias error is not significant. All seasons and altitude ranges are similar, although the number of data points decreases markedly with increasing elevation.

Table IX shows a corresponding error distribution using the density departure tables in FM 6-16 for region 6. Since the older tables are in-

MONTHS-11 TO 2
ELEVATION- 0 TO 200 METERS
2000 HOURS TO 760 HOURS

170 POINTS OUT OF RANGE

TABLE VIII												
	1	2	3	4	5	6*	7	8	9	10	11	TOTAL
-15	0	0	0	3	5	6	7	8	9	10	11	72
-14	0	1	1	5	10	8	4	5	1	0	1	43
-13	0	2	2	8	12	8	4	2	4	4	3	59
-12	2	4	9	10	14	14	6	9	5	5	2	74
-11	0	2	5	19	15	21	9	7	8	4	9	103
-10	2	6	15	29	27	25	24	19	13	12	17	189
-9	3	8	16	39	34	35	27	20	12	15	15	227
-8	10	14	36	43	47	47	20	37	30	24	20	344
-7	8	37	59	83	63	84	36	54	43	44	33	548
-6	29	64	67	71	67	68	72	74	64	36	41	693
-5	62	84	111	122	125	122	81	94	71	61	60	1010
-4	146	143	125	135	114	135	110	118	99	90	88	1320
-3	175	174	143	114	123	135	131	143	117	121	101	1507
-2	239	218	181	135	154	137	157	146	114	147	100	1753
-1	250	184	158	141	119	151	161	169	141	149	126	1771
0	365	213	156	144	146	121	119	157	144	151	100	1854
1	208	209	141	137	144	142	154	144	124	151	110	1831
2	206	174	156	141	143	120	147	144	114	135	74	1558
3	145	157	119	120	109	104	119	161	130	123	70	1330
4	92	134	127	115	106	114	122	105	114	96	64	1199
5	47	94	108	105	124	85	117	107	94	87	51	1040
6	41	72	84	100	93	67	86	84	76	67	44	850
7	17	57	75	83	69	74	63	73	104	60	40	757
8	17	47	61	53	69	56	64	77	63	47	38	597
9	6	32	44	46	47	64	75	46	40	43	30	544
10	6	20	45	45	40	48	46	52	46	54	41	464
11	2	15	30	29	41	41	43	33	44	27	26	330
12	2	6	16	26	32	30	24	24	21	26	23	214
13	0	5	13	12	14	32	26	24	16	12	11	171
14	0	3	7	10	16	20	14	21	26	14	11	140
15	1	2	5	7	13	17	16	12	11	11	7	94
2197	2196	2197	2194	1993	2191	2175	2155	1993	1995	1427	TOTALS	
0.2	0.3	0.4	0.5	0.5	0.5	0.5	0.5	0.4	0.4	0.5	AVERAGE ERROR(ARS)	
0.0	0.0	0.1	0.0	0.1	0.1	0.1	0.1	0.1	0.1	0.1	AVERAGE ERROR(ALG)	

22813 DATA POINTS

* Values in this column represent density errors X 10

DISTRIBUTION OF ERRORS
NEW VIETNAM TABLES

TABLE VIII

tended for use in all seasons, there is no differentiation for summer and winter, and the tables include all data for all months.

DISTRIBUTION OF ERRORS -- OLD BALLISTIC DENSITY DEPARTURE TABLES

POINTS- 1 TO 12
ELEVATION- 0 TO 200 METERS
2000 METERS TO 700 HOURS

4334 PRINTS OUT OF RANGE

	ZONES											TOTAL
	1	2	3	4	5	6	7	8	9	10	11	12
-20	0	0	1	2	3	14	25	27	12	1	0	87
-19	0	0	1	1	1	19	23	47	14	2	0	114
-18	1	2	2	5	10	33	49	82	34	6	0	244
-17	0	1	2	6	12	59	77	114	46	0	0	331
-16	0	0	3	6	19	74	124	144	80	7	0	457
-15	0	2	3	18	36	118	179	194	119	7	0	544
-14	2	4	7	27	72	157	211	255	153	12	0	619
-13	0	3	16	40	83	192	305	334	235	24	0	744
-12	3	11	34	90	171	274	368	352	289	34	0	1432
-11	2	14	48	104	173	259	344	451	342	75	1	1814
-10	8	24	69	165	172	303	474	452	497	104	0	2204
-9	2	34	113	176	215	366	516	525	414	140	0	2574
-8	15	64	141	247	293	431	515	520	545	187	0	2871
-7	18	122	215	315	359	474	527	434	514	270	0	3261
-6	14	154	293	409	394	524	457	379	509	324	0	3402
-5	49	234	373	431	452	464	373	386	442	404	0	3534
-4	97	341	444	490	591	474	404	312	337	444	0	3162
-3	134	394	470	544	499	384	371	283	321	410	0	1884
-2	454	500	504	480	399	389	297	244	290	404	1	4047
-1	344	567	444	443	448	327	294	219	247	501	0	4034
0	682	561	479	490	384	302	220	148	141	371	0	3849
1	400	544	494	427	343	251	201	105	152	347	0	3332
2	247	444	497	370	350	213	123	84	111	304	2	2401
3	117	444	423	319	270	189	72	74	63	263	1	2240
4	745	484	315	244	229	179	54	56	48	212	2	2412
5	509	461	286	252	243	155	39	34	38	142	0	2131
6	672	312	244	225	191	77	38	24	25	120	5	1947
7	349	230	217	164	119	74	29	14	24	109	2	1384
8	431	184	144	110	101	41	18	16	17	71	8	1140
9	145	137	119	74	63	34	11	14	6	34	15	484
10	142	44	100	57	41	34	4	3	10	40	21	541
11	50	63	46	45	40	14	5	4	3	34	27	444
12	45	40	50	44	37	12	5	4	4	14	20	334
13	24	34	27	9	10	9	2	0	2	14	17	294
14	15	25	19	25	13	1	0	1	0	9	123	231
15	12	24	22	15	4	4	2	0	1	4	144	241
16	11	15	13	9	7	1	1	1	0	5	218	241
17	4	1	4	4	1	1	0	0	0	4	237	254
18	5	1	6	2	1	0	0	0	0	3	213	204
19	0	1	3	0	0	1	0	0	0	2	339	444
20	3	1	1	1	1	0	0	0	0	2	374	381

4915 4914 4914 4901 4896 4884 4884 4892 4114 3750 4334 TOTALS
 * Values in this column represent density errors X10
 0.4 0.4 0.4 0.5 0.5 0.4 0.4 0.4 0.7 0.4 2.2 AVERAGE ERROR(ACC)
 0.3 0.1 0.0 0.1 0.2 0.4 0.7 0.4 0.7 0.2 2.2 AVERAGE ERROR(SIG)

TORRE DATA PRINTS

TABLE IX

It is obvious that the absolute errors are greater and that there is a significant bias in many zones. Note especially line 11, which has an average error of more than 2%! It can be seen that fewer than 2000 of the 4334 values in line 11 have errors less than 2.0%, all the others are off range as far as the table goes. It is all the more surprising that line 10 is excellent, with mean errors as low as lines 1 and 2, and no significant bias!

If one wishes to compare the tables on a yearly basis, the seasonal errors can be combined to give an annual figure. Table X shows the breakdown into four categories of error.

Cases with errors from 0 to 0.5% can be classified excellent, those from 0.6% to 1.0% can be called fair; those from 1.0% to 1.5%, poor; and greater than 1.5% would be disastrous.

TABLE X

ANNUAL DISTRIBUTION OF DENSITY ERRORS (SEA-LEVEL STATIONS)

<u>Percent Error</u>	<u>New Tables</u>		<u>Old Tables</u>	
	<u>Day</u>	<u>Night</u>	<u>Day</u>	<u>Night</u>
0 - 0.5	72.6%	77.1%	50.1%	53.5%
0.6 - 1.0	22.4%	19.2%	28.1%	28.5%
1.1 - 1.5	4.3%	3.3%	12.5%	10.5%
> 1.5	0.7%	0.4%	9.3%	7.5%

We see that the number of cases of very poor assessment of ballistic density when the new tables are used is less than one-tenth the number with the old tables; the probability of obtaining an excellent ballistic density has increased by more than 20% and an acceptable density profile is obtained in 95 of every 100 cases. There is no doubt that the tables offer an excellent back-up system for obtaining ballistic densities in Vietnam in the absence of a direct radiosonde observation.

The lack of a direct measurement may be due to outage of equipment, communication difficulties, or it may be that it is the period between scheduled balloon releases, usually every six hours in Vietnam.

Earlier studies have shown that the variation of ballistic density in a 1-to-8-hour period is not significant [5]. Should the tables be used when the latest metro message is 3, 4, 5, or 6 hours old?

To answer this question one must examine the diurnal variation of density. The special 3-hourly data from Vietnam (mentioned earlier) are ideal for this analysis. Even though the period is short, about one week, the changes in density throughout the day will give some clue as to those periods in which the temporal variation is greatest. Figures 14 and 15 show the density for the various ballistic zones throughout the day for the station near sea level and the elevated location. The shape of both the curves is the same although the amplitude of the density fluctuation is greater at the lower station. About 2100Z (0500LST), well before sunrise, the surface and ballistic densities decrease rapidly, fluctuating between 0300Z and 0900Z (1100-1700LST) to a minimum a little before 0900Z. The decrease over this 6-hour period is greater than 3.0% at sea level and more than 2.0% at elevated stations.

[5] Lowenthal, Marvin J., "Applications of Accuracy of Upper Air Data to Artillery," ECOM-3122, May 1969.




FIG.14 DIURNAL VARIATION OF BALLISTIC DENSITY
 VIETNAM STATION 142080 ELEVATION 30 METERS
 GMT = GREENWICH MERIDIAN TIME (Z)




FIG.15 DIURNAL VARIATION OF BALLISTIC DENSITY
 VIETNAM STATION 140080 ELEVATION 770 METERS
 GMT = GREENWICH MERIDIAN TIME (Z)

Thus, the ballistic density obtained from a sounding made at 0000Z and used at 0500Z (before a new measurement is customarily made) would be in error by more than 1%. Similarly, a sounding made at 0600Z and used at 1100Z would contain an error almost as great, whereas the ballistic densities from a 1200Z sounding used at 1700Z and from a 1800Z sounding used at 2300Z would be in error by less than 1%. While these values are for the tropics, similar curves, although with different amplitudes, obtain for the other climatic regions. This means that the time of day is the determining factor for the time variability of ballistic density for periods of less than 6 hours.

To decrease the error in ballistic density in the interval between soundings, the curves of Figures 14 and 15 may be used to correct the earlier measurement. Even though the daily curves of ballistic density variation may not coincide absolutely, the trend is the same. Thus a diminishing of the 0000Z ballistic density by 1% for use at 0300Z would give a much better estimate of the density than the use of the uncorrected 0000Z value. A table could be constructed giving such corrections for each measurement hour of the day and the time of usage.

This could be supplied to the Fire-Direction Center to decrease the range error due to improper ballistic density corrections. A sample of such a table for one location in Vietnam is shown in Table XI. While both location and altitude are different from Figures 14 and 15, the trend is the same.

Corrections for Ballistic Density -- Line 1
Station 144075 (Vietnam)

Sounding Time (LST)	00	01	02	03	04	05	06	07	08	09	10	11	12	13	14	15	16	17	18	19	20	21	22	23	
Using Time	00	+2	+4	+6	+8	+9	+1.0	+9	+7	+3	-3	-9	-1.7	-2.3	-2.9	-3.2	-3.3	-3.1	-2.7	-2.2	-1.7	-1.1	-	-	-
01	-2	+2	+4	+5	+7	+8	+7	+5	+1	-5	-1.2	-1.9	-2.6	-3.1	-3.4	-3.5	-3.3	-2.9	-2.4	-1.9	-1.3	-0.7	-0.1	-0.5	-1.1
02	-4	+2	+4	+5	+6	+5	+3	-1	-7	-1.3	-2.1	-2.7	-3.3	-3.6	-3.7	-3.5	-3.1	-2.6	-2.1	-1.6	-1.1	-0.6	-0.1	-0.5	-1.1
03	-6	-4	-2	+2	+3	+4	+3	+1	-3	-8	-1.5	-2.3	-2.9	-3.5	-3.8	-3.8	-3.7	-3.3	-2.8	-2.3	-1.7	-1.2	-0.6	-0.1	-0.5
04	-8	-5	-4	-2	+1	+2	+2	-1	-4	-1.0	-1.7	-2.4	-3.1	-3.6	-4.0	-4.0	-3.8	-3.5	-3.0	-2.5	-1.9	-1.4	-0.8	-0.3	-0.7
05	-9	-7	-5	-3	-1	+1	0	-2	-6	-1.2	-1.8	-2.6	-3.2	-3.8	-4.1	-4.2	-4.0	-3.6	-3.1	-2.6	-2.1	-1.5	-1.0	-0.4	-0.8
06	-1.0	-8	-6	-4	-2	-1	-1	-3	-7	-1.2	-1.9	-2.6	-3.3	-3.8	-4.2	-4.2	-4.1	-3.7	-3.2	-2.7	-2.1	-1.6	-1.0	-0.5	-0.9
07	-9	-7	-5	-3	-2	0	+1	+2	-2	-6	-1.2	-1.9	-2.6	-3.2	-3.8	-4.1	-4.0	-3.6	-3.1	-2.6	-2.1	-1.5	-1.0	-0.4	-0.8
08	-7	-5	-3	-1	+1	+2	+3	+2	-4	-1.0	-1.6	-2.4	-3.0	-3.6	-3.9	-4.0	-3.8	-3.4	-2.9	-2.4	-1.8	-1.3	-0.7	-0.2	-0.6
09	-3	-1	+1	+3	+5	+6	+7	+6	+4	-6	-2	-2.0	-2.7	-3.2	-3.5	-3.6	-3.4	-3.0	-2.5	-2.0	-1.5	-1.0	-0.5	-0.1	-0.5
10	+2	+5	+7	+8	+1.0	+1.2	+1.2	+1.0	+6	-7	-1.4	-2.1	-2.6	-2.9	-3.0	-2.8	-2.5	-2.0	-1.4	-0.9	-0.4	+0.1	+0.6	+1.1	+1.6
11	+9	+1.2	+1.3	+1.5	+1.7	+1.8	+1.9	+1.6	+1.3	+7	-7	-1.4	-1.9	-2.2	-2.3	-2.1	-1.8	-1.3	-0.7	-0.2	+0.3	+0.8	+1.3	+1.8	+2.3
12	+1.7	+1.9	+2.1	+2.3	+2.4	+2.6	+2.6	+2.4	+2.0	+1.4	+7	-7	-1.2	-1.5	-1.6	-1.4	-1.0	-0.5	0	+0.5	+1.0	+1.5	+2.0	+2.5	+3.0
13	+2.3	+2.6	+2.7	+2.9	+3.1	+3.2	+3.3	+3.3	+3.0	+2.7	+2.1	+1.4	+7	-5	-9	-9	-7	-4	+1	+7	+1.2	+1.7	+2.2	+2.7	+3.2
14	+2.9	+3.1	+3.3	+3.5	+3.6	+3.8	+3.8	+3.8	+3.6	+3.2	+2.6	+2.0	+1.2	+5	-3	-4	-2	+2	+7	+1.2	+1.7	+2.2	+2.7	+3.2	+3.7
15	+3.2	+3.4	+3.6	+3.8	+4.0	+4.1	+4.2	+4.1	+3.9	+3.5	+2.9	+2.3	+1.5	+9	+3	-1	+1	+5	+1.0	+1.5	+2.0	+2.5	+3.0	+3.5	+4.0
16	+3.3	+3.5	+3.7	+3.9	+4.0	+4.2	+4.2	+4.2	+4.0	+3.6	+3.0	+2.3	+1.6	+9	+4	+1	+2	+5	+1.1	+1.6	+2.1	+2.6	+3.1	+3.6	+4.1
17	+3.1	+3.3	+3.5	+3.7	+3.8	+4.0	+4.1	+4.0	+3.8	+3.4	+2.8	+2.1	+1.4	+7	+2	-1	-2	+4	+9	+1.4	+1.9	+2.4	+2.9	+3.4	+3.9
18	+2.7	+2.9	+3.1	+3.3	+3.5	+3.6	+3.7	+3.6	+3.4	+3.0	+2.5	+1.8	+1.0	+4	-2	-5	-5	+4	+9	+1.4	+1.9	+2.4	+2.9	+3.4	+3.9
19	+2.2	+2.4	+2.6	+2.8	+3.0	+3.1	+3.2	+3.1	+2.9	+2.5	+2.0	+1.3	+5	-1	-7	-1.0	-1.1	-0.9	-0.5	+0.6	+1.1	+1.6	+2.1	+2.6	+3.1
20	+1.6	+1.9	+2.1	+2.2	+2.4	+2.6	+2.6	+2.4	+2.0	+1.4	+7	0	-7	-1.2	-1.5	-1.6	-1.4	-1.1	-0.6	+0.5	+1.0	+1.5	+2.0	+2.5	+3.0
21	+1.1	+1.3	+1.5	+1.7	+1.9	+2.0	+2.1	+2.0	+1.8	+1.4	+9	+2	-6	-1.2	-1.8	-2.1	-2.1	-2.0	-1.6	-1.1	-0.6	+0.5	+1.0	+1.5	+2.0
22	+7	+9	+1.1	+1.2	+1.4	+1.6	+1.6	+1.5	+1.4	+1.0	+4	-3	-1.0	-1.7	-2.2	-2.5	-2.6	-2.4	-2.1	-1.6	-1.1	-0.6	+0.5	+1.0	+1.5
23	+3	+5	+7	+9	+1.0	+1.2	+1.3	+1.2	+1.0	+6	0	-7	-1.4	-2.1	-2.6	-2.9	-3.0	-2.8	-2.4	-1.9	-1.4	-0.9	-0.4	+0.1	+0.6

Apply appropriate correction to measured Ballistic Density to get corrected value for use at firing time.

TABLE XI

CONCLUDING REMARKS

In this report, the development of ballistic density departure tables has been traced over the last three decades from their earliest beginnings, where elevation of the battery was the only parameter considered, through the promulgation of climatic regions where geography and time of day were added, to the latest tables that are constructed for a limited geographical area far smaller than the previous regions in FM 6-16.

In addition, the possibilities for minimizing interdiurnal density variations have been pointed out, as well as impossibility of predicting ballistic densities at elevated locations with sufficient accuracy for Artillery purposes from measurements at sea level.

The new tables constructed from individual radiosondings have been shown to be much more accurate than the older ones derived mainly from mean values at selected locations.

Use of the tables can effect a considerable savings in expendables, as wind measurements can be made every two hours by tracking a balloon alone (without the more costly radiosonde), since the density can be corrected by the tables and associated techniques.

APPENDIX

COMPUTER DETERMINATION OF METRO MESSAGES FROM RADIOSONDE DATA

By

Hope S. Perlman
U. S. Army Electronics Command

Computer reduction of radiosonde flights to give thermodynamical quantities is common where a large number of flights must be handled or where checking of manual operations is required in data repositories. Less common is the computer determination of NATO or computer meteorological messages since ballistic meteorology is less extensively researched than pure synoptic meteorology.

Secondly, the recent change in the format of the computer met message has necessitated a change in the reported quantities. That, in turn, requires a program change in any computer routine that produces the artillery met messages.

This paper outlines the procedure for computer determination of ballistic and computer met messages, with especial detail given to the evaluation of the two new elements of the computer message - mean virtual temperature and pressure at the midpoint of the zone.

Procedures

Since data for research purposes is most often obtained from World Data Centers, the analysis begins with the data in the usual card (or tape) format of the mandatory pressure levels - the customary way in which data are archived. It must be remembered that heights on the cards are expressed above mean sea level (MSL). Since artillery met data are concerned with heights above ground, the elevation of the station must be subtracted from all given heights of the standard pressure levels. That is the reason for Steps 1 and 3(c) below. A complete computer program is available upon request.

Method of Solution

1. Read in surface height (S_z) of station.
2. Read in complete data for one flight. This will include n sets of pressure (P), height (Z), temperature in °C(T) and relative humidity in % (H).

3. Adjust data and number of levels (n) as follows:
 - a. If $H < 0$, set it equal to zero.
 - b. Eliminate any level where temperature is missing.
 - c. Subtract S_z from all values of Z .
4. Calculate virtual temperature, in °K, (T_v) at each level, as follows:

$$T_v = \frac{P(T+273)}{P - (.0037812)(H)(6.105)\exp\left(\frac{25.22T}{T+273} - 5.31 \ln \frac{T+273}{273}\right)} \quad (1)$$

5. Print out adjusted data (P, Z, T, H, T_v) for each level.
6. Set up ballistic zone top heights (ZLT) and standard temperatures (ZLTMS) and densities (ZLDNS) at these heights.
7. Print all surface information.
8. For each zone (ZL) compute ZZTM, zone temperature and ZZPR, pressure at the top of the zone, using intermediate data level values of P, Z and T_v , indexed from 0, the bottom of the zone, to d , the top of the zone.
 - a. Obtain a starting value of T_{vd} , virtual temperature at the top of the zone, by interpolating linearly, using values of $T_{vd+1}, T_{vd-1}, Z_{d+1}, Z_d$, and Z_{d-1} .
 - b. ZZTM is calculated by selecting a starting value (1/2 the sum of the maximum and minimum of the T_v values within the zone), and moving it toward the maximum or minimum until

$$1/2 \sum_{i=0}^{d-1} (Z_{i+1} - Z_i) (T_{vi} + T_{vi+1} - 2ZZTM) \sim 0^* \quad (2)$$

In practice, the process is continued until two successive values of ZZTM differ by less than .1°K.

- c. P_d is found by solving the transcendental equation

$$29.27 \text{ ZZTM} \ln \frac{(P_0)}{(P_d)} = Z_n - Z_0 \quad (3)$$

* The rationale for Equation (2) is found in Appendix I.

- d. A new value of T_{vd} is computed by interpolating linearly, using T_{vd+1} , T_{vd-1} , P_{d+1} , P_d , and P_{d-1} .
- e. Steps b, c, and d are repeated until two successive values of ZZTM, calculated in step d differ by $.2^\circ\text{K}$ or less.
- f. $ZZPR = P_d$ (4)

9. For each zone level, compute

- a. $ZZTMP_{ZL}$ zone temperature (% of standard)

$$ZZTMP_{ZL} = \frac{100(ZZTM_{ZL})}{ZLTMS_{ZL}} \quad (5)$$

- b. $ZZPR2_{ZL}$, pressure at the midpoint of the zone, is found by solving the following transcendental equation:

$$29.27(ZZTM_{ZL}) \ln \frac{(ZZPR_{ZL-1})}{(ZZPR2_{ZL})} = 1/2 (Z_d - Z_o) \quad (6)$$

- c. $ZZDN_{ZL}$, zone density

$$ZZDN_{ZL} = 348.384 ZZPR2_{ZL} / ZZTM_{ZL} \quad (7)$$


- d. $ZZDNP_{ZL}$, zone density in percent of standard

$$ZZDNP_{ZL} = \frac{100(ZZDN_{ZL})}{ZLDNS_{ZL}} \quad (8)$$

- 10. For each zone, calculate ballistic weighted temperature ($ZZBT_{ZL}$) and ballistic weighted density ($ZZBD_{ZL}$), using temperature weighting factors (TWF) and density weighting factors (DWG) for that zone.

APPENDIX I

CALCULATION OF ZONE TEMPERATURE



Given: T_{v0}, Z_0, T_{v1}, Z_1 ---- T_{vd}, Z_d , where the zero subscript denotes the bottom of the zone and the d subscript the top of the zone.

Find: ZZTM, the zone temperature, such that the areas to the right of ZZTM equal the areas to the left of ZZTM, or ZZTM is correct to $.1^\circ\text{K}$.

Analysis:

1. Between any successive reading of T_{vi}, Zi and $T_{vi+1}, Zi+1$ the figure is either a trapezoid or 2 triangles.

- a. If a trapezoid, Area = $1/2 h (a+b)$

$$= 1/2 (Z_{i+1} - Z_i) [(T_{vi+1} - ZZTM) + (T_{vi} - ZZTM)]$$

$$= 1/2 (Z_{i+1} - Z_i) (T_{vi} + T_{vi+1} - 2ZZTM)$$

and will be positive or negative depending on whether T_{vi} and T_{vi+1} are greater or less than ZZTM.

- b. If two triangles, the area equals area of triangle

$$(T_{vi}, Zi) (T_{vi+1}, Zi+1) (T_{vi+1}, Zi)$$

minus area of rectangle

$$(ZZTM, Zi) (ZZTM, Zi+1) (T_{vi+1}, Zi+1) (T_{vi+1}, Zi)$$

$$\text{or } 1/2 (Z_{i+1} - Z_i) (T_{Vi} - T_{Vi+1}) - (Z_{i+1} - Z_i) (ZZTM - T_{Vi+1})$$

∴ the area of the trapezoid equals the area of the two triangles, and is

$$1/2(Z_{i+1} - Z_i) (T_{Vi} + T_{Vi+1} - 2ZZTM)$$

2. ZZTM is calculated by selecting a starting value of ZZTM (half the sum of the maximum and minimum values of the T_v 's) and moving it ΔT toward T_v min until

$$1/2 \sum_{i=0}^{d-1} (Z_{i+1} - Z_i) (T_{Vi} + T_{Vi+1} - 2ZZTM) \sim 0.$$

ATMOSPHERIC SCIENCES RESEARCH PAPERS

1. Miers, B. T., and J. E. Morris, Mesospheric Winds Over Ascension Island in January, July 1970, ECOM-5312, AD 711851.
2. Webb, W. L., Electrical Structure of the D- and E-Region, July 1970, ECOM-5313, AD 714365.
3. Campbell, G. S., F. V. Hansen and R. A. Lise, Turbulence Data Derived from Measurements on the 32-Meter Tower Facility, White Sands Missile Range, New Mexico, July 1970, ECOM-5314, AD 711852.
4. Pries, T. H., Strong Surface Wind Gusts at Holloman AFB (March-May), July 1970, ECOM-5315, AD 711853.
5. D'Arcy, E. M., and B. F. Engebos, Wind Effects on Unguided Rockets Fired Near Maximum Range, July 1970, ECOM-5317, AD 711854.
6. Matonis, K., Evaluation of Tower Antenna Pedestal for Weather Radar Set AN/TPS-41, July 1970, ECOM-3317, AD 711520.
7. Monahan, H. H., and M. Armendariz, Gust Factor Variations with Height and Atmospheric Stability, August 1970, ECOM-5320, AD 711855.
8. Stenmark, E. B., and L. D. Drury, Micrometeorological Field Data from Davis, California; 1966-67 Runs Under Non-Advection Conditions, August 1970, ECOM-6051, AD 726390.
9. Stenmark, E. B., and L. D. Drury, Micrometeorological Field Data from Davis, California; 1966-67 Runs Under Advection Conditions, August 1970, ECOM-6052, AD 724612.
10. Stenmark, E. B., and L. D. Drury, Micrometeorological Field Data from Davis, California; 1967 Cooperative Field Experiment Runs, August 1970, ECOM-6053, AD 724613.
11. Rider, L. J., and M. Armendariz, Nocturnal Maximum Winds in the Planetary Boundary Layer at WSMR, August 1970, ECOM-5321, AD 712325.
12. Hansen, F. V., A Technique for Determining Vertical Gradients of Wind and Temperature for the Surface Boundary Layer, August 1970, ECOM-5324, AD 714366.
13. Hansen, F. V., An Examination of the Exponential Power Law in the Surface Boundary Layer, September 1970, ECOM-5326, AD 715349.
14. Miller, W. B., A. J. Blanco and L. E. Traylor, Impact Deflection Estimators from Single Wind Measurements, September 1970, ECOM-5328, AD 716993.
15. Duncan, L. D., and R. K. Walters, Editing Radiosonde Angular Data, September 1970, ECOM-5330, AD 715351.
16. Duncan, L. D., and W. J. Vechione, Vacuum Tube Launchers and Boosters, September 1970, ECOM-5331, AD 715350.
17. Stenmark, E. B., A Computer Method for Retrieving Information on Articles, Reports and Presentations, September 1970, ECOM-6050, AD 724611.
18. Hudlow, M., Weather Radar Investigation on the BOMEX, September 1970, ECOM-3329, AD 714191.
19. Combs, A., Analysis of Low-Level Winds Over Vietnam, September 1970, ECOM-3346, AD 876935.
20. Rinehart, G. S., Humidity Generating Apparatus and Microscope Chamber for Use with Flowing Gas Atmospheres, October 1970, ECOM-5332, AD 716994.
21. Miers, B. T., R. O. Olsen, and E. P. Avara, Short Time Period Atmospheric Density Variations and a Determination of Density Errors from Selected Rocket-sonde Sensors, October 1970, ECOM-5335.
22. Rinehart, G. S., Sulfates and Other Water Solubles Larger than 0.15μ Radius in a Continental Nonurban Atmosphere, October 1970, ECOM-5336, AD 716999.
23. Lindberg, J. D., The Uncertainty Principle: A Limitation on Meteor Trail Radar Wind Measurements, October 1970, ECOM-5337, AD 716996.
24. Randhawa, J. S., Technical Data Package for Rocket-Borne Ozone-Temperature Sensor, October 1970, ECOM-5338, AD 716997.

25. Devine, J. C., The Fort Huachuca Climate Calendar, October 1970, ECOM-6054.
26. Allen, J. T., Meteorological Support to US Army RDT&E Activities, Fiscal Year 1970 Annual Report, November 1970, ECOM-6055.
27. Shinn, J. H., An Introduction to the Hyperbolic Diffusion Equation, November 1970, ECOM-5341, AD 718616.
28. Avara, E. P., and M. Kays., Some Aspects of the Harmonic Analysis of Irregularly Spaced Data, November 1970, ECOM-5344, AD 720198.
29. Fabrici, J., Inv. of Isotopic Emitter for Nuclear Barometer, November 1970, ECOM-3349, AD 376461.
30. Levine, J. R., Summer Mesoscale Wind Study in the Republic of Vietnam, December 1970, ECOM-3375, AD 721585.
31. Petriw, A., Directional Ion Anemometer, December 1970, ECOM-3379, AD 720573.
32. Randhawa, J. S., B. H. Williams, and M. D. Kays, Meteorological Influence of a Solar Eclipse on the Stratosphere, December 1970, ECOM-5345, AD 720199.
33. Nordquist, Walter S., Jr., and N. L. Johnson, One-Dimensional Quasi-Time-Dependent Numerical Model of Cumulus Cloud Activity, December 1970, ECOM-5350, AD 722216.
34. Avara, E. P., The Analysis of Variance of Time Series Data Part I: One-Way Layout, January 1971, ECOM-5352, AD 721594.
35. Avara, E. P., The Analysis of Variance of Time Series Data Part II: Two-Way Layout, January 1971, ECOM-5353.
36. Avara, E. P., and M. Kays., The Effect of Interpolation of Data Upon the Harmonic Coefficients, January 1971, ECOM-5354, AD 721593.
37. Randhawa, J. S., Stratopause Diurnal Ozone Variation, January 1971, ECOM-5355, AD 721309.
38. Low, R. D. H., A Comprehensive Report on Nineteen Condensation Nuclei (Part II), January 1971, ECOM-5358.
39. Armendariz, M., L. J. Rider, G. Campbell, D. Favier and J. Serna, Turbulence Measurements from a T-Array of Sensors, February 1971, ECOM-5362, AD 726390.
40. Maynard, H., A Radix-2 Fourier Transform Program, February 1971, ECOM-5363, AD 726389.
41. Devine, J. C., Snowfalls at Fort Huachuca, Arizona, February 1971, ECOM-6056.
42. Devine, J. C., The Fort Huachuca, Arizona 15 Year Base Climate Calendar (1956-1970), February 1971, ECOM-6057.
43. Levine, J. R., Reduced Ceilings and Visibilities in Korea and Southeast Asia, March 1971, ECOM-3403, AD 722735.
44. Gerber, H., et al., Some Size Distribution Measurements of AgI Nuclei with an Aerosol Spectrometer, March 1971, ECOM-3414, AD 729331.
45. Engebos, B. F., and L. J. Rider, Vertical Wind Effects on the 2 75-inch Rocket, March 1971, ECOM-5365, AD 726321.
46. Rinehart, G. S., Evidence for Sulfate as a Major Condensation Nucleus Constituent in Nonurban Fog, March 1971, ECOM-5366.
47. Kennedy, B. W., E. P. Avara, and B. T. Miers, Data Reduction Program for Rocketsonde Temperatures, March 1971, ECOM-5367.
48. Hatch, W. H., A Study of Cloud Dynamics Utilizing Stereoscopic Photogrammetry, March 1971, ECOM-5368.
49. Williamson, L. E., Project Gun Probe Captive Impact Test Range, March 1971, ECOM-5369.
50. Henley, D. C., and G. B. Hoidale, Attenuation and Dispersion of Acoustic Energy by Atmospheric Dust, March 1971, ECOM-5370, AD 728103.
51. Cionco, R. M., Application of the Ideal Canopy Flow Concept to Natural and Artificial Roughness Elements, April 1971, ECOM-5372, AD 730638.
52. Randhawa, J. S., The Vertical Distribution of Ozone Near the Equator, April 1971, ECOM-5373.
53. Ethridge, G. A., A Method for Evaluating Model Parameters by Numerical Inversion, April 1971, ECOM-5374.

54. Collett, E., Stokes Parameters for Quantum Systems, April 1971, ECOM-3415, AD 729347.
55. Shinn, J. H., Steady-State Two-Dimensional Air Flow in Forests and the Disturbance of Surface Layer Flow by a Forest Wall, May 1971, ECOM-5383, AD 730681.
56. Miller, W. B., On Approximation of Mean and Variance-Covariance Matrices of Transformations of Joint Random Variables, May 1971, ECOM-5384, AD 730302.
57. Duncan, L. D., A Statistical Model for Estimation of Variability Variances from Noisy Data, May 1971, ECOM-5385.
58. Pries, T. H., and G. S. Campbell, Spectral Analyses of High-Frequency Atmospheric Temperature Fluctuations, May 1971, ECOM-5387.
59. Miller, W. B., A. J. Blanco, and L. E. Traylor, A Least-Squares Weighted-Layer Technique for Prediction of Upper Wind Effects on Unguided Rockets, June 1971, ECOM-5388, AD 729792.
60. Rubio, R., J. Smith and D. Maxwell, A Capacitance Electron Density Probe, June 1971, ECOM-5390.
61. Duncan, L. D., Redundant Measurements in Atmospheric Variability Experiments, June 1971, ECOM-5391.
62. Engebos, B. F., Comparisons of Coordinate Systems and Transformations for Trajectory Simulations, July 1971, ECOM-5397.
63. Hudlow, M. D., Weather Radar Investigations on an Artillery Test Conducted in the Panama Canal Zone, July 1971, ECOM-5411.
64. White, K. O., E. H. Holt, S. A. Schleusener, and R. F. Calfee, Erbium Laser Propagation in Simulated Atmospheres II. High Resolution Measurement Method, August 1971, ECOM-5398.
65. Waite, R., Field Comparison Between Sling Psychrometer and Meteorological Measuring Set AN/TMQ-22, August 1971, ECOM-5399.
66. Duncan, L. D., Time Series Editing By Generalized Differences, August 1971, ECOM-5400.
67. Reynolds, R. D., Ozone: A Synopsis of its Measurements and Use as an Atmospheric Tracer, August 1971, ECOM-5401.
68. Avara, E. P., and B. T. Miers, Noise Characteristics of Selected Wind and Temperature Data from 30-65 km, August 1971, ECOM-5402.
69. Avara, E. P., and B. T. Miers, Comparison of Linear Trends in Time Series Data Using Regression Analysis, August 1971, ECOM-5403.
70. Miller, W. B., Contributions of Mathematical Structure to the Error Behavior of Rawinsonde Measurements, August 1971, ECOM-5404.
71. Collett, E., Mueller Stokes Matrix Formulation of Fresnel's Equations, August 1971, ECOM-3480.
72. Armendariz, M., and L. J. Rider, Time and Space Correlation and Coherence in the Surface Boundary Layer, September 1971, ECOM-5407.
73. Avara, E. P., Some Effects of Randomization in Hypothesis Testing with Correlated Data, September 1971, ECOM-5408.
74. Randhawa, J. S., Ozone and Temperature Change in the Winter Stratosphere, November 1971, ECOM-5414.
75. Miller, W. B., On Approximation of Mean and Variance-Covariance Matrices of Transformations of Multivariate Random Variables, November 1971, ECOM-5413.
76. Horn, J. D., G. S. Campbell, A. L. Wallis (Capt., USAF), and R. G. McIntyre, Wind Tunnel Simulation and Prototype Studies of Barrier Flow Phenomena, December 1971, ECOM-5416.
77. Dickson, David H., and James R. Oden, Fog Dissipation Techniques for Emergency Use, January 1972, ECOM-5420.
78. Ballard, H. N., N. J. Beyers, B. T. Miers, M. Izquierdo, and J. Whitacre, Atmospheric Tidal Measurements at 50 km from a Constant-Altitude Balloon, December 1971, ECOM-5417.
79. Miller, Walter B., On Calculation of Dynamic Error Parameters for the Rawinsonde and Related Systems, January 1972, ECOM-5422.

80. Richter, Thomas J., Rawin Radar Targets, February 1972, ECOM-5424.
81. Pena, Ricardo, L. J. Rider, and Manuel Armendariz, Turbulence Characteristics at Heights of 1.5, 4.0, and 16.0 Meters at White Sands Missile Range, New Mexico, January 1972, ECOM-5421.
82. Blanco, Abel J., and L. E. Traylor, Statistical Prediction of Impact Displacement due to the Wind Effect on an Unguided Artillery Rocket During Powered Flight, March 1972, ECOM-5427.
83. Williams, B. H., R. O. Olsen, and M. D. Kays, Stratospheric-Ionospheric Interaction During the Movement of a Planetary Wave in January 1967, March 1972, ECOM-5428.
84. Schleusener, Stuart A., and Kenneth O. White, Applications of Dual Parameter Analyzers in Solid-State Laser Tests, April 1972, ECOM-5432.
85. Pries, Thomas H., Jack Smith, and Marvin Hamiter, Some Observations of Meteorological Effects on Optical Wave Propagation, April 1972, ECOM-5434.
86. Dickson, D. H., Fogwash I An Experiment Using Helicopter Downwash, April 1972, ECOM-5431.
87. Mason, J. B., and J. D. Lindberg, Laser Beam Behavior on a Long High Path, April 1972, ECOM-5430.
88. Smith, Jack, Thomas H. Pries, Kenneth J. Skipka, and Marvin Hamiter, Optical Filter Function for a Folded Laser Path, April 1972, ECOM-5433.
89. Lee, Robert P., Artillery Sound Ranging Computer Simulations, May 1972, ECOM-5441.
90. Lowenthal, Marvin J., The Accuracy of Ballistic Density Departure Tables 1934-1972, April 1972, ECOM-5436.
91. Cantor, Israel, Survey of Studies of Atmospheric Transmission from a 4π Light Source to a 2π Receiver, April 1972, ECOM-5435.
92. Barr, William C., Accuracy Requirements for the Measurement of Meteorological Parameters Which Affect Artillery Fire, April 1972, ECOM-5437.