AD-750 878

STUDIES OF ATMOSPHERIC PROCESSES

Frederick P. Boynton, et al

Wayne State University

Prepared for:

Air Force Cambridge Research Laboratories Advanced Research Projects Agency

15 June 1972

DISTRIBUTED BY:

National Technical Information Service U. S. DEPARTMENT OF COMMERCE 5285 Port Royal Road, Springfield Va. 22151 AFCRL-72-0453

STUDIES OF ATMOSPHERIC PROCESSES

by

Frederick P. Boynton, Edward R. Fisher, Ralph H. Kummler

Pieter K. Rol

Research Institute for Engineering Sciences College of Engineering Wayne State University Detroit, Michigan 48202

Contract No. F19628-72-C-0007

Semi-Annual Technical Report No. 2

15 June 1972

The views and conclusions contained in this document are those of the authors and should not be interpreted as necessarily representing the official policies, either expressed or implied, of the Advanced Research Projects Agency of the U.S. Government.

> Contract Monitor: Alva T. Stair Optical Physics Laboratory

Approved for public release; distribution unlimited.

Sponsored by

Reproduced by NATIONAL TECHNICAL INFORMATION SERVICE US Department of Commerce age out of the VA 22151

Advanced Research Projects Agency ARPA Order No. 1856

Monitored by

AIR FORCE CAMBRIDGE RESEARCH LABORATORIES AIR FORCE SYSTEMS COMMAND UNITED STATES AIR FORCE BEDFORD, MASSACHUSETTS 01730 ARPA Order No. - 1856 Contract No. F19628-72-C-0007 Program Code No. - 2E40 Principal Investigator and Phone No. Dr. Adolf R. Hochstim/313 577-3867 Name of Contractor- Wayne State University Effective Date of Contract - 12 July 1971

Project Scientist and Phone No. Dr. Alva T. Stair/617 861-4911

Contract Expiration Date - 16 October 1972

1 :





	ROL DATA - R & D
(Security eleastification of itig, body of obstract and indexing ORIGINATING ACTIVITY (Corporate author)	annotation must be entered when the overall report is cleasified) 22. REPORT SECURITY CLASSIFICATION
Wayne State University, College of)	Engineering UNCLASSIFIED
Research Institute for Engineering	Sciences Z. GROUP
Detroit, Michigan 48202	
REPORT TITLE	
STUDIES OF ATMOSPHERIC PROCESSES	
DESCRIPTIVE NOTES (Type of report and inclusive dctee)	
SCIENCILLC. Incerim.	
Frederick P. Boynton	Ralph H Kummler
	Kerph H. Kunnter
Edward R. Fisher	Pieter K Rol
REPORT DATE	31 5
June 15, 1972	94. ORIGINATOR'S REPORT NUMBER(S)
ARPA Order No. 1856	
F19628-72-C-0007 6. Project no.	SemiAnnual Technical Report No. 2
8692	
^{c.} DoD Element 62301D	yp. OTHER REPORT NO(S) (Any other numbers that may be assigned this report)
, DoD Subelement n/a	AFCRL-72-0453
0. DISTRIBUTION STATEMENT	
supported by the Derense Advanced kesearch Projects Agency	Laboratories (OP) L.G. Hanscom Field Bedford, Massachusetts 01730
The research described in thi the non-equilibrium flow and coup	s report is aimed at characterizing ed chemistry associated with the
interaction of rocket exhaust eff Infrared chemiluminescence emitted C ₂ H ₄ has been quantitatively measu ratio of the rate coefficient	from the reaction between 0 and red under lean conditions. The for the reactions
interaction of rocket exhaust eff Infrared chemiluminescence emitted C ₂ H ₄ has been quantitatively measu ratio of the rate coefficient CHO + O- CO	from the reaction between 0 and red under lean conditions. The for the reactions + OH* and CHO + 02- CO + HO2
interaction of rocket exhaust eff Infrared chemiluminescence emitted C ₂ H ₄ has been quantitatively measu ratio of the rate coefficient CHO + O- CO is found to be 1.7 while the rate	the the upper atmosphere. I from the reaction between 0 and ared under lean conditions. The is for the reactions + OH* and CHO + 0_2 - CO + HO ₂ coefficient for the reaction
interaction of rocket exhaust eff] Infrared chemiluminescence emitted C_2H_4 has been quantitatively measured ratio of the rate coefficient CHO + O- CO is found to be 1.7 while the rate CH_D + O- CF	from the reaction between 0 and ared under lean conditions. The is for the reactions + OH* and CHO + O_2 - CO + HO ₂ coefficient for the reaction HO + OH
interaction of rocket exhaust eff] Infrared chemiluminescence emitted C_2H_4 has been quantitatively measured ratio of the rate coefficient CHO + O- CO is found to be 1.7 while the rate $CH_2O + O- CH$ is found to be less than 3 x 10 ⁻¹⁴ than 20%.	the the upper atmosphere. I from the reaction between 0 and ared under lean conditions. The is for the reactions + OH* and CHO + O_2 - CO + HO ₂ coefficient for the reaction HO + OH Cc/sec with a quantum yield of less
interaction of rocket exhaust eff Infrared chemiluminescence emitted C_2H_4 has been quantitatively measured ratio of the rate coefficient CHO + O- CO is found to be 1.7 while the rate $CH_2O + O- CH$ is found to be less than 3 x 10 ⁻¹⁴ than 20%. A compilation of all available included herein for use in highly The molecular beam measurement between O atoms and plume species	the first in the upper atmosphere. I from the reaction between 0 and ared under lean conditions. The is for the reactions + OH* and CHO + O_2 - CO + HO ₂ coefficient for the reaction HO + OH CC/sec with a quantum yield of less the cross section information is rarefied plume calculations. Its on the interaction potential is described.
interaction of rocket exhaust eff Infrared chemiluminescence emitted C_2H_4 has been quantitatively measured ratio of the rate coefficient CHO + O- CO is found to be 1.7 while the rate $CH_2O + O- CH$ is found to be less than 3 x 10 ⁻¹⁴ than 20%. A compilation of all available included herein for use in highly The molecular beam measurement between O atoms and plume species	the the upper atmosphere. I from the reaction between 0 and ared under lean conditions. The is for the reactions + OH* and CHO + O_2 - CO + HO ₂ coefficient for the reaction HO + OH fcc/sec with a quantum yield of less the cross section information is rarefied plume calculations. Its on the interaction potential is described.
interaction of rocket exhaust eff Infrared chemiluminescence emitted C_2H_4 has been quantitatively measure ratio of the rate coefficient CHO + O- CO is found to be 1.7 while the rate $CH_2O + O- CH$ is found to be less than 3 x 10 ⁻¹⁴ than 20%. A compilation of all available included herein for use in highly The molecular beam measurement between O atoms and plume species	UNCLASSIFIED

UNCLASSIFIED

Security Classification			1		1		
KEY WORDS		NK A	ROLE	С.В. М.Т.	ROLE WT		
	NOLS	-	1				
	•				Į į		
chemiluninescence				1			
combustion flows						ł	
rareified flows	}					}	
vibrational cross sections	l	1					
plume chemistry		1	1		1	ļ	
		1					
			}				
		ļ				ļ	
		1					
				1	1	1	
				1	1	1	
	1		1				
	ļ				1	1	
				1	1		
					{		
					{		
			1				
					ļ		
			1			1	
		l l					
		}				1	
						1	
		}					
		1		l l		1	
		1				1	
					1		
		ł			-	1	
	ļ	ļ	{		1		
		Į	L L	ļ			
	r			l	ł		
1							
د د بالا الاستان المراجعة المراجعة اليونية الورية الورية المراجع المراجع المراجع المراجع المراجع الم		UNCI.A	SSTRTI	20			
		Sec	urity Clas	sificatio	'n		
		Sec	urity Clas	●111C#[10			

ł,

- **3**4-

See. In

AFCRL-72-0453

STUDIES OF ATMOSPHERIC PROCESSES

1 saliditto

by

Frederick P. Boynton, Edward R. Fisher, Ralph H. Kummler

Pieter K. Rol

Research Institute for Engineering Sciences College of Engineering Wayne State University Detroit, Michigan 48202

Contract No. F19628-72-C-0007

Semi-Annual Technical Report No. 2

15 June 1972

The views and conclusions contained in this document are those of the authors and should not be interpreted as necessarily representing the official policies, either expressed or implied, of the Advanced Research Projects Agency of the U.S. Government.

> Contract Monitor: Alva T. Stair Optical Physics Laboratory

Approved for public release; distribution unlimited.

Sponsored by

Advanced Research Projects Agency ARPA Order No. 1856

Monitored by

AIR FORCE CAMBRIDGE RESEARCH LABORATORIES AIR FORCE SYSTEMS COMMAND UNITED STATES AIR FORCE BEDFORD, MASSACHUSETTS 01730 ARPA Order No. - 1856
Program Code No. - 2E40
Name of Contractor- Wayne State University
Effective Date of Contract - 12 July 1971

Contract No. F19628-72-C-0007

Principal Investigator and Phone No. Dr. Adolf R. Hochstim/313 577-3867

Project Scientist and Phone No. Dr. Alva T. Stair/617 861-4911

Contract Expiration Date - 16 October 1972



IV.

TABLE OF CONTENTS

Ρ	age	

takes to

Abs	tract	1
Pre	face	2
I.	Fluid Mechanical Studies	3
II.	Vibrational Cross sections	5
III.	Flow Discharge System Measurements	11
IV.	Molecular Beam Experiments	23
v.	Rotational Excitation	27

- Vi

ABSTRACT

The research described in this report is aimed at characterizing the non-equilibrium flow and coupled chemistry associated with the interaction of rocket exhaust effluents in the upper atmosphere. Infrared chemiluminescence emitted from the reaction between 0 and C_2H_4 has been quantitatively measured under lean conditions. The ratio of the rate coefficients for the reactions

$$CHO + O \rightarrow CO + OH^*$$

and
$$CHO + O_2 \rightarrow CO + HO_2$$

is found to be 1.7 while the rate coefficient for the reaction

$$CH_2O + O \rightarrow CHO + OH$$

is found to be less than 3×10^{-14} cc/sec with a quantum yield of less than 20%.

A compilation of all available cross section information is included herein for use in highly rarefied plume calculations.

The molecular beam measurements on the interaction potential between 0 atoms and plume species is described.

PREFACE

The research described in this report falls into four main categories, identified in the contractual work statement as follows:

- a) Calculate the continuum flow structure and rate chemistry of high altitude plumes.
- b) Calculate the vibrational excitation cross-sections for plume relevant species using available interaction potential data.
- c) Measure the quantum yields in the infrared region for the chemi-excitation involving O or OH and plume species.
- d) Measure the inter-molecular potential between atomic oxygen and various plume species as an input to the calculation of vibrational excitation cross sections.

Item (a) is discussed in Section I, item (b) in Section II, item (c) in Section III, and item (d) in Section VI.

I. Fluid Mechanical Studies

In this portion of the Wayne State effort, emphasis has been focused on two specific problem areas: development of a 1-D unsteady continuum fluid description as a bridge between the 2-D steady continuum MULTITUBE code and a full Monte Carlo rarefied flow description (under development at TRW), and examination of the methodology for including chemistry and excitation processes in a 1-D Monte Carlo rarefied flow description.

The development of the 1-D unsteady continuum flow code has been described in the previous semi-annual technical report dated 15 December, 1971.⁽¹⁾ To date the coding on this program has been completed and the calculation successfully checked out for the case of expansion into a vacuum. Current effort is focused on studying shock formation, i.e. studying the development of shock structure through a thick shoc description. This phase of the fluid study will be carried on by \mathbb{F} . Boynton elsewhere.⁽²⁾

The Monte Carlo code development has several purposes in addition to accounting for chemistry and excitation processes. These include an examination of multicomponent effects in a simple geometry and as a means of gaining familiarity with Carlo analysis for advisory the structure of Monte а purposes in chemistry input. The early development of the Monte Carlo code is described in the semi-annual report dated 15 December, 1971, so will not be further described here. Work in this last 6 months has established the validity of the 1-D unsteady Monte Carlo code for shock wave formation. A routine for the inclusion of vibrational excitation processes has been developed and included into the M-C analysis. In addition, this excitation subroutine has been forwarded to TRW for inclusion with the full Monte Carlo development underway at TRW. The current problem in the M-C code development is that when the excitation probability is small, the code is slow in running on the IBM 360/67 and CDC 7600 machines.

Literature Cited

- F.P. Boynton, E.R. Fisher, R.H. Kummler and P.K. Rol, Semi-Annual Technical Report No.115 December (1971).
- 2) Physical Dynamics, Inc., San Francisco, California.

II. Vibrational Cross Sections

In the high altitude rarefied flow regime, highly non-thermal collisions characterize the interaction between ambient atmospheric species and exhaust constituents. In order to characterize the excitation processes associated with these high relative energy collisions, cross section information for various plume and atmospheric species must be generated. The close coupling code of Marriott⁽¹⁾ has been applied to some of these cross section calculations over the past years. In this reporting period, the results of Marriott's past cross section calculations, in addition to some new results of plume interest, have been correlated into a form convenient for evaluation of the excitation cross section at any collision energy above threshold.

No experimental information is available on the excitation cross sections for any of the collision processes calculated. Therefore, in order to attempt to assess the validity of these cross section calculations, rate coefficients (or relaxation times) have been generated by integrating the cross sections over a Boltzmann distribution of relative energies. Where thermalized data are available these calculated relaxation times have been compared and conclusions drawn.

Table IIa

State Notation for the Polyatomic Species

Notation	<u>~1</u>	<u>v</u> 2	<u>~3</u>
0	0	0	0
1	0	1	0
2	0	2	0
3	1	0	0
4	0	0	1

A full description of the theoretical close coupling code is currently being included in a technical report describing this work⁽²⁾ and is also available elsewhere so only the results will be summarized here. The species considered in this tabulation include CO, CO_2 , H_2O , H_2 and He. The state notation for the polyatomic species is given in Table II a. It should be noted that the v_2 bending mode of CO_2 is doubly degenerate and the cross sections do not account for this, i.e. the total excitation cross section from ground to the $v_2 = 1$ state is twice the cross section listed (see Marriott for a further discussion of this).

The functional form assumed for correlating the energy dependence of the cross section is

$$\sigma(E) = C3_{ln}(E/E_{T})^{C2} (E_{T}/E)^{C4}$$
 II-1

where E = relative collision energy (eV)

 E_{m} = threshold energy

and $\sigma(E) = excitation cross section (cm²).$

Table IIb lists all the coefficients to equation (II-1) for the reactions studied.

These cross section functions can be integrated over a Boltzmann distribution of relative energies, yielding rate coefficients with the inclusion of degeneracy factors. The complete tabulation of rate coefficients as determined from the data in Table IIb will be available in a forthcoming report complete with the comparison to available rate coefficient data.⁽²⁾ The conclusions drawn from these comparisons are, that at high temperatures the agreement between the calculated rate coefficients and measurements is good in all cases for which data exist except $CO_2(v_2) + H_2$. At lower temperatures, i.e. below $1000^{\circ}K$, the calculated rate coefficients are generally smaller than the data suggest reflecting the neglect of vibration-rotation (VR) and vibration-vibration (VV) processes in the calculation of the cross sections. These latter processes, particularly VR processes, are expected to be important in effecting thermalized energy transfer processes involving H₂ and H₂O. For the case of $CO_2(v_2) + H_2$, the calculated

مستنبط سارات المستنوب

REACTION	ET	<u>C2</u>	<u>C3</u>	<u>C4</u>
$\begin{array}{llllllllllllllllllllllllllllllllllll$	2.89E-01 2.660E-01 5.310E-01 2.660E-01 5.310E-01 2.660E-01 1.980E-01 3.950E-01 4.410E-01 4.650E-01 1.980E-01 2.430E-01 2.430E-01 2.430E-01 3.950E-01 4.560E-02 7.000E-02 2.440E-02 1.980E-01 3.950E-01 4.410E-01 4.560E-02 7.000E-02 2.440E-02 1.980E-01 2.680E-01 3.950E-01 3.950E-01 3.950E-01 3.950E-01 3.950E-01 3.950E-01 3.950E-01 3.950E-01 4.650E-01 1.980E-01 2.680E-01 1.980E-01 2.680E-01 1.980E-01 2.680E-01 1.980E-01 2.680E-01 1.980E-01 2.680E-01 1.980E-01 2.680E-01 1.980E-01 2.680E-01 1.980E-01 2.680E-01 1.980E-01 2.680E-01 1.980E-01 2.680E-01 1.980E-01 2.680E-01 1.980E-01 2.680E-01 1.980E-01 2.680E-01 1.980E-01 2.680E-01 2.680E-01 1.980E-01 2.680E-01 1.980E-01 2.910E-01	1.093E01 3.658E00 3.659E00 3.779E00 1.306E01 1.175E01 6.830E300 3.937E00 4.387E00 2.628E00 3.488E00 2.398E00 4.588E00 5.631E00 3.992E00 4.180E00 9.229E00 5.160E00 3.355E00 3.260E00 3.134E00 3.682E01 3.977E00 3.728E00 6.307E00 5.153E00 1.157E00 1.098E01 8.663E00 1.481E01 5.116E00 4.923E00 2.119E01 1.451E01 8.291E00 7.518E00 4.145E00 5.473E00	1.110E-23 3.263E-20 8.189E-23 1.299E-19 1.449E-18 2.189E-20 4.082E-22 3.755E-16 1.630E-16 2.205E-18 1.183E-18 1.939E-16 5.263E-17 2.619E-17 5.143E-18 3.589E-18 3.589E-18 3.589E-18 3.589E-18 3.589E-18 3.589E-18 3.589E-18 3.589E-18 3.589E-18 3.589E-18 3.589E-18 3.589E-18 2.754E-17 5.112E-20 2.065E-20 1.676E-18 2.756E-17 5.328E-18 1.993E-16 1.067E-16 1.419E-16 1.708E-19 9.932E-21 1.103E-19 1.689E-23 3.386E-19 1.793E-16 9.282E-18 2.392E-17 1.289E-16 4.420E-16 1.142E-16 9.133E-17 2.600E-17	$\begin{array}{c} 0.0\\ -2.147E00\\ -4.857E00\\ -8.742E-01\\ 3.183E-00\\ 1.764E-00\\ -2.357E-00\\ 1.380E-00\\ 1.734E-00\\ -3.184E-01\\ 4.102E-01\\ 7.175E-01\\ 1.374E-00\\ 1.999E-00\\ 3.068E-01\\ 5.583E-01\\ 2.230E-00\\ 1.287E-00\\ 1.287E-00\\ 1.287E-00\\ 1.287E-00\\ 1.400E-00\\ -1.709E-00\\ 1.144E-00\\ 9.232E-01\\ 2.017E-01\\ 1.569E-00\\ 6.524E-02\\ 4.090E-00\\ 3.680E-00\\ 3.393E-01\\ 8.457E-00\\ 5.192E-00\\ 1.948E-00\\ 2.180E-00\\ 9.704E-01\\ 1.244E-00\\ 9.704E-01\\ 1.244E-00\\ 9.704E-01\\ 1.244E-00\\ 3.772E-01\\ 1.721E-00\\ \end{array}$
••- ···				

Coefficients to Cross Section vs. Energy Function

Table con't.

.

REACTION	ET	<u>C2</u>	<u>C3</u>	<u>C4</u>
CO2 + H2 Q(12)	8.280E-02	2.267E00	8.757E-16	9.479E-01
CO2 + H2 Q(13)	8.090E-02	1.395200	2.3595-17	4.9065-02
CO2 + H2 Q(14)	2.0906-01	5.4/0EUU 5.107E01	0 254E 20	4.3/96-01
CO2 + H2 Q(32)	2.000E-03	7 10/EUL	0.234 <u>6</u> -30	1.603E-00
CO2 + H2 Q(24)	1.200E-01	1 301E00	9.000E-10	1.0836-00
CO2 + H2 Q(34)	1.200E-01 9.290E-02	4.391600	J.0316-10	J 172E 00
CO2 + H20 Q(01)	1 660F-01	5 237E00	2 4098 17	1 2175.00
CO2 + H20 Q(02)	$1.640E_01$	4 624E00	2.400E-17 1 307E-19	2 6795-01
CO2 + H20 Q(03)	2 910F-01	4 696F00	1.970E-10	2.079E-01
CO2 + H20 O(12)	8 280E-01	6 947E00	5.245E-17	1 795F_00
CO2 + H20 O(12)	8 090E-02	6 495E00	8 598E-18	1.795E-00
CO2 + H20 O(14)	2.090E-01	1 164E01	7 792E - 16	5 282E-00
CO2 + H20 O(32)	2.000E-01	1 030E01	6 608E-21	1 440E-00
CO2 + H20 O(24)	1.260E-01	1 488E01	1.483E-15	6 078E-00
CO2 + H20 O(34)	1.280E-01	4.601E00	9.341E-19	-4.290E-01
CO2 + CO2 O(01)	8.280E-02	1.955E01	4,993E-19	4.957E-00
CO2 + CO2 O(02)	1.660E-01	1.144E01	3.059E-20	1.324E-00
CO2 + CO2 O(03)	1.640E-01	1.630E01	4.223E 20	3.603E-00
CO2 + CO2 O(04)	2.910E-01	1.926E01	1.209E-17	7.240E-00
$CO2 + CO2 \tilde{Q}(12)$	8.280E-02	2.804E0C	4.183E-18	0.0
CO2 + CO2 O(13)	8,090E-02	1.064E01	2.919E-19	2.411E-00
CO2 + CO2 Q(14)	2.090E-01	2.067E01	1.880E-15	8.969E-00
$CO2 + CO2 \overline{Q}(32)$	2.000E-03	1.048E00	2.062E-17	0.0
$CO2 + CO2 \overline{Q}(24)$	1.260E-01	1.968E01	7,164E-16	7.333E-00
CO2 + CO2 Q(34)	1.280E-01	8.858E00	6.543E-20	2.766E-01

results are larger than measurements suggest even without the inclusion of VR processes. Apparently, this can only be interpreted by assuming that the interaction potential used in the calculation is not quite correct. This point is being currently investigated.

In summary, the cross section information listed in Table II b is the only source of high energy excitation data available for use in the rarified plume - atmosphere chemistry description. These data are expected to be reliable for this application even though the thermalized comparisons show some disagreement with data. This is due to the expectation that at high relative collision energies, VR and VV processes will be relatively unimportant compared to direct VT excitation.

Current work on the close coupling code is aimed at including structure in the collision partner in order to characterize VV processes for species of plume interest.

Literature Cited

- (1) R. Marriott, <u>Proc. Phys. Soc.</u>, <u>83</u>, 159 (1964).
 R. Marriott, <u>Proc. Phys. Soc.</u>, <u>84</u>, 877 (1964).
 R. Marriott, <u>Proc. Phys. Soc.</u>, <u>86</u>, 1041 (1965).
 R. Marriott, <u>Proc. Phys. Soc.</u>, <u>88</u>, 83 (1966).
 R. Marriott, <u>Proc. Phys. Soc.</u>, <u>88</u>, 617 (1966).
 R. Marriott, RIES Report 71-21 (1971).
 R. Marriott, and G. Kindt, RIES Report 71-22 (1971).
 R. Marriott, RIES Report 70-17 (1970).
 - Sections and Rate Coefficients for VT Processes Involving CO_2 , H_2O , H_2 and CO," report in preparation.

III. Flow Discharge System Measurements

Prior to quantitative model predictions on the radiative properties of exhaust products in the upper atomosphere, several important chemical reaction rate coefficients must be determined. In particular, our studies have focussed on simple exothermic oxidation reactions involving 0 and 0H attack on various hydrocarbon fuel species and accompanying vibrational relaxation processes.

The interest simple exothermic reactions is directed toward the release of the available energy of the reaction into radiative modes of the product molecules. Measurements in our laboratory have been made from 2500 A to 25μ in an attempt to determine the reaction mechanism and available radiative modes.

In order to determine which of the various hydrocarbon reations was worth studying in detail, i.e. determining quantum yields at 2.7μ , a brief review of hydrocarbon combustion chemistry was performed. A simplified flow diagram for the decomposition of straight chain hydrocarbons is shown in Fig. III a. We note that 0 atom attack on C_nH_{2n+2} produces OH in a nearly thermoneutral reaction in addition to the C_nH_{2n+1} radical. This radical is rapidly attacked by 0 atoms to produce C_nH_{2n} and OH with sufficient exothermicity to produce vibrationally excited OH^{*}. H_2O^* can subsequently be produced by the reaction OH + OH. This cycle continues, as outlined in the figure until C_2H_4 is formed. This then decomposes as shown in Fig. VIb upon further O atom attack.





12.



The attack of C atoms on C_2H_4 has been selected as a prototype reaction in the degradation of straight chain hydrocarbons by O atoms. From Figure IIIb, the primary products are CHO + CH₃. The rapid reaction of O + CHO is sufficiently exothermic to produce OH^{*}. At low concentrations of C_2H_4 , OH^{*} is expected to be the dominant 2.7 μ radiator. This is discussed at length in a recent thesis performed at Wayne State University under the direction of R. Kummler.⁽¹⁾ The work performed during that thesis study will be published as a technical report under the plume program.

To summarize the results of that study, experimental infrared measurements at 2.7 $_{\mu}$ were taken for the reaction system illustrated in Figure IIIb at low concentrations of $C_{2}H_{4}$. Under these conditions, as shown in the computer simulation of Figure IIIc, the OH^{*} is considerably higher in concentration than $H_{2}O^{*}$ (both of which radiate at 2.7 $_{\mu}$). The kinetic mechanism in the computer model is shown in Table IIIa. Under these lean conditions, strong 2.7 $_{\mu}$ emission is observed from the primary radiation step

$$CHO + O \rightarrow CO + OH^*$$
.

The primary CHO loss step is found to be the reaction

$$CHO + O_2 \rightarrow CO + HO_2$$
.

The ratio of rate coefficients for these two competitive CHO reactions is experimentally found to be 1.7. $^{(1)}$ In addition, the rate coefficient for the slightly exothermic reaction

$$CH_{2}O + O \rightarrow CHO + OH$$

is found to be less than 3×10^{-14} cc/sec with a quantum yield of than than 20%. ⁽¹⁾ A full report of this study is being prepared.



ENDUS2, 300 . LOLX, PLOT NO. 105

15

KINFTICS OF THE REACTIONS. TIME = 3.65499E-03 SEC.

. .

NO.			REACT	TION		CONSTANT	RATE	
1.C2H4	+0			=CH3	+CHO	8.4E-13	6.CE	14
3. HCHO	+0			=CHO	+OH1	3.0E-15	1. 3E	12
5.080	+0			=C02	+H	3.0E-14	7.4E	11
7.CH0	+8			=C0	+82	4.0E-12	1.1E	13
9.080	+02			=C0	+H02	1.0E-12	6.9E	14
11.OH	+0 H			=820	+0	2.5E-12	1. 1E	11
13.CH3	+02	+	M	=CH302	+ M	8.0E-32	1.6E	12
15. H20*	+ 8			= H2O	+ M	1.CE-12	1.7E	12
17. HCHO	+0			=CHC	+OH	6.0E- 14	2.6E	13
19. OH	+H CHO			= H 20	+CHC	1.4E-11	1.5E	13
21.08	+H2			=H20	+ H	6.6E-15	5.2E	07
23. HO2	+0			=OH	+02	1.0E-11	6.9E	14
25.08	+ HO 2			= H20"	+02	1.0E-12	1.7E	11
27. CHO*	+0			=C0	+O用*	1.CE-11	8.0E	10
29.0	+H02			=OH •	+02	3.0E-14	2.1E	12
31.CH302	+ HV			=CH302	+ HV	1.0E-20	7.3E-	11

NO.			REAC	r Ion		CONSTANT	RATE
2.083	+0			=HCHO	+H	3.0E-11	6.3E 14
4.CHO	+0			=C 0	+OH	1.0E-13	2.5E 12
6.08	+0			=02	+H	5.0E-11	9.1E 14
8.08	+ HV			=OH	+ HV	3.3E 00	4.4E 10
10.08	+ 0			=OH	+ 0	5.0E-11	5.7E 13
12.08	+0H			= H20*	+0	2.5E-13	1.1E 10
14.H	+02	+	M	=H02	+ 8	3.0E-32	2.4E 13
16. #20*	+ HV			=820	+ HV	3.5E 01	1.7E 09
18.CHO	+0			=C0	+OH *	2.0E-12	4.9E 13
20. OH	+HCHO			= H2O*	+CHO	1.4E-12	1.5E 12
22.08	+802			=#20	+02	1.0E-11	1.7E 12
24.8	+H02			=OH •	+OH	3.0E-15	2.3E 10
26.0284	+0			=CH3	+C HO*	1.0E-16	7.2E 10
28.H	+802			=OH	+0 H	1.0E-11	7.7E 13
30.00	+0H			=C 02	+H	1.5E-13	1.7E 11
32.820	+C02			=120	+C 02	1.0E-20	4.2E 00

Figure IIIa. Kinetic reactions and rate coefficients at 300° K for the computer (Simulation of Figure IIIb).

Experimental

All studies were performed in the discharge flow system shown schematically in Figure IIId.Molecular oxygen (American Cryogenics, purity 99.6%) was discharged in an Evenson or Broida Cavity at 2450 MHZ using a Raytheon PGM 10x2 diathermy system. The discharge was pulsed at 238 Hz for simultaneous infrared, visible and ultraviolet measurements. The discharged oxygen was passed into the flow tube reactor through a teflon tube with a radially-multiperforated exit stream which permits excellent mixing with a second gas and prevents discharge region radiation from reaching the observation The stainless steel reactor is lined with a teflon ports. sleeve which largely prevents wall recombination of the atomic species. The inside diameter of the reactor is one inch and typical gas velocities are 30 to 40 m/sec through a Heraeus Engelhard E - 225 mechanical pump. Six ports permit radial viewing through KBr windows using an RCA C31025C uncooled photomultiplier on a Jarrell Ash scanning quarter meter monochrometer with a Keithley 602 Electrometer recorded on a Honeywell Electronik 194 recorder. Signal to noise ratios were such that no damping or time integration of signals was necessary in the u.v. and visible. Measurements in the infrared were made using a Santa Barbara uncooled PbS detector (to 3_{μ}), Ge:Au detector (to 8_{μ}) and Ge:Cu detector (to 25_{μ}). All infrared measurements were made with a PAR HR8 Phase Sensitive Lock in Amplifier.

The hydrocarbons employed as reactant gases were methane, ethane, propane, ethylene, propylene, trans-butene, acetylene and formaldehyde. Linde research grade methane and ethane were employed. Union Carbide chemically pure propane was used. The propylene and trans-butene from Cryogenic Sales had purities of 95% and 99% respectively. Several runs were made with "house" methane with no perceptibly different results.



Figure IIId. Schematic of discharge flow tube reactor and detection system. Legend:

- M = Monochromator Jarrell Ash, quarter meter, scanning.
- P = Photomultiplier RCA C31025C (uncooled).
- E = Electrometer, D.C. Amplifier Keithley 602.
- Pi = Picoampere source Keithley 602.
 - R = Recorder Honeywell Electronik 194.
- Hi = High voltage C.C. power supply Power
 Designs Pacific model 2K-10.

Summary of Wayne State Measurements

A summary review of all the flow discharge measurements to date is presented in Table IIIb. It should be mentioned that reaction (1) in the table is the generation reaction for producing OH, and therefore the reactions initiated with OH, i.e. reactions (2), (3), and (4), have IR signals in excess of reaction (1). Some of these preliminary survey measurements are being further quantified in a way similar to the work on $O + C_2H_4$. Table IIIb. Wayne State University: Summary of Data to June 1972

د مند المعاد مورد . ت. دند

and the second second

And a state of the state of the

CHEMILUMINESCENT FLOW TUBE MEASUREMENTS

React	ion System	Visible	<u>2,7µ</u>	<u>5-10µ</u>	Comments
1. н (ол ог	+ $NO_2 \rightarrow NO$ + OH H + $H_2 \rightarrow H_2O$ + H H + OH $\rightarrow H_2O$ + O)	Meinel bands	large signal	large signal	maximum found in 2.7µ signal, no maximum in 5-10µ
2. 01	H + CO	nothing	nothing	?	
3. OI 4. OI	н + сн ₂ 0 н + с ₂ н ₂	? nothing	large signal small signal	medium signal medium signal	signals greater than (1) above
5. 0	+ c ₂ H ₄	Meinel bands	large signal	medium signal	maximum found in 2.7µ signal
6. 0	+ C ₄ H ₈	Meinel bands	large signal	?	
7. 0	+ c ₃ H ₆	Meinel bands	?	?	
8. 0	+ CH ₂ 0	Meinel bands (3500-4500	large signal	?	maximum found in 2.7µ signal
9. 0	+ c ₂ H ₂	Meinel bands	small signal	?	
LO. O	+ c ₂ ^H 6	Meinel bands	-		
L1. O	+ $C_n H_{2n+2}$	Meinel bands			
L2. 0	+ NH ₃	HNO [*]			

Literature Cited

(1) M. Malki, "Infrared Chemiluminescence from the Reaction of Oxygen Atoms on Ethylene", MS Thesis, Wayne State University, June (1972).

IV. Molecular Beam Experiments

The calculation or measurement of the excitation of exhaust species by atmospheric O atoms is recognized as an important problem. To date, direct experimental measurement of these excitation cross sections has not proved feasible. In this study, the excitation cross sections are calculated using the close coupled code and the intermolecular potential acting between O atoms and plume species. This potential is being experimentally measured using an Amdur-type molecular beam apparatus.

This experiment proceeds by measuring the attenuation of a fast (50 - 5000eV) O atom beam in passing through a scattering gas. The elastic scattering cross section is then determined from

$$\varrho_n(I/I_n) = -nS_{\ell} \qquad (VII.1)$$

where I = attenuated beam flux
I_o = incident beam flux
n = scattering gas density
i = path length in scattering gas
and S = incomplete scattering cross section.

Thus, from a plot of $p_n(I/I_0)$ vs.n for a constant p, the cross section can be determined. For a mixture of 2 atomic species in the incident beam, i.e., O(3P) and O(¹D), the (I/I₀) vs. n curve can be resolved into two linear parts, yielding two cross sections. S is an incomplete cross section since the finite width of the detector and the atomic beam cause part of the scattered particles to reach the detector.

The molecular beam apparatus used in this experiment is shown in Figure IV a. Oxygen atomic ions are produced in the duo-plasmatron source, focused and accelerated through a magnet to separate out both neutrals and molecular ions. The oxygen atomic ion beam is then collimated and energy selected prior to being neutralized by charge exchange with water vapor. The neutralized oxygen atomic



beam is then passed through the scattering chamber and the attenuation measured.

The beam apparatus, donated to Wayne State by NASA, originally had problems with the beam handling system. The original unipotential lens was replaced by a series of lenses and two sets of deflection plates to guide the beam through the small orifices in the scattering chamber. The system now is operational and preliminary measurements should be available within the month.

Measured incomplete scattering cross sections depend critically on the beam geometry. This fact, not initially recognized in the MIT experiments, is displayed in Figure IV b. Measurements made too close to the cusp in the scattering cross section can yield large errors to the sensitivity in the cross section to variations in both the beam width (d) and the detector width (o). The potential determined for the case in the figure is $V(R) = K/r^{2.5}$.

The repulsive part of the interaction potential determined from these beam experiments is in the range from leV to about 30eV. No difficulties are currently anticipated in these measurements.



Figure IVb: Calculated dependence of relative beam width (d) to detector width (p) on scattering cross section.

V. Rotational Excitation

The rotational excitation of simple molecules plays a critical role in determining the spectral spread of near infrared emission from vibrationally excited molecules. Therefore, a review of rotational excitation models was undertaken to provide background in estimating the spectral signature expected from vibrational emission of CO, CO_2 and H_2O in non-equilibrium flow environments. This review has been compiled into a report which is in the final stages of typing.⁽¹⁾

In addition, based on this review of rotational excitation processes, short numerical codes have been written to calculate the effective rotational relaxation times of simple molecules, the population distribution of rotational states under nonequilibrium conditions and collision-limited steady state emission from rotational states. The codes are based on Miller's statistical model of rotational excitation⁽²⁾ and employ the rigid rotor approximation.

Literature Cited

- F.P. Boynton, "Recent Developments in Theories of Rotational Excitation of Simple Molecules", report in preparation.
- (2) W.H. Miller, J. Chem. Phys. <u>52</u>, 543 (1970).

Contributors to the research described in this report:

Section I

F.P. Boynton III, Research Assistant, Wayne State University.

Section II

- E.R. Fisher, Associate Professor, Research Institute for Engineering Sciences, Wayne State University.
- R. Marriott, Professor, Research Institute for Engineering Sciences, Wayne State University.

Section III

- E.R. Fisher, Associate Professor, Research Institute for Engineering Sciences, Wayne State University.
- R.H. Kummler, Associate Professor, Dept. of Chemical Engineering and Material Sciences, Wayne State University.

Section IV

- P.K. Rol, Professor, Research Institute for Engineering Sciences, Wayne State University.
- P. Foreman, Research Associate.

Section V

F.P. Boynton III, Research Assistant, Wayne State University.

PREVIOUS CONTRACT AND REPORTS

The research continued under this contract was previously supported under Contract No. FO 4701-69-C-0230, supported by the Advanced Research Projects Agency of the Department of Defense, and monitored by Space and Missile Systems Organization, Norton Air Force Base, California.

The following reports have been sponsored in full or in part by this contract and the previous contract:

"Numerical Calculations of Viscous High-Altitude Exhaust Plume Flow Fields," F.P. Boynton, RIES-70-18.

"Transport Coefficients and Rates of Collisional Energy Transfer of Electrons and Ions in the Upper Atmosphere," A.R. Hochstim, RIES-70-8.

"Quasi-Steady Motions of Weakly Ionized Inhomogeneous Plasma Clouds in a Strong Magnetic Field," J.Alex Thomson, RIES-70-2.

"Energy Deposition in Artificial Ionospheric Heating Experiments,"

J. A. Thomson, J. Geoph. Res. <u>75</u>, 6446 (1970), RIES-70-4. "A Calculation of the Vibrational Excitation of Carbon Monoxide by Collision with Atomic (xygen in the ¹D State", R. Marriott, RIES-70-17.

"Relaxation Processes in Expanding Flows of Combustion Products," F.P. Boynton and E.R. Fisher, RIES-71-32.

"A Hydrodynamic Analog of Ionospheric Plasma," J. Alex Thomson, invited paper presented at the Fall Meeting of the A.G.U., San Francisco, December 1971.

"Alkali Quenching in High Temperature Environments," E.R. Fisher and G.K. Smith, Chem. Phys. Letters, <u>13</u>, 448 (1972).

"Studies of Atmospheric Processes," F.P. Boynton, E.R. Fisher, R.H. Kummler and P.K. Rol, Semi-Annual Technical Report No. 1, Contract No. F19628-72-C-0007, December 1971.