

AD-751 304

INVESTIGATION OF LASER PROPAGATION
PHENOMENA

Stuart A. Collins, Jr., et al

Ohio State University

Prepared for:

Advanced Research Projects Agency
Rome Air Development Center


August 1972

DISTRIBUTED BY:

NTIS

National Technical Information Service
U. S. DEPARTMENT OF COMMERCE
5285 Port Royal Road, Springfield Va. 22151

RADE-TR-72-247
Technical Report
August 1972



INVESTIGATION OF LASER PROPAGATION PHENOMENA
(3432-1)


AD751304

The Ohio State University
ElectroScience Laboratory

Department of Electrical Engineering
Columbus, Ohio 43212

Sponsored by
Defense Advanced Research Projects Agency
ARPA Order No. 1279

Approved for public release;
distribution unlimited.



The views and conclusions contained in this document are those of the authors and should not be interpreted as necessarily representing the official policies, either expressed or implied, of the Defense Advanced Research Projects Agency or the U. S. Government.

NATIONAL TECHNICAL
INFORMATION SERVICE

Rome Air Development Center
Air Force Systems Command
Griffiss Air Force Base, New York

22 R

UNCLASSIFIED

Security Classification

DOCUMENT CONTROL DATA - R&D		
<i>(Security classification of title, body of abstract and indexing annotation must be entered when the overall report is classified)</i>		
1. ORIGINATING ACTIVITY (Corporate author) The Ohio State University ElectroScience Laboratory Department of Electrical Engineering Columbus, Ohio		2a. REPORT SECURITY CLASSIFICATION Unclassified
3. REPORT TITLE INVESTIGATION OF LASER PROPAGATION PHENOMENA		2b. GROUP
4. DESCRIPTIVE NOTES (Type of report and inclusive dates) Quarterly Technical Report		
5. AUTHOR(S) (Last name, first name, initial) Collins, S.A., Jr. Reinhardt, G.W.		
6. REPORT DATE August 1972	7a. TOTAL NO. OF PAGES 15	7b. NO. OF REFS 7
8a. CONTRACT OR GRANT NO. F30602-72-C-0305	9a. ORIGINATOR'S REPORT NUMBER(S) ElectroScience Laboratory 3432-1	
b. ARPA Order No. 1279		
c. Program Code No. 9E20		
d.	9b. OTHER REPORT NO(S) (Any other numbers that may be assigned this report) RADC-TR-72-247	
10. AVAILABILITY/LIMITATION NOTICES Approved for public release; distribution unlimited.		
11. SUPPLEMENTARY NOTES Monitored by: Raymond P. Urtz, Jr. (OCSE) RADC, GAFB, NY 13440		12. SPONSORING MILITARY ACTIVITY Advanced Research Projects Agency Washington, D.C. 20301
13. ABSTRACT This is the first quarterly technical report under a contract entitled "Investigation of Laser Propagation Phenomena". Work during the quarter was concentrated on temporal spectra of phase difference fluctuations of atmospherically degraded spherical waves and on averaging times of optical random data. The temporal spectra derivation considered the case of a horizontal wind transverse to the optical path but with the phase measurement points oriented so that their separation vector makes an arbitrary angle with the horizontal. The spectra also included outer scale effects. The averaging time discussion outlined a technique for determining the averaging time of random data while the data is being recorded but in a time much shorter than the total data duration.		

DD FORM 1473
1 JAN 64

UNCLASSIFIED

Security Classification

ia

UNCLASSIFIED

Security Classification

14. KEY WORDS	LINK A		LINK B		LINK C	
	ROLE	WT	ROLE	WT	ROLE	WT
Averaging time						
Temporal spectra						
Turbulence						
Outer scale						
Optical phase difference						

INSTRUCTIONS

1. **ORIGINATING ACTIVITY:** Enter the name and address of the contractor, subcontractor, grantee, Department of Defense activity or other organization (*corporate author*) issuing the report.
- 2a. **REPORT SECURITY CLASSIFICATION:** Enter the overall security classification of the report. Indicate whether "Restricted Data" is included. Marking is to be in accordance with appropriate security regulations.
- 2b. **GROUP:** Automatic downgrading is specified in DoD Directive 5200.10 and Armed Forces Industrial Manual. Enter the group number. Also, when applicable, show that optional markings have been used for Group 3 and Group 4 as authorized.
3. **REPORT TITLE:** Enter the complete report title in all capital letters. Titles in all cases should be unclassified. If a meaningful title cannot be selected without classification, show title classification in all capitals in parenthesis immediately following the title.
4. **DESCRIPTIVE NOTES:** If appropriate, enter the type of report, e.g., interim, progress, summary, annual, or final. Give the inclusive dates when a specific reporting period is covered.
5. **AUTHOR(S):** Enter the name(s) of author(s) as shown on or in the report. Enter last name, first name, middle initial. If military, show rank and branch of service. The name of the principal author is an absolute minimum requirement.
6. **REPORT DATE:** Enter the date of the report as day, month, year, or month, year. If more than one date appears on the report, use date of publication.
- 7a. **TOTAL NUMBER OF PAGES:** The total page count should follow normal pagination procedures, i.e., enter the number of pages containing information.
- 7b. **NUMBER OF REFERENCES:** Enter the total number of references cited in the report.
- 8a. **CONTRACT OR GRANT NUMBER:** If appropriate, enter the applicable number of the contract or grant under which the report was written.
- 8b, 8c, & 8d. **PROJECT NUMBER:** Enter the appropriate military department identification, such as project number, subproject number, system numbers, task number, etc.
- 9a. **ORIGINATOR'S REPORT NUMBER(S):** Enter the official report number by which the document will be identified and controlled by the originating activity. This number must be unique to this report.
- 9b. **OTHER REPORT NUMBER(S):** If the report has been assigned any other report numbers (*either by the originator or by the sponsor*), also enter this number(s).

10. **AVAILABILITY/LIMITATION NOTICES:** Enter any limitations on further dissemination of the report, other than those imposed by security classification, using standard statements such as:

- (1) "Qualified requesters may obtain copies of this report from DDC."
- (2) "Foreign announcement and dissemination of this report by DDC is not authorized."
- (3) "U. S. Government agencies may obtain copies of this report directly from DDC. Other qualified DDC users shall request through _____."
- (4) "U. S. military agencies may obtain copies of this report directly from DDC. Other qualified users shall request through _____."
- (5) "All distribution of this report is controlled. Qualified DDC users shall request through _____."

If the report has been furnished to the Office of Technical Services, Department of Commerce, for sale to the public, indicate this fact and enter the price, if known.

11. **SUPPLEMENTARY NOTES:** Use for additional explanatory notes.

12. **SPONSORING MILITARY ACTIVITY:** Enter the name of the departmental project office or laboratory sponsoring (paying for) the research and development. Include address.

13. **ABSTRACT:** Enter an abstract giving a brief and factual summary of the document indicative of the report, even though it may also appear elsewhere in the body of the technical report. If additional space is required, a continuation sheet shall be attached.

It is highly desirable that the abstract of classified reports be unclassified. Each paragraph of the abstract shall end with an indication of the military security classification of the information in the paragraph, represented as (TS), (S), (C), or (U).

There is no limitation on the length of the abstract. However, the suggested length is from 150 to 225 words.

14. **KEY WORDS:** Key words are technically meaningful terms or short phrases that characterize a report and may be used as index entries for cataloging the report. Key words must be selected so that no security classification is required. Identifiers, such as equipment model designation, trade name, military project code name, geographic location, may be used as key words but will be followed by an indication of technical context. The assignment of links, rules, and weights is optional.

UNCLASSIFIED

Security Classification

16

INVESTIGATION OF LASER PROPAGATION PHENOMENA

**S. A. Collins, Jr.
G. W. Reinhardt**

**Contractor: Ohio State University
Contract Number: F30602-72-C-0305
Effective date of Contract: 1 April 1972
Contract Expiration Date: 31 March 1973
Amount of Contract: \$75,000.00
Program Code Number: 9E20**

**Principal Investigator: Dr. Stuart A. Collins, Jr.
Phone: 614 422-5045**

**Project Engineer: Edward K. Damon
Phone: 614 422-5953**

**Contract Engineer: Raymond P. Urtz, Jr.
Phone: 315 330-3443**

**Approved for public release;
distribution unlimited.**

**This research was supported by the
Defense Advanced Research Projects
Agency of the Department of Defense
and was monitored by Raymond P. Urtz,
Jr. RADC (OCSE), GAFB, NY 13440 under
Contract F30602-72-C-0305.**

ABSTRACT

This is the first quarterly technical report under a contract entitled "Investigation of Laser Propagation Phenomena". Work during the quarter was concentrated on temporal spectra of phase difference fluctuations of atmospherically degraded spherical waves and on averaging times of optical random data. The temporal spectra derivation considered the case of a horizontal wind transverse to the optical path but with the phase measurement points oriented so that their separation vector makes an arbitrary angle with the horizontal. The spectra also included outer scale effects. The averaging time discussion outlined a technique for determining the averaging time of random data while the data is being recorded but in a time much shorter than the total data duration.

CONTENTS

	Page
INTRODUCTION	1
TEMPORAL PHASE DIFFERENCE SPECTRA	1
AVERAGING TIMES	7
SUMMARY	14
BIBLIOGRAPHY	15

INTRODUCTION

This is the first quarterly technical report under Contract F30602-72-C-0305 titled "Investigation of Laser Propagation Phenomena". This effort is aimed at providing theoretical support to the RADC Laser Propagation Program in the areas of atmospheric propagation phenomena and microturbulence statistics. The efforts are in direct support of the experimental program being conducted at RADC. Two immediate aims are the theoretical description of the temporal spectrum of phase difference fluctuations and theoretical development of methods of determining appropriate averaging times for atmospherically degraded light beams. Work done during the past quarter has concentrated on these two topics. This work is only summarized in this report; the material will be covered in detail in two technical reports to be disseminated at a later date.

The work on temporal spectra is of interest because it provides information on time scales of phase and arrival angle useful to systems designers. More generally it utilizes a simple method of examining experimental data, i.e., the determination of temporal spectra, to give a check on our knowledge of the light-beam turbulence interaction.

In the present report the phase difference temporal spectrum of turbulence degraded light received at two pinholes is considered. Specifically the case when there is a horizontal wind blowing across the beam and the vector between the two pinholes is not necessarily parallel with the ground but makes an arbitrary angle with the horizontal is examined. Outer scale effects are also included.

The averaging time work is important because it establishes the duration of data necessary in order for time averages measured experimentally to agree with the theoretically developed predictions using ensemble averages and a variety of assumptions. In the present report the emphasis is on determining, using a computer-centered technique, the appropriate averaging time while the data is being recorded.

The work performed on these two topics during the past quarter will be summarized respectively in the next two sections.

TEMPORAL PHASE DIFFERENCE SPECTRA

An analytical expression for the spherical wave phase difference power spectrum has been determined for a horizontal transverse wind and arbitrary angle between the line connecting the phase points and the horizontal using the Von Karmen index spectrum (i.e., including the

finite outer scale effects). This extends previous work (3163-3) where the line between the phase points and wind direction were both horizontal.

In this section the derivation of the analytical expression for the frequency spectrum is summarized. The details of the derivation will be presented in a future report. The final equation of this derivation is evaluated for several limiting cases: small and large normalized separations; and small, intermediate and large normalized frequencies.

The physical picture for the calculation is one of a spherically divergent beam propagating a horizontal distance L to a pair of points in a plane perpendicular to the propagation direction. A wind of constant velocity is blowing across the beam. The difference of the electromagnetic phase of the two points is then measured.

Figure 1 illustrates the coordinate system, where $\phi_1(\rho_1, t)$ and $\phi_2(\rho_2, t)$ are the phases at the two points on a spherical wave front separated by distance $\bar{\rho} = (\bar{\rho}_2 - \bar{\rho}_1)$ and a distance L from the spherical wave source. The time-lagged difference in phase between the electromagnetic fields at the phase points is Fourier transformed to yield the desired spectrum.

The spherical wave phase difference spectrum is calculated for any angle between the line connecting the phase points and transverse wind velocity component (v) using the Rytov approximation.

Two cases are of particular interest. The first case is for very small normalized frequencies where the spectrum approaches a constant amplitude except when the wind direction and separation are exactly parallel. The second case occurs when the normalized frequency is large. There the spectrum is orientation independent in contrast to a pronounced orientation dependence for small normalized frequencies.

The phase difference spectrum calculations and experimental measurements are especially useful in the low frequency region where outer scale effects become dominant. In this case phase variance measurements with phase point separation on the order of several outer scale lengths (L_0) are replaced with more easily obtained temporal correlation measurements having a time lag comparable to that required for the wind to travel the distance of several outer scale lengths. This technique allows an independent determination of outer scale and directly shows the spectrum of the time variation of the relative phase. This time variation is valuable input for any calculation involving the time variation of quantities dependent on the wave or phase structure functions.

The summary of the derivation starts with an expression for the time lagged phase difference spectrum,




Fig. 1. Phase difference spectrum experiment.

$$(1) \quad W_{\delta S}(\omega) = \mathcal{F}\{ \langle (\phi_2(\bar{\rho}_2, \tau+t) - \phi_1(\bar{\rho}_1, \tau+t))(\phi_2(\bar{\rho}_2, t) - \phi_1(\bar{\rho}_1, t)) \rangle \}$$

where we use $\mathcal{F}$ to indicate the one dimensional Fourier transform, and brackets indicate the ensemble average. $\bar{\rho}_1$ and $\bar{\rho}_2$ are the phase point coordinates.

Equation (1) is related (Clifford, 1971) to the phase structure function (Carlson, 1967)

$$(2) \quad D_S(\rho) = 8\pi^2 k^2 \int_0^\infty \kappa d\kappa [1 - J_0(\kappa\rho)] \int_0^L d\eta C_n^2(\eta) \left(\frac{L}{\eta}\right)^2 \\ \times \cos^2\left(\frac{1}{2} \frac{(L-\eta)\kappa^2}{k\eta}\right) \phi_n^{(0)}\left(\frac{\kappa L}{\eta}\right)$$

and approximately to the wave structure function

$$(3) \quad D_S(\rho) = \left\{ \begin{matrix} 1. \\ .5 \end{matrix} \right\} D_W(\rho) = \left\{ \begin{matrix} 8 \\ 4 \end{matrix} \right\} \pi^2 k^2 \int_0^\infty d\kappa \kappa (1 - J_0(\kappa\rho)) \\ \times \int_0^L d\eta C_n^2(\eta) \left(\frac{L}{\eta}\right)^2 \phi_n^{(0)}\left(\frac{\kappa L}{\eta}\right)$$

by employing Taylor's frozen turbulence hypothesis

$$(4) \quad \phi(\rho(\eta), \tau+t) = \phi(\rho(\eta) - v\tau\left(\frac{\eta}{L}\right)\hat{x}, t),$$

where η is the longitudinal mean position variable, ($0 < \eta < L$), and $\hat{x}$ is the unit vector in the x (wind) direction. The index of refraction spectrum is

$$(5) \quad \phi_n^{(0)}(\kappa) = \left(\left(\frac{1.071}{L_0} \right)^2 + \kappa^2 \right)^{-11/6}$$

where κ is the spatial frequency and L_0 is the outer scale.

The resultant integrals consist of several terms that simplify into Delta functions. The remaining integral is written in terms of the normalized variables ρ , β and R , where

$$(6a) \quad \Omega \equiv \frac{2\pi f \rho}{V}$$

$$(6b) \quad R \equiv \frac{1.071 \rho}{L_0}$$

$$(6c) \quad \beta^2 = \left(1 + \frac{R^2}{\Omega^2}\right)$$

It is

$$(7) \quad \frac{W_{\delta s}(\Omega) \left(\frac{V}{\rho}\right)}{8.77 \left\{ \begin{matrix} 1. \\ .5 \end{matrix} \right\} k^2 L C_n^2 \rho^{5/3}} = \frac{2\Gamma(11/6)}{\sqrt{\pi} \Gamma(4/3)} \Omega^{-8/3} \beta^{-8/3}$$

$$\times \int_0^\infty dk \left[1 - \frac{\sin(\{\cos \theta + \kappa \beta \sin \theta\} \Omega)}{2\{\cos \theta + \kappa \beta \sin \theta\} \Omega} - \frac{\sin(\{\cos \theta - \kappa \beta \sin \theta\} \Omega)}{2\{\cos \theta - \kappa \beta \sin \theta\} \Omega} \right] (1+\kappa^2)^{-11/6}.$$

The integral in Eq. (7) can be evaluated for certain specific cases, depending on frequency, wind velocity, phase point orientation, and outer scale.

Table 1 summarizes the cases for arbitrary orientation angle. The limiting cases are divided into two groups depending on the relative size of $|\Omega \cos \theta \pm (\Omega^2 + R^2)^{1/2} \sin \theta|$ and $\sqrt{6}$. A general equation is given for each group. These groups are further simplified into two cases depending on the relative magnitude of the normalized separation and normalized frequency. Table 2 presents the results for large and small separation when the line connecting the phase points is horizontal. Table 3 gives corresponding equations when the wind is perpendicular to the line connecting the phase points.

Figure 2 presents a graph of the limiting cases when the normalized separation, R , is 0.1. This could represent the case for a near earth propagation measurement when the outer scale, $L_0 \approx 2$ meters and the phase point separation is ≈ 20 cm.




Fig. 2. Limiting form of normalized phase difference spectra versus normalized frequency for several orientation angles.

This figure illustrates the fact that small fluctuations in wind direction could easily mask the ω^2 dependence of the phase difference frequency spectrum for very small ω when the angle θ is nominally 0° .

Referring to Tables 1, 2 and 3, there is a break point in each of the phase difference spectra at $\omega = R$. This allows a direct determination of the outer scale in terms of break point frequency, phase point separation and wind velocity. Furthermore once L_0 is known, one can determine C_n^2 from the amplitude of the phase difference spectrum in one or more of the frequency regions where the limiting forms in Tables 1, 2 or 3 apply.

In summary the derivation of an expression for the phase difference spectrum was outlined and a final integral form was given. Tables were presented showing the results of evaluating the integral for limiting cases in each frequency range.

AVERAGING TIMES

This section will present a survey of the recent work on averaging time determination. A procedure for determination of averaging time while the data is being taken is presented. The developments follow previous work on averaging time in which data was tested after the fact to see if sufficiently long data sections had been obtained.

Here, as in previous work (3163-3) two equivalent averaging time criteria are considered; ergodicity, for which the normalized difference between time and ensemble averages is chosen as the indicator; and stationarity, for which the normalized difference between the data variance and the mean of the variances of many data subsections is chosen as the indicator.

Previously the data was examined after the recording process was completed. The examination indicated whether or not data sufficient for a given precision had been taken. A much more desirable approach is presented here; that of determining while the data is being taken what a sufficient duration will be for a given criterion and indicator.

The method presented assumes the availability of an on-line digital computer fast enough to handle simple arithmetic on the incoming data. An equivalent scheme presented previously (Report 3163-3) used electronic analog techniques. Both schemes rely heavily on developments in the literature: (Lumley and Panofsky, 1964, Bendat and Piersol, 1966, and Charnok and Robinson, 1957).

In the remainder of the section, previous work will be summarized to provide formulae for the real time procedure. Then the basic approach is to determine the autocorrelation integral scale from which the averaging time immediately follows.

TABLE 1

THE ONE SIDED PHASE DIFFERENCE SPECTRUM FOR
ANY ORIENTATION OF PHASE POINTS AND WIND VELOCITY

$$\Omega \equiv \frac{2\pi f \rho}{v}$$

$$R \equiv \frac{1.071 \rho}{L_0}$$

Small normalized separation ($\theta \neq 0^\circ$) $|\Omega \cos \theta \pm (\Omega^2 + R^2)^{1/2} \sin \theta| \ll \sqrt{6}$
and small normalized frequency

$$\begin{aligned} \frac{W_{\delta S}(\Omega) \left(\frac{v}{\rho}\right)}{8.77 \left\{ \begin{smallmatrix} 1 \\ .5 \end{smallmatrix} \right\} k_L^2 C_{n\rho}^2 5/3} &= (\Omega^2 + R^2)^{-4/3} \left[\frac{\Omega^2 \cos^2 \theta}{6} + \frac{(\Omega^2 + R^2) \sin^2 \theta}{4} \right] \text{General case,} \\ &= R^{-8/3} \left[\frac{\Omega^2 \cos^2 \theta}{6} + \frac{R^2 \sin^2 \theta}{4} \right] \quad R \gg \Omega, \\ &= \Omega^{-2/3} \left[\frac{\cos^2 \theta}{6} + \frac{\sin^2 \theta}{4} \right] \quad \Omega \gg R. \end{aligned}$$

Large normalized separation ($\theta \neq 0^\circ$) $|\Omega \cos \theta \pm (\Omega^2 + R^2)^{1/2} \sin \theta| \gg \sqrt{6}$
or large normalized frequency

$$\begin{aligned} \frac{W_{\delta S}(\Omega) \left(\frac{v}{\rho}\right)}{8.77 \left\{ \begin{smallmatrix} 1 \\ .5 \end{smallmatrix} \right\} k_L^2 C_{n\rho}^2 5/3} &= (\Omega^2 + R^2)^{-4/3} \quad \text{General case,} \\ &= R^{-8/3} \quad R \gg \Omega, \\ &= \Omega^{-8/3} \quad \Omega \gg R. \end{aligned}$$

TABLE 2

PHASE DIFFERENCE SPECTRUM WHEN THE WIND DIRECTION IS PARALLEL
TO THE LINE CONNECTING THE PHASE POINTS

$$\Omega = \frac{2\pi f \rho}{v}$$

$$R = \frac{1.071 \rho}{L_0}$$

$$\theta = 0^\circ$$

Small normalized separation:

$$\underline{R \ll \sqrt{6}}$$

$$\frac{W_{\delta s}(\Omega) \left(\frac{v}{c} \right)}{8.77 \left\{ \begin{matrix} 1. \\ .5 \end{matrix} \right\} k^2 L C_{n^0}^2 \frac{5}{3}}$$

$$= \frac{1}{6} R^{-8/3} \Omega^2$$

$$R \gg \Omega$$

$$= \frac{1}{6} \Omega^{-2/3}$$

$$\sqrt{6} \gg \Omega \gg R$$

$$= \Omega^{-8/3}$$

$$\Omega \gg \sqrt{6}$$

Large normalized separation:

$$\underline{R \gg \sqrt{6}}$$

$$\frac{W_{\delta s}(\Omega) \left(\frac{v}{c} \right)}{8.77 \left\{ \begin{matrix} 1. \\ .5 \end{matrix} \right\} k^2 L C_{n^0}^2 \frac{5}{3}}$$

$$= \frac{1}{6} R^{-8/3} \Omega^2$$

$$6^{3/14} R^{4/7} \gg \Omega$$

$$= \Omega^{-8/3}$$

$$\Omega \gg 6^{3/14} R^{4/7}$$

TABLE 3

PHASE DIFFERENCE SPECTRUM WHEN THE WIND DIRECTION IS
PERPENDICULAR TO THE LINE CONNECTING THE PHASE POINT

$$\Omega \equiv \frac{2\pi f \rho}{v}$$

$$R \equiv \frac{1.071 \rho}{L_0}$$

$$\theta = 90^\circ$$

Small normalized separation

$$\underline{\underline{R \ll \sqrt{6}}}$$

$$\frac{W_{\delta s}(\Omega) \left(\frac{v}{\rho} \right)}{8.77 \left\{ \begin{matrix} 1. \\ .5 \end{matrix} \right\} k_L^2 C_{n^0}^2 5/3}$$

$$= \frac{1}{4} R^{-2/3}$$

$$R \gg \Omega$$

$$= \frac{1}{4} \Omega^{-2/3}$$

$$2 \gg \Omega \gg R$$

$$= \Omega^{-8/3}$$

$$\Omega \gg 2$$

Large normalized separation

$$\underline{\underline{R \gg \sqrt{6}}}$$

$$\frac{W_{\delta s}(\Omega) \frac{v}{\rho}}{8.77 \left\{ \begin{matrix} 1. \\ .5 \end{matrix} \right\} k_L^2 C_{n^0}^2 5/3}$$

$$= R^{-8/3}$$

$$R \gg \Omega$$

$$= \Omega^{-8/3}$$

$$\Omega \gg R$$

The analytical background will now be outlined. Consider a recording of a random data function, $f(t)$. It is shown in several references: (Lumley and Panofsky, 1964, Bendat and Pierson, 1966, and Davenport and Root, 1958) that ϵ , the ensemble average mean square difference between the time and ensemble averages normalized to the mean value squares $\langle f \rangle^2$, is related to the averaging time T by

$$(8) \quad T = \frac{2C(0)}{\epsilon \langle f \rangle^2} \int_0^T \left(1 - \frac{t}{T}\right) \rho(t) dt$$

where

$C(0)$ = ensemble variance of $f(t)$

$\rho(t)$ = autocorrelation of $f(t)$.

It is generally pointed out that if the autocorrelation becomes substantially zero in a time much less than T , then the second term under the integral is negligible and T is given by

$$(9) \quad T = \frac{2C(0)}{\epsilon \langle f \rangle^2} I$$

where I is the integral scale of ρ ,

$$(10) \quad I = \int_0^\infty \rho(t) dt .$$

Another situation has been considered in the literature (Charnok and Robinson, 1957) in which there are N samples in the complete data section. The data section is divided into m subsections of $s = N/m$ elements.

The variance of each subsection is computed and $\overline{C_s(0)}$ the mean variance of each of the sections is examined. Then by definition

$$(11) \quad \overline{C_s(0)} = \frac{1}{m} \sum_{a=0}^{m-1} \frac{1}{s} \sum_{i=as+1}^{as+s} \left(f_i - \frac{1}{s} \sum_{i=as+1}^{as+s} f_i \right)^2 .$$

By subtracting the complete section mean, $\langle f \rangle$, from both terms in Eq. (11) expanding, rearranging, and writing the resultant expression in terms of the complete section variance

$$C(0) = \frac{1}{m} \sum_{a=0}^{m-1} \frac{1}{s} \sum_{i=as+1}^{as+s} (f_i - \langle f \rangle)^2 ,$$

we get

$$(12) \quad C(0) - \overline{C_s(0)} = \frac{1}{m} \sum_{a=0}^{m-1} \left(\frac{1}{s} \sum_{i=as+1}^{as+s} - \langle f \rangle \right)^2 .$$

Equation (12), as has been indicated previously (3163-3), gives a measure of stationarity. If the section length is sufficiently long and the function is stationary, then each subsection mean will be nearly equal to the complete section mean and the expression in Eq. (12) will be small. The question is how long must each subsection be in order for this to happen. A plot of $C(0) - \overline{C_s(0)}$ versus subsection length s , will indicate the section length for which the difference becomes small, and thus the subsection length for which the function may be regarded as essentially stationary. This requires, however, at least twice as much data as the minimum amount for stationarity (3163-3).

By further rearranging Eq. (12) and defining covariances $C_0(k)$

$$(13) \quad C_0(k) = \frac{1}{m} \sum_{a=0}^{m-1} \frac{1}{(s-k)} \sum_{i=as+1}^{as+s-k} (f_i - \langle f \rangle) (f_{i+k} - \langle f \rangle)$$

Equation (12) can be written in the form

$$(14) \quad \frac{C(0) - \overline{C_s(0)}}{C(0)} = \frac{1}{s} \sum_{k=-(s+1)}^{s+1} \left(1 - \frac{|k|}{s} \right) \frac{C_0(k)}{C_M(0)}$$

Taking the limit of the summation as the number of terms becomes large, identifying $C_0(k)/C(0)$ with the autocorrelation $\rho(k)$, and putting $T_s = s\delta t$ and $t_k = k\delta t$ where δt is the sampling interval gives

$$(15) \quad \frac{C(0) - \overline{C_s(0)}}{C(0)} = \frac{2}{T_s} \int_0^{T_s} \left(1 - \frac{t_k}{T_s} \right) \rho(t_k) \delta t_k$$

The integral in Eq. (15) is identical with that in Eq. (8). If (t_k) becomes small while t_k/T_s is still small, then the integral is identified with the integral scale in Eq. (10) and Eq. (15) becomes

$$(16) \frac{C(0) - \overline{C_s(0)}}{C(0)} = \frac{2I}{T_s}$$

Equation (16) gives the method of finding the integral scale as the data is being taken. The procedure is to plot the function $T_s(C(0) - \overline{C_s(0)})/2C(0)$ versus subsection duration, T_s , as data is being received for particular values of T_N , the total record duration. For small values of N the function $T_s(C(0) - \overline{C_s(0)})/2C(0)$ is poorly defined. As N becomes sufficiently large the curve asymptotically approaches the value of the integral inner scale.

Once the integral scale is computed, then Eq. (9) can be used to give the averaging time.

The approach of Eq. (16) to the proper integral scale value is limited by two factors. The first is the requirement that the term t_k/T_s in the parentheses under the integral in Eq. (15) be negligible. A data section of only a few integral scales is sufficient for that, and since the complete data section will be the order of a hundred integral scales in duration the requirement is satisfied. The other requirement is that the data sampling rate be sufficiently fast so that the transition from summation to integration in Eq. (15) is sufficiently precise. The digitizing rate can be chosen sufficiently fast to satisfy this requirement, for typical cases when $I \sim 1$ second.

The net result is that the value of the function approaches the value of the integral scale in a time much less than the total data section. The application of Eq. (9) then gives the predicted averaging time.

An experimental check of the above procedure was carried out using optical phase difference data. The data was recorded at the RADC Laser Propagation Range in Verona, N.Y. Figure 3 shows a trace of $T_s(C(0) - \overline{C_s(0)})/2C(0)$ versus data length, T_s , for $T_N = 10$ sec. There it is seen that the ordinate value peaks and then drops to a lower level. The points represent the values $T_N/2, T_N/3, T_N/4$ secs, etc., corresponds to $m = 2, 3, 4$, etc. The value at the peak is chosen as a representative conservative value of the integral scale. This value is roughly 0.8 secs which would lead to an averaging time in the range of 80 seconds. Other corroborating studies (3163-3) show that the 80 seconds range is indeed the appropriate one. Further the value was predicted using the first eighth of the data.




Fig. 3. Real time determination of the integral scale for phase difference data.

In a real time procedure the computer would plot curves equivalent to that of Fig. 3 for successively larger intervals (T_N) until a consistent peak is recognized.

To summarize, a real time computer method of predicting averaging time has been outlined and applied to a single set of random optical data. The process predicts an integral scale time from which the averaging time is then obtained. The process will be outlined in more detail and further demonstrated in a report to be released subsequently.

SUMMARY

During the past quarter, work has concentrated on temporal spectra of phase difference fluctuations of atmospherically degraded spherical waves and on averaging times of optical random data. The temporal spectra considered the case of a horizontal wind transverse to the optical path but with phase points oriented so that their separation vector makes an arbitrary angle with the horizontal. The spectra also included outer scale effects. A final integral was given along with the expressions and graphs for particular limiting cases.

The averaging time discussion outlined a technique for determining the averaging time of random data while the data is being recorded but in a time much shorter than the total data duration. This method assures that a sufficient but economical duration of data is recorded. The procedure assumes the availability of an on-line digital computer.

BIBLIOGRAPHY

- (3163-3), S. A. Collins, "Investigation of Laser Propagation Phenomena," Report 3163-3, April 1972, ElectroScience Laboratory, The Ohio State University Department of Electrical Engineering; prepared under Contract F30602-71-C-0132 for Rome Air Development Center, Air Force Systems Command, Griffiss Air Force Base, New York. (RADC-TR-72-123)
- (Bendat and Piersol, 1966), J. S. Bendat and Piersol, A.G., Measurement and Analysis of Random Data, John Wiley and Sons, New York, 1966.
- (Carlson, 1967) F.P. Carlson, "Propagation in Stationary and Locally Stationary Random Media," Ph.D. Dissertation, University of Washington, 1967.
- (Charnok and Robinson, 1957) H. Charnock, and Robinson, G.D., Spectral Estimates from Subdivided Meteorological Series, a paper of the Meteorological Research Committee (London), M.R.P. No. 1062, S.C. III/240, 27 August 1957.
- (Clifford, 1971) S. F. Clifford, JOSA 61 1285 (1971).
- (Davenport and Root, 1958) W. B. Davenport and Root, W.L., Random Signals and Noise, McGraw-Hill Book Company, New York, 1958.
- (Lumley and Panofsky, 1964) Lumley, J.L. and Panofsky, H.A., The Structure of Atmospheric Turbulence, John Wiley and Sons, New York, 1964.