AD 756585

1.

Report Number 7

ORGANOTINS FOR ANTIMALARIAL SCREENING

Final Report

by Henry G. Kuivila

5 January 1973

Supported by

U. S. Army Medical Research and Development Command

Washington, D. C. 20315

Contract_DA-49-193-MD-3018_

Research Foundation of the State University of New York

Albany, New York 12224

This document has been approved for public release

and sale; its distribution is unlimited.

The findings in this report are not to be construed as an official Department of the Army position unless so designated by other authorized documents. **DISCLAIMER NOTICE**

THIS DOCUMENT IS BEST QUALITY PRACTICABLE. THE COPY FURNISHED TO DTIC CONTAINED & SIGNIFICANT NUMBER OF PAGES WHICH DO NOT REPRODUCE LEGIBLY.

TABLE OF CONTENTS

| I. | General | Summary |
|------|---------|---------------------------------|
| II. | Synthet | ic results |
| | A. | Sulfones |
| | В. | Allenyltins |
| | C. | Bi- and Tricyclic Organotins |
| | D. | Nitrogen-containing Organotins |
| | E. | Organotin-Schiff Base Complexes |
| III. | Experi | mental, |
| | A. | Allyltin Compounds |
| | В. | Sulfones |
| | C. | Allenyltins |
| | D. | Bi- and Tricyclic Organotins |
| | E. | Nitrogen-containing Organotins |

LIST OF TABLES

| Table II | The Physical Properties and Analyses of Allylalkyltin Compounds |
|-----------|--|
| Table III | THE UV SPECTRAL DATA OF ALLYLTIN COMPOUNDS 28 |
| Table IV | The Physical Properties and Analyses of Sulfones |
| Table V | Allenyl and Propargyltins |
| Table VI | ANALYTICAL DATA ON PYRIDYLTINS |

1

ORGANOTINS FOR ANTIMALARIAL SCREENING

Summary of Final Report

When new strains of malaria were discovered in Southeast Asia which did not respond to treatment by the usual antimalarials quinine and chloroquine, an extensive screening program for new antimalarials was initiated at the Walter Reed Army Institute of Research. It was discovered that certain organotins displayed activity toward some of these parasites. The author of this report was invited to submit a proposal for the synthesis of compounds which might show promise. A long range proposal was submitted and formed the basis of the research conducted under this contract.

It was proposed to prepare for screening organotins in four general categories: 1) preparation of a variety of relatively simple organotins for general screening; 2) introduction of organotin groups into quinine and its derivatives; 3) introduction of organotin groups into 4-aminoquinoline; 4) introduction of organotin groups into sulfones which had shown some antimalarial activity.

During the course of the contract it was possible to carry on research on three of these categories. However, successful synthetic results were obtained only in the first of these. A total of forty three compounds was submitted for testing, and test results were returned of thirty six of these. These included: sulfones, alkylallyltins; alkylallenyltins; alkylpyridyltins; and bis-salicylidenediaminoethane-alkyltin halide complexes. Some showed activity in mice, but were inactive or toxic in birds; and <u>vice versa</u>. None was active in both, and many were toxic in one or both.

ORGANOTINS FOR ANTIMALARIAL SCREENING

002

144 SAN 20

I. General Summary

When new strains of malaria were discovered in Southeast Asia which did not respond to treatment by the usual antimalarials, quinine and chloroquine, an extensive screening program for new antimalarials was initiated at the Walter Reed Army Institute of Research. It was discovered that certain simple organotins displayed activity toward some of these parasites. The author of this report was instead to submit a proposal for the synthesis of compounds suitable for antimalerial screening. A long range proposal was submitted and formed the basis of the research conducted under this contract. Euringthe two years of the contract it was possible to complete only a portion of the longer range goals.

It was proposed to carry on research under two broad categorics: 1) modification of known antimalarials by the introduction of organotin functions; 2) proparation of a variety of simple organotins in the hope that some would be found to be active.

In the first category one of the more premising objectives involved the modification of quinine to form compounds of structures 1 and 2. Apparently simple routes for their preparation are available, but no success was realized.



It was also hoped to modify the aminoquinoline group of antimalarials by introduction of an organotin group to form compounds such as <u>3</u>. However, time did no permit experimentation in this direction. The third class, whose synthesis was attempted unsuccessfully, were crotyl sulfones containing



the second s

organotin functions: (R₃SnCH=CHCH₂)₂SO₂ and CH₃CH(SnR₃)CH=CHSO₂CH₂CH=CHCH₃. However, forty three more simple compounds were prepared and submitted for screening. These are listed in Table I. The table includes all of the screening results made available to us. It will be noted that a number of the compounds are toxic to mice or chicks, or both. However, a few (7-9,11) are active in mice and, in the cases of 8 and 9 are inactive in chicks. In summary, it may be said that none of the compounds tested showed any real promise as antimalarials.

Table I.

ORGANOTINS SUBMITTED FOR ANTIMALARIAL SCREENING

Ę

のであるという

÷x,

Į

Submitter code 0378

| | - | | scree | ning r | results ⁸ | L | |
|-----|--------------------------------------|------------|------------|--------|----------------------|----|----|
| | ~ • | , <u>,</u> | мъ | BC | P ^d | l | C, |
| 1. | phenyl propargyl sulfone, WR-49464-B | | t | | | | |
| 2. | phenyl crotyl sulfone, WR,-84820-A | | i | | i | | |
| 3. | phenyl allyl sulfone, WR-84818-A | | i | | i | | |
| 4. | ethy] propargyl sulfone, WR-84819-A | • | 1 | | i | | |
| 5. | tetrallyltin, WR-73750-B | | 1 | | | | |
| 6. | allyltrimethyltin, -WR-80280-A | | t | | | | |
| 7. | triethylallyltin, WR-80281-A | ł | 3 | | | | |
| 8. | allyltri-n-propyltin, WR-80282-A | i | 8 | i | | | |
| 9. | allyltri-n-butyltin, WR-80283-A | ł | 1 | i | | | |
| 10. | diallyldimethyltin, WR-80284-A | : | L | | | | |
| 11. | diallyldi-n-propyltin, WR-80285-A | ŧ | 1 . | | | | |
| 12. | diallyldi-n-bùtyltin, WR-80286-A | 4 | Ĺ | | | | |
| 13. | diallyldiethyltin, WR-80287-A | | | | | | |
| 14. | triallylmethyltin, WR-92109-A | | | | | | |
| 15. | triallyl∋thyltin, WR92089-A | | | | | | |
| 16. | trially1-n-propyltin, WR-92090-A | | | i | | | |
| 17. | trially1-n-butyltin, WR-92091-A | | | | | | |
| 18. | triallylphenyltin, WR-92092-A | | | | | | |
| 19. | diallyldiphenyltin, WR-59215-C | | | i | | | |
| 20. | allyltrichenyltin, WR-27343-C | | | | | | |
| 21. | tri-n-butylallenyltin, WR-27343-C | t | | i | | | |
| 22. | triphenylallenyltin, AF58635 | | | i | ao, | cs | |
| 23. | tripropylallenyltin, AF58662 | t | | t | | • | |
| 24. | triphenylallenyltin, AF58671 | t | | t | | | |
| 25. | di-n-butyldiallenyltin, AF58680 | t | | 4 | | | |

| Table | Ι | (contd | 1 |
|-------|---|--------|---|
|-------|---|--------|---|

| | Table I (co | ntd) | |
|-------------|--|----------------|----|
| | | M ^a | Bp |
| 26. | di-n-propyldiallenyltin, AF58706 | i | t |
| 27. | p-tri-n-butylstannyl-N,N-dimethyl- | t | i |
| 28. | 2-trimethylstannyl-5(or 6)- trichloromethyl-6(or 5)-chloro- bicyclo/2.2.1/heptane, AF58617 | t | |
| 2 9. | 2-trimethylstannyl-5(or 6)-thio- phenylbicyclo/2.2.1/heptane, AF58626 | t | |
| 30. | 2(or 3)-trimethylstannyl-5(or 6)- tri-n-butylstannylbicyclo/2.2.1/ heptane, AE72425 | i | i |
| 31. | bis-2(or 3),5(or 6)-trimethylstannyl- bicyclo/2.2.1/heptane, AE72416 | t | 8 |
| 32. | 3-trimethylstannyltricyclo/2.2.1.0 ²⁻⁶ / heptane, ** | | |
| 33. | penta-2,3-dien-2-yltrimethyltin,AE7244 | 3 t | i |
| 34. | penta-2, 3-dien-2-yltri-n-butyltin, AE?2 | 434 a | t |
| 35. | tri-n-probyl-3-pyridyltin, ** | t | |
| 36. | tri- <u>n</u> -butyl-3-pyridyltin, 🌫 | t | |
| 37. | n-butyl-tris-(3-pyridyl)-tin, ** | | |
| 38. | triphenylproparzyltin, ** | t | i |
| LO. | $R, R' = CH_{2}, AC94921$ | t | |

3 41. R, R' = C_2H_5 , AC94930 t 42. R, R' = $\underline{n}-C_{3}H_{7}$, AC94949 $R = CH_{3}$; $R' = C1_{3}$, AC94958 1 43.

^a Mica; ^b White leghorn chicks; ^c <u>Flasmodium gallinaceum</u> infected <u>Aedes aegypti</u>

** No accesion number available.

bis-salićylidenedčumino. Uhane-alkyltin halide vomplexes N=CH 9 5-H=NCH 5

5

P^C

ps

1

1

i

i

SYNTHETIC RESULTS

A. <u>Sulfones</u>. - Sulfones with α -hydrogen have been routinely converted to monoanions by a variety of basic reagents including methyl magnesium bromide¹; <u>n</u>-butyllithium²; potassium <u>t</u>-butoxide³; sodium hydride⁴; and lithium amide⁵.

Field and McMariand indicated that <u>n</u>-butyllithium was the most effective metalation reagent, and benzene as solvent afforded better yields than ether or tetrahydrofuran. We have attempted to prepare organotin substituted sulfones by the following reaction sequence.

 $C_6H_2SO_2CH_2CH = CHR + BuLi + C_6H_5SO_2CHCHCHR^+ LI^- + BuH$ R_3SnCI $C_6H_5SO_2CHCHCHR + LICI$

ŚńRa

However, the products sought did not form, presumably due to the low nucleophilicity of the sulfanyially! anion, and the relative weekness of the fin-carbon bond adjacent to the sulfanyi group.

An alternate attempt to precare organotin substituted subsciences involved the addition of triethyltin hydrice to phenyl propargyl sulfone. Whereas simple acetylunes undergo facile addition of organotin hydrides, the sulfone was impervious to triethyltin hydride.

B. <u>Allenyltins</u>. - The groups bonded to the terminal carbon atoms in allenes exist in two planes perpendicular to each other. The allene molecule is, therefore, dissymmetric if the groups present on either of the terminal carbon atoms are not the same.

One of the main areas of interest in allene chemistry is the syntheses and resolution of optically active compounds. Recently, the resolution and absolute configuration of 1,3-dimethylallene has been carried out by Waters and Caserio⁶

With a view to study the stereochemistry of allenyltin

derivatives, compounds III and IV have been prepared by the method described earlier?



Amalgamated magnesium, prepared by stirring magnesium turnings in an etherial solution of mercuric chloride, was used for preparing the Grignard reagent from 2-brono-3-pentyne. Still, the yield of allenyitin compounds was considerably low ($\sim 25-30$ %), probably due to coupling reactions. The use of cyclic Grignard reactor did not prove of any advantage.

The allonyitin derivatives prepared by the above method were contaminated with substantial amounts of unreacted trialkyltin chloride. However, they could be easily purified by treating aqueous KF and removing white precipitate of trialkyltin fluoride. After careful fractionation, the allenyitins were obtained in $\sim 90\%$ purity.

<u>The Propargyl/Allenyltin System</u>. - This system was studied because of the structural similarity of the compounds to the allyltins prepared and submitted earlier by Mr. Kawakami; it has not previously been studied in any great detail, though Le Quan and Cadiot^{®,9} have published some results on it, while Cochran and Kuivila¹⁰ have studied substituted allenyltins.

Two methods were used to prepare these compounds. Firstly, propargyl bromide was treated with n-butyllithium (hopefully to give the propargyllithium), and the product reacted with the appropriate trialkyltin halide.

 $BrCH_2C\equiv CH + BuLi \rightarrow LiCH_2C\equiv CH \xrightarrow{R_3SnCl} R_3SnCH_2C\equiv CH$ The second method used was that described in the literature, namely treatment of the Grignard reagent derived from propargyl bromide with the appropriate trialkyltin halide.

 $[BrMgCH_2C\equiv CH] + R_3SnCI \rightarrow R_3SnCH_2C\equiv CH + MgBrCI$

in all cases studied (R=Et,Bu,Ph) the products obtained from the reaction scheme shown in eq. I were neither allenyl nor propargyltins, but propynyltins R₃SnC=C-CH₃, as evidenced by I.R. and P.M.R. spectroscopy. This is probably due to rearrangement of the lithium derivative: it was found that on adding propargyl bromide to butyllithium in ether at low temperatures, a yellow color first built up in the solution, but within a few seconds disappeared. This behavior was observed at various temperatures and at various raises of addition of propargyl bromide and, also, if the butyllithium was added to the propargyl bromide.

The results obtained when the Grignard route was used were rather more complex. Gaudemar and co-workers have suggested that the Grignard reagent derived from propargyl bromide is entirely allenic^{9,10} this conclusion is not borne out by the work reported here.

Allenyltins are characterized by a band in the infrared at ca. 1930 cm⁻¹ (C=C=C), while propargyltins have a band at ca. 2110 cm⁻¹ (C=C) and propynyltins a band at ca. 2160 cm⁻¹ (C=C).

The product obtained when the Grignard prepared from propargy!

bromide and magnesium in ether was treated with Bu₃SnCl was a mixture of tributy|propargy|- and tributy|alleny|-tin, and on standing for about a day, the propargy! compound isomerized to the alleny! compound. When the preparation was repeated using magnesium amalgam rather than magnesium metal, no trace of the propargy! isomer was observed in the product.

Similarly, triethyltin chloride and the Grignard (magnesium amalgam) gave only the allenyl isomer, but this (though in an impure state) isomerized on standing to the propynyl isomer (Et₃SnC C-CH₃).

A more calcultive series of investigations was carried out in the case of triphenyltin chloride; here, it had been reported that the propargyl and allenyl isomers could be isolated and were both stable. (This is the normal trend in organotins, aryltin compounds generally being more stable than alkyltins.) Here the Grignard was prepared in three different ways:-

- a) Propargyl bromide/Mg amaigam/ether
 Here the product mixture contained triphenylallen yltin and triphenylpropargyltin in the ratio 2:1,
 with a trace of the propynyl derivative.
- b) Propargyl bromide/Mg metal/ether

The product mixture contained the allenyl and propargyl isomers in the ratio 1:6, and the propargyl isomer was obtained in a pure state, the physical properties agreeing with those quoted by Le Quan and Cadiot.

c) Propargyl bromide/Mg metal/THG
 The product mixture contained no propargyltin,
 but included allenyl and propynyltins.

Thus, it appears that this particular system is extremely labile, and that under only slightly differing conditions various product mixtures may be obtained. The isolated triphenylpropargyltin will be submitted but, at the present time, we do not plan to carry out any further studies on this particular system.

c. <u>Bl- and Tricyclic Organotins.</u> - Although ionic additions to norbornadione has been studied extensively!! the analogous free radical additions have received comparatively little attention. <u>p</u>-Theocresol has been shown to add to norbornadiene, giving a mixture of <u>exo-5-</u> norbornen-2-yl <u>p</u>-tolyl sulphide and 3-nortricyclyl-<u>p</u>-tolyl sulphide!⁷ <u>p</u>-Tolyl sulphonyl chloride⁷ perfluoropropyl iodide¹⁴ and bromotrichloronothene⁴, however, gave the corresponding 3,5-disubstituted nortricyc+ lenes. Carbon tetrachloride, chloroform and diethylphosphite, on the other hand, gave the corresponding 3-nortricyclyl derivative as the major product!⁵

The present investigation was undertaken as a part of the extensive study of free radical additions of organotin hydrides to unsaturated compounds being carried out in these laboratories in order to provide more organotins for screening.

Trimethyltin hydride reacted radily with norbornadiene on photolysing an equimolar mixture in a pyrex sealed ampoule to give a

のないである



The ratio of the components was as shown above. The same mixture was obtained when the reaction was carried out thermally or photolytically in a quartz tube.

On a 1,2,3-tris-β-cyanoethoxypropane (TCEP) column, compounds I and II show the same retention times whereas on an Apiezon column, compounds II and III show same retention time. Compound I was identified by independent synthesis from 3-bromonortricyclene¹⁶ and trimethyltin chloride via the Grignard method. Compounds II and III have previously been characterized by Francis Pelczar in these laboratories¹⁷

The free radical chain nature of such additions has been well substantiated and the following equations account for the formation of Compounds 1 - 111 ($Sn=Snie_3$).



$$\overbrace{(V11)}^{\underline{Sn}} \rightleftharpoons \overbrace{(V11)}^{\underline{Sn}}$$
(3)

 $(V) + HSn \longrightarrow 11$ (4)

 $(VI) + HSn \longrightarrow III$ (5) $VII + HSn \longrightarrow I$

11

When mixture (A) (b.p. $35^{\circ}/0.02 \text{ mm}$) was photolysed with excess trimethyltin hydride (~ 2 moles), Compounds I and IV remained unchanged but II and III reacted further and gave a mixture (B) of diadducts (b.p. $82^{\circ}/0.02 \text{ mm}$). Mixture (B) shows only three peaks on SE30 column and four peaks on aplezon column. This mixture possibly contains Compounds VIII - XI. The tormation of compounds XII and XIII is most unlikely on steric grounds.



The interaction of mixture (A) with trimethyltin hydride was carried out under various conditions, photolytically as well as thermally and in every case mixture (B) was obtained. Similar results were observed when the reactions were carried out between norbornadiene and excess (~ 2.5 moles) trimethyltin hydride.

Addition of tributyltin hydride to mixture (A) was comparatively slower. However, in this case also, compounds I and IV remained unchanged but II and III gave a mixture (C) of bisadducts quantitatively (b.p. 135°/0.02 mm).



Tributyltin hydride reacted with norbornadiene in equimolar ratio to give a mixture (D) of monoadducts (b.p. 105°/0.02 mm). This mixture possibly contains the <u>exo</u> and <u>endo</u> norbornyl as well as the nortricyclyl derivatives. No attempt was made to isolate the different

components or to determine their relative ratio in the mixture. On S.E.30 column, the mixture shows only two not well separated peaks; the small peak may be due to nortricyclic compound and the big peak due to exo and endo isomers.

Mixture (D) reacted with excess trimethyltin hydride very _ slowly to give a mixture (E) of diadducts, (b.p. 135°/0.02 mm).



Mixtures (C) and (E) seem to consist of the same isomers. They have identical boiling points and their infra-red spectra are virtually identical. However, the relative ratio of the isomers is not the same in the two mixtures as shown by g.l.p.c. on S.E. 30 column.

As expected, the attack of trifluoroacetic acid on mixture (B) was found to be extremely slow. On stirring the mixture with excess trifluoroacetic acid (\sim 6 moles) in pentane for four days, a white solid gradually precipitated out with evolution of methane. The infra-red spectrum of the solid showed that all methyl groups were not cleaved.

. D. Nitrogen-containing Organotins.

Simple Quinolines. - In the preparation of 2-triethyltin-4methyl quinoline from 2-chloro-4-methyl quinoline with butyllithium in anhydrous ether, followed by the addition of triethyltin chloride, a compound, m.p. 57-58.5°, was obtained. The nmr spectrum indicated quinoline protons, methylprotons, and ethyl protons at τ 7.5, τ 2.5 and τ 1.0 with the relative integrated intensities in the ratio of 28:19:90 (5:3:15), respectively. However, the elemental analysis did not give a satisfactory result, indicating there was impurities present in the compound. Therefore, further purification is necessary before submitting for anti-malarial screening.

For the preparation of 3-trialkyltin quinoline, the final products were viscous liquids. Distillation or recrystallization failed to lead to the desired compound. It appears that the halogenmetal exchange reaction goes to completion as shown by the color test for lithium quinoline. In addition to the desired 3-trialkyltin quinoline, the coupling product, diquinolyl might be formed as side product which could contaminate the desired compounds.

B. 7-Chloroquinolines. - Attempts have been made to prepare 4-tributylstannyl 7-chloroquinoline by three methods: a) treatment of the dichloro compound with n-butyl lithium at -40° to -50°, followed by treatment of the lithium derivative with tributyltin chloride, b) treatment of the dichlorocompound with tributyltin lithium, and c) preparation of 4-lithio 7-chloroquinoline from the dichlorocompound and lithium metal, followed by treatment with tributyltin chloride.

Neither method b) nor c) was successful. Method a) produced

a mixture of several components with no major component under the conditions first used (addition of butyl lithium at -45° to a THF suspension of 4,7-dichloroquinoline, followed by addition of tributyltin chloride at room temperature), but a change in conditions (diethyl ether as solvent, addition of tin halide at -40°, addition of excess ammonium chloride solution before workup) produced a mixture with two major components; distillation, followed by G.L.C. analysis, showed the more volatile of these to be unreacted tributyltin chloride. The other component could not be obtained in anything approaching a pure state.

However, the recovery of tributyltin chloride in such a high yield (>60%) indicates that in this case the problem is complicated by coupling of two quinoline nuclei, involving some quinoly! Lithium as an intermediate.

In the case of 2-chloroquinoline and of other quinoline derivatives studied by Dr. K. C. Yeh, where buty! lithium is added to a solution of the quinoline derivative rather than a suspension, the reaction appears to be quite straightforward.

C. Quinines. - It was found that quinine could be readily converted into its tin alkoxide by azeotropic dehydration of a mixture of bistributyltin oxide and quinine, using toluene as solvent,



but that the compound so formed, in common with other tin alkoxides, was too readily hydrolyzed by moisture in the air to be of any volue. It dld, however, react like simple alkoxides with acceptor molecules such as phenyl isocyanate and chloral:

「日本」というと言語でいた。

 $Bu_3Sn OR + PhNCO \rightarrow Bu_3Sn NPh CO R$

 $Bu_3Sn OR + CCI_3CHO \rightarrow Bu_3Sn \cdot O \cdot CH(CCI_3) OR$

The products of these reactions are also very readily hydrolysed; however, the hydrolysis products have been isolated and will be submitted when satisfactory analytical data have been obtained.

A second approach used was to attempt to convert the ethylenic group in both quinine and cinchonine to an acetylenic group by bromination and subsequent dehydrobromination and then to add trialkyltin hydrides to this group. However, we were unable to repeat earlier work reported in the literature⁴ which indicated that this had been done.

Attempts to replace the hydroxyl group of quinine by bromine, which could then perhaps be replaced by a trialkyltin group, were also not successful.

D. Alkylpyridines. - Gilman¹⁸ has reported the preparation of triphenyl-2- and -3- pyridyltin. His method has been used and extended to prepare the following compounds. The purity after distillation of the compounds, as calculated from G.L.C. traces, is given in parenthesis.

| Bu₃Sn 2-py | (87%) | b.p. 103-105°/.075 mm |
|---------------------------------------|-------|-------------------------|
| Bu₃Sn 3-py | (97%) | b.p. 106.5-108°/.075 mm |
| Bu ₂ Sn(2-py) ₂ | (95%) | b.p. 130-132°/.1 mm |
| Bu ₂ Sn(3-py) ₃ | (91%) | b.p. 154-158°/.075 mm |
| Pr₃Sn 2-py | (95%) | b.p. 96°/.15 mm |
| Pr₃Sn 3-py | (88%) | b.p. 91-93°/.1 mm |

These compounds will be submitted when purification procedures (where necessary) have been carried out and when satisfactory analytical data become available.

The reaction scheme used in c) and d) may be summarized as

follows, where **TR** is n-butyl or n-propyl R^t may be pyridyl or quinolyl, X is halido:

 $R'X + BuLi \xrightarrow{-40^{\circ}} R'Li + BuX$

 $n R^{\dagger}Li + (4-n) R_{4-n} SnCi_{n} \longrightarrow (4-n) R_{n} SnR^{\dagger}n + nLiCi_{n}$

N-Tri-n-butyIsulfanilamide. - This compound was prepared during the course of Dr. Mitchell's Ph.D. research. The preparation has been repeated as this compound is an organotin derivative of a known antibacterial and contains a relatively unreactive tin-nitrogen bond; the compound can be handled in air for short periods without any appreciable hydrolysis occurring. E. <u>Organotin Schiff Base Complexes</u>. - So far, our efforts have involved the ligand, bissalycilidenediaminoethane (BSDE·2H) and two other ligands were checked as possible chelating agents, but they turned out to be hopeless. The preparative procedure was a deprotonation reaction of BSDE·2H by means of Bu_2Sn0 , $Bu_2Sn(OMe)_2$ and $Bu_2Sn(NEt_2)_2$. In all cases, reactions gave H_20 , CH_30H and $NH(C_2H_5)_2$ but the products always had wide melting ranges. In one typical case (Bu_2Sn0) the product was run through a column packed with neutral alumina and T.H.F. The solid product obtained after evaporation of T.H.F. was recrystallized several times from n-heptane, and yellow crystals with m.p. of 154-156°C were obtained. Starting from 5 g. of crude product, only 0.5 g. of crystals were recovered.

In order to check the coordination ability of nitrogen atoms in BSDE 2H molecule, reactions between organotin chlorides and the ligand were tried. Four simple addition compounds have thus been prepared: $R_2SnCl_2 \cdot BSDE \cdot 2H$ (R = CH₃, C_2H_5 and $n-C_3H_7$) and $C_2H_5SnCl_3$ BSDE 2H. A reaction between trimethyltin chloride and ESDE $\cdot 2H$ was carried out but no immediate precipitation of adduct was observed. Thus, trialkyltin chloride does not have enough acceptor capacity toward this ligand which seems to function well as a bidentate ligand. No hexa-coordinated trialkyltin chloride adduct has been reported. All of these compounds showed good C, H, N, Cl, and Sn analyses. Therefore, if the Sn-O bond formation occurred, these nitrogen atoms would be able to coordinate to tin atom if the conformation were favorable.

Some features of n.m.r. spectra of the BSDE2H ligand, $(CH_3)_2SnCl_2\cdot BSDE\cdot 2H$ and Bu_2SnO ($Bu_2Sn(BSDE)$) merit comment. There are

11

four typical protons: -CH=N-, aromatic ring, =N-CH₂- and alkyl protons attached to tin in order from the low field to the high field. They are summarized in Table 1.

TABLE I

τ VALUE (p.p.m.) OF BSDE2H AND ITS TIN COMPLEXES

| Compound | -CH=N- | | +N-CH2 | - | Sn-CH ₃ | Solvent |
|---|--------|-----|--------|-----|--------------------|---------------------------------|
| BSDE2H | 1.68 | | 6.15 | | - | CH2CI2 |
| (CH ₃) ₂ SnCl ₂ •BSDE2H | 1.65 | | 6.03 | | 8,84 | CDCI3 |
| Bu2Sn(BSDE) | 1.67 | (1) | 6.10 | (2) | - | CDCI3 |
| | 2.00 | (9) | 6.17 | (8) | | |
| Bu ₂ Sn(BSDE) | 1.62 | (4) | 6.08 | (4) | - | CH ₂ Cl ₂ |
| | 1.98 | (5) | 6.19 | (5) | | |

In all these compounds, good relative intensity ratios were obtained. An interesting thing is that $(CH_3)_2SnCl_2BSDE \cdot 2H$ adduct shows only one set of -CH=N- and =N-CH₂- protons but $Bu_2Sn(BSDE)$ complex shows two sets of these protons whose intensity changes depending on a small change of the solvent. One assumption in the case of $Bu_2Sn(BSDE)$ is that the intramplecular motion sometimes favors the coordination of both nitrogen atoms which must be easier at low temperature. We assume a chelate structure by two nitrogen atoms forming a six-coordinated tin atoms in the case of the adducts, since only one set of -CH=N- and -N-CH₂- protons were observed.

As shown in Table 1, =N-CH₂- protons of $(CH_3)_2SnCl_2BSDE \cdot 2H$ are shifted to downfield by 0.12 p.p.m. from those of the ligand BSDE2H. This might be because nitrogen atoms coordinate to tin. Among two sets of =N-CH₂- protons in Bu₂Sn(BSDE), the low field ones might be attached to the nitrogen coordinated to tin. Among two sets of -CH=Nproton; the low field one could be attributed to the one adjacent to the coordinated nitroge, and the high field one could occur because of the breaking down of a strong hydrogen bond which exists in the ligand itself.

BIBLIOGRAPHY

1. L. Field and J. W. McFarland, J. Amer. Chem. Soc., 75, 5582 (1953).

- 2. W. E. Truce and R. D. Buser, ibid., 76, 3577 (1954).
- 3. H. D. Becker and G. A. Russell, J. Org. Chem., 28, 1896 (1963).
- 4. E. J. Corey and M. Chaykovsky, J. Am. Chem. Soc., 87, 1345 (1965).
- 5. L. Field and J. W. McFarland, *ibid.*, 75, 5582 (1953).
- 6. W. L. Water and M. C. Caserio, personal communication.
- 7. J. C..Cochran, Ph. D. dissertation, University Of New Hampshire, 1966.

8. M. Le Quan and P. Cadiot, Comtes rendus, 254, 133 (1962).

9. M. L. Quan and P. Cadiot, Bull. Soc. Chim. Fance, 1965 (1), 45.

10. H. G. Kuivila and J. C. Cochran, J. Am. Chem. Soc., <u>89</u>, 7152 (1967).

11. P. von R. Schleyee, ibid., 80, 1700 (1958).

12. S. J Cristol, G. D. Brindell and J. A. Reeder, ibid., 80, 635 (1958).

13. S. J. Cristol and J. A. Reeder, J. Org. Chem., <u>26</u>, 2182 (1961).

14. N. Ol Brace, <u>ibid</u>., <u>27</u>, 3027 (1962).

VICE LEBE

- 15. D. J Trecker and J. P. Henry, J. Am. Chem. Soc., <u>85</u>, 3204 (1963).
- 16. J. D. Roberts, E. R. Trumbull, Jr., W. Bennett and R. Arrstrong, <u>ibid.</u>, <u>72</u>, 3116 (1950).
- 17. F. Pelczar, Ph. D. Thesis, University of New Hampshire, 1968.
- 18. H. Gilman and T. N. Goreau, J. Org. Chem., <u>17</u>, 1470 (1952).

III. EXPERIMENTAL

A. Allyltin Compounds

Allyltri-n-propyltin

Into a dry 250 ml three-neckec flask equipped with a sealed stirrer. a reflux condenser and a pressure-equalized dropping funnel were placed 7.3 g. (0.30 g. atoms) of dry magnesium turnings and 100 ml of anhydrous ether. The flask was cooled in an ice bath, a few drops of ethylene dibromide were added and a solution of 11.5 g. (0.15 mole) of allyl chloride in an equal volume of anhydrous ether was added dropwise over a period of 1 hour. After the addition was completed, the reaction mixture was stirred for 30 minutes. The Grignard reagent was transferred by means of nitrogen pressure into a 500 ml flask similarly equipped. The residue in the first flask was washed with 20-30 ml of dry ether and the wash liquid was added to the second flask. Then a solution of 28.3 gr (22.3 ml, 0.10 mole) of tri-n-propyltin chloride in 20 ml. of ether was added to the Grignard reagent in the course of 1 hour during which the reaction mixture warmed to room temperature. The reaction mixture was heated to reflux for 3 hours, was cooled and hydrolyzed with saturated aqueous ammonium chloride solution until it showed two clear layers. The organic layer was separated and the inorganic layer was washed with ether. The combined organic layer and ether washings were washed by dry ammonia to remove any organotin chlorides. The white precipitate which was thought to be n-propyltin chlorideammonia adduct was filtered off and the filtrate was concentrated and distilled under reduced pressure to give 16.1 g. (59%) of allyltri-n-propyltin, b.p. 114-115/13 mm.

The other allyltin compounds were all prepared in similar ways. Yields varied from 30% to 70%. Generally speaking, allylmagnesium bromide gave, better yield than the chloride.

Allytrimethyltin

The Grignard reagent was prepared from 7.4 g. (0.35 g. atoms) of magnesium turnings and 11.5 g. (0.15 mole) of allyl chloride in 120 ml. of ether. A solution of 19.9 g. (0.10 mole) of trimethyltin chloride in 50 ml. of ether was added. Thus, 14.4 g. (70%) of allytrimethyltin, b.p. 123° C, was obtained.

<u>Allyltriethyltin</u>

The Grignard reagent was prepared from 3.2 g. (0.13 g. atoms) of magnesium turnings and 8.8 g. (0.115 moles) of allyl chloride in 50 ml. of annydrous ether. A solution of 24.5 g. (0.10 mole) of triethyltin chloride in 20 ml. of ether was added. After treatment similar to the preparation of Allyltri-n-propyltin, 9.2 g. (37%) of allytriethyltin, b.p. $71^{\circ}C/10$ mm was obtained.

<u>Allyltri-n-butyltin</u>

The Grignard reagent was prepared from 3.2 g. (0.13 g. atoms) of magnesium turnings and 8.8 g. (0.115 moles) of allyl chloride in 50 ml. of ether. A solution of 32.6 g (0.10 moles) of tri-<u>n</u>-butyltin chloride in 20 ml. of ether was added. Thus, 11.9 g. (36%) of allyltrin-butyltin, b.p. $73-76^{\circ}C/0.09$ mm. was obtained. Allyltriphenyltin (H. Gilman and J. Fisch, J. Org. Chem., 20, 763 (1955)

To 0.1 mole_of allyImagnesium bromide in 190 ml of dry ether was added 27.0 g.(0.0700 mole) of dry powdered triphenyltin chloride by means of a rubber tube connecting a solid vessel of the flask. The addition required 30 minutes and a lively reaction ensured. After stirring for two hours, the reaction mixture was worked up in the usual manner. (See the procedure for diallyIdiphenyltin.) After drying, the ether layer and removing the solvent, there remained a pale yellow oil. This was refluxed with 100 ml. of petroleum ether (b.p. 77-115°) and then refrigerated. Filtration of the ether and concentration of the mother liquor gave a total yield of 24.0 g. (87.5%), melting at 73-75°. The analytical sample was obtained from 95% ethanol as colorless prisms and melted at 73.5-74.5°.

Diallyldimethyltin

The Grignard reagent was prepared from 5.4 g. (0.22 moles) of magnesium turnings and 15.3 g. (0.20 moles) of allyl chloride in 100 ml. of ether. A solution of 20.0 g. (0.09 mole) of dimethyltin dichloride in 10 ml. of T.H.F. and 40 ml. of ether was added. Thus, 7.5 g. (36%) of diallyldimethyltic, b.p. $62-63^{\circ}C/15$ mm. was obtained.

Diallyldiethyltin

The Grignard reagent was prepared from 6.4 g. (0.26 g. atoms) of magnesium turnings and 17.6 g. (0.23 moles) of allyl chloride in

100 ml. of ether. A solution of 24.8 g. (0.10 moles) of diethyltin cichloride in 10 ml. of T.H.F. and 49 ml. of ether was added. Thus, 7.4 g. (28%) of diallyidiethyltin, b.p. $49-52^{\circ}$ C/mm. was obtained.

<u>Ciallyldi-n-propyltin</u>

The Grignard reagent was prepared from 14.6 g. (0.60 g. atoms) of magnesium turnings and 23.0 g. (0.30 moles) of allyl chloride in 230 ml. of ether. A solution of 27.6 g. (0.10 mole) of di-<u>p</u>-propyltin dichloride in 40 ml. of ether was added. Thus, 7.3 g. (25%) of diallyldi-<u>n</u>-propyltin, b.p. $46-48^{\circ}$ C/mm. was obtained.

Diallyldi-n-butyltin

The Grignard reagent was prepared from 14.6 g. (0.60 g. atoms) of magnesium turnings and 23.0 g. (0.30 moles) of allyl chloride in 230 ml. of ether. A solution of 30.4 g. (0.10 mole) of di-n-butyltin dichloride in 50 ml. of ether was added. Thus, 16.4 g. (52% of diallyldi-n-butyltin, b.p. $70-72^{\circ}C/0.1$ mm. was obtained.

Diallyldiphenyltin (H. Gilman and J. Eisch, J. Org. Chem., 20, 763 (1955)

Allylmagnesium bromide was prepared in an 85% yield by the addition of 40.8 g. (0.336 mole) of allyl bromide in 190 ml. of dry ether to 48.9 g. (2.02 g. atoms) of magnesium turnings suspended in dry ether over the course of 2 hours. The reagent was analyzed by titration of an aliquot with standard acid. To 0.10 mole of the allylmagnesium bromide so prepared were added 17.2 g. (0.050 mole) of diphenyltin dichloride in 50 ml. of dry ether over 50 minutes. The addition was accompanied by smooth reflux and the formation of a white, gelatinous precipitate. After 1.5 hours of stirring Color Test I was negative. The reaction mixture then was hydrolyzed by pouring into cold ammonium chloride solution. The separated ether layer was dried over sodium sulfate and the ether was removed. The residue was distilled under a vacuum and a colorless fraction taken at $173-174^{\circ}$ (5.5 mm.), n_{D}^{20} 1.6025. This weighed 12.5 g. and was a 70.3% yield.

Triallylmethyltin

The Grignard reagent was prepared from 37.9 g. (1.56 g. atoms) of magnesium turnings and 82.3 g. (0.68 miles) of allyl bromide in 470 ml. of ether. A solution of 55.1 g. (0.15 mole) of methyltin tribromide in 20 ml. of THF. and 50 ml. of ether was added. Thus, 20.0 g. (52%) of triallylmethyltin, b.p. 100-101/23 mm. was obtained.

<u>Triallylethyltin</u>

The Grignard reagent was prepared from 26.3 g. (1.08 g. atoms) of magnesium turnings and 49.4 g. (0.45 moles) of allyl bromide in 330 ml of ether. A solution of 25.4 g. (0.10 mole) of ethyltin trichloride in 20 ml. of other was added. Thus, 11.9 g (44%) of triallylethyltin, b.p. 60-61/0.6 mm. was obtained.

Triallyl-n-propyitin

The Grignard reagent was prepared from 53.4 g. (2.28 g. atoms)

of magnesium turnings and 72.7 g. (0.95 moles) of allyl chloride in 600 ml. of ether. A solution of 42.3 g. (0.16 moles) of μ -propyltin trichloride in 50 ml. of ether was added. Thus, 29.5 g. (66%) of triallyl- μ -propyltin, b.p. 51-52°C/0.2 mm. was obtained.

Triallyl-n-butyltin

The Grignard reagent was prepared from 21.9 g. (0.90 g. atoms) of magnesium, turnings and 54.5 g. (0.45 moles) of allyl bromide in 300 ml. of ether. A solution of 28.2 g. (0.10 moles) of n-butyltin trichloride in 30 ml. of ether was added. Thus, 19.7 g. (67%) of triallyl-<u>n</u>-butyltin, b.p. $60-61^{\circ}$ C/0.16 mm. was obtained.

Trially]phenyltin

The Grignard reagent was prepared from 29.8 g. (1.22 g. atoms) of magnesium turnings and 61.7 g. (0.51 moles) of allyl bromide in 360 ml. of ether. A solution of 30.2 g. (0.10 moles) of phenyltin tii-chloride in 50 ml. of ether was added. Thus, 15.5 g. (49%) of triallyl-phenyltin, b.p. $80-82^{\circ}C/(07 \text{ mm. was obtained.})$

<u>Tettaallyltin</u>

The Grignard reagent was prepared from 73.0 g. (3.0 g. atoms) of magnesium turnings and 181.4 g. (1.5 moles) of allyl bromide in 1 liter of ether. A solution of 180.4 g. (0.5 mole) of diallyltin dibromide (K. Sisido and Y. Takeda, J. Org. Chem., 26, 2301 (1961) in 100 ml of ether was added. Thus, 115 g. (80%) of tetraallyltin, b.p. $66-68^{\circ}C/0.04$ mm. was obtained.

Table II

The Physical Properties and Analyses of Allylalkyltin Compounds

ş

| Compound | Yield.X | b.p. | | היזה- מיזרב- | 2H fcund | calc'd | \$S found | n calc'd |
|--------------------------------|---------|---------------------------|----------------|-----------------|-------------|--------------|---------------|----------------|
| All.Sn | 80 | 50-51 _{/0-} 1 | 51.04 | 50.93 | 7.23 | 7.12 | 42.6 | 41.94 |
| 4 | | | | | | | | |
| All ₂ EtSn | 44 | 60-61 _{/0.6} | 48.52 | i48•75 | 7.37 | 7.444 | 44.0 | 43.80 |
| All ₃ PrSn | | 51-52/0.2 | 20°5 20°5 | 50.57 50.57 | 7.94 | 7.78 8.09 | 112.0 39.8 | 41.05 39.69 |
| All ₃ BuSn | . 49 | 60-61/0.15 80-82 /0.17 | در.1ر 56.28 | 56 .48 | 6.37 | 6.32 | 37.5 | 37.20 |
| 3 | УÇ | 62-63 | 41.49 | 41.61 | 7.04 | 6.99 | 51.7 | 51.40 |
| AII2 ^{me} 25 n | 2 8 | ¢1/ | 46.22 | 46.36 | 7.70 | 1.79 | 45.7 | 45.85 |
| A11264231 | 25 | 46-48 / 1 | 50.50 | 50.22 | 8.56 | 8.43 | 41.5 | 41.35 |
| All2r23 All2Eu2Snb | | 70-72/0.1 | 53.58 | 53.37 | 9.09 | 8.96 | 37.2 | 37.67 |
| | | 123 | 35_53 | 35.19 | 7.06 | 6.89 | 58.0 | 57.93 |
| All'te ₃ 5n | 27 | 71 | 43.96 | 43.77 | 8.36 | 8.16 | 48.6 | 48.06 |
| Allet35n | 5 | 114-115 /13 | 50.11 | 49.87 | 9.26 | 9.07 | 41.1 | 41.06 |
| AllBu ₃ Sn | 36 | 73-76/0.09 | 54.48 | 54.41 | 09.6 | 9.74 | 36.4 | 35.85 |

27

,

۰.

:

TABLE III

THE UV SPECTRAL DATA OF ALLYLTIN COMPOUNDS

| | n-Ce | H14 | _ CH3(| CN |
|------------------------|-----------|------------------------|-------------|------------------------|
| Compounds | λmax (mµ) | emax XIO ⁻⁴ | λmax (mµ) · | emax XI0 ⁻⁴ |
| All4Sn | 217 | 3.16 | 220 | 3.14 |
| All ₃ MeSn | 211 | 2.49 | 217 | 2,50 |
| Et | 210 | 2.77 | 217 | . 2.96 |
| Pr | . 208 | 2.98 | 215 | 2,88 |
| Bu | 209 | 3.05 | 214 | 2.74 |
| All2Me2Sn | 208 | 1.99 | 212 | 1.83 |
| Et | 208 | 1.94 | 212 | 1.99 |
| Pr | 208 | 1.89 | 211 | 1.80 |
| Bu | 203 | 2.00 | 212 | 2.03 |
| AT IMa ₃ Sn | <210 | *** ** | 210 | 1.07 |
| Et | <200 | * ** | 203 | 1.29 |
| Pr | . <200 | | 204 | 1.12 |
| Bu | <200 | 4 0 6 0 | · <200 | |

representation and the second and the second s

B. PREPARATION OF SUBFONES

Phouvi Propartyl Sulfone .

Following the procedure of Overberger, 44.5 g. (0.4 mole) of phunyl propargyl sulfide in 300 ml of glacial acatic acid was cooled to 0° and 120 ml of 30% hydrogen peroxide was added during one hour. The solution was then heated at 80-90° for several hours and allowed to stand overnight. The solid sulfone was removed by filtration and washod with alcohol and water. Recrystal lization from absolute ethanol obtained 43 g. (60%) of phenyl propargyl sulfone, mp 89-89.5°. Anal. Calco. for CallaSO2: C. 59.99; H,4.48

Found: C, 59.64; H, 4.71

Phényl Grótyl Sulfone (A.C. Cope, D.E. Morrison and L. Field; J. Am. Chom. Soc., <u>72</u>, 60 (1950))

Phonyl crojyl sulfone, bp 98° (0.05 mm), was prepared in 40% yield as described by two method of Cope. A mixture of 51 g. (0.3 mole) of phonyl crofyl sulfide in 200 ml of glacial acetic acid was cooled to 0° in ice-bath and 100 ml of 30% hydrogen peroxide was added during one hour. The solution was then heated at 70-80° for several hours, allowed to stand overhight, diduted with an equal volume of water and extracted with 200 ml of chloroform. The extract was washed with water, dried over anhydrous magnesium sulfate, and the solvent removed under reduced pressure. The residue was vacuum distilled and 23.5 g. (40%) of phenylcrotyl sulfone was obtained. Phonyl Ally Sulfone (A.C. Cope, D.E. Morrison and L. Field, loc. cit.)

Phenyl allyl sulfone, bp 86-88° (0.95 mm) was prepared in 45% yield from phenyl allyl sulfide and 30% hydrogen peroxide in glacial acetic acid as described above for the preparation of phenyl crotyl sulfone.

Ethyl Propargyl Sulfone

Ethyl propargyl sulfone, bp 63-65° (0.05 mm) was prepared in: 30-40% yield from ethyl propargyl sulfille and 30% hydrogen peroxide in glacial acetic acid as described above for the preparation of phenyl crotyl sulfone. Carbon hydrogen-analysis-are in progress.

Ethyl Crotyl Sulfone

A SUBLEY C. I DESCRIPTION

Étnyl crótyl sulfone, op 54° (0.9 mm) was prepared in 30-40% viela from étnyl crotyl sulfide and 30% hydrogen peroxide in glačiák acutic acid as described above for the preparation of phonyl crotyl sulfone.

| • • • • • • • • • • • • • • • • • • • | * 2012/07/2000/07 ** *** | uuuu 'i'i'a' au | .j. | 1 | | | ((*)) ((((((((((((((((| • • | and a poleo. | | • | 31, | li. | | |
|---------------------------------------|--------------------------|----------------------|----------------------------------|------------------|---------------------|------------------------|--|-----|--------------|---|-------|----------|--------|------------------|----|
| | • | · | Found Calc'd | 4.71 4.48 | in Progress | | | - | | | | e) () | , , | | |
| چند بی الجند | | es of Sulfones | KC Found Calle ¹ d | 59°64 59°99 | Analyses | · | | | • | | • | | • | • | |
| | Table IV | coporties and Analys | bp/mp | 89-89.5 | 63-65° /0.05 | 98 <mark>0/0.05</mark> | 86-88 <mark>0%05</mark> | | | • | | | | ب مربع مربع ۲ | |
| | , | The Physical P | Yield % | 60 | 30-40 | 40 | 45 | | | | | | | | 5. |
| · | | | Compounds | Phenyl propargyl | Ethyl propargyl | Phenyl crotyl | Phenyl allyl | 2. | | | · | • | • | • • • | |

•

`1`

•

. د •,

ý. F

0

ς.

, <u>t</u> . $\langle \cdot \rangle_{\mathcal{C}}$

Attempted preparation of debenzonesulfonyltriethyltin. - The method used for the metalation of sulfones with <u>n</u>-butyliithium in benzeno has been described by Truce and Buser? In a 500-mi three-necked round bottomed flask, fitted with a condenser, a dropping funnel and a mechanical stirrer, was placed 6 g.(0.03 mole) of phenyl allyl sulfone and 100 mi of anhydrous benzene. <u>n</u>-Butyliithium in hexane (1.9 g., 0.03 mole) was added slowly to the mixture with cooling, while a stream of hitrogen was passed through the system. The mixture was then refluxed approximately three hours and allowed to stir at room temperature for one hour. Triethyltin chloride (7.2 g., 0.03 mole) was added to the mixture and again if was refluxed for 72 hours. Filtration followed by vacuum distillation, led to recovery of approximately 60% of triethyltin chloride and starting sulfone.

Attempted addition of triethyltin hydrida to athyl sulfone

by photolysis. - A two-nocked Pyrex tube was evocuated; heated, flushed with argon and cooled. Ethyl propargyl sulfond (4 g., 0.03 mole), 10 ml of anhydrous ether, and triethyltin hydride (7.2 g., 0.03 mole) were introduced under a stream of argon and stoppered. The Pyrex tube was then irradiated with 100 W medium pressure mercury lamp at 25-30° for 216 hours. After removal of diethyl ether, distillation yielded only unreacted statting materials.

NEGRETALICAL DECOMPOSIT

たちたいとうである

l'anna

sulfone by thermo-method (AlBN). ~ A 250 ml three-necked flask was

Attempted addition of triethyltin hydride to ethyl propargy

evacuated, heated and filled with argon. Ethyl propargyl sulfone (4 g., 0.03 mole), 50 ml of anhydrous benzene, and 7.2 g. (0.03 mole) of triethyltin chloride were introduced to the flask. The mixture was removed under reduced pressure. Distillation yielded unreacted triethyltin chloride and ethyl propargyl sulfone, which were characterized by I.R. spectra and g.l.p.c. retention times.

Note of the second

C. PREPARATION OF ALLENYLTINS

Preparation of Triethylallenyltin

In a 3-necked 500 ml. round-bottomed flask fitted with a stirrer, reflux condenser and dropping funnel were placed magnesium (2.4g.; 0.1 mole) and dry ether (70 ml.). Propargyl bromide (11.9g., 0.1 mole) in ether (15 ml.) was added slowly with stirring: the reaction was initiated by iodine. After all the propargyl bromide had been added, the mixture was stirred for two hours, then cooled to -60° . Triethyltin chloride (14.5 g., 0.06 mole) was added, and the mixture stirred and allowed to warm to -15° . Saturated ammonium chloride solution was added slowly to hydrolyse the mixture, which was then allowed to warm to room temperature; the ether layer was separated and dried over magnesium sulphate, which was then filtered off. The ether was removed and the remaining pale yellow oil distilled, the major fraction coming over at $55-56^{\circ}/$ 1.5 rm.

The coloriess distillate was shown by L.R. to contain allenyltin but no propargyltin and by G.L.C. to consist of only one component.

This procedure was used for the other allenyl and propargyltins. <u>Preparation of 3-Chloro-1-phenyl-1-butyne</u>

A solution of 20.0g. (0.137 mole) of 1-phenyl-1-butyn-3-ol in 25 ml. anhydrous ether and a solution of 23.8g. (0.20 mole) of thionyl chloride in 25 ml. anhydrous ether were added simultaneously, dropwise, to 250 ml. of refluxing ether containing 20.7g. (0.15 mole) of potassium carbonate. After the addition, the reaction mixture was allowed to

TABLE ፗ

Allenyl and Propargyltins

| • • • | Compound or Mixture | 5'Yield % | VCEC or VC=C=C (cm ⁻¹) | m.p. °C | b.p. °C/mm. |
|-------------|---|---------------------|---------------------------------------|------------|----------------|
| | (Me ₃ SnCH=C=CH ₂) | | 2110 s | | 133-136 |
| | (Me3SnC=C-CH3) | 12 | 2165 m | | . . |
| | Et ₃ SnCH=C=CH ₂ | . 54 | 1928 s | Y | 55-58/1.5 |
| | Pr ₃ SnCH=C=CH ₂ | 46 | 1928 s | • | 86-88/1.5 |
| •. | Bu ₃ SnCH=C=CH ₂ | 75 | 1928 s | | 112/1.25 |
| | Ph₃SnCH=C∺GH₂ | 95 | 1928 s | 61-62 | |
| | Ph₃SnCH₂-C≡CH | 90 | 2110 w | 81.5-83.5 | |
| | $\left(Bu_2 Sn(CH=C=CH_2)_2 \right)$ | | 1928 s | | |
| | Bu2Sn(CH2CECH) | 62 | 2105 m | | 100/1.5 |
| | Bu2Sn(C≡CCH3)2 | | 2163 m | • | |
| | Ph3Sn(Me)C=C=CHMe | 25 | | 47-48 | |
| | Me ₃ Sn(Ph)C=C=CHMe | 40 | | | 101-103/0.25 |

ŀ

í

reflux for two hours, filtered and the filtrate washed with 50 ml. of water, five 50 ml. portions of 5% aq. sodium bicarbonate solution and finally with 50 ml. of water. The ether solution was dried over magnesium sulfate and distilled to give 3-chloro-1-pheny1-1-butyne (13.1g., 58%) at 75-77°/0.25 mm. (authentic infrared spectrum).

Preparation of 2-Bromo-3-pentyne

A mixture of 3-pentyn-2-ol (100 g., 1.19 moles), dry pyridine (20 ml) and dry ether (500 ml) was placed in a flask equipped with a dropping funnel, stirrer and a reflux condenser with a CaCl₂ guard tube. Phosphorus tribromide (150 g., 0.56 mole) was added dropwise to above with stirring and cooling (0°). After an initial period of five minutes, the cooling bath was removed and the ether was allowed to reflux while the rest of the phosphorus tribromide was added (30 minutes). The mixture was then heated at 55° with stirring for three hours, after which it was cooled (0°) and ice was slowly added to hydrolyze excess phosphorus tribromide. The ether layer was washed with saturated NaHCO₃ solution and saturated NaCl solution, dried over MgSo4 and then distilled. The bromide (119 g., 68%) was obtained at 41-44°/20 mm.

Preparation of Penta-2,3-dion-2-yltrimethyltin

2-Bromo-3-pentyne (14.7 g., 1 mole) in dry ether (100 ml) was added very slowly (ten hours) to activated magnesium turnings (3.0 g.) in refluxing ether (50 ml). The r-sulting Grignard compound was slightly dark in color. Trimethyltin chloride (19.8 g., 1 mole) in ether (40 ml) was then added to it and the mixture was refluxed for two hours. It was hydrolyzed by saturated ammonium chloride (just enough to precipitate the magnesium salts as a solid crust) and the etherial layer was treated with acqueous pofassium fluoride solution. The white precipitate obtained was filtered off, and the ether layer was dried over MgSo4. Distillation afforded the allenyltin compound (5.8 g., 25%) as a colorless liquid, b.p. 46-47°/15 mm.

Preparation of Penta-2,3-dien-2-yl tributyltin

This compound (10.7 g., 30%) was obtained as a colorless liquid, b.p. 85°/0.02 mm, by the same procedure as described in the previous experiment.

Preparation of Penta-2, 3-dien-2-yltriphenyltin

Triphenyltin chloride (37.2g., 1 mole) in ether (50 ml.) was added to the Grignard reagent, obtained from 2-bromo-3-pentyne (14.7g., 1 mole) and magnesium (2.5 g.). The mixture was refluxed for two hours and hydrolysed by saturated ammonium chloride solution. The crude product obtained from the etherial layer was crystallized from methanol; m.p. $47-48^\circ$ (yield 25%).

Preparation of 1-Phenyl-1,2-dien-1-yltrimethyltin

This was obtained as a colorless liquid, b.p. 101-103°/0.25 mm. in 40% yield from 3-chloro-1-phenyl-1-butyne and trimethyltin chloride by the same method as in the previous experiment.

D. PREPARATION OF BI- AND TRI-CYCLIC ORGANOTINS

General----- -- Trimethyltin

hydride and tributyltin hydride were prepared as reported earlier⁶ 3-Bromonortricyclene was obtained from norbornylene and N-bromosuccinimide as described in the literature? Norbornadiene was distilled before use.

Photolysis was carried out in sealed pyrex or quartz ampules. A 100 w. mercury lamp was used as the light source. All experiments were carried out in argon atmosphere.

Interaction between Norbornadiene and Trimethyltin Hydride.

A. Equimolar Ratio. - A mixture of trimethyltin hydride (3.2 g.) and norbornadiene (1.8 g.) was sealed in a pyrex ampoule and was photolyzed for six hours at 55°. The ampoule was broken and the product was distilled to give a colorless liquid (mixture A) at 35°/0.02 mm. (4.5 g., 90%). The infrared spectrum was identical to that obtained by Mr. Pelczar.

> (1) The above experiment was repeated in a quartz tube. The photolysis was carried out for 27-1/2 hours. Identical results were obtained. The mixtures obtained in both

30

(2) The mixture A (3.8 g.) from the above experiment was sealed in a quartz tube and photolyzed for eight days at 70°. The tube was broken and the mixture redistilled.

infrared spectrum and g.l.p.c. showed that the mixture remained unchanged.

B. With excess Trimethyltin Hydride.

(1) A mixture of norbornadiene (1.36 g.) and trimethyltin hydride (6.5 g.) was heated under reflux (bath temp. 90°) with stirring for 17 hours. Excess trimethyltin hydride was removed at 20 mm. The product on distillation gave a small forerun (0.22 g., 5.8%) at $32^{\circ}/0.02$ mm and, finally, the main fraction (4.0 g., 65%) at $92^{\circ}/0.02$ mm, leaving a small residue.

Anal. Calc. for $C_{13}H_{40}Sn_2$: C, 37.02; H, 6.64. Found: C, 37.10; H, 6.60.

(2) A mixture of norbornadiene (1.8 g.) and trimethyltin. hydride (8.0 g.) was scaled in a pyrex ampoule and photolyzed for 30 hours at 55°. Excess trimethyltin hydride was removed at 20 mm. Distillation afforded a smail fraction (0.8 g.) at 35°/0.02 mm. Mixture B (6.8 g.) was finally obtained at 82°/0.02 mm.

(3) Similar results were obtained when the above experiment was repeated in a quartz tube.

Interaction between Mixture A and Trimethyltin Hydride.

•

(1) Trimethyltin hydride (3.5 g; 1.48 mole) was added -21mixture A (3.66 g; 1 mole) in a pyrex ampoule which was sealed in argon atmosphere and photolyzed for 30 hours at 65°. After removing excess trimethyltin hydride and distilling the product, a forerun (0.4 g.) was obtained at 35°/0.02 mm and then mixture B (3.90 g.) distilled at 82°/0.02 mm.

(2) The above experiment was repeated twice, the photolysis being carried out for ten hours and two hours, respectively. Exactly similar results were obtained. (3) Trimethyltin hydride (1.26 g., 0.5 mole) and mixture A (3.74 g.) were mixed in a pyrex ampoule and photolyzed for two hours at 65°. After the usual workup, the product was distilled to give first fraction (0.52 g.) at 35°/0.02 mm and the second fraction (3.06 g.) at 82°/0.02 mm. leaving a small residue. First fraction was unreacted

Interaction between Mixture A and Tributyltin Hydride. - Tributyltin hydride (5.7 g., 1 mole) was added to mixture A (5.0 g.) in a pyrex tube which was sealed and photolyzed for six hours at 65°. Distillation of the product gave a first fraction (0.32 g.; b.p. 35-45°/ 0.05 mm.), a second fraction (2.39 g., b.p. 45-52°/0.05 mm.) and finally a third fraction (mixture C; 6.43 g., 52%) at 140-2°/0.05 mm.

mixture A with high ratio of Compounds 1 and 1V.

Anal. Calc. for C₂₂H₅₂Sn₂: C, 48.23; H, 8.40. Found: C, 48.26; H, 8.44.

G.I.p.c. showed that fractions one and two were unreacted mixture A and tributyltin hydride. These were combined and photolyzed again in a pyrex tube for 30 hours at 65°. Distillation gave a small fraction (0.4 g.) at $35^{\circ}/0.02$ mm. and mixture C (1.1 g.) at $135^{\circ}/0.02$ mm.

-22-

Preparation of Nortricyclyl trimethyltin (Compound 1). -

3-Bromonortricyclene (14.4 g.) was added slowly to magnesium turnings (2.22 g.) in dry ether (100 ml.) with stirring. The resulting Grignard compound was refluxed for two hours. It was then cooled to 0° and a solution of trimethyltin chloride (16.6 g.) in 50 ml. dry ether was added to it in two hours. The mixture was again refluxed for two hours, 'cooled to 0° and then hydrolyzed by saturated ammonium chloride solution. The etherial layer was separated, washed with water, dried over magnesium sulphate and distilled. Nortricyclic trimethyltin (13.2 g., 62%) was obtained as a coloriess liquid at 35°/0.02 mm.

<u>Attempted Reaction between Nortricyclyl Trimethyltin and Tri-</u> <u>methyltin Hydride</u>. - A mixture of nortricyclictrimethyltin (4.5 g.) and trimethyltin hydride (5.0 g.) was photolyzed in a sealed pyrex tube for 18 hours at 65°. After removing unreacted trimethyltin hydride, the residual liquid was found to be unchanged nortricyclic trimethyltin contaminated with a little hexamethylditin.

Interaction between Norbornadione and Tributyltin Hydride. -A mixture of norbornadiene (1.8 g., 1 mole) and tributyltin hydride (5.8 g., 1 mole) was photolyzed in a pyrex ampoule for 18 hours at 65°. Distillation of the product yielded a mixture (D) of monoadducts (6.5 g., 85%) at 102°/0.02 mm.

Anal. Calc. for C19H36Sn: C, 59.58; H, S.40. Found: C, 59.49; H, 9.37.

Interaction between Mixture (D) and Trimethyltin Hydride. - Trimethyltin hydride (3.8 g., 1.5 mole) was added to mixture (D) (5.8 g., 1 mole) in a pyrex tube. The photolysis was carried out for 30 hours at 65°. Distillation afforded four fractions having b.p. 115-125°/0.02 mm; 125-130°/0.02 mm; 130-135°/0.02 mm and 135°/0.02 mm. G.I.p.c. showed

that the last fraction (3.1 g., $37\frac{1}{2}$) was a mixture (E) of the bisadducts whereas the first three fractions were mixtures of unreacted mixture D and the product.

Interaction between Mixture (B) and Trifluoroacetic Acid. -Trifluoroacetic acid (2.5 g., 6 mole) was added to mixture (B) (1.42 g., I mole) in dry pentane (50 ml.). The mixture was stirred at room temperature for four days. Precipitation of a white solid started after 7-8 hours and gradually increased with slow evolution of methane. The white ppt. was filtered off and washed with more pentane and dried at 0.02 mm. The infrared spectrum shows that all methyl groups have not been cleared.

Preparation of 2(or 3)-Trimethylstannyl-5(or 6)-thiophenylvicycloheptane

Reduiros: C, 52.33; H, 6.54; S, 8.72; Sn 32.36.

Proparation of 2(or 3)-Trimethylstannyl-5(or 6)trichloromethyl-6(or 5)chlorobicycloheotane

Carbon tetrachloride (4.60g., 1.5 mole) was added to the mixture of trimethylstannylbicycloheptene (5.1g., 1 mole) in a pyrex tube which was sealed and photolysed for six hours. Distillation under reduced pressure afforded the addition compound as a colorless liquid (6.5g., 80%) b.p. 113-14°/0.02 mm.

Found: C, 32.3; H, 4.40; Cl, 34.37; Sn, 27.98%. C₁₁H₁₈SnCl₄ Requires: C, 32.14; H, 4.38; Cl, 34.57; Sn, 28.89%

E. NITROGEN-CONTAINING ORGANOTINS

Preparation of 3-bromoquinoline. - In a three-necked flask, fitted with a condenser, a mechanical stirrer and a dropping funnel was heated, evacuated and filled with nitrogen gas. Redistilled quinoline, 39 g. (0.3 mole) and 320 ml. of anhydrous carbon tetrachloride were placed in the flask, 48 g. (0.3 mole) of Bromine was added slowly to the mixture. The mixture was heated for reflux about one hour thereafter, a solution of 23.7 g. (0.3 mole) of anhydrous pyridine in 20 ml. of carbon tetrachloride was added slowly to the refluxing mixture over a period of two hours. During the terminal 18 hours reflux period, the original orange suspension became tan-colored. Filtration of the cooled suspension, removal of solvent from the filtrate and fractional distillation afforded 36 g. (60% yield) of 3-bromoguinoline, b.p. 94°/2.25 mm.

Preparation of trialkyltin quinoline. - In a 3-necked roundbottomed flask, fitted with a condenser, a mechanical stirrer and a dropping funnel was evacuated and filled with nitrogen gas. n-Butyllithium (1.6 g. 0.025 mole) in hexane was placed in the flask together with 100 ml. of anhydrous ether. The flask was cooled by means of dry-ice-acetone bath to -50°. 3-Bromoquinoline (5.2 g. 0.025 mole) was introduced very quickly to the mixture. The color of the mixture became dark red. Color test for quinoline lithium was positive after fifteen minutes. Equivalent amount of trialkyltin chloride was added to the mixture; it turned to a yellow color after twenty minutes. The mixture was hydrolyzed with saturated ammonium chloride solution after several hours stirring. The ether layer was dried; after removal of ether, the liquid became viscous. Preparation of 2-chloro-4-methyl quinoline. - In a 3-necked round-bottomed flask, fitted with a condenser, a mechanical stirrer and a dropping funnel was placed 29.4 g. (0.2 mole) of 2-hydroxy-4-methyl quinoline. About 60 ml. of phosphorous oxychloride was added slowly to the flask with cooling. After refluxing for twelve hours, the mixture was hydrolyzed with cold water. It was then extracted with convenient amounts of ether and dried over anhydrous magnesium sulfate. Removal of ether yielded 65% of corresponding 2-chloro-4-methyl quinoline, m.p. 53-54°.

Preparation of 2-triethyltin-4-methyl quinoline. - The same procedure as previously described for the preparation of 3-trialkyltin quinoline was employed.

Preparation of 3-(tributyistannyloxy) quinine. - Quinine (3.2 g., 0.01 mole) and bistributyitin oxide (3 g., 0.005 mole) in toluene (60 ml.) were heated under reflux for 1.5 hrs., with a Dean and Stark separator to remove water. The toluene was removed by pumping at a rotary pump, leaving an extremely viscous oil, the infrared spectrum of which showed no -OH band at 3120 cm⁻¹. On exposing the infrared plates to air for two minutes, the oil became a semi-solid, which the infrared spectrum showed to consist largely of quinine and bistributyitin carbonate.

Addition of acceptors to the alkoxide.

(a) <u>Phenyl isocyanate</u>. - To .01 mole of the tin alkoxide dissolved in carbon tetrachloride (10 ml.) was added phenyl isocyanate (1.19 g., .01 mole). The mixture was stirred rapidly for 30 minutes, heat being evolved on initial mixing. Volatile components were removed at the pump, leaving a viscous oil; the latter was readily hydrolyzed by atmospheric moisture, and on ethanolysis, followed by treatment with pentane, gave a highly viscous oil containing no tributyltin compounds. This oil was sublimed in vacuo, yielding a white powder, melting slowly over the range 70-110°, v_{max} 1730s (C=O) cm⁻¹.

(b) <u>Chloral</u>. - To .01 mole of the tin alkonide, dissolved in carbon tetrachloride (10 ml.) was added chloral (1.47 g., .01 mole). Immediately polychloral was formed, but on stirring for several hours dissolved. Volatile components were removed, leaving a readily hydrolyzed highly viscous oil. On methanolysis this yielded a white solid, m.p. 137-147°, the infrared spectrum of which showed no -OH bands.

<u>Preparation of N¹-tributyIstannyIsulfanilamide</u>. - Sulfanilamide (3.4 g., .02 mole) and bistributyItin oxide (5.96 g., .01 mole) in nylene (70 ml.) were heated under reflux for 2.5 hours with a Dean and Stark separator to remove water. The solvent was removed at a rotary pump, leaving a white crystalline solid. This was further purified by washing rapidly in air with dry pentane. An attempt to recrystallize a sample from toluene in air, taking no precautions to exclude atmospheric moisture, resulted in hydrolysis of the sample.

Preparation of Alkylpuridyltins. - The method of preparation of all of these compounds is essentially the same; tributyl 3-pryidyltin will be guoted as an example.

To 3-bromopyridine (3.16 g., .02 mole) in ether (25 ml.) cooled to -40° , was added a solution of n-butyl lithium in hexane (13.4 ml. of 1.6 M). A brown precipitate was thrown down. Tributyltin chloride (6.5 g., .02 mole) was added, and the mixture stirred for two hours, during which time it was allowed to warm to room temperature and the brown color disappeared.

A saturated solution of ammonium chloride (40 ml.) was added; the ether layer separated and dried over magnesium sulfate. The ether was removed, leaving a pale orange oil, shown by G.L.C. to consist of two components, the minor one being tributyltin chloride. Distillation gave a pale orange oil, b.p. 106.5-108°/.075 mm; the G.L.C. of this indicated it to be 97% pure tributyl 3-pyridyltin.

Some of the other pyridyltins were obtained as brown oils; the brown color was removed by passing an ether solution through a short alumina column.

Table VI-

ANALYTICAL DATA ON PYRIDYLTINS

| | ۶ | Ċ | % | H | ۶ | N |
|---|--------------|--------------|------------|------------|--------------|-------|
| | calc. | found | calc. | found | calc. | found |
| tri- <u>n</u> =butyl=3-pyridyltin | 55•և | 55•3 | 8.4 | 8.3 | 3.8 | 3.4 |
| di- <u>n</u> -butylbis-(3-pyridyl)tin | 55.5 | 55.6 | 6.7 | 6.9 | 7.2 | 7.2 |
| tri- <u>n</u> -propyl-3-pyridyltin | 51.5 | 51.4 | 7.8 | 7.8 | 4.3 | 4.4 |
| di- <u>n</u> -butvlbis-(2-pyridyl)tin | 55•5 | 55•7 | 6.7 | 7.0 | 7.2 | 7.0 |
| tri- <u>n</u> -propyl-3-pyridyltin di- <u>n</u> -butvlbis-(2-pyridyl)tin | 51.5 55.5 | 51.4 55.7 | 7.8 6.7 | 7.8 7.0 | 4•3 . 7•2 | |

a All isolated as oils.

he a have a starter to

Preparation of Dimethyltin Dichloride - Bissalicylidendiamino-

ethane Adduct. - Dimethyltin dichloride was stirred and dissolved in benzene to which bissalicylidenediaminoethane in benzene was added. Yellow precipitates appeared immediately. They were filtered off and had m.p. of 149 to 153 C. Recrystallization from benzene-heptane mixture raised the m.p. to 153 to 154 C. The crystalls were dried under reduced pressure at 80 C. Found: C, 44.52; H, 4.59; N, 5.83; Cl, 14.70; Sn, 24.6

Calcd. for C18H22N2O2Cl2Sn

C, 44.30; H, 4.55; N, 5.74; Cl, 14.53; Sn, 24.32

Diethyltin dichloride - Bissalicylidenediaminoethane Adduct. -Diethyltin dichloride (2.48 g., 0.01 mole) in 10 ml of benzene was added to bissalicylidenediaminoethane (2.68 g., 0.01 mole) in 30 ml of benzene with stirring. Yellow precipitates came out immediately. They were dissolved by adding 60 ml of benzene and 50 ml of n-heptane. The solution was kept to stand overnight to yield yellow crystals, m.p. 161.5 -162.5 C.

Found: C, 46.77; H, 5.15; N, 5.42; Cl, 13.64; Sn, 22.9 Calcd. for $C_{20}H_{26}N_2O_2Cl_2Sn$

C, 46.55; H, 5.08; N, 5.43; CI, 13.74; Sn, 23.00

<u>Di-n-propyltin dichloride - Bissalicylidenediaminoethane</u> <u>Adduct. - Di-n-propyltin dichloride (2.76 g., 0.01 mole) in 10 mi of</u> benzene was added to the ligand (2.68 g., 0.01 mole) in 30 ml of benzene with stirring. Yellow precipitates came out immediately which were recrystallized from 90 ml of benzene and 50 ml of n-heptane to yield yellow crystals, m.p. 109-110 C.

Found: C, 48.81; H, 5.62; N, 5.15; Cl, 13.29; Sn, 21.4 Calcd. for C₂₂H₃₀il₂O₂Cl₂Sn

C, 48.57; H, 5.56; N, 5.15; CI, 13.03; Sn, 21.81

Ethyltin Trichloride - Bissalicylidenediaminoethane Adduct. -Ethyltin trichloride (2.54 g., 0.01 mole) in 10 mi of benzene was added to the ligand (2.68 g., 0.01 mole) in 30 ml of benzene with stirring. Yellow precipitates came out immediately. The solubility of this complex was so low that recrystallization was not tried. The precipitates were filtered off and washed with n-pentane several times to give yellow powder, m.p. 191-3° C.

Found: C, 41.24; H. 4.09; N, 5.43; Cl, 20.21; Sn, 22.6 Calcd. for $C_{18}\overline{H_{21}}N_2O_2CI_3Sn$

C, 41.38; H, 4.05; N, 5.36; C1, 20.36; Sn, 22.72

Preparation of Di-n-buty (bissaly cilethylenedilminato) tin.

Bu₂SnO (0.023 mole, 5.7 g.) and BSDE2H (0.02 mole, 5.4 g.) were mixed in 100 ml of toluene and heated to reflux for five hours, filtered when hot and concentrated to give 9.7 g. of yellow precipitates, m.p. 100-200 C. 7.2 g. of these precipitates was passed through a column using neutral alumina and T.H.F. The solid left when the solvent was evaporated was recrystallized from benzene-heptane mixture to give 3.8 g. of yellow needle crystals, m.p. 80-200 C. The n.m.r. spectra showed fairly good intensity ratio of our kinds of protons. 3.8 g. of crystals were fractionally recrystallized several times and ended up with less than I g. of yellow needle crystals, m.p. 152-154 C.

Calcd. for $C_{24}H_{32}N_2O_2Sn$

C, 57.75; H, 6.48; N, 5.61; Sn, 23.77

DISTRIBUTION LIST

- 20 copies Defense documentation Center Attn: DDCIR Cameron Station Alexandria, Virginia 22314
- l copy U. S. Army Combat Development Agency Medical Service Agency Brooke Army Mdical Center Fort Sam Houston, Texas 78234

4 Copies Commanding General U. S. Army Medical Research and Development Command Office of the Surgeon General Attn : MEDDH-SI Washington, D. C. 2031).

16 covies Director

Walter Reed Army Institute of Research Walter Reed Army Medical Centor Attn: Division of Medicinal Chemistry Washington, D. C. 20012

| Creating characterises and in the base of instants assured as and and and the de array report is called and and and and and and and and and an | Ceremp damingene of this, bed of sharest and indense security most is a carlied when the secure report is cardingly of New York allowed when the secure report is cardingly of New York allowed when the secure report is cardingly of New York allowed when the secure report is cardingly of New York allowed when the secure report is cardingly of New York allowed when the secure report is cardingly of New York allowed when the secure report is cardingly of New York allowed when the secure report is cardingly of New York allowed when the secure report is cardingly of New York allowed when the secure report is a secure r | | DOCUMENT CONT | ROL DATA - R | & D | |
|--|--|---|--|---|--|--|
| L. CARENT HE ACTIVITY CORPORE Setting of New York Andread Setting of New York Andread Research Second Research Second Setting Print Research Second Secon | One Definition of the State University of New York Definition of the State University of New York Albary, New York 12221 Definition of the State University of New York Definition of the State University of New York Albary, New York 12221 Definition of the State University of New York Definition of the State University of New York Albary, New York 12221 Definition of the State University of New York Definition of the State University of New York ORGANOTING FOR ANTIMALARIAL SCREENING Definition of the State University of New York Definition of New York Organization of the State University of New York January 1973 Definition of New York Aduption of State University of New York Definition of New York Definition of New York Aduption of State University of The State University of New York Definition of New York Definition of New York Aduption of State University of The Sta | (Security classification of title, bo | dy of abstract and indexing | annotation must be | enfered when the | overall report is classified) |
| Albany, New York 12221 Albany, New York 12221 ORGANOTINS FOR ANTIMALARIAL SCREENING DEGAMOTINS FOR ANTIMALARIAL SCREENING description of report and inclusive dates Final report covering period 15 June 1966- 31 October 1968 description of the covering period 15 June 1966- 31 October 1968 description of the covering period 15 June 1966- 31 October 1968 description of the covering period 15 June 1966- 31 October 1968 description of the covering period 15 June 1966- 31 October 1968 description of the covering period 15 June 1966- 31 October 1968 description of the second of the covering period 15 June 1966- 31 October 1968 description of the second of the seco | Albany, New York 1222). AREONY TILE ORANOTINS FOR ANTIMALARIAL SCREENING THE report covering period 15 June 1966- 31 October 1968 ANTHON (MUTIA) FOR ANTIMALARIAL SCREENING THE report covering period 15 June 1966- 31 October 1968 ANTHON (MUTIA) Henry C. Kuivila Report no.7 A CONTRACT OF SAME THE ALA-19-135-MD-3018 A CONTRACT OF SAME THE A CONTRACT OF SAME THE SA | The Research Toundation | or) of the State Univ | ersitu | 24. REPORT SE | CURITY CLASSIFICATION |
| Albany, New York 12224 - REFORMENT ORGANOTING FOR ANTIMALARIAL SCREENING - DECEMBENT - DECEM | Albany, New York 12224 Abany, New York 12224 ABANY, New York 12224 AREANY UNCENTRY FOR ANTIMALARIAL SCREENING Description of the contract of the series | of New York | | erervy | | · |
| A SECONT THE ORGANOTINS FOR ANTIMALARIAL SCREENING Support notes (Type of report and Machadin adms) Final report accounting period 15 June 1966-31 October 1966 Support account adms, adde should be adms) Henry C. Kuivila A Secont Total report account adms adms adms A should be administrate adms) Henry C. Kuivila A Secont Total administrate administration adms) Henry C. Kuivila A Secont Total administrate administration administra | AREANT NIE ORGANOTINS FOR ANTIMALARIAL SCREENING | Albany, New York 1222 | | | ZO. GROOP | |
| ORGANOTING FOR ANTIMALARIAL SCREENING | ORGANOTINS FOR ANTIMALARIAL SCREENING | . REPORT TITLE | | | | |
| ORGANCTINS FOR ANTIMALARIAL SCREENING | ORGANOTINS FOR ANTIMALARIAL SCREENING Instant report covering period 15 June 1966- 31 October 1968 Instant on the state state states Years of the states <t< td=""><td></td><td></td><td></td><td></td><td></td></t<> | | | | | |
| Beschervice works for a disperient of initial data of the second se | Pinal report covering period 15 June 1966- 31 October 1968 Turnenne (Parl Lease and Annue Lease Annue Lea | ORGANOTINS FOR ANT | IMALARIAL SCREENI | NG | | |
| Description and and a measure datases Pinary of a seried in the intervience of t | - Description was added failed to be added by the second of the second second of the second of the second second second of the second second second of th | | | | | •, • |
| Final report covering period is one 1900-31 October 1900 Henry G. Kuivila Autonomy divinance added multicly faituated Autonomy divind faituated for testing and test results were faituated for testi | That report covering period is only food 5 of covering for the covering of the covering and the covering of the covering of the covering and th | 4. DESCRIPTIVE NOTES (Type of report and | inclusive dates) | 1066 27 0 | | |
| Henry 0. Kuivila Image: Status 1973 Ima | Henry O. Kuivila Image: Stanuary 1973 Conventor on service of the servic | Final report covering | ng period 15 June | 1900- 31 00 | ctoper 1900 | · · |
| 1 1 TOTAL NO. OF FARTS 13 1 13 13 1 14 CONTRACT ON SAME YOU. 13 1 1 CONTRACT CONTRACT SATEMAN OF THOSE AND ON THE AND ON THOSE AND ON THOS | S January 1973 48 pages 13 Conversion statution 10 10 DA-197-193-ND-3018 Report no. 7 A remeasure no. 10 10 Conversion statution 10 10 Conversion statution 10 10 10 Conversion statution 10 10 10 Conversion statution 10 10 10 10 Conversion statution 10 10 10 10 10 Conversion statution 10 10 10 10 10 10 Conversion statution 10 </td <td>Henry G. Kuivila</td> <td></td> <td></td> <td></td> <td></td> | Henry G. Kuivila | | | | |
| S January 1973 1 18 Datas 18 As CONTRACTOR SHART NO. DA-16-13-MD-3018 He ORIGINATOR'S REFORT NUMBERING Report no. 7 * PROJECT NO. He ORIGINATOR'S REFORT NUMBERING Report no. 7 * CONTRACTOR SHART DISTRIBUTION OF this document is unlimited. Social Stream (Stream) * Control Sesarch & Development Command 1. Coop Commanding General, U. S. Army Medical Research & Development Command 1. Coop Commanding General, U. S. Army Medical Service Agency. 16 copies Walter Reed Army Medical The Sufficiency Notice Center, Medicinal Chemistry. U. S. Army Medical Research and Development Command. U. S. Army Medical Research and Development Command. * Austract Is unimated. Chemistry. U. S. Army Medical Research and Development Command. * Austract Is unimated to for command to for command to for command development Command * Austract U. S. Army Medical Research and Development Command * Austract Is used to prepare for antimalarial screening organotins in four general screening (2) introduction of organotin groups into unimate its derivatives; 3) introduction of organotin groups into unimate its derivatives; 3) introduction of organoting roups into unimate its derivatives. During the course of the contract it was nossible to carry on research on three of these categories. However, successful synthetic results were obtained only in the first of these. A total of forty three compounds was aubmitted for testing, and test results were returned on thirty six of thes | S January 1973 1 TOTAL NO. OF PAGES 13 4 CONTRACT ON STANT NO. DA-15-39-SUD-3018 4 Seport no. 7 5 Annuary 1973 5 Seport no. 7 6 CONTRACT ON STANT NO. DA-15-39-SUD-3018 5 Seport no. 7 6 Contract on Stant Not State S | | | | | • |
| 5. Allowary 1973 18. Converse 18. No. or market 10. DA-19-193-ND-3018 48. pages 11. Production of Reany No. DA-19-193-ND-3018 12. Converse Reany No. DA-19-193-ND-3018 13. Products No. 7 No. or market 14. Converse Reany No. DA-19-193-ND-3018 15. Products No. 7 No. or market 16. Converse Reany No. No. or market 17. Superclasser No. No. or market 18. Converse Reany No. No. or market 19. Constantion Command (Medical Service Army Medical Research & Development Command) I. S. Army Medical Research and Development Command 10. Superclasser Name Center, Medicinal Is someonic organotins in four general screening; 2) introduction of organotin groups into Undersearch and Development Command 11. was proposed to prepare for antimalarial screening organotins in four general screening; 2) introduction of organotin groups into Undersearch and Development Command 11. threduction of organotin groups ito sulfones which had shown some antimalarial activit | 1 10. or fairs 18 2 24 8 pages 18 2 10. or fairs 19. or fairs 3 10. or fairs 10. or fairs 4 10. or fairs 10. or fairs 4 10. or fairs 10. or fairs 5. or fairs 10. or fairs 10. or fairs 4 10. or fairs 10. or fairs 5. or fairs 10. or fairs 10. or fairs 5. or fairs 10. or fairs 10. or fairs 5. or fairs 10. or fairs 10. or fairs 2 10. or fairs 10. or fairs 3. or fairs 10. or fairs 10. or fairs 4. or fairs 10. or fairs 10. or fairs 4. or fairs 10. or fairs 10. or fairs 5. or fairs 10. or fairs 10. or fairs 1. or fairs 10. or fairs 10. or fairs 2. or fairs 10. or fairs 10. or fairs 3. or fairs 10. or f | | | | | |
| 5 January 1973 48 pages 18 DA-49-193-MD-3018 Report no. 7 A preserve. Report no. 7 A Development Distribution of this document is unlimated. 20 conies Defense Documentation Center; L copies Commanding General, U. S. Army Medical Research & Development Command; I copy Commanding General, U. S. Army Medical Research & Development Command; I copy Commanding General, U. S. Army Medical Research & Development Command; I copy Commanding General, U. S. Army Medical Center, Medicinal The user proposed to prepare for antimalarial screening organctins in four general screening; 2) introduction of organotin groups into L-aminoquinoling; L) introduction of organoting roups into L-aminoquinoling; L) introduction of these categories. Mewer, success'ul synthetic results were obtained only in the first of these. A total of forty three compounds was submitted for testing, and test results were returned on thirty six of these. These included: unsaturated fulfones; alkylallylting; alkylallenyltins; alkylallenyltins; alkylatinglicyloheeylyltins; and bis-salicylideediaminoethane-alkyltin halide complexes. Sone showed activit | 5 January 1973 1 48 pages 18 • CONTACT OB SAMATION DA-L9-193-MD-3018 • CONSISTING STATEMENT OF SAME AND | REPORT DATE | | 74. TOTAL NO. C | OF PAGES | 75. NO. OF REFS |
| Cheritater en enant No. DA-L(9-19)-ND-3018 Project No. 7 Report no. 7 Statistic Present Nousers and Annual Statistic Action of this document is unlimated. Coroles Defense Documentation Center; is copies Commanding General, U. S. Army Combe The Contract & Development Command; 1 copy Commanding General, U. S. Army Medical Research & Development Command; 1 copy Commanding General, U. S. Army Medical The Contract Nousers Nousers are not not the second The Context Nousers are not not the second and Development Command Center, Medicinal Center, Medical Research and Development Command Composition of organotin groups into chain(contex) coroles of the contract it was possible to carry on research on three of these categories, Sone s | L CONTRACT ON SAAT NO. DA-16-133-KD-3018 * MOUNCET NO. 7 * OTHER REPORT NOUNCEARS Report NO. 7 * OTHER REPORT NOUNCEARS Contense Decementation Center; L copies Commanding General, U. S. Army Comb Medical Research & Development Command J Cocy Commanding General, U. S. Army Medical BrailDevelopment Command, Medical Service Agency. 16 copies Walter Reed Army Medical T-SUPERLEMENTARY NOTES Center, Medicinal Chemistry. T superlement Command, I cocy Commanding General, U. S. Army Medical T-SUPERLEMENTARY NOTES Center, Medicinal Chemistry. T superlement Command, Service Agency. 16 copies Walter Reed Army Medical T-SUPERLEMENTARY NOTES Center, Medicinal Chemistry. T superlement Command, Service Agency. 16 copies Walter Reed Army Medical T-SUPERLEMENTARY NOTES Center, Medicinal Chemistry. T superlement Command, Service Agency. 16 copies Walter Reed Army Medical T-SUPERLEMENTARY NOTES Center, Medicinal Chemistry. T superlement Command, Service Agency. 16 copies Walter Reed Army Medical T-SUPERLEMENTARY NOTES Center, Medicinal Chemistry. T superlement Command, Service Agency. 16 copies Walter Reed Army Medical Chemistry. T superlement Command, Service Agency. 16 copies Walter Reed Army Medical Chemistry. T superlement Command, Service Agency of the search and Development Command Chemistry. During the course of the contract it was nossible to carry on research on three of these categories. However, success'ul synthetic results were obtained only in the first of these. A total of forty three compounds was submitted for testing, and test results were returned on thirty six of these. These included: unsaturated fulfones; alkylallylitins; alkylallenyltins; alkylpyridines; bicycloheptyltins; and bia-salicylidenediaminethane- alkyltin halde complexes. Some showed activity in mice, but were inactive or toxic in chicks; and vice versa. None was active in both, and many were toxic in one or both. D rows. 1473 Second 200 and 200 a | 5 January 1973 | ů. | 48 pag | es | 18 |
| Report no. 7 k endustry. Actively and the first of the contract it was nossible to carry on research on three of these cateories. However, successful synthetic results were obtained only in the first of these. At the of forty three compoundings was submitted for testing, and test results were returned on thirty six of these. These included: unsaturated fulfones; alkylallyltins; alkylalleyltins; alkyltin halide complexes. Some showed activity in mice, but were inactive or toxic in chicks; and vice versa. None was active in both, and many were toxic in one or both. | Hardy-Ly-symuly-Joho Keport no. 7 Hardy-Ly-symuly-Joho Array Medical Service Account is unlimited. Coordies Defense Documentation Center; L copies Commanding General, U. S. Army Combe Jum Development Command, Medical Service Account is unlimited. Co cordination Command, Medical Service Account Command in General, U. S. Army Combe Jum Development Command, Medical Research and Development Command. Chemistry. Chemistry. Chemistry. Chemistry is a proposed to prepare for antimalarial screening organotins in four general categories: 1) preparation of a variety of relatively simple organotine for general screening; 2) introduction of organotin groups into L-aminoquinoline; 1, introduction of organoting roups into L-aminoquinoline; 1, introduction of organoting roups into L-aminoquinoline; 1, introduction of organoting groups into L-aminoquinoline; 1, introduction of organoting roups into L-aminoquinoline; 1, introduction of these. A total of forty three compounds was submitted for testing, and test results were returned on thirty six of these. These included: unsaturated fulfones; alkylaylallyltins; alkylalleyltins; alkylalleyltins; alkylayling is alkylawide. Some showed activity in mice, but were inactive or toxic in chicks; and vice versa. None was active in both, and many were toxic in one or both. D Page 1473 Statest Statest Sta | DA LO TOT ID TOT | | M. ORIGINATOR | S REPORT NUM |)ER(\$) |
| It was proposed to prepare for antimalarial screening organotins in four general categories: 1) preparation of organotin groups into quinine its derivatives; 3) introduction of organotin groups into quinine its derivatives; 3) introduction of organotin groups into quinine its derivatives; 3) introduction of organotin groups into quinine its derivatives; 3) introduction of organotin groups into quinine its derivatives; 1) introduction of organotin groups into quinine its derivatives; 3) introduction of organotin groups into quinine its derivatives; 3) introduction of organotin groups into quinine its derivatives; 3) introduction of organoting groups into quinine its derivatives; 3) introduction of organoting groups into quinine its derivatives; 3) introduction of organoting groups into quinine its derivatives; 3) introduction of organoting groups into quinine its derivatives; 3) introduction of organoting groups into quinine its derivatives; 3) introduction of organoting groups into quinine its derivatives; 3) introduction of organoting groups into quinine its derivatives; 3) introduction of organoting groups into the second derivatives; 4) introduction of organoting groups into the derivative; 4) introduction of organoting groups into the derivative; 5) introduction of organoting groups into the derivative of these. These included in unsurvated fulfones; alkylallyllting; alkylallenyllting; alkylallenyl | A DIVER REPORT HOUS (Any other manham but may be assigned as our mouse of the presence of the open of the presence of the presence of the open of the presence of the open of the presence of the open of the presence of the presence of the open of the presence of the presence of the open of the presence of the open of the presence of the presence of the presence of the presence of the open of the presence of the open of the presence of | UA-449-193-MU-3010 | | Report | no. 7 | |
| Definition of the contract it was to solve to carry on research on three of the contract it was to solve the solve to the solve to the contract of the contract it was to solve the contract. During the course of the contract it was to solve to the contract was a solved to three the contract of the contract it was a contract. During the course of the contract it was to solve to the contract was a solved to three the contract. During the course of the contract it was to solve to the contract it was to solve to the contract it was a contract. During the course of the contract it was to solve to the contract was a solved to the contract. These include: unsaturated fulfones; alkylallyltins; alkylallenyltins; alkylallen | A STREM REPORT NOID (Any other manhaters that may be dealed in the strength of th | O, FROJECT NO. | | 1 | | |
| A Minimum Annu Andrew Market Market Strategy and Strat | A Service of the contract it was possible to carry on research or drives of these attents. However, successful synthetic results were obtained only in the first of these. A total of forty three compounds was submitted for testing, and test results were returned on thirty six of these. These included: unsaturated fulfones; alkylalythins; alkylapyridines; bioyclohepithins; and bis-salicylidenediamineethane-alkylation of ortex. None was active in both, and many were toxic in one or both. | 6. | · | B. OTHER STOC | RT NO(1) /Any A | het numbers that may be seed as |
| 4 So correst before bounder of this document is unlimated. 20 cooles Defense Documentation Center; 4 copies Commanding General, U. S. Army Combe TamlDevelopment Command, Medical Service Arency. 16 copies Walter Reed Army Medical The service and the service arency. 16 copies Walter Reed Army Medical The service arency. 18 service Walter Reed Army Medical The service arency. 18 service arency. 18 service arency and Development Command the service arency of the service arency of relatively simple organotins for general screening; 2) introduction of organoting groups into quinine its derivatives; 3) introduction of organoting roups into quinine its derivatives; 3) introduction of organoting roups into quinine its derivatives; 3) introduction of organoting roups into animoquinoline; b) introduction of organoting groups into animoquinoline; b) introduction of organoting roups into areninoquinoline; b) introduction of organoting sources which had shown some antimal- arial activity. During the course of the contract it was possible to carry on research on three of these categories. However, success ² and of forty three compounds was submitted for testing, and test results were returned on thirty six of these. These included: unsaturated fulfones; alkylallyltins; alkylallenyltins; alkylpyridines; bicycloheptyltins; and bis-salicylidenediamineethane- alkyltin halide complexes. Some showed activity in mice, but were inactive or toxic in chicks; and vice versa. None was active in both, and many were toxic in one or both. | A summation erargement Distribution of this document is unlimited. 20 conies Defense Documentation Center; & copies Commanding General, U. S. Army Combe Tas Development Command, Medical Service Azency. 16 copies Walter Reed Army Medical Tester Tester Tester Center, Medicinal U. S. Army Medical Research and Development Command Tester Tester Tester Tester Center, Medicinal U. S. Army Medical Research and Development Command Tester | ν. | • | this report) | and the second sec | |
| Distribution statement Distribution of this document is unlimited. 20 conies Defense Documentation Center; i copies Commanding General, U. S. Army Medical Research & Development Command; l covy Commanding General, U. S. Army Medical TableVelopment Command, Medical Service Arency. 16 copies Malter Reed Army Medical Chemistry. Is stonadawa Multian Activity U. S. Army Medical Research and Development Command A servator A servator It was proposed to prepare for antimalarial screening organotins in four general categories: 1) preparation of a variety of relatively simple organotins for general screening; 2) introduction of organotin groups into quinine its derivatives; 3) introduction of organotin groups into quinine its derivatives; 3) introduction of organotin groups into u-aminoquinoline; 1) introduction of organotin groups into h-aminoquinoline; 1) introduction of organoting swhich had shown some antimalarial activity. During the course of the contract it was possible to carry on research on three of these categories. However, successful synthetic results were obtained only in the first of these. A total of forty three compounds was sublitued for testing, and test results were returned on thirty six of these. These included: unsaturated fulfones; alkylallyltins; alkylallenyltins; alkylpyridines; bicycloheptyltins; and bis-salicylidenediaminethane-alkyltin halide complexes. Some showed activity in mice, but were inactive or toxic in one or both. Differentiation one or both. | Distribution statement Distribution of this document is unlimited. Cooles Defense Documentation Center; L copies Commanding General, U. S. Army Medical Research & Development Command; L coy Commanding General, U. S. Army Medical Research & Development Command, Medical Service Agency. 16 copies Malter Reed Army Medical Topies Malter Reed Army Medical Service Agency. 16 Copies Malter Reed Army Medical Chemistry. Chemistry. Chemistry. Chemistry. Center, Medicinal Command: Loop Commanding General, U. S. Army Medical Research and Development Command Assynact Assynact Assynact The was proposed to prepare for antimalarial screening organotins in four general categories: L) preparation of a variety of relatively simple organotins for general screening; 2) introduction of organotin groups into quinine its derivatives; 3) introduction of organoting roups into L-aminoquinoline; L) introduction of organoting roups into has shown some antimalarial activity. During the course of the contract it was possible to carry on research on three of these categories. However, successful synthetic results were obtained only in the first of these. A total of forty three compounds was submitted for testing, and test results were returned on thirty six of these. These included: unsaturated fulfones; alkylallyltins; alkylallenyltins; alkylallenyltins; alkylpyridines; bicycloheptyltins; and bis-salicylidenediaminethane-alkyltin halide complexes. Some showed activity in mice, but were inactive or toxic in one or both. Difference in one or both. | 4 | | 1 | | |
| <pre>general categories: 1) preparation of a variety of relatively simple organotins for general screening; 2) introduction of organotin groups into 4-aminoquinoline; 4) introduction of organotin groups into sulfones which had shown some antimal- arial activity.</pre> During the course of the contract it was possible to carry on research on three of these categories. However, successful synthetic results were obtained only in the first of these. A total of forty three compounds was submitted for testing, and test results were returned on thirty six of these. These included: unsaturated fulfones; alkylallyltins; alkylallenyltins; alkylpyridines; bicycloheptyltins; and bis-salicylidenediaminethane- alkyltin halide complexes. Some showed activity in mice, but were inactive or toxic in chicks; and vice versa. None was active in both, and many were toxic in one or both. DD roots 1473 Stelects Dor form 12.1. And 54. Number 13. Security ClassificeUoa | general categories: 1) preparation of a variety of relatively simple organotins for general screening; 2) introduction of organotin groups into 4-aminoquinoline; 4) introduction of organotin groups into sufferences which had shown some antimal- arial activity. During the course of the contract it was possible to carry on research on three of these categories. 'However, success'ul synthetic results were obtained only in the first of these. A total of forty three compounds was submitted for testing, and test results were returned on thirty six of these. These included: unsaturated fulfones; alkylallyltins; alkylallenyltins; alkyloyridines; bicycloheptyltins; and bis-salicylidenediaminoethane- alkyltin halide complexes. Some showed activity in mice, but were inactive or toxic in chicks; and vice versa. None was active in both, and many were toxic in one or both. | S. ABSTRACT | are for entimelar | L | ng organoti | ns in four |
| During the course of the contract it was nossible to carry on research on three of these categories. However, successful synthetic results were obtained only in the first of these. A total of forty three compounds was submitted for testing, and test results were returned on thirty six of these. These included: unsaturated fulfones; alkylallyltins; alkylallenyltins; alkylpyridines; bicycloheotyltins; and bis-salicylidenediaminoethane- alkyltin halide complexes. Some showed activity in mice, but were inactive or toxic in chicks; and vice versa. None was active in both, and many were toxic in one or both. | During the course of the contract it was possible to carry on research on three of these categories. However, successful synthetic results were obtained only in the first of these. A total of forty three compounds was submitted for testing, and test results were returned on thirty six of these. These included: unsaturated fulfones; alkylallyltins; alkylallenyltins; alkylpyridines; bicycloheptyltins; and bis-salicylidenediamincethane- alkyltin halide complexes. Some showed activity in mice, but were inactive or toxic in chicks; and vice versa. None was active in both, and many were toxic in one or both. | It was proposed to prepa | NTC TAT MITATURY CIT | lai screeni | ng utganuut | |
| obtained only in the first of these. A total of forty three compounds was submitted for testing, and test results were returned on thirty six of these. These included: unsaturated fulfones; alkylallyltins; alkylallenyltins; alkylpyridines; bicycloheptyltins; and bis-salicylidenediamincethane- alkyltin halide complexes. Some showed activity in mice, but were inactive or toxic in chicks; and vice versa. None was active in both, and many were toxic in one or both. D. rows. 1473 Concerts poor commuter. Jan 44, Which is Security Classification | obtained only in the first of these. A total of forty three compounds was submitted for testing, and test results were returned on thirty six of these. These included: unsaturated fulfones; alkylallyltins; alkylallenyltins; alkylpyridines; bicycloheptyltins; and bis-salicylidenediaminoethane- alkyltin halide complexes. Some showed activity in mice, but were inactive or toxic in chicks; and vice versa. None was active in both, and many were toxic in one or both. D. MON. 1473 Structs DD FORM 1473, 1 JAN 44, WHICH 15 Security Classification | It was proposed to prepa general categories: 1); for general screening; derivatives; 3) introduction (1) introduction of organ arial activity. | preparation of a 2) introduction uction of organot notin groups into | variety of s of organotis in groups is sulfones w | relatively n groups in nto 4-amino hich had sh | simple organotins to quinine its quinoline; own some antimal- |
| D rotu 1473 ASPLACES DO FORM 1475. 1 JAN 64, WHICH 15 Security Classification | Inese included: unsaturated fulfones; alkylallyltins; alkylallenyltins; alkylallenyltins; alkylapyridines; bicycloheptyltins; and bis-salicylidenediaminoethane-alkyltin halide complexes. Some showed activity in mice, but were inactive or toxic in chicks; and vice versa. None was active in both, and many were toxic in one or both. D round to the second se | It was proposed to prepa general categories: 1) ; for general screening; derivatives; 3) introduction (1) introduction of organ arial activity. During the course of the on three of these categories | preparation of a 2) introduction action of organot notin groups into me contract it wa pries. However | variety of : of organoti: in groups in sulfones with s possible - successful | relatively n groups in nto 4-amino hich had sh to carry o synthetic | simple organotins to quinine its quinoline; own some antimal- n research results were |
| alkyltin halide complexes. Some showed activity in mice, but were inactive or toxic in chicks; and <u>vice versa</u> . None was active in both, and many were toxic in one or both. D. root at 1473 Applaces DD Form 1475. 1 Jan 64, WHICH 19 Security Classification | alkyltin halide complexes. Some showed activity in mice, but were inactive or toxic in chicks; and vice versa. None was active in both, and many were toxic in one or both. D. MOV1473 ALPLACES DO FORM 1473. I JAN 64, WHICH IS Security Classification | It was proposed to prepa general categories: 1) ; for general screening; derivatives; 3) introduction h) introduction of organ arial activity. During the course of the on three of these categories obtained only in the fin submitted for testing, a | preparation of a 2) introduction action of organot notin groups into the contract it wa pries. However, rst of these. A and test results | s possible successful total of for | to carry of synthetic relatively n groups in nto 4-amino hich had sh to carry of synthetic rty three c ed on thirt | simple organotins to quinine its quinoline; own some antimal- n research results were ompounds was y six of these. |
| D Form 1473 ASPLACES DO FORM 1475. 1 JAN 64, WHICH 18 Security Classification | D rota 1473 Alphaces do ronm lars. I Jan 64, Which is Security Classification | It was proposed to prepa general categories: 1) ; for general screening; derivatives; 3) introduction (1) introduction of organ arial activity. During the course of the on three of these categories obtained only in the fin submitted for testing, a These included: unsature | preparation of a 2) introduction action of organot notin groups into me contract it wa pries. However, rst of these. A and test results ated fulfones; al | s possible successful total of for sulfones with successful total of for were returned kylallylting | to carry of subtractions of subtractions of the subtraction of the sub | simple organotins to quinine its quinoline; own some antimal- n research results were ompounds was y six of these. enyltins; |
| D FORM 1473 AEPLACES DO FORM 1475, 1 JAN 64, WHICH 15 1 NOV 1473 OBSOLETE FOR ARMY USE. Security Classification | D FORM 1473 AEPLACES DD FORM 1473. 1 JAN 64, WHICH 18 NOV 4.1473 OBSOLETE FOR ARMY USE. Security Classification | It was proposed to prepa general categories: 1) ; for general screening; derivatives; 3) introduction h) introduction of organ arial activity. During the course of the on three of these categories obtained only in the first submitted for testing, a These included: unsatures alkyloyridines; bicyclob alkyltin balide completed | preparation of a 2) introduction action of organot notin groups into the contract it wa pries. However, rst of these. A and test results ated fulfones; all heptyltins; and b | s possible successful total of for successful total of for were return kylallyltin is-salicylic | to carry of subtractions of the second secon | simple organotins to quinine its quinoline; own some antimal- n research results were ompounds was y six of these. enyltins; ethane- were inactive |
| D FORM 1473 AEPLACES DO FORM 1473. I JAN 64, WHICH IS D NOV 00 1473 OBSOLETE FOR ARMY USE. | D 1000 4. 1473 AEPLACES DO FORM 1475, 1 JAN 44, WHICH IS OBSOLETE FOR ARMY USE. Security Classification | It was proposed to prepa general categories: 1) ; for general screening; derivatives; 3) introduction h) introduction of organ arial activity. During the course of the on three of these category obtained only in the first submitted for testing, a These included: unsature alkylpyridines; bicyclou alkyltin halide complexed or toxic in chicks; and were toxic in one or both | preparation of a 2) introduction action of organot notin groups into me contract it wa pries. However, rst of these. A and test results ated fulfones; al heptyltins; and b as. Some showed vice versa. Non th. | al screeni variety of : of organoti: in groups in sulfones wi successful total of for were return kylallyltin is-salicylic activity in e was active | to carry o synthetic rty three c ed on thirt s; alkylall denediamino mice, but e in both, | simple organotins to quinine its quinoline; own some antimal- n research results were ompounds was y six of these. enyltins; ethane- were inactive and many |
| D FORM 1473 REPLACES DD FORM 1473, 1 JAN 64, WHICH 18 D 1 NOV 45 1473 OBSOLETE FOR ARMY USE. Security Classification | D FORM 1473 AEPLACES DD FORM 1473. 1 JAN 44, WHICH IS 0850LETE FOR ARMY USE. Security Classification | It was proposed to prepa general categories: 1) ; for general screening; derivatives; 3) introduction arial activity. During the course of the on three of these category obtained only in the first submitted for testing, a These included: unsature alkylpyridines; bicyclow alkyltin halide complexed or toxic in chicks; and were toxic in one or both | preparation of a 2) introduction action of organot notin groups into the contract it wa pries. However, rst of these. A and test results ated fulfones; al heptyltins; and b as. Some showed vice versa. Non th. | al screen variety of s of organotis in groups in sulfones with successful total of for were return kylallyltins is-salicylic activity in e was active | relatively n groups in nto 4-amino hich had sh to carry o synthetic rty three c ed on thirt s; alkylall denediamino mice, but e in both, | simple organotins to quinine its quinoline; own some antimal- n research results were ompounds was y six of these. enyltins; ethane- were inactive and many |
| D FORM 1473 AEPLACES DO FORM 1475, I JAN 64, WHICH IS OBSOLETE FOR ARMY USE. Security Classification | D FORM 1473 AEPLACES DD FORM 1473. 1 JAN 64, WHICH IS NOV 4.1473 OBSOLETE FOR ARMY USE. Security Classification | It was proposed to prepa general categories: 1) ; for general screening; derivatives; 3) introduction i) introduction of organ arial activity. During the course of the on three of these category obtained only in the fin submitted for testing, a These included: unsature alkylpyridines; bicyclou alkyltin halide complexed or toxic in chicks; and were toxic in one or both | preparation of a 2) introduction action of organot notin groups into the contract it wa pries. However, rst of these. A and test results ated fulfones; al heptyltins; and b bs. Some showed vice versa. Non th. | s possible sulfones wi sulfones wi successful total of for were return kylallyltin is-salicylic activity in e was active | relatively n groups in nto 4-amino hich had sh to carry o synthetic rty three c ed on thirt s; alkylall denediamino mice, but e in both, | simple organotins to quinine its quinoline; own some antimal- n research results were ompounds was y six of these. enyltins; ethane- were inactive and many |
| D PORM 1473 AEPLACES DO FORM 1475, 1 JAN 64, WHICH IS 1 NOV 0. 1473 OBSOLETE FOR ARMY USE. Security Classification | D FORM 1473 AEPLACES DD FORM 1475, I JAN 44, WHICH IS NOV 44 1473 OBSOLETE FOR ARMY USE. Security Classification | It was proposed to prepa general categories: 1) ; for general screening; derivatives; 3) introdu h) introduction of organ arial activity. During the course of the on three of these category obtained only in the first submitted for testing, a These included: unsature alkylpyridines; bicyclou alkyltin halide complexed or toxic in chicks; and were toxic in one or both | preparation of a 2) introduction action of organot notin groups into the contract it wa pries. However, rst of these. A and test results ated fulfones; al heptyltins; and b as. Some showed vice versa. Non th. | al screen variety of : of organoti: in groups in sulfones w successful total of for were return kylallyltin is-salicylic activity in e was active | relatively n groups in nto 4-amino hich had sh to carry o synthetic rty three c ed on thirt s; alkylall denediamino mice, but e in both, | simple organotins to quinine its quinoline; own some antimal- n research results were ompounds was y six of these. enyltins; ethane- were inactive and many |
| DD FORM 1473 REPLACES DD FORM 1473, I JAN 64, WHICH IS NOV 45 1473 OBSOLETE FOR ARMY USE. Security Classification | D 1000 4. 1473 AEPLACES DO FORM 1473, 1 JAN 44, WHICH IS Security Classification | It was proposed to prepa general categories: 1) i for general screening; derivatives; 3) introduction (h) introduction of organ arial activity. During the course of the on three of these categories obtained only in the fin submitted for testing, a These included: unsature alkylpyridines; bicyclou alkyltin halide complexe or toxic in chicks; and were toxic in one or both | preparation of a 2) introduction action of organot notin groups into the contract it wa pries. However, rst of these. A and test results ated fulfones; all heptyltins; and b as. Some showed vice versa. Non th. | al screen variety of s of organotis in groups in sulfones with successful total of for were return kylallylting activity in e was active | relatively n groups in nto 4-amino hich had sh to carry o synthetic rty three c ed on thirt s; alkylall denediamino mice, but e in both, | simple organotins to quinine its quinoline; own some antimal- n research results were ompounds was y six of these. enyltins; ethane- were inactive and many |
| DD PORM 1473 AEPLACES DO FORM 1473. 1 JAN 64, WHICH IS NOV 00 1473 OBSOLETE FOR ARMY USE. Security Classification | D PORM 1473 AEPLACES DD PORM 1473. I JAN 64, WHICH IS BEOLETE POR ARMY USE. Security Classification | It was proposed to prepa general categories: 1) i for general screening; derivatives; 3) introduction arial activity. During the course of the on three of these categories obtained only in the first submitted for testing, a These included: unsature alkylpyridines; bicyclow alkyltin halide complexed or toxic in chicks; and were toxic in one or both | preparation of a 2) introduction action of organot notin groups into the contract it wa pries. However, rst of these. A and test results ated fulfones; all heptyltins; and b as. Some showed vice versa. Non th. | al screen variety of s of organotis in groups in sulfones with successful total of for were return kylallyltins is-salicylic activity in e was active | relatively n groups in nto 4-amino hich had sh to carry o synthetic rty three c ed on thirt s; alkylall denediamino mice, but e in both, | simple organotins to quinine its quinoline; own some antimal- n research results were ompounds was y six of these. enyltins; ethane- were inactive and many |
| Security Classification | Security Classification | It was proposed to prepa general categories: 1) ; for general screening; derivatives; 3) introdu h) introduction of organ arial activity. During the course of th on three of these category obtained only in the first submitted for testing, a These included: unsature alkylpyridines; bicyclou alkyltin halide complexed or toxic in chicks; and were toxic in one or both | preparation of a 2) introduction action of organot notin groups into the contract it wa pries. However, rst of these. A and test results ated fulfones; al heptyltins; and b as. Some showed vice versa. Non th. | al screen variety of s of organotis in groups is sulfones wi successful total of for were return kylallyltins is-salicylic activity in e was active | relatively n groups in nto 4-amino hich had sh to carry o synthetic rty three c ed on thirt s; alkylall denediamino mice, but e in both, | simple organotins to quinine its quinoline; own some antimal- n research results were ompounds was y six of these. enyltins; ethane- were inactive and many |
| | | It was proposed to prepa general categories: 1) for general screening; derivatives; 3) introdu h) introduction of organ arial activity. During the course of th on three of these category obtained only in the fin submitted for testing, a These included: unsature alkylpyridines; bicyclol alkyltin halide complexe or toxic in chicks; and were toxic in one or both | preparation of a 2) introduction action of organot notin groups into the contract it wa pries. However, rst of these. A and test results ated fulfones; al heptyltins; and b as. Some showed vice versa. Non th. | All screening variety of s of organotis in groups in sulfones with successful total of for were returned kylallylting activity in e was active | relatively n groups in nto 4-amino hich had sh to carry o synthetic rty three c ed on thirt s; alkylall denediamino mice, but e in both, | simple organotins to quinine its quinoline; own some antimal- n research results were ompounds was y six of these. enyltins; ethane- were inactive and many |
| · · · | | It was proposed to prepa general categories: 1) i for general screening; derivatives; 3) introduction h) introduction of organ arial activity. During the course of the on three of these categories obtained only in the first submitted for testing, a These included: unsature alkyloyridines; bicyclow alkyltin halide complexed or toxic in chicks; and were toxic in one or both D | preparation of a 2) introduction action of organot notin groups into the contract it wa pries. However, rst of these. A and test results ated fulfones; al heptyltins; and b as. Some showed vice versa. Non th. | All screen variety of s of organotis in groups in sulfones with successful total of for were return kylallyltins is-salicylic activity in e was active | relatively n groups in nto 4-amino hich had sh to carry o synthetic rty three c ed on thirt s; alkylall denediamino mice, but e in both, | simple organotins to quinine its quinoline; own some antimal- n research results were ompounds was y six of these. enyltins; ethane- were inactive and many |
| | | It was proposed to prepa general categories: 1) for general screening; derivatives; 3) introduction h) introduction of organ arial activity. During the course of the on three of these categories obtained only in the first submitted for testing, a These included: unsature alkylpyridines; bicyclou alkyltin halide complexed or toxic in chicks; and were toxic in one or both D PORM 1473 SECONT | preparation of a 2) introduction action of organot notin groups into the contract it wa pries. However, rst of these. A and test results ated fulfones; al heptyltins; and b bas. Some showed vice versa. Non th. | All Screeni variety of s of organotis in groups in sulfones wi successful total of for were return kylallyltin is-salicylic activity in e was active | relatively relatively n groups in nto 4-amino hich had sh to carry o synthetic rty three c ed on thirt s; alkylall denediamino mice, but e in both, | simple organotins to quinine its quinoline; own some antimal- n research results were ompounds was y six of these. enyltins; ethane- were inactive and many |
| | | It was proposed to prepa general categories: 1) for general screening; derivatives; 3) introdu l) introduction of organ arial activity. During the course of the on three of these categories obtained only in the first submitted for testing, a These included: unsature alkylpyridines; bicyclow alkyltin halide complexed or toxic in chicks; and were toxic in one or both D: MON1473 | preparation of a 2) introduction action of organot notin groups into the contract it was pries. However, rst of these. A and test results ated fulfones; all heptyltins; and b as. Some showed vice versa. Non th. | All screen variety of s of organotis in groups in sulfones wi successful total of for were return kylallyltins is-salicylic activity in e was active | relatively relatively n groups in nto 4-amino hich had sh to carry o synthetic rty three c ed on thirt s; alkylall denediamino mice, but e in both, Secunty | simple organotins to quinine its quinoline; own some antimal- n research results were ompounds was y six of these. enyltins; ethane- were inactive and many Classification |

مر المراجع مر المراجع المراجع المراجع المراجع المراجع المراجع المراجع مراجع المراجع المراجع المراجع المراجع المراجع ا

Person con

| Secu | rity Classifics | tion | | | | | | | | |
|------------------------------------|-----------------|----------|-----|-------------------------|----------|-----|--------------|-----|----------|----|
| 14. | | KEY WO | RDS | | LIN | K A | LIN | KB | LIN | ĸc |
| | • | | | • | ROLE | | ROLE | WT | ROLE | |
| antima | ania]e | | | • | | | • | • | | |
| a1191/14. | tal 1410 | | | | | | | ĺ | | |
| organo | tins | ••• | | | | | | | | |
| allylt | ins | | | | | | | 4 | j | |
| alleny | ltins | | | | | | | | | |
| nvrfdv | tins | 2. 1. 10 | | | [· | l · | • • | · · | | |
| <i>p</i> y 1 2 4 <i>y</i> 1 | | | | | | | ¹ | ŀ. | | |
| | | | | | | Ì | | | | |
| | . | | | • •••••• ••• | {. | - | | | | |
| | - | • | | \$ 1. • • | . | | | 1 | ì | |
| | | • | • | | ł | ł | | | - | |
| | | | | | [| ł | | | | |
| • | | | • | . 1 | | | | | | |
| | | • | × | · · · | : | | | | | |
| • • • • • | , , | • | | • • • • | | | | | | |
| : | • • | ~ | | | | | } | | | ŀ |
| | | • | | | | | Į | | | |
| | · | • | | • | | | | | • | • |
| | | | | | ł | | | | | |
| • | | ●, | | • • • • • * * * | . | [| | | · | |
| | | | с. | • . | ł | · · | | • • | | |
| • | • • • • | - | | · · · · · | [| į . | • | | · · | |
| •· · · | •• | 5 D D D | . • | | | | , , | , . | : | |
| | | • | | | | ļ | | | | |
| | | | | · · · | | | | | | |
| | | · | | | | | | | | |
| | ; | • | • . | 4 * | | | | | | |
| | • | | | • | | | | • | | |
| | • | • | | | | | | | | |
| | | | | |] | | | | | |
| | | | | | | ł | | | | |
| | | | | | | | | | | |
| | | | | | 1 | 1 | i i | | 1 1 | |

с.

÷ ×*. -

التوادينين ويتعاد

Security Classification

2

,

والمتعادية والمتعاد و