

**Best
Available
Copy**

AD-763 503

THRESHOLD BEHAVIOR OF INELASTIC COLLISION
CROSS SECTIONS

James N. Bardsley, et al

Pittsburgh University

Prepared for:

Army Research Office
Advanced Research Projects Agency

May 1973

DISTRIBUTED BY:

NTIS

National Technical Information Service
U. S. DEPARTMENT OF COMMERCE
5285 Port Royal Road, Springfield Va. 22151

Unclassified

Security Classification

DOCUMENT CONTROL DATA - R & D

(Security classification of title, body of abstract and indexing annotation must be entered when the overall report is classified)

1. ORIGINATING ACTIVITY (Corporate author)		2a. REPORT SECURITY CLASSIFICATION	
University of Pittsburgh Pittsburgh, Pennsylvania 15203		Unclassified	
3. REPORT TITLE		2b. GROUP	
Threshold Behavior of Inelastic Collision Cross Sections			
4. DESCRIPTIVE NOTES (Type of report and inclusive dates)			
Scientific paper, to be published July, 1973			
5. AUTHOR(S) (First name, middle initial, last name)			
Bardsley, James N. Nesbet, Robert K.			
6. REPORT DATE	7a. TOTAL NO. OF PAGES	7b. NO. OF REFS	
May 1973	10 13	19	
8a. CONTRACT OR GRANT NO	8b. ORIGINATOR'S REPORT NUMBER(S)		
DA-31-124-ARO-D-440			
b. PROJECT NO			
P-6563-P			
c.	9b. OTHER REPORT NO(S) (Any other numbers that may be assigned this report)		
ARPA Order No. 826, Am. 9			
d.			
10. DISTRIBUTION STATEMENT			
Approved for public release; distribution unlimited			
11. SUPPLEMENTARY NOTES		12. SPONSORING MILITARY ACTIVITY	
		Advanced Research Projects Agency 1400 Wilson Blvd. Arlington, Virginia 22209	
13. ABSTRACT			
The behavior of scattering cross sections near an excitation threshold is examined for the case of long range potentials. The theory is confirmed by computations of e-Li scattering. Recent observations of threshold structure in e-Na collisions are ascribed to a 1D resonance of Na^-. A true cusp can occur only in the 1P partial cross section.			

Reproduced by
NATIONAL TECHNICAL
INFORMATION SERVICE
U S Department of Commerce
Springfield VA 22151

DD FORM 1 NOV 66 1473

Unclassified

Security Classification

AD 763503

Unclassified

Security Classification

14	KEY WORDS	LINK A		LINK B		LINK C	
		ROLE	WT	ROLE	WT	ROLE	WT
	Electron Scattering Scattering Cross Sections Alkali-metal atoms Lithium Sodium						

Un

THRESHOLD BEHAVIOR OF INELASTIC SCATTERING CROSS SECTIONS

by

J. N. Bardsley[†] and R. K. Nesbet

IBM Research Laboratory
San Jose, California

[†]Permanent address: Physics Department
University of Pittsburgh
Pittsburgh, Pennsylvania 15213

Sponsored by

The Advanced Research Projects Agency ARPA Order No. 826

Approved for public release;
distribution unlimited

July 1973

The views and conclusions contained in this document are those of the authors and should not be interpreted as necessarily representing the official policies, either expressed or implied, of the Advanced Research Projects Agency or the U.S. Government.

The effects of long range potentials on the threshold behavior of elastic scattering cross sections were studied some time ago, principally by O'Malley, Spruch and Rosenberg¹ and by Levy and Keller.² However, there has been little analysis of the effects of such potentials on inelastic collisions. The purpose of this paper is to extend the analysis of Levy and Keller² to inelastic scattering, and to relate the results to recent experiments involving alkali atoms. One particular feature that we will examine is the appearance of Wigner cusps at excitation thresholds. Our analysis will be applicable to any atom except hydrogen, for which special problems arise due to the ℓ -degeneracy. These special problems have been studied by Gaillitis and Damburg³ and others.

The analysis of Levy and Keller² is based on the variable phase method, which has been extended to multi-channel problems by several authors.^{4,5} For each channel α , the associated single-particle function $\phi_{\alpha\beta}(r)$ is written

$$\phi_{\alpha\beta}(r) = w_{0\alpha}(r)\delta_{\alpha\beta} + w_{1\alpha}(r)t_{\alpha\beta}(r) \quad (1)$$

The index β denotes the incident channel; $w_{0\alpha}(r)$ and $w_{1\alpha}(r)$ are independent functions which have the asymptotic form appropriate for the channel α . If there is no unscreened Coulomb interaction in the asymptotic region, these can be expressed in terms of spherical Bessel functions. For open channels we take

$$w_{0\alpha}(r) = k_{\alpha}^{-1/2} r j_{\ell_{\alpha}}(k_{\alpha} r) \quad ; \quad (2)$$

$$w_{1\alpha}(r) = -k_{\alpha}^{\frac{1}{2}} r n_{\ell_{\alpha}}(k_{\alpha} r) \quad (3)$$

The scattering information is contained in the matrix $t_{\alpha\beta}(r)$. The limit of $t_{\alpha\beta}(r)$, as $r \rightarrow \infty$, is the reactance matrix, $K_{\alpha\beta}$, whose eigenvalues are the tangents of the eigenphases.

By substitution in the Schrödinger equation we obtain an integral equation for $t_{\alpha\beta}(r)$, of the form⁴

$$t_{\alpha\beta}(r) = -2 \int_0^r \sum_{\gamma} \sum_{\delta} (\delta_{\alpha\gamma} w_{0\gamma}(r') + t_{\alpha\gamma}(r') w_{1\gamma}(r')) V_{\gamma\delta}(r') (w_{0\delta}(r') \delta_{\delta\beta} + w_{1\delta}(r') t_{\delta\beta}(r')) dr' \quad (4)$$

where $V_{\gamma\delta}(r)$ is the interaction potential, expressed in matrix form. The K-matrix Born Approximation⁶ can be obtained by neglecting the terms involving $t(r')$ on the RHS of Eq. (4). Substituting for $w_{0\beta}(r)$, we obtain

$$K_{\alpha\beta} \equiv t_{\alpha\beta}(\infty) = -k_{\alpha}^{\frac{1}{2}} k_{\beta}^{\frac{1}{2}} \int_0^{\infty} 2V_{\alpha\beta}(r) j_{\ell_{\alpha}}(k_{\alpha} r) j_{\ell_{\beta}}(k_{\beta} r) r^2 dr \quad (5)$$

As shown for single channel scattering by Levy and Keller,² the dominant term in the threshold behavior is given by Eq. (5).

Let us suppose that we have one or more old channels labelled α , β , ..., and one or more new channels labelled p , q , We will assume that the new channels share a common threshold so that $k_q = k_p$, and also to determine the dependence of the elements of the reactance matrix

upon k_p . Let us first consider K_{pq} which corresponds to elastic scattering in the new channels. For any short range interaction the integral is dominated, as $k_p \rightarrow 0$, by small r and we find

$$K_{pq}^S = -k_p^{\ell_p + \ell_q + 1} \left\{ \int_0^\infty 2V_{pq}(r) r^{\ell_p + \ell_q + 2} dr + O(k_p^2) \right\}. \quad (6)$$

However, if the interaction contains a long range component, say Cr^{-s} , this leads to a contribution of the form

$$K_{pq}^L = -2Ck_p \int_0^\infty r^{2-s} j_{\ell_p}(k_p r) j_{\ell_q}(k_p r) dr. \quad (7)$$

If $s < (\ell_p + \ell_q + 3)$, this integral is well defined and gives

$$K_{pq}^L = \frac{-\pi C}{2^{s-1}} k_p^{s-2} \frac{\Gamma(s-1) \Gamma\left(\frac{\ell_p + \ell_q - s + 3}{2}\right)}{\Gamma\left(\frac{\ell_p - \ell_q + s}{2}\right) \Gamma\left(\frac{\ell_p + \ell_q + s + 1}{2}\right) \Gamma\left(\frac{\ell_q - \ell_p + s}{2}\right)}. \quad (8)$$

Since, given the inequality above, $(s-2) < (\ell_p + \ell_q + 1)$, this term is dominant for small k . Thus we confirm the conclusion of Levy and Keller that the threshold behavior of the matrix elements K_{pq} can be changed by the presence of long range forces. If, however, $s \geq (\ell_p + \ell_q + 3)$ the potential must be modified at small r , and the dominant term remains $k_p^{\ell_p + \ell_q + 1}$, since the integral is made finite by the short range behavior of the potential.

Let us next consider the term $K_{\alpha p}$, corresponding to inelastic scattering. For any short range interaction we obtain a result similar to

Eq. (6) with a dominant term proportional to $k_p^{\ell_p + 1/2}$. From our long range component, Cr^{-s} , we find that if $s < (\ell_\alpha + \ell_p + 3)$

$$\begin{aligned}
 K_{\alpha p}^L &= -2C k_\alpha^{1/2} k_p^{1/2} \int_0^\infty r^{2-s} j_{\ell_\alpha}(k_\alpha r) j_{\ell_p}(k_p r) dr \\
 &= -\frac{\pi C}{2^{s-1}} k_\alpha^{s-1/2} k_p^{s-1/2} \frac{\Gamma\left(\frac{\ell_\alpha + \ell_p - s + 3}{2}\right)}{\Gamma(\ell_p + 3/2) \Gamma\left(\frac{\ell_\alpha - \ell_p + s}{2}\right)} \\
 &\quad \times \left[1 + O\left(\frac{k_p^2}{k_\alpha^2}\right) \right].
 \end{aligned} \tag{9}$$

For small k_p this has the same form as $K_{\alpha p}^S$. Thus we conclude that the long range interactions do not change the power of the dominant term for K-matrix elements in which the initial and final wave numbers are different. In this conclusion we confirm the analysis of Gelman,⁷ but differ from the result of Mott and Massey⁸ who state that $K_{\alpha p} \propto k_p^{s-1/2}$ when $s < \ell_p + 2$.

In order to examine the physical cross sections we must construct the T-matrix, given by $K(1 - iK)^{-1}$. By analytic continuation of the T-matrix, the behavior of the K-matrix below the threshold can be deduced.^{9,10} It is well known^{9,10} that if an element $K_{\alpha p}$ is proportional to $k_p^{1/2}$, then the elastic scattering T-matrix element $T_{\alpha\alpha}$ will, in general, have infinite slope at the threshold. This can lead to a cusp in $T_{\alpha\alpha}$ and in the corresponding partial wave cross section $\sigma_{\alpha\alpha}$. From our previous analysis we see that $K_{\alpha p} \propto k_p^k$ if, and only if, $\ell_p = 0$, that is, if the angular momentum of the scattered electron is zero.

Let us apply these ideas to electron scattering by alkalis in the ground 2S state, at energies close to the threshold for excitation of the lowest 2P state. For collisions in which the 2P state is excited the initial and final values of the electronic angular momentum, l_α and l_p , are linked through the selection rule

$$l_p = l_\alpha \pm 1 \quad . \quad (10)$$

Thus l_p can be zero only if $l_\alpha = 1$, so that cusps can appear at the excitation threshold only for incident p-waves.

Although through this type of analysis we can show that cusps may exist, we are unable to predict their magnitude except through an explicit calculation. We have, therefore, performed numerical computations of the reactance matrix and partial wave cross sections for electron scattering by lithium. We have used a variational method, used previously by Sinfaillam and Nesbet¹¹ and by Oberoi and Nesbet,¹² which has been described in detail by Lyons *et al.*¹³ The basis set of square integrable functions included 8 s-wave functions, 7 p-wave functions, 5 d-wave functions and 4 f-wave functions, supplemented by continuum functions with asymptotic forms given by Eq. (2).

At energies just above the inelastic threshold each of the elements of the reactance matrix varied with energy in a manner consistent with our previous analysis. However, some of the elements could be fitted by a simple power law only over a very small range of energies ($\sim 10^{-3}$ eV). The calculated partial wave cross sections for elastic scattering are shown in Fig. 1. The s- and d-wave contributions pass smoothly through

the threshold energy. There is a very clear cusp in the 1P contribution, which falls from $19 \pi a_0^2$ at threshold to $9.5 \pi a_0^2$ at an energy 0.02 eV above threshold. The total cross section decreases by approximately 6% over this energy range. The 1P contribution should have an infinite slope on both sides of the threshold energy. The finite slope that we find at energies below threshold is probably due to an inadequacy in our trial wave function. The basis set does not include any terms which have exactly the proper asymptotic form required for closed channels. We do, however, have continuum functions with the proper asymptotic form for the open channels, and so obtain the correct analytic form of the cross sections just above threshold.

The 3P contribution to elastic scattering also has an infinite slope at the threshold energy. However, the cross section derivative ($d\sigma/dE$) is negative on both sides of the threshold, so there is not a cusp, but a point of inflexion. In spite of the infinite derivative the magnitude of the change in the 3P contribution near threshold is negligible. This is because the elastic scattering phase shift, $\eta_{\alpha\alpha}$, is very close to $(n + \frac{1}{2})\pi$ (we find $\sin \eta_{\alpha\alpha} = 0.999$). In this situation a change of 10% in the K -matrix element $K_{\alpha\alpha}$ leads to a change of only 0.04% in the cross section.

From comparison with previous calculations^{11,14} of elastic electron-alkali scattering at lower energies and from exploratory calculations at higher energies it is clear that there are resonances in the 1P and 1D cross sections. These are presumably associated with the $(1s)^2(2s)(2p)^1P$ and $(1s)^2(2p)^2^1D$ states of Li^- . The effect of the resonances can be seen, for example, in the fact that the 1P and 1D contributions to

inelastic scattering are considerably greater than the 3P and 3D contributions. The presence of the 1P resonance enhances the effect of the cusp in that channel, and the maximum value of the 1P partial cross section occurs at the threshold energy. The 1D resonance causes the 1D partial cross section also to peak at an energy very close to the threshold.

The existence of these resonances was previously noted by Burke and Taylor.¹⁵ These authors also studied the sum of the scattering eigenphases, and found a discontinuity of slope at the threshold energy in each partial wave. We have confirmed this feature and find that for s-waves and d-waves this discontinuity is consistent with a smooth variation of the elastic scattering partial cross section.

Observations of cusps in e-Na differential scattering cross sections have been reported by Andrick *et al.*¹⁶ and by Gehera and Reichert.¹⁷ The structure observed by Andrick *et al.*¹⁵ is weak near 60° , and strong near 90° . However, the threshold cusps should vanish at 90° , since they are confined to incident p-waves. Since the qualitative features of e-Li scattering and e-Na scattering are very similar^{11,14} it seems plausible that the observed structure is due predominantly to a 1D resonance of Na^- . For many scattering angles the p-wave cusps should be masked by the 1D resonance. However, the fact that there is no observed structure near 60° must still be explained.

The effects of the cusps may be more easily observed in photodetachment of the alkali negative ions, since the final state must have 1P symmetry. From the calculations of Moores and Norcross¹⁸ it is clear that the detachment cross section peaks very close to the threshold

energy for the production of excited neutral atoms. Using Na^- and K^- , Patterson et al.¹⁹ have observed such peaks and from their observations have derived accurate values for the electron affinities of these alkalis.

REFERENCES

1. T. F. O'Malley, L. Spruch and L. Rosenberg, J. Math. Phys. 2, 491 (1961).
2. B. R. Levy and J. B. Keller, J. Math. Phys. 4, 54 (1963).
3. M. Gailitis and R. Damburg, Proc. Phys. Soc. 82, 192 (1963).
4. C. Zemach, Nuovo Cimento 33, 939 (1964); J. J. Matese and R.J.W. Henry, Phys. Rev. A5, 222 (1972).
5. R. K. Nesbet, Comments on Atomic and Molecular Physics 3, 143 (1972).
6. M. J. Seaton, Proc. Phys. Soc. (London) 77, 174 (1961).
7. S. Geltman, Topics in Atomic Collision Theory (Academic Press; New York, 1969), pp. 152-167.
8. N. F. Mott and H.S.W. Massey, Theory of Atomic Collisions, 3rd edition, (Clarendon Press: Oxford, 1965), p. 403.
9. M. H. Ross and G. L. Shaw, Ann. Phys. (New York) 13, 147 (1961).
10. B. H. Bransden, Atomic Collision Theory (Benjamin; New York, 1970), pp. 155-60; see also ref. 8, pp. 377-85.
11. A. L. Sinfailam and R. K. Nesbet, Phys. Rev. A6, 2118 (1972); Bull. Am. Phys. Soc. 17, 1133 (1972).
12. R. S. Oberoi and R. K. Nesbet, Bull. Am. Phys. Soc. 17, 1133 (1972).
13. J. D. Lyons, R. K. Nesbet, C. C. Rankin and A. C. Yates, to be published.
14. D. W. Norcross, J. Phys. B4, 1458 (1971).
15. P. G. Burke and A. J. Taylor, J. Phys. B2, 869 (1969).
16. D. Andrick, M. Eyb and M. Hofmann, J. Phys. B5, L15 (1972).
17. W. Gehenn and E. Reichert, Zeit. f. Phys. 254, 28 (1972).
18. D. W. Norcross and D. L. Moores, in Atomic Physics III, Proc. of the 3rd Int. Conf. on Atomic Physics, Boulder, 1972 (Plenum Press, New York).
19. T. A. Patterson, H. Hotop and W. C. Lineberger, Bull. Am. Phys. Soc. 17, 1128 (1972).




Figure 1. Partial wave contributions to the elastic scattering cross section in electron collisions with lithium atoms at energies close to the excitation threshold, E_{th} , which is calculated to be 1.8411 eV.