

*File Copy*

BRL R 1652

# BRL

AD763895

REPORT NO. 1652

CROSS SECTIONS FOR FORWARD-ANGLE ELASTIC  
SCATTERING OF 7 TO 10 MeV NEUTRONS FROM  
CARBON, NITROGEN, AND OXYGEN

by

William P. Bucher  
Clinton E. Hollandsworth  
Andrus Niiler  
David McNatt  
James E. Youngblood

June 1973

Approved for public release; distribution unlimited.

USA BALLISTIC RESEARCH LABORATORIES  
ABERDEEN PROVING GROUND, MARYLAND

Destroy this report when it is no longer needed.  
Do not return it to the originator.

Secondary distribution of this report by originating  
or sponsoring activity is prohibited.

Additional copies of this report may be obtained  
from the National Technical Information Service,  
U.S. Department of Commerce, Springfield, Virginia  
22151.

The findings in this report are not to be construed as  
an official Department of the Army position, unless  
so designated by other authorized documents.

BALLISTIC RESEARCH LABORATORIES

REPORT NO. 1652

JUNE 1973

CROSS SECTIONS FOR FORWARD-ANGLE ELASTIC SCATTERING OF 7 to 10 MeV  
NEUTRONS FROM CARBON, NITROGEN, AND OXYGEN

William P. Bucher  
Clinton E. Hollandsworth  
Andrus Niiler  
David McNatt  
James E. Youngblood

Approved for public release; distribution unlimited.

This work was supported by the Defense Nuclear Agency under  
DNA Subtask W99QAXPC101  
and in part by  
RDT&E Project No. 1W062117AD52

ABERDEEN PROVING GROUND, MARYLAND

B A L L I S T I C   R E S E A R C H   L A B O R A T O R I E S

REPORT NO. 1652

WPBucher/CEHollandsworth/  
ANiiler/DMcNatt/JEYoungblood/rr  
Aberdeen Proving Ground, Md.  
June 1973

CROSS SECTIONS FOR FORWARD-ANGLE ELASTIC SCATTERING OF 7 TO 10 MeV  
NEUTRONS FROM CARBON, NITROGEN, AND OXYGEN

ABSTRACT

Absolute differential cross sections for the elastic scattering of 7.40, 7.55, 8.00, 8.55, 9.00, and 9.50 MeV neutrons from carbon, nitrogen, and oxygen in the angular range from  $2.5^\circ$  to  $15^\circ$  have been determined. Measurements were made relative to the scattering from lead by use of a special-purpose collimator. The cross section scale was then established by absolute measurements for lead in a ring geometry experiment. The results for nitrogen are consistent with recent measurements at other laboratories which suggest that the discrepancy in previous nitrogen cross section data around 7-8 MeV neutron energy arose from an underestimate of the elastic scattering cross section. The data presented in this report represent the first direct measurements of neutron scattering cross sections for light elements at very small scattering angles.




## TABLE OF CONTENTS

	Page
ABSTRACT. . . . .	3
LIST OF FIGURES . . . . .	7
I. INTRODUCTION. . . . .	9
II. EXPERIMENTAL PROCEDURE. . . . .	9
III. RESULTS AND DISCUSSION. . . . .	12
A. Carbon. . . . .	12
B. Nitrogen. . . . .	17
C. Oxygen. . . . .	17
IV. SUMMARY . . . . .	17
ACKNOWLEDGEMENTS. . . . .	21
REFERENCES. . . . .	22
DISTRIBUTION LIST . . . . .	25



## LIST OF FIGURES

Figure	Page
1. Schematic Diagram of Small-Angle Collimator . . . . .	11
2. Forward-angle Scattering Cross Sections for Carbon. . . . .	16
3. Forward-angle Scattering Cross Sections for Nitrogen. . . . .	18
4. Forward-angle Scattering Cross Sections for Oxygen. . . . .	19



## I. INTRODUCTION

A technique, which utilizes a neutron collimator assembly designed to provide shielding along the entire path length of scattered neutrons from scatterer to detector, has been developed for measurements of the forward-angle elastic scattering of fast neutrons.

In a previous report<sup>1\*</sup> we described the experimental method and presented results for scattering of 7.55 and 9.50 MeV neutrons from carbon, nitrogen, and oxygen. These experiments were motivated by the desire to provide experimental data bearing on the resolution of an existing discrepancy<sup>2</sup> in the nitrogen cross sections in the 7 to 10 MeV range of neutron energies. The results of air transport calculations<sup>3</sup> for neutrons are sensitive to errors in the input cross sections of the magnitude and nature of those implied by the discrepancy. Our results<sup>1</sup> for N at 7.55 MeV indicated that the 0° cross section was approximately 40% higher than the minimum value given by the Wick limit and implied that previous measurements of the elastic scattering probably yielded an underestimate of the total cross section for elastic scattering.

The new measurements reported here were initiated primarily to provide additional data concerning the discrepancy. Since initiating the present work, new measurements at large angles ( $15^\circ \leq \theta \leq 160^\circ$ ) between 7 and 10 MeV neutron energy have been reported.<sup>4,5</sup> The results of these recent measurements are consistent with our results and produce integrated elastic cross sections approximately 20% higher than the earlier elastic scattering data<sup>6</sup> in the 7 to 9 MeV range of neutron energies.

## II. EXPERIMENTAL PROCEDURE

The measurements were carried out at the BRL tandem Van de Graaff facility. Deuterons from a direct-extraction diode source were pulsed at a repetition rate of 2 MHz and klystron bunched. The overall time resolution for the detection of elastically scattered neutrons was approximately two nanoseconds. After acceleration, the pulsed deuteron beam was used to produce neutrons through the  $D(d,n)^3\text{He}$  reaction in a  $\text{D}_2$  gas cell with a thin (3.5 or 5 $\mu$ ) Mo entrance window. The cell length was varied from 2 cm to 3.8 cm during the course of these measurements. Neutron production was monitored by observation of the neutron time-of-flight spectrum in a scintillation detector positioned approximately 1.5 meters from the source.

The experimental arrangement is shown in Figure 1. A scattering run with the sample in the scattering position (see Figure 1) is followed by a background run, for the same number of monitor counts, with the sample in the attenuating position. The scattering samples are in the form of discs 26 cm in diameter. The liquid samples,  $\text{N}_2\text{H}_4$  and  $\text{H}_2\text{O}$  are confined

---

\* References are listed on page 22.

in thin-walled stainless steel containers. Scattered neutrons are detected in NE-213 scintillators. Standard neutron time-of-flight techniques, including neutron-gamma ray discrimination, are used for background suppression. A general description of the experimental techniques has been given in a previous report.<sup>1</sup>

All scattering measurements with the collimator of Figure 1 are made relative to the scattering from lead. The absolute cross sections for lead are then determined by ring geometry measurements.<sup>7</sup> This approach is particularly feasible due to the high scattering cross section for lead, approximately 7 barns/sr. at 3°, a factor of ten or more higher than those of carbon, nitrogen, or oxygen. The excellent counting statistics of the lead data taken with the collimator introduce only a small uncertainty in the final data.

In the ring geometry measurements, the signal to background is sufficiently high to allow good counting statistics. However, it is necessary to correct these data for systematic errors due to air scattering by performing similar measurements with polyethylene (CH<sub>2</sub>) rings. The final data for lead are also corrected for multiple scattering and, to the second order, for the finite sizes of the source, scatterer, and detector. The angles for which measurements were made correspond closely to those measured with the collimator so that errors due to either interpolation or extrapolation are small.

The lead data at five energies (7.55, 8.00, 8.55, 9.00, and 9.50 MeV) were compared with the predictions of the Perey-Buck optical model<sup>8</sup> using the equivalent local potentials given by Wilmore and Hodgson.<sup>9</sup> While the agreement is found to be reasonably good, it is noted that the measured angular dependence at each energy is slightly more forward peaked than that predicted by these optical model parameters. However, the energy dependence is correct; this is verified by the fact that when the data are expressed in terms of fractional deviation from the optical model values the resulting plot is seen to be energy independent. Thus, optical model predictions can be used to correlate the data from various energies and thereby yield more accurate results than by considering the data at each energy separately. This represents an improvement in the normalization used in our previously reported<sup>1</sup> C, N, and O cross section data at 7.55 and 9.50 MeV, and for this reason the results are revised and included in the following tables and graphs along with the present results.

Because measurements were made for all three elements at each incident neutron energy, the energy spread frequently encompassed regions of structure in the total cross sections for one or more elements. Further, the mean neutron energy was uncertain to about 30 keV. To minimize uncertainties in the final results, an auxiliary measurement was performed; the transmission of each sample was measured under conditions identical to those of the differential scattering measurements, e.g., energy, energy spread, and same sample discs (the center channel of the




Figure 1. Schematic Diagram of Small-Angle Collimator.

collimator is used for this purpose).

### III. RESULTS AND DISCUSSION

The results of measurements of the forward-angle elastic scattering from carbon, nitrogen, and oxygen at  $7.40 \pm 0.13$ ,  $7.55 \pm 0.06$ ,  $8.00 \pm 0.12$ ,  $8.55 \pm 0.07$ ,  $9.00 \pm 0.10$ , and  $9.50 \pm 0.05$  MeV are presented in Tables I-III. The angular resolutions of the measurements were  $\pm 0.7$ ,  $\pm 0.9$ ,  $\pm 1.2$ , and  $\pm 1.5$  degrees for the scattering angles 2.6, 6.6, 10.4, and 14.2 degrees, respectively. The uncertainty in the mean neutron energy is estimated to be 30 keV.

The data are corrected for multiple scattering effects<sup>10</sup> and finite geometry effects are taken into account. The final uncertainty in the oxygen and nitrogen data was magnified by the use of compounds as scattering samples ( $H_2O$  and  $N_2H_4$ ). As an independent check of these data, scattering measurements were also carried out with a  $CH_2$  sample. The results inferred for carbon are in good statistical agreement with that obtained by the use of a graphite sample (or conversely, the correct values for the hydrogen cross section are inferred with the C and  $CH_2$  data). Included in the assigned standard deviations, in addition to the counting statistics, are the uncertainties in the differential cross section for lead (used to normalize the data) and in the transmission of each sample averaged over the energy spread (used to infer the final results from the data).

#### A. Carbon

Differential cross sections for the scattering of 7.40 to 9.50 MeV neutrons from carbon are shown in Figure 2. In this figure (and in succeeding ones for other elements), the energy-averaged Wick's limit<sup>11</sup> is indicated by an arrow along the vertical axis. Total cross sections used to calculate this value were taken from ENDF/B files, and the square of the cross section was then averaged over the appropriate energy interval. In Table I this limit is given as well as the extrapolated  $\sigma(0^\circ)$  as calculated from a least squares fit of the experimental data to the equation  $\ln\sigma(\theta) = a - m(1 - \cos\theta)$ , where  $a$  and  $m$  are constant. The solid curves give the prediction of ENDF/B evaluation 1165 for carbon,<sup>12</sup> appropriately averaged over the experimental energy spread. With the exception of the 8.55 and 9.00 MeV data, which agree well with the evaluation, the measurements are higher than the evaluated curve at all energies. As noted by Perey and Kinney,<sup>13</sup> the carbon evaluation was based on rather meager experimental data. This fact coupled with the strongly resonant nature of fast neutron scattering from carbon suggest that an adequate evaluation of neutron interactions with carbon cannot be expected without several additional experimental measurements.

Our measurements at 7.55 and at 7.40 MeV are in excellent agreement with results ORNL for  $7.54 \pm 0.06$  MeV at large angles. The measured cross

Table I. Cross Sections for Forward-Angle Scattering From Carbon.

$\theta_{\text{c.m.}}$	$\frac{7.40 \pm 0.13}{\sigma_{\text{c.m.}} \text{ (mb/sr)}}$	% error	$\theta_{\text{c.m.}}$	$\frac{7.55 \pm 0.06}{\sigma_{\text{c.m.}} \text{ (mb/sr)}}$	% error
2.9°	668	7	2.8°	690	8
7.4°	641	5	7.2°	604	5
11.3°	583	4	11.7°	596	4
15.4°	518	4	15.4°	534	4
$\sigma(0^\circ)^a$	678		$\sigma(0^\circ)$	658	
$W_L^b$	545		$W_L$	559	
$\sigma(0^\circ)/W_L$	1.24		$\sigma(0^\circ)/W_L$	1.18	
$\theta_{\text{c.m.}}$	$\frac{8.00 \pm 0.12}{\sigma_{\text{c.m.}} \text{ (mb/sr)}}$	% error	$\theta_{\text{c.m.}}$	$\frac{8.55 \pm 0.07}{\sigma_{\text{c.m.}} \text{ (mb/sr)}}$	% error
2.9°	998	8	2.9°	360	8
7.4°	880	5	7.4°	396	5
11.7°	821	5	11.4°	320	5
15.3°	756	4	15.4°	302	4
$\sigma(0^\circ)$	953		$\sigma(0^\circ)$	393	
$W_L$	688		$W_L$	244	
$\sigma(0^\circ)/W_L$	1.39		$\sigma(0^\circ)/W_L$	1.61	
$\theta_{\text{c.m.}}$	$\frac{9.00 \pm 0.10}{\sigma_{\text{c.m.}} \text{ (mb/sr)}}$	% error	$\theta_{\text{c.m.}}$	$\frac{9.50 \pm 0.05}{\sigma_{\text{c.m.}} \text{ (mb/sr)}}$	% error
2.9°	335	7	2.8°	456	7
7.4°	329	5	7.2°	446	5
11.4°	311	5	11.7°	414	4
15.4°	269	4	15.4°	354	4
$\sigma(0^\circ)$	347		$\sigma(0^\circ)$	475	
$W_L$	288		$W_L$	374	
$\sigma(0^\circ)/W_L$	1.20		$\sigma(0^\circ)/W_L$	1.27	

<sup>a</sup>Zero degree differential cross section determined by extrapolation of experimental data.

<sup>b</sup>Minimum value of zero degree cross section determined from Wick's Limit.

Table II. Cross Sections for Forward-Angle Scattering from Nitrogen.

$7.40 \pm 0.13$			$7.55 \pm 0.06$		
$\theta_{c.m.}$	$\sigma_{c.m.} \text{ (mb/sr)}$	% error	$\theta_{c.m.}$	$\sigma_{c.m.} \text{ (mb/sr)}$	% error
2.9°	545	13	2.8°	575	14
7.3°	507	9	7.1°	526	11
11.1°	492	6	11.6°	538	8
15.2°	433	8	15.2°	402	7
$\sigma(0^\circ)^a$	546		$\sigma(0^\circ)$	609	
$W_L^b$	486		$W_L$	426	
$\sigma(0^\circ)/W_L$	1.12		$\sigma(0^\circ)/W_L$	1.43	
$8.00 \pm 0.12$			$8.55 \pm 0.07$		
$\theta_{c.m.}$	$\sigma_{c.m.} \text{ (mb/sr)}$	% error	$\theta_{c.m.}$	$\sigma_{c.m.} \text{ (mb/sr)}$	% error
2.9°	606	12	2.9°	438	13
7.3°	526	9	7.3°	388	10
11.6°	458	8	11.2°	358	8
15.1°	389	8	15.2°	313	9
$\sigma(0^\circ)$	594		$\sigma(0^\circ)$	427	
$W_L$	444		$W_L$	360	
$\sigma(0^\circ)/W_L$	1.34		$\sigma(0^\circ)/W_L$	1.19	
$9.00 \pm 0.10$			$9.50 \pm 0.05$		
$\theta_{c.m.}$	$\sigma_{c.m.} \text{ (mb/sr)}$	% error	$\theta_{c.m.}$	$\sigma_{c.m.} \text{ (mb/sr)}$	% error
2.9°	413	13	2.8°	476	10
7.3°	398	10	7.1°	401	10
11.2°	335	8	11.6°	418	7
15.2°	262	9	15.2°	378	8
$\sigma(0^\circ)$	436		$\sigma(0^\circ)$	455	
$W_L$	382		$W_L$	414	
$\sigma(0^\circ)/W_L$	1.14		$\sigma(0^\circ)/W_L$	1.10	

<sup>a</sup>Zero degree differential cross section determined by extrapolation of experimental data.

<sup>b</sup>Minimum value of zero degree cross section determined from Wick's Limit.

Table III. Cross Sections for Forward-Angle Scattering From Oxygen.

$\theta_{\text{c.m.}}$	$\frac{7.40 \pm 0.13}{\sigma_{\text{c.m.}} \text{ (mb/sr)}}$	% error	$\theta_{\text{c.m.}}$	$\frac{7.55 \pm 0.06}{\sigma_{\text{c.m.}} \text{ (mb/sr)}}$	% error
2.8°	262	22	2.8°	325	16
7.2°	289	14	7.2°	271	14
11.1°	295	11	11.5°	249	14
15.1°	248	11	15.1°	164	14
$\sigma(0^\circ)^a$	299		$\sigma(0^\circ)$	331	
$W_L^b$	328		$W_L$	258	
$\sigma(0^\circ)/W_L$	0.91		$\sigma(0^\circ)/W_L$	1.28	
$\theta_{\text{c.m.}}$	$\frac{8.00 \pm 0.12}{\sigma_{\text{c.m.}} \text{ (mb/sr)}}$	% error	$\theta_{\text{c.m.}}$	$\frac{8.55 \pm 0.07}{\sigma_{\text{c.m.}} \text{ (mb/sr)}}$	% error
2.8°	317	18	2.8°	246	21
7.2°	252	15	7.2°	314	12
11.4°	239	13	11.2°	231	12
14.9°	164	15	15.1°	241	10
$\sigma(0^\circ)$	315		$\sigma(0^\circ)$	290	
$W_L$	195		$W_L$	278	
$\sigma(0^\circ)/W_L$	1.62		$\sigma(0^\circ)/W_L$	1.04	
$\theta_{\text{c.m.}}$	$\frac{9.00 \pm 0.10}{\sigma_{\text{c.m.}} \text{ (mb/sr)}}$	% error	$\theta_{\text{c.m.}}$	$\frac{9.50 \pm 0.05}{\sigma_{\text{c.m.}} \text{ (mb/sr)}}$	% error
2.8°	344	15	2.8°	375	13
7.2°	328	11	7.0°	348	11
11.2°	332	8	11.5°	309	9
15.1°	218	10	15.1°	249	11
$\sigma(0^\circ)$	388		$\sigma(0^\circ)$	384	
$W_L$	360		$W_L$	385	
$\sigma(0^\circ)/W_L$	1.08		$\sigma(0^\circ)/W_L$	1.00	

<sup>a</sup>Zero degree differential cross section determined by extrapolation of experimental data.

<sup>b</sup>Minimum value of zero degree cross section determined from Wick's Limit.

# CARBON




Figure 2. Forward-angle Scattering Cross Sections for Carbon.

sections for  $15^\circ$  scattering at 7.54 and 7.55 MeV agree within 4%. Our 8.55 MeV results for carbon are somewhat lower than the ORNL data at 8.56 MeV; however, in general we obtain results consistent with those of Perey and Kinney (see succeeding discussions).

## B. Nitrogen

Our nitrogen results are shown in Figure 3. The solid curves were computed from the recent evaluation of nitrogen data by Young and Foster, ENDF/B MAT 1133.<sup>14</sup> The evaluation is in good agreement with our measurements around 7 and 8 MeV; whereas, the evaluated curve tends to be slightly higher than the measured one for the higher energies. Our results at 7.55 (and at 7.40) join smoothly with the large-angle data of Perey and Kinney.<sup>15</sup>

## C. Oxygen

Differential cross sections for forward-angle scattering of 7 to 10 MeV neutrons from oxygen are compared with the Young and Foster evaluation<sup>16</sup> of oxygen data (MAT 1134) in Figure 4. The data are well described by the evaluation except for the 9.50 MeV data which are somewhat lower than the evaluated curve. Our measurements near  $15^\circ$  at 7.55 and 8.55 are consistent with the measurements of Kinney and Perey<sup>17</sup> at 7.54 and 8.56 for approximately the same energy spreads.

## IV. SUMMARY

The data presented in this report represent the first direct measurements of neutron cross sections for scattering from light elements at very small angles. These results are of special importance in the case of nitrogen because of the discrepancy in the nitrogen cross sections discussed in Ref. 2. Briefly stated, the elastic scattering cross section obtained by subtracting the measured nonelastic cross sections from the total was approximately 200 mb higher than the elastic cross section obtained by direct measurement.<sup>6</sup>

In their evaluation,<sup>14</sup> Young and Foster assigned the 200 mb of cross section to the elastic scattering cross section. Their choice was influenced in part by our previous measurements<sup>1</sup> of small-angle scattering which indicated that the differential cross section at zero degrees exceeded Wick's limit by about 40% around 7-8 MeV. The data of Bauer, *et al.*<sup>6</sup> would not predict forward cross sections in excess of Wick's limit. A second consideration was that the assignment of the 200 mb to the elastic cross section assumed that only one measurement was in error, that of Bauer *et al.* Conversely, the assignment of this 200 mb contribution to the nonelastic cross section would have implied that several independent measurements of the nonelastic cross section were in error. Our results at small angles and the large-angle data from ORNL




Figure 3. Forward-angle Scattering Cross Sections for Nitrogen.




Figure 4. Forward-angle Scattering Cross Sections for Oxygen.

and TNC tend to confirm the choice made by Young and Foster in their evaluation. Furthermore, the results presented here demonstrate that, even with energy resolutions typical of those used in elastic scattering measurements, the zero degree cross sections for light elements appreciably exceed, in some instances, the value given by an energy-averaged Wick's limit. (See Reference 1 for discussion of energy averaging of Wick's inequality.)

#### ACKNOWLEDGEMENTS

We wish to thank L. Stewart, P. Young, and D. G. Foster, Jr. for several helpful discussions. In addition, we gratefully acknowledge the contributions of R. Juckens, W. B. Beverly, and J. A. Morrissey to this work.

## REFERENCES

1. W. P. Bucher, C. E. Hollandsworth, R. D. Lamoreaux, A. Niiler, R. R. Sankey, and D. Eccleshall, Small-Angle Elastic Scattering of 7.55 and 9.50 MeV Neutrons From C, N, and O. BRL Report No. 1562, 1971. (AD #734877)
2. J. K. Dickens and F. G. Perey, "The  $^{14}\text{N}$  (n,xy) Reaction for  $5.8 \leq E_n \leq 8.6$  MeV", Nucl. Sci. Eng. 36, 280 (1969).
3. Norman E. Banks, Neutron Cross Section Sensitivity Studies For Oxygen, Nitrogen, and Dry Air. BRL Report No. 1583, 1972. (AD #901087)
4. F. G. Perey and W. E. Kinney, "On A Possible Resolution of the Nitrogen Nonelastic Neutron Cross Section Discrepancy Below 9 MeV", Nucl. Sci. Eng. 46, 428 (1971).
5. D. O. Nellis and P. S. Buchanan, Neutron Scattering and Gamma-Ray Production Cross Sections for N, O, Al, Si, Ca, and Fe". DNA 2716, 1972.
6. P. W. Bauer, J. D. Anderson, H. F. Lutz, C. Wong, J. W. McClure and B. A. Pohl, "Elastic Scattering of 7 to 14 MeV Neutrons from Nitrogen", Nucl. Phys. A93, 673 (1967).
7. W. P. Bucher, C. E. Hollandsworth, and A. Niiler, BRL Report (to be published).
8. F. Perey and B. Buck, "A Non-local Potential Model For the Scattering of Neutrons by Nuclei", Nucl. Phys. 32, 353 (1962).
9. D. Wilmore and P. E. Hodgson, "The Calculation of Neutron Cross Sections from Optical Potentials", Nucl. Phys. 55, 673 (1964).
10. W. Beverly, Muse: A Multiple Scattering Effect Code for Infinite Slabs, BRL Report No. 1588, 1972. (AD #903008L)
11. G. C. Wick, Atti Acad. d'Itac. 13, 1203 (1943).
12. E. L. Slaggie and J. T. Reynolds, C<sup>12</sup> Fast Neutron Cross Sections and Legendre Moments Below 15 MeV, Knolls Atomic Power Laboratory Report KAPL 3099, 1966; See also, J. T. Reynolds, C. J. Slavik, C. R. Lubitz, and N. C. Francis, "Coupled-Channel Calculation of C<sup>12</sup> Neutron Scattering", Phys. Rev. 176, 1213 (1968).
13. F. G. Perey and W. E. Kinney, Carbon Neutron Elastic and Inelastic Scattering Cross Section from 4.5 to 8.5 MeV, Oak Ridge National Laboratory Report, ORNL-4441, 1969.

14. P. G. Young and D. G. Foster, Jr., An Evaluation of the Neutron and Gamma-Ray Production Cross Sections for Nitrogen, Los Alamos Scientific Laboratory Report, LA-4725, 1972.
15. F. G. Perey and W. E. Kinney, Private Communication, 1972.
16. D. G. Foster, Jr. and P. G. Young, A Preliminary Evaluation of the Neutron Cross Section of Oxygen, Los Alamos Scientific Laboratory Report, LA 4780, 1972.
17. W. E. Kinney and F. G. Perey, Neutron Elastic and Inelastic Scattering Cross Sections for Oxygen in the Energy Range 4.34 to 8.56 MeV. Oak Ridge National Laboratory Report, ORNL-4780, 1972.



DISTRIBUTION LIST

<u>No. of</u> <u>Copies</u>	<u>Organization</u>	<u>No. of</u> <u>Copies</u>	<u>Organization</u>
12	Commander Defense Documentation Center ATTN: TIPCR Cameron Station Alexandria, Virginia 22314	1	Commander U.S. Army Materiel Command ATTN: AMCRD-TP 5001 Eisenhower Avenue Alexandria, Virginia 22304
5	Director Defense Nuclear Agency ATTN: Tech Library (1 cy) RARP (4 cys) Washington, DC 20305	1	Commander U.S. Army Materiel Command ATTN: AMCRD-WN-RE 5001 Eisenhower Avenue Alexandria, Virginia 22304
1	Commander U.S. Army Materiel Command ATTN: AMCDL 5001 Eisenhower Avenue Alexandria, Virginia 22304	1	Commander U.S. Army Aviation Systems Command ATTN: AMSAV-E 12th and Spruce Streets St. Louis, Missouri 63166
1	Commander U.S. Army Materiel Command ATTN: AMCDMA, MG J.R. Guthrie 5001 Eisenhower Avenue Alexandria, Virginia 22304	1	Director U.S. Army Air Mobility Research and Development Laboratory Ames Research Center Moffett Field, California 94035
1	Commander U.S. Army Materiel Command ATTN: AMCRD, MG S. C. Meyer 5001 Eisenhower Avenue Alexandria, Virginia 22304	2	Commander U.S. Army Electronics Command ATTN: AMSEL-RD AMSEL-DL Fort Monmouth, New Jersey 07703
1	Commander U.S. Army Materiel Command ATTN: AMCRD, Dr. J.V.R. Kaufman 5001 Eisenhower Avenue Alexandria, Virginia 22304	1	Commander U.S. Army Missile Command ATTN: AMSMI-R Redstone Arsenal, Alabama 35809
1	Commander U.S. Army Materiel Command ATTN: AMCRD-TE 5001 Eisenhower Avenue Alexandria, Virginia 22304	1	Commander U.S. Army Tank Automotive Command ATTN: AMSTA-RHFL Warren, Michigan 48090

DISTRIBUTION LIST

<u>No. of Copies</u>	<u>Organization</u>	<u>No. of Copies</u>	<u>Organization</u>
2	Commander U.S. Army Mobility Equipment Research & Development Center ATTN: Tech Docu Cen, Bldg. 315 AMSME-RZT Fort Belvoir, Virginia 22060	2	HQDA (DARD-MSN-N; DARD-ARP Dr. R. Watson) Washington, DC 20310
1	Commander U.S. Army Munitions Command ATTN: AMSMU-RE Dover, New Jersey 07801	1	Commander U.S. Army Research Office (Durham) ATTN: RDRD-P, Dr. Mace Box CM, Duke Station Durham, North Carolina 27706
2	Commander U.S. Army Weapons Command ATTN: AMSWE-RE AMSWE-RDF Rock Island, Illinois 61202	4	Commander US Naval Air Systems Command ATTN: AIR-604 (3 cys) AIR-52023, Nuc Appl Br Washington, DC 20360
1	Director U.S. Army Advanced Materiel Concepts Agency 2461 Eisenhower Avenue Alexandria, Virginia 22314	4	Commander U.S. Naval Ordnance Systems Command ATTN: ORD-0632 ORD-035 ORD-5524 ORD-0523 Washington, DC 20360
2	Commander U.S. Army Harry Diamond Laboratories ATTN: AMXDO-TD/002 AMXDO-NP Washington, DC 20438	1	Director Argonne National Laboratory ATTN: Dr. A. B. Smith Reactor Physics Division 9700 South Cass Avenue Argonne, Illinois 60440
1	Commander U.S. Army Materials and Mechanics Research Center ATTN: AMXMR-ATL Watertown, Massachusetts 02172	1	Brookhaven National Laboratory National Neutron Cross Sec Ctr ATTN: Dr. Murray Goldberg P. O. Box 150 Upton, Long Island, New York 11973
1	Commander U.S. Army Natick Laboratories ATTN: AMXRE, Dr. D. Sieling Natick, Massachusetts 01762		

DISTRIBUTION LIST

<u>No. of</u> <u>Copies</u>	<u>Organization</u>	<u>No. of</u> <u>Copies</u>	<u>Organization</u>
3	Director Los Alamos Scientific Lab ATTN: Dr. Philip Young Dr. Donald Harris Dr. Leona Stewart P. O. Box 1663 Los Alamos, New Mexico 87544	1	Ohio University ATTN: Dr. R. O. Lane Dept of Physics Athens, Ohio 45701
2	Gulf Energy & Environmental Systems Inc. ATTN: Mgr, Tech Info Svcs for Dr. Martin Fricke Mr. Charles Preskitt P. O. Box 1111 San Diego, California 92112	1	University of California Lawrence Livermore Laboratory ATTN: Technical Info Division P. O. Box 808 Livermore, California 94550
2	Union Carbide Corporation Nuclear Division Oak Ridge National Laboratory ATTN: Dr. Francis Perey Mr. C. E. Clifford P. O. Box X Oak Ridge, Tennessee 37830		<u>Aberdeen Proving Ground</u>  Ch, Tech Lib Marine Corps In Ofc CDC In Ofc Dir, USAMSAA ATTN: Dr. Sperrazza

Unclassified

Security Classification

DOCUMENT CONTROL DATA - R & D

(Security classification of title, body of abstract and indexing annotation must be entered when the overall report is classified)

1. ORIGINATING ACTIVITY (Corporate author) US Army Ballistic Research Laboratories Aberdeen Proving Ground, Md. 21005		2a. REPORT SECURITY CLASSIFICATION NA	
		2b. GROUP	
3. REPORT TITLE CROSS SECTIONS FOR FORWARD-ANGLE ELASTIC SCATTERING OF 7 TO 10 MeV NEUTRONS FROM CARBON, NITROGEN, AND OXYGEN			
4. DESCRIPTIVE NOTES (Type of report and inclusive dates)			
5. AUTHOR(S) (First name, middle initial, last name) W. P. Bucher, C. E. Hollandsworth, A. Niiler, D. McNatt, J. E. Youngblood			
6. REPORT DATE JUNE 1973		7a. TOTAL NO. OF PAGES 27	7b. NO. OF REFS 17
8a. CONTRACT OR GRANT NO.		8a. ORIGINATOR'S REPORT NUMBER(S) BRL Report No. 1652	
b. PROJECT NO. DNA Subtask W99QAXPC101 RDT&E Proj. No. 1W062117AD52		8b. OTHER REPORT NO(S) (Any other numbers that may be assigned to this report)	
c.			
d.			
10. DISTRIBUTION STATEMENT Approved for public release; distribution unlimited.			
11. SUPPLEMENTARY NOTES		12. SPONSORING MILITARY ACTIVITY Defense Nuclear Agency and US Army Materiel Command Washington, DC	
13. ABSTRACT Absolute differential cross sections for the elastic scattering of 7.40, 7.55, 8.00, 8.55, 9.00, and 9.50 MeV neutrons from carbon, nitrogen, and oxygen in the angular range from 2.5° to 15° have been determined. Measurements were made relative to the scattering from lead by use of a special-purpose collimator. The cross section scale was then established by absolute measurements for lead in a ring geometry experiment. The results for nitrogen are consistent with recent measurements at other laboratories which suggest that the discrepancy in previous nitrogen cross section data around 7-8 MeV neutron energy arose from an underestimate of the elastic scattering cross section. The data presented in this report represent the first direct measurements of neutron scattering cross sections for light elements at very small scattering angles.			

14. KEY WORDS	LINK A		LINK B		LINK C	
	ROLE	WT	ROLE	WT	ROLE	WT
<p>Neutron scattering</p> <p>cross sections</p> <p>elastic scattering</p> <p>small-angles</p> <p>Carbon</p> <p>Nitrogen</p> <p>Oxygen</p>						