

WOODARD RESEARCH CORPORATION

ARO-11545.2-L

AD \_\_\_\_\_

INVESTIGATION OF THE PROLONGED RECOVERY TIME  
FOLLOWING BOTULINUM INTOXICATION - EFFICACY OF  
INTRAPERITONEAL VERSUS INTRAVENOUS INJECTION  
OF BOTULINUM ANTITOXIN

FINAL REPORT

by

James F. Novotny, Ph.D.

August 10, 1973

Supported By

U.S. ARMY RESEARCH OFFICE - DURHAM  
Box CM, Duke Station, Durham, North Carolina 27706

Contract No. DAHC 19-72-C-0011 P-00001  
Woodard Research Corporation  
Herndon, Virginia 22070

Reproduced by  
NATIONAL TECHNICAL  
INFORMATION SERVICE  
US Department of Commerce  
Springfield, VA. 22151

D D C  
R  
SEP 10 1973  
B

DOD DISTRIBUTION STATEMENT

Distribution of this document is unlimited.

AD 766955

AD-766 955

INVESTIGATION OF THE PROLONGED RECOVERY  
TIME FOLLOWING BOTULINUM INTOXICATION.  
EFFICACY OF INTRAPERITONEAL VERSUS INTRA-  
VENOUS INJECTION OF BOTULINUM ANTITOXIN

James F. Novotny

Woodard Research Corporation

Prepared for:

Army Research Office -Durham

10 August 1973

DISTRIBUTED BY:

**NTIS**

National Technical Information Service  
U. S. DEPARTMENT OF COMMERCE  
5285 Port Royal Road, Springfield Va. 22151

## DOCUMENT CONTROL DATA - R &amp; D

(Security classification of title, body of abstract and indexing annotation must be entered when the overall report is classified)

1. ORIGINATING ACTIVITY (Corporate author)		2a. REPORT SECURITY CLASSIFICATION	
Woodard Research Corporation 12310 Pinecrest Road Herndon, Virginia 22070		Unclassified	
3. REPORT TITLE		2b. GROUP	
Investigation of the Prolonged Recovery Time Following Botulinum Intoxication - Efficacy of Intraperitoneal versus Intravenous Injection of Botulinum Antitoxin			
4. DESCRIPTIVE NOTES (Type of report and inclusive dates)			
Final Report, July 10, 1972-July 10, 1973			
5. AUTHOR(S) (First name, middle initial, last name)			
James F. Novotny			
6. REPORT DATE	7a. TOTAL NO. OF PAGES	7b. NO. OF REFS	
10 August 1973	23	0	
8a. CONTRACT OR GRANT NO.	8c. ORIGINATOR'S REPORT NUMBER(S)		
DAHC 19-72-C-0011 P-00001	N/A		
b. PROJECT NO.	9d. OTHER REPORT NO(S) (Any other numbers that may be assigned this report)		
	N/A		
c.			
d.			
10. DISTRIBUTION STATEMENT			
Distribution of this document is unlimited.			
11. SUPPLEMENTARY NOTES		12. SPONSORING MILITARY ACTIVITY	
N/A		U.S. Army Research Office-Durham	
13. ABSTRACT			
<p>Studies to provide information concerning botulinum intoxication were conducted in an attempt to determine what parameters influence the prolonged recovery time following intoxication and if the routes of antitoxin treatment could influence the rates of recovery and decrease mortality in test species. (U)</p> <p>In order to study prolonged recovery times, mice were injected via the intramuscular route with sublethal concentrations of botulinum type A toxin. Proteolytic enzyme solutions were injected into the test site to determine if the removal of bound proteinacious toxin from neuromuscular junctions could influence paralytic recovery times. Parameters of observation were gait of animal, production of lesion at site of injection and activity of paralyzed and treated mice. Trypsin and protease at 1 mg concentrations appeared to markedly influence recovery whereas chymotrypsin, papain, ficin, pepsin and CaCl<sub>2</sub> did not significantly affect decrease of paralysis. (U)</p> <p>The efficacy of intraperitoneal versus intravenous injection of botulinum antitoxin was studied by initiating a state of paralysis with resulting death of groups of mice and prophylactically administering specific antitoxin by either the intravenous or intraperitoneal routes. There appeared to be no effect of route of antitoxin administration on protective action of antisera therapy. (U)</p>			

14. KEY WORDS	LINK A		LINK B		LINK C	
	ROLE	WT	ROLE	WT	ROLE	WT
Type A Cl. botulinum toxin Botulinum paralysis Botulinum therapy Enzyme therapy Botulinum prophylaxis Swiss-Webster mice Fort Detrick Inbreed mice						

WOODARD RESEARCH CORPORATION

INVESTIGATION OF THE PROLONGED RECOVERY TIME  
FOLLOWING BOTULINUM INTOXICATION - EFFICACY OF  
INTRAPERITONEAL VERSUS INTRAVENOUS INJECTION  
OF BOTULINUM ANTITOXIN

SUMMARY

Studies to provide information concerning botulinum intoxication were conducted in an attempt to determine what parameters influence the prolonged recovery time following intoxication and if the routes of antitoxin treatment could influence the rates of recovery and decrease mortality in test species.

In order to study prolonged recovery times, mice were injected via the intramuscular route with sublethal concentrations of botulinum type A toxin. Proteolytic enzyme solutions were injected into the test site to determine if the removal of bound proteinacious toxin from neuromuscular junctions could influence paralytic recovery times. Parameters of observation were gait of animal, production of lesion at site of injection and activity of paralyzed and treated mice. Trypsin and protease at 1 mg concentrations appeared to markedly influence recovery whereas chymotrypsin, papain, ficin, pepsin and  $\text{CaCl}_2$  did not significantly affect decrease of paralysis.

The efficacy of intraperitoneal versus intravenous injection of botulinum antitoxin was studied by initiating a state of paralysis with resulting death of groups of mice and prophylactically administering specific antitoxin by either the intravenous or intraperitoneal routes. There appeared to be no effect of route of antitoxin administration on protective action of antisera therapy.

## WOODARD RESEARCH CORPORATION

.1.

### INVESTIGATION OF THE PROLONGED RECOVERY TIME FOLLOWING BOTULINUM INTOXICATION

The course of botulinal intoxication following the binding of toxin to a specific site in the nerve fiber was studied by injecting mice with a sublethal paralyzing dose of botulinum type A toxin. Toxin which was not bound immediately following injection was neutralized intraperitoneal inoculation with specific antitoxin.

As reported in previous reports, testing criteria to determine the extent and severity of paralysis consisted of use of test mouse motion in an activity cage. As the mouse moves through beams of invisible light, activity is measured as counts per period of time.

The results of activity of mice following administration of toxin by the intramuscular, intraperitoneal and stomach intubation routes are presented in Table 1. Significant paralysis was observed following intramuscular injection of  $1/2$  LD<sub>50</sub> of toxin.

Following these initial experiments, tests were performed to determine if injection of various proteolytic enzymes could destroy botulinum toxin in vivo. In the previous experiments four mice were placed in the activity cage. Three mice per group were used in these experiments. One group of three received either two LD<sub>50</sub> of toxin in the left hind leg or PBS diluent. One hour later all animals received 0.1 ml of antitoxin and four hours later all animals received 0.1 ml of freshly prepared proteolytic enzyme containing 10 mg/ml in the

WOODARD RESEARCH CORPORATION

- 2 -

TABLE 1  
ACTIVITY OF MICE FOLLOWING ADMINISTRATION  
OF DIFFERENT AMOUNTS OF TOXIN - ACTIVITY  
PRESENTED AS PERCENT OF CONTROL GROUP

Route of Injection	Percent LD <sub>50</sub>	Day			
		1	4	11	15
Intra-muscular	1/2	45.96	34.16	64.91	82.48
	1/4	82.72	89.11	107.45	88.73
	1/8	94.68	91.62	101.94	89.18
Intraperitoneal	1/2	86.99	191.10	105.11	111.06
	1/4	285.11	436.84	159.56	158.07
	1/8	162.58	284.47	142.56	187.63
Stomach Intubation	1/2	29.84	12.20	106.79	137.62
	1/4	67.91	119.75	94.93	129.00
	1/8	90.97	136.03	173.28	158.79

same hind leg. Chymotrypsin, protease, trypsin and CaCl<sub>2</sub> (control) were used as saline solutions. The animals were observed for walk and stance, ability to walk down a vertical rod, and presence of enzyme-induced lesion or sore at the site of injection for two weeks. The results are presented in Tables 2, 3 and 4.

Tables 5 and 6 present activity cage results of mice treated with toxin to induce paralysis and then injected with proteolytic enzymes. This data demonstrated marked paralysis in all toxin-treated animals except those receiving protease and trypsin. Papain and ficin resulted in extensive tissue damage at the site of injection.

WOODARD RESEARCH CORPORATION

- 3 -

TABLE 2

VERTICAL ROD, GAIT AND STANCE, AND SORES AT THE  
INJECTION SITE AS A RESULT OF ENZYME TREATMENT

Treatment	Toxin Animals					Diluent Animals				
	Day					Day				
	1	3	7	11	16	1	3	7	11	16
Control										
Vert			+	++	+					+
Gait										
Sore										
Vert			+	++	+					
Gait	+									
Sore										
Vert				+						
Gait	+			+						
Sore										
Pepsin										
Vert	+	++	+							
Gait	++	+								
Sore										
Vert			+	+	+					
Gait	+	+	+							+
Sore										
Vert			+	+	+					
Gait	+	+								
Sore										
Trypsin										
Vert			+	+						
Gait	+			+						+
Sore	+			+						+
Vert	+	+	+	+	++					
Gait	+		+							++
Sore										
Vert		++	+	±						
Gait	+	+	+						++	+
Sore			+							+



WOODARD RESEARCH CORPORATION

- 4 -

TABLE 2 (Contd)

VERTICAL ROD, GAIT AND STANCE, AND SORES AT THE  
INJECTION SITE AS A RESULT OF ENZYME TREATMENT

Treatment	Toxin Animals					Diluent Animals				
	Day					Day				
	1	3	7	11	16	1	3	7	11	16
α Chymotrypsin										
Vert				+						
Gait	+					+				
Sore	+		+			+			+	
Vert										
Gait						+				
Sore	+		+							
Vert				+						
Gait	+					+				
Sore	+		+							
Papain										
Vert		++	+	+		+	+++		*	
Gait	++	+	+			+++	+++			
Sore	+		+			++	++			
Vert										
Gait	+					+				
Sore	+					+				
Vert	+	++		+	+					
Gait	++	+								
Sore	+		+							+
Ficin										
Vert		+++	+	+	+				+	
Gait	++	++								
Sore	+		+	+		+				+
Vert	+	+	+	+	+			++		
Gait	++							+		
Sore	+		+	+		+			+	+
Vert		+		+	+			+++	+	+
Gait	++	+	++			+++	+++	+	+	+
Sore	+		+	+		+	+	+	+	+

\* Leg badly crippled



WOODARD RESEARCH CORPORATION

- 6 -

TABLE 3

VERTICAL ROD, GAIT AND STANCE, AND SORES AT THE  
INJECTION SITE AS A RESULT OF ENZYME TREATMENT

NEOSTIGMINE INDUCED HYPERSENSITIVITY

Treatment	Toxin Animals			Diluent Animals		
	Vert	Gait	Sore	Vert	Gait	Sore
Control	+	+	+	-	-	-
Pepsin	+	+		-		
Trypsin	-					
α Chymotrypsin	-	-		-		
Papain	+	-		-		
Ficin	-					
Protease	-	-		-		
CaCl <sub>2</sub>	+	-				
Uninjected Control	-					







## WOODARD RESEARCH CORPORATION

- 10 -

TABLE 5  
PERCENT ACTIVITIES OF MICE PARALYZED  
WITH BOTULINUM TOXIN AND TREATED  
PROPHYLACTICALLY WITH PROTEOLYTIC ENZYMES

Treatment	Day				
	1	3	7	11	16
Control	50.1	48.6	73.1	68.1	49.4
Pepsin	37.2	41.2	46.1	39.1	53.0
Trypsin	88.1	80.4	81.9	86.2	115.8
Chymotrypsin	54.4	54.7	85.9	61.0	79.9
Papain	43.2	38.9	46.6	51.0	52.8
Ficin	74.6	57.0	68.8	86.2	71.5
Protease	66.8	121.8	110.7	121.4	94.2
CaCl <sub>2</sub>	43.2	33.9	51.5	53.1	57.5

## WOODARD RESEARCH CORPORATION

- 11 -

TABLE 6  
PERCENT ACTIVITIES OF MICE TREATED WITH  
DIFFERENT CONCENTRATIONS OF PROTEOLYTIC ENZYMES  
FOLLOWING BOTULINAL PARALYSIS

Treatment	Dosage Level mg	Percent Decrease of Control				
		Day				
		1	4	7	11	18
Control	-	61.6	63.8	80.7	82.5	78.2
Chymotrypsin	1	55.0	77.0	72.8	50.5	54.3
CaCl <sub>2</sub>	1	91.8	188.3	236.1	179.2	177.3
Protease	1	62.6	54.4	90.9	84.1	90.4
Protease	0.33	109.4	89.9	222.4	210.9	184.4
Protease	0.11	52.9	73.7	73.8	56.4	79.2
Trypsin	1	49.2	51.1	75.3	52.4	62.3
Trypsin	0.33	144.5	245.9	591.7	380.7	292.3
Trypsin	0.11	50.7	81.3	47.6	76.3	87.3



EFFICACY OF INTRAPERITONEAL VERSUS INTRAVENOUS INJECTION  
OF BOTULINUM ANTITOXIN

Studies conducted were designed to compare the difference in efficacy between intraperitoneal (IP) and intravenous (IV) injection of botulinum antitoxin following stomach intubation of type A botulinum toxin into Swiss-Webster mice. Mice were inoculated with 0.25 ml of toxin solution by the per os route by inserting a round end needle into the stomach of each mouse and expressing the toxin. Botulinum antitoxin (Wellcome Lot K 9710) was serially diluted in saline and 0.1 ml of each dilution was injected either by the IP or IV route into each toxin treated mouse. The results of initial studies are presented in Table 7. Marked differences in protection were not noted.

A second group of experiments was conducted in which greater dilutions of antitoxin were used in order to determine if the low death rates initially observed were due to high antitoxin concentrations. Again, a significant difference between the two routes was not observed. This data is presented in Table 8. In addition, species sex did not alter the results which were obtained as shown in Table 9.

A chance observation within the laboratory indicated mice which are fasted become more susceptible to botulinal intoxication. Table 10 presents data obtained when mice were fasted for 24 hours and inoculated with 10-fold dilutions of botulinum type A toxin by stomach intubation. A total of three trials was conducted. Two of three trials provides expected dose-response results.



## WOODARD RESEARCH CORPORATION

- 14 -

TABLE 8

INTRAPERITONEAL VERSUS INTRAVENOUS PROPHYLAXIS  
OF BOTULINAL INTOXICATION IN MICE  
USING DILUTE CONCENTRATIONS OF ANTITOXIN

Route of Antitoxin Admn.	Dilution of Antitoxin	Number of Deaths Post Injection (Hours)									Dead/ Total
		4	20	28	44	52	68	76	92	106	
IV	10 <sup>-5</sup>	0	1	1	0	0	0	0	2	0	4/20
	10 <sup>-4</sup>	0	0	0	0	0	0	0	0	0	0/20
	10 <sup>-3</sup>	0	1	0	0	0	0	0	0	0	1/20
	10 <sup>-2</sup>	0	1	0	0	0	0	0	0	0	1/20
IP	10 <sup>-5</sup>	0	2	0	0	0	0	0	0	0	2/20
	10 <sup>-4</sup>	0	0	0	0	0	0	0	0	0	0/20
	10 <sup>-3</sup>	0	0	0	0	0	0	0	0	0	0/20
	10 <sup>-2</sup>	0	0	0	0	0	0	0	0	0	0/20
Control	-	0	1	0	1	0	0	0	0	0	2/20

WOODARD RESEARCH CORPORATION

- 15 -

TABLE 9  
THE EFFECT OF SEX ON THE INTRAPERITONEAL  
VERSUS INTRAVENOUS PROPHYLAXIS  
OF BOTULINAL INTOXICATION IN MICE

Route of Antitoxin Admn.	Dilution of Antitoxin	Sex	Number of Deaths Post Inj. (Hours)					Dead/Total
			8	24	32	48	96	
IV	10 <sup>-5</sup>	20M*	0	1	0	3	0	4/20
	10 <sup>-4</sup>	20M	0	0	0	0	0	0/20
	10 <sup>-3</sup>	10M	0	1	0	0	0	1/10
		10F**	0	0	0	0	0	0/10
	10 <sup>-2</sup>	10M	0	1	0	0	1	2/10
		10F	0	0	0	1	0	1/10
IP	10 <sup>-5</sup>	20M	0	0	0	1	0	1/20
	10 <sup>-4</sup>	10M	0	1	0	1	0	2/10
		10F	0	1	0	1	0	2/10
	10 <sup>-3</sup>	10M	0	0	0	0	1	1/10
		10F	1	0	0	0	0	1/10
	10 <sup>-2</sup>	10M	0	0	0	1	0	1/10
	10F	0	0	0	0	0	0/10	
Control	-	3M	0	1	0	0	0	1/3
		12F	0	3	0	1	0	4/12

\* Males  
 \*\* Females

## WOODARD RESEARCH CORPORATION

- 16 -

TABLE 10MORTALITY OF MICE RECEIVING BOTULINUM TOXIN  
FOLLOWING REMOVAL FROM FEED FOR 24 HOURS

Botulinum Toxin Dilution	Death-Survivor Ratio		
	Trial 1	Trial 2	Trial 3
1:2	4/10	9/10	3/10
1:4	3/10	7/10	2/10
1:8	3/10	5/10	2/10
1:16	0/10	2/10	3/10
1:32	0/10	-	0/10
1:64	0/10	-	0/10

Following experiments to determine the potency of botulinum toxin by the stomach intubation route in fasted mice, tests were conducted in which mice intoxicated by this route were treated with antitoxin by the intraperitoneal route in order to determine if this treatment can indeed lessen the severity of symptoms or number of deaths. Three trials were conducted in which mice were inoculated by the stomach tube route and were injected by the intraperitoneal route at two hours with 0.1 ml of antitoxin (Wellcome Batch K 9710). The mice were observed daily for five days and deaths and/or surviving animals were recorded. The results of these studies are presented in Table 11.

## WOODARD RESEARCH CORPORATION

- 17 -

TABLE 11  
MORTALITY OF MICE INJECTED WITH BOTULINUM TOXIN  
AND TREATED INTRAPERITONEALLY WITH  
BOTULINAL ANTITOXIN

Botulinum Toxin Dilution	Death-Survivor Ratio		
	Trial 1	Trial 2	Trial 3
10 <sup>-1</sup>	2/20	-	-
10 <sup>-2</sup>	3/20	-	-
10 <sup>-3</sup>	4/20	4/20	4/10
10 <sup>-4</sup>	7/20	2/20	1/10
10 <sup>-5</sup>	-	7/20	1/10
10 <sup>-6</sup>	-	10/20	1/10
10 <sup>-7</sup>	-	-	2/10

Antisera therapy via this route appeared to be extremely variable and at the recommendation of the scientific technical representative, Dr. Carl LaManna, experiments in this area of study were terminated.

James F. Novotny, Ph.D.  
Microbiologist

Submitted: August 10, 1973

WOODARD RESEARCH CORPORATION

In conducting the research described in this report, the Investigator(s) adhered to the "Guide for Laboratory Animal Facilities and Care," as promulgated by the Committee on the Guide for Laboratory Animal Resources, National Academy of Sciences, National Research Council.

Contract No. DAHC 19-72-C-0011 P-00001

Woodard Research Corporation  
12310 Pinecrest Road  
Herndon, Virginia 22070