

AD-770 782

PRESENCE OF ERABUTOXINS A AND B IN
VENOM OF THE SEA SNAKE LATICAUDA SEMI-
FASCIATA FROM TAIWAN

C.S. Kao, et al

Naval Medical Research Unit No. 2
Taipei, Taiwan


1973

DISTRIBUTED BY:

NTIS

**National Technical Information Service
U. S. DEPARTMENT OF COMMERCE
5285 Port Royal Road, Springfield Va. 22151**

DISCLAIMER NOTICE



THIS DOCUMENT IS BEST QUALITY AVAILABLE. THE COPY FURNISHED TO DTIC CONTAINED A SIGNIFICANT NUMBER OF PAGES WHICH DO NOT REPRODUCE LEGIBLY.



BIOCHEMISTRY DEPARTMENT

R. QUENTIN BLACKWELL, PH.D., HEAD

A 26

ADMINISTRATIVE INFORMATION

THIS WORK WAS SUPPORTED BY
THE BUREAU OF MEDICINE AND
SURGERY, DEPARTMENT OF THE
NAVY, WASHINGTON, D. C.

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

.....

R. H. WATTEN
CAPT MC USN
COMMANDING OFFICER

III

UNCLASSIFIED

Security Classification

AD 770782

DOCUMENT CONTROL DATA - R & D		
<i>Security classification of title, body of abstract and indexing annotation must be entered when the overall report is classified</i>		
1. ORIGINATING ACTIVITY (Corporate author) U. S. NAVAL MEDICAL RESEARCH UNIT No. 2 Box 14, APO SAN FRANCISCO 96263		2b. REPORT SECURITY CLASSIFICATION UNCLASSIFIED
3. REPORT TITLE SHORT COMMUNICATION PRESENCE OF ERABUTOXINS A AND B IN VENOM OF THE SEA SNAKE <u>LATICAUDA SEMIFASCIATA</u> FROM TAIWAN		2c. GROUP
4. DESCRIPTIVE NOTES (Type of report and inclusive dates) TECHNICAL REPORT		
5. AUTHOR(S) (First name, middle initial, last name) C.-S. KAO, C.-S. LIU AND R. Q. BLACKWELL		
6. REPORT DATE 1973	7a. TOTAL NO. OF PAGES 3	7b. NO. OF REFS 7
8a. CONTRACT OR GRANT NO.	9a. ORIGINATOR'S REPORT NUMBER(S) NAMRU-2-TR-571	
b. PROJECT NO.	9b. OTHER REPORT NO(S) (Any other numbers that may be assigned this report)	
c.		
d.		
10. DISTRIBUTION STATEMENT DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED		
11. SUPPLEMENTARY NOTES PUBLISHED IN TOXICON 11: 383-385, 1973		12. SPONSORING MILITARY ACTIVITY BUREAU OF MEDICINE AND SURGERY DEPARTMENT OF THE NAVY WASHINGTON, D. C. 20372
13. ABSTRACT		

Reproduced by
**NATIONAL TECHNICAL
 INFORMATION SERVICE**
 U.S. Department of Commerce
 Springfield VA 22151

I

UNCLASSIFIED
Security Classification

10 KEY WORDS	LINK A		LINK B		LINK C	
	ROLE	WT	ROLE	WT	ROLE	WT
ERABUTOXINS A AND B IN TAIWAN <u>L. SEMIFASCIATA</u>						
<i>II</i>						

DD FORM 1 NOV 66 1473 (BACK)
(PAGE 2)

UNCLASSIFIED
Security Classification

SHORT COMMUNICATION

PRESENCE OF ERABUTOXINS a AND b IN VENOM OF THE SEA SNAKE *LATICAUDA SEMIFASCIATA* FROM TAIWAN*

C.-S. KAO, C.-S. LIU and R. Q. BLACKWELL†

Department of Biochemistry, U.S. Naval Medical Research Unit No. 2, Taipei, Taiwan,
Republic of China

(Accepted for publication 5 March 1973)

TWO PRINCIPAL toxins from venom of the sea snake *Laticauda semifasciata* have been purified and their amino acid compositions studied previously. TAMIYA and ARAI (1966), working with venom obtained from sea snakes collected in Okinawan waters, called the two toxins Erabutoxins a and b; amino acid compositions and sequences of the two compounds were reported by SATO and TAMIYA (1971). TU *et al.* (1971) obtained two similar toxins, called Toxins a and b, from the same species of *Laticauda* collected in Philippine waters; results of their amino acid analyses are somewhat different from those of Sato and Tamiya. In the present paper we report results of studies made on the two principal toxins obtained from venom of *Laticauda semifasciata* collected in Taiwan. As summarized below, our results are in complete agreement with those of Sato and Tamiya.

Sea snakes used in this study were collected in the vicinity of Pescadore Islands in the Formosan Strait west of Taiwan and near Green Island and Orchid Island which are near the southeast end of Taiwan. Venom was collected by milking live snakes and was pooled for analysis. Methods used for venom fractionation and further chemical characterizations were essentially those reported in our previous study of *Hydrophis cyanocinctus* (LIU *et al.*, 1973).

Protein fractions obtained by chromatographic separation of venom components are shown in Fig. 1. Fractions IV, V and VI had relatively high lethality as measured by screening tests in mice. For example, 10 µg quantities of each fraction, injected intraperitoneally (in 0.2 ml of saline) into 20 g mice caused death in all animals within 30 min; five mice were used to test each fraction at the 10 µg level. Higher amounts of all three fractions also killed all animals tested, while studies were not made at lower levels.

Fraction IV was not investigated further; from its chromatographic behavior, it appears to be identical with the minor fraction, Erabutoxin c, described by TAMIYA (1972) and TAMIYA and ABE (1972). Fractions V and VI were desalted with a Sephadex G-15 column and purified by column chromatography on CM 52 carboxymethylcellulose using an ammonium acetate buffer system. Portions of purified fractions were lyophilized, carboxy-

* This work was supported by the Bureau of Medicine and Surgery, Department of the Navy, Washington, D.C. The opinions and assertions contained herein are those of the authors and are not to be construed as official or reflecting the views of the U.S. Navy Department or the U.S. Naval Service at large.

† Publications Office, U.S. Naval Medical Research Unit No. 2, Box 14, APO San Francisco 96263, U.S.A.




FIG. 1. RESULTS OF TYPICAL SEPARATION OF PROTEIN COMPONENTS FROM LYOPHILIZED VENOM OF *Laticauda semifasciata*.

Seventy-seven milligrams of dry venom were dissolved in 2 ml of the initial buffer (pH 6.3, 0.01 M sodium phosphate containing 0.01 M sodium chloride) and applied to the column packed with Whatman carboxymethylcellulose, CM-52. Gradient elution was achieved with 500 ml of initial buffer and 500 ml of final buffer (pH 6.3, 0.01 M sodium phosphate containing 0.5 M sodium chloride). Fractions V and VI were isolated for further study.

TABLE 1. COMPARISON OF AMINO ACID COMPOSITIONS OF TOXINS FROM *Laticauda semifasciata*

Amino acid residue	Fraction V		Erabutoxin a		Toxin a		Fraction VI		Erabutoxin b		Toxin b	
	nmoles	Molar ratio	Molar ratio*	Molar ratio†	nmoles	Molar ratio	Molar ratio*	Molar ratio*	Molar ratio†			
CM-Cys	215	8.0	8	8	150	8.0	8	8	8			
Asp	135	5.0	5	5	75	4.0	4	4	4			
Thr	125	4.6	5	6	90	4.8	5	5	5			
Ser	205	7.6	8	7	145	7.7	8	6	6			
Glu	225	8.3	8	8	150	8.0	8	8	8			
Pro	90	3.3	4	4	65	3.5	4	4	4			
Gly	140	5.2	5	5	95	5.1	5	6	6			
Val	50	1.9	2	2	34	1.8	2	3	3			
Ile	80	3.0	4	4	55	3.0	4	4	4			
Leu	27	1.0	1	1	16	0.9	1	1	1			
Tyr	32	1.2	1	1	17	0.9	1	1	1			
Phe	55	2.0	2	2	33	1.8	2	2	2			
Lys	110	4.1	4	4	75	4.0	4	5	5			
His	32	1.2	1	1	42	2.2	2	1	1			
Arg	80	3.0	3	3	50	2.7	3	2	2			

* SATO and TAMIYA (1971).

† TU *et al.* (1971).

methylated, hydrolyzed for 24 hr at 108°C with constant boiling hydrochloric acid containing mercaptoethanol (1/2000 v/v) (KEUTMANN and POTTS, 1969), and analyzed for their amino acid composition. Results are given in Table 1 and compared with results obtained

by previous workers. Except for isoleucine, where recovery was incomplete, presumably because of incomplete hydrolysis of the isoleucylisoleucine bond (SATO and TAMIYA, 1971), amino acid compositions of Fractions V and VI agree completely with those given by SATO and TAMIYA (1971) for Erabutoxins a and b. Compositions of Toxins a and b reported by TU *et al.* (1971) had several differences.

Peptide maps of our tryptic digests of Fractions V and VI were identical except for one peptide from each which had different mobilities. Those peptides were analyzed for amino acid composition with the following results. The peptide from Fraction V had 2 cysteinyl and 3 seryl residues and one each of asparaginyl (determined as aspartyl after hydrolysis), threonyl, glutamyl, prolyl, glycyl, tyrosyl, and lysyl. Composition of the peptide from Fraction VI was the same except for absence of asparaginyl and presence of one histidyl group. These results match those obtained by SATO and TAMIYA (1971) for that section of the erabutoxin molecules between amino acid residues 16 and 27. The sole difference between Erabutoxins a and b reported by SATO and TAMIYA (1971) was at position 26 which was occupied by asparaginyl in Erabutoxin a and by histidyl in Erabutoxin b.

From these results it was concluded that the two toxic fractions isolated from venoms collected from specimens of *Laticauda semifasciata* in Taiwan are identical to those reported by Sato and Tamiya in *L. semifasciata* from Okinawa and different from Toxins a and b reported by TU *et al.* (1971) from *L. semifasciata* from the Philippines.

Acknowledgement—We thank Dr. J.-J. GUNNING for providing the sample of lyophilized crude venom and Miss HELEN HSIN for amino acid analyses.

REFERENCES

- KEUTMANN, H. T. and POTTS, J. T., JR. (1969) Improved recovery of methionine after acid hydrolysis using mercaptoethanol. *Anal. Biochem.* **29**, 175.
- LIU, C.-S., HUBER, G. S., LIN, C.-S. and BLACKWELL, R. Q. (1973) Fractionation of toxins from *Hydrophis cyanocinctus* venom and determination of amino acid composition and end groups of Hydrophitoxin a. *Toxicon* **11**, 73.
- SATO, S. and TAMIYA, N. (1971) The amino acid sequence of Erabutoxins, neurotoxic proteins of sea-snake (*Laticauda semifasciata*) venom. *Biochem. J.* **122**, 453.
- TAMIYA, N. (1972) Contents of toxic components in *Laticauda semifasciata* venom. *J. formosan Med. Ass.* **71**, 389.
- TAMIYA, N. and ABE, H. (1972) The isolation, properties and amino acid sequence of Erabutoxin c, a minor neurotoxic component of the venom of a sea snake *Laticauda semifasciata*. *Biochem. J.* **130**, 547.
- TAMIYA, N. and ARAI, H. (1966) Studies on sea-snake venoms. Crystallization of Erabutoxins a and b from *Laticauda semifasciata* venom. *Biochem. J.* **99**, 624.
- TU, A. T., HONG, B.-S. and SOLIE, T. N. (1971) Characterization and chemical modifications of toxins isolated from the venoms of the sea snake, *Laticauda semifasciata*, from Philippines. *Biochemistry* **10**, 1295.