

UNCLASSIFIED

AD NUMBER
AD810492
NEW LIMITATION CHANGE
TO Approved for public release, distribution unlimited
FROM Distribution authorized to U.S. Gov't. agencies and their contractors; Administrative/Operational Use; MAR 1967. Other requests shall be referred to Department of the Army, Fort Detrick, Attn: Technical Releases Branch, Frederick, MD 21701.
AUTHORITY
Army Biological Defense Research Lab. ltr dtd 28 Sep 1971

THIS PAGE IS UNCLASSIFIED

AD 810 492

AD

TECHNICAL MANUSCRIPT 373


EFFECT OF INPUT MULTIPLICITY
AND TISSUE CELL CONCENTRATION UPON
THE GROWTH OF RIFT VALLEY FEVER VIRUS

Michael D. Orlando
Richard D. DeLauter
Jean M. Riley

MARCH 1967

DEPARTMENT OF THE ARMY
Fort Detrick
Frederick, Maryland

DISCLAIMER NOTICE



**THIS DOCUMENT IS BEST
QUALITY AVAILABLE. THE COPY
FURNISHED TO DTIC CONTAINED
A SIGNIFICANT NUMBER OF
PAGES WHICH DO NOT
REPRODUCE LEGIBLY.**

Reproduction of this publication in whole or in part is prohibited except with permission of the Commanding Officer, Fort Detrick, ATTN: Technical Releases Branch, Technical Information Division, Fort Detrick, Frederick, Maryland, 21701. However, DDC is authorized to reproduce the publication for United States Government purposes.

DDC AVAILABILITY NOTICES

Qualified requesters may obtain copies of this publication from DDC.

Foreign announcement and dissemination of this publication by DDC is not authorized.

Release or announcement to the public is not authorized.

DISPOSITION INSTRUCTIONS

Destroy this publication when it is no longer needed. Do not return it to the originator.

The findings in this publication are not to be construed as an official Department of the Army position, unless so designated by other authorized documents.

DEPARTMENT OF THE ARMY
Fort Detrick
Frederick, Maryland 21701

TECHNICAL MANUSCRIPT 373

EFFECT OF INPUT MULTIPLICITY AND TISSUE
CELL CONCENTRATION UPON THE GROWTH
OF RIFT VALLEY FEVER VIRUS

Michael D. Orlando

Richard D. DeLauter

Jean M. Riley

Product Development Division
AGENT DEVELOPMENT AND ENGINEERING LABORATORY

Project 1B533601D426

March 1967

ABSTRACT

The growth of Rift Valley fever (RVF) virus in L (Earle) cells obtained from two sources and the effect of input multiplicity and tissue cell concentration upon the growth of the virus are described. The titers obtained in the suspension cultures with the cells from two laboratories were significantly different. With both monolayer culture system and suspension culture system, an input multiplicity of 2.5 resulted in the greatest proliferation of virus. Optimal viral yield was obtained in suspension cultures with 40×10^6 tissue cells per 100 ml of suspension.

I. INTRODUCTION

Although many papers have been presented on Rift Valley fever (RVF) virus since its isolation by Daubney, et al.¹ work on the growth of RVF virus in tissue culture systems has been meager. Easterday and Murphy² described the growth of RVF virus in five established cell lines under a limited set of conditions. Coackley³ described the alteration in virulence during serial passage in lamb testis cells. Tribble and Boyle⁴ studied the growth of RVF virus in L (Earle) cells in suspension cultures of 30-ml volume in Nagle's defined medium. However, the effect of input multiplicity (ratio of inoculum to tissue cells) and tissue cell concentration were not reported.

This report describes the growth of RVF virus in L cells obtained from two sources and the effect of input multiplicity and tissue cell concentration upon that growth.

II. MATERIALS AND METHODS

A. VIRUS STRAIN

The pantropic Van Wyk strain of RVF virus described by Kashula⁵ was used. Infected lamb serum stored at -70 C was the working seed stock. It had a titer of about $10^{8.3}$ mouse intracerebral median lethal doses (MICLD₅₀) per ml.

B. TISSUE CULTURE SYSTEMS

The established L cell line was obtained from two laboratories. In laboratory 1, the cell line had been maintained in suspension cultures⁶ on Eagle's minimum essential medium (double strength) with Earle's balanced salt solution supplemented with bovine serum (10%), cysteine (260 µg/ml), and ascorbic acid (50 µg/ml). In laboratory 2, the cell line had been maintained as monolayers with medium 199 (described by Morgan et al.⁷) supplemented with 10% calf serum.

Cultures were maintained in our laboratory as either monolayers or suspension cultures at 36 C. The monolayer cultures were prepared in roller bottles* (650 ml volume) rotated at 7 rpm. The total cell concentration in the roller bottle was 2.8×10^7 cells with 80 ml of growth medium. Suspension cultures were prepared in 100-ml volumes

* Strumia blood plasma bottles obtained from Arthur Thomas Co., Philadelphia, Pa.

in centrifuge bottles (250-ml) fitted with Teflon-coated, suspended bar magnets and removable vents.* Medium 199 supplemented with 10% calf serum was used as the growth medium. Penicillin (250 units per ml), dihydrostreptomycin (250 µg/ml), and kanamycin (50 µg/ml) were added to the growth medium during the viral infectivity studies only.

C. INFECTION OF TISSUE CULTURES

The roller bottles and suspension cultures were infected with sufficient inoculum in a total volume of 10 ml of growth medium to obtain input multiplicities of 1, 2.5, 5, and 10.

The suspension cultures were started with total cell populations of 20, 40, 80, 120, 160, and 200 million cells. The amount of inoculum varied with the total cell population so that a constant input multiplicity of 2.5 was maintained. The culture systems were incubated at 36 C.

D. ASSAY

Tenfold serial dilutions were made in medium 199 plus 10% calf serum. Ten- to 14-gram Swiss-Webster mice were injected intracerebrally with 0.03-ml amounts of appropriate virus dilutions. Ten mice were injected per dilution, and deaths were recorded during the next 6 days. Deaths prior to 24 hours were assumed to be traumatic. The LD₅₀ values were calculated by the method of Reed and Muench.⁸

III. RESULTS

The results of the study (means of four replicates) on the effect of input multiplicity are presented in Table 1. In the roller bottle (monolayer system) the effect of input multiplicity can readily be seen by a comparison of the output factor (yield titer divided by inoculum titer) as follows:

<u>Input Multiplicity</u>	<u>Output Factor</u>
1	40
2.5	40
5	8
10	5

* Personal communication, J. Rosensteel and Wm. F. Daniels.

TABLE 1. EFFECT OF INPUT MULTIPLICITY UPON THE GROWTH OF RIFT VALLEY FEVER VIRUS

Input Multiplicity	Titers, MICLD ₅₀ per ml						
	Roller Bottles ^{a/}		Suspended Cultures ^{b/}				
	at Indicated Time, hr		at Indicated Time, hr				
	0	72	0	24	48	72	96
1	6.00	7.60	6.60	c/	6.44	6.30	5.34
2.5	6.40	7.99	6.90	6.51	6.51	7.17	5.64
5	6.70	7.65	7.20	7.50	6.80	6.74	5.82
10	7.00	7.69	7.50	7.26	6.70	6.73	6.27

- a. Results are the means of the titers obtained from cells from both laboratories (no significant difference between cell sources).
 b. Results are those obtained with cells from laboratory 2 only.
 c. Samples lost.

On the basis of the output factor, the use of either the 1- or 2.5-input multiplicity produces the maximum growth ratio, but on the basis of the largest yield per unit volume (MICLD₅₀/ml) the 2.5-input multiplicity is optimal.

The results of the study on input multiplicity in suspension cultures were disappointing. Apparently the only input multiplicity that resulted in a proliferation of the virus occurred with the 2.5 ratio after 72 hours (an increase of only 0.27 log).

The results of the study (means of four replicates) on the effect of tissue cell concentration are summarized in Table 2 on the basis of titers obtained after 72 hours of incubation. There were no significant differences in the titers for each cell source. However, the titers obtained with cells from the two laboratories were significantly different. At each cell concentration, the titers obtained with the cells from laboratory 2 were greater than those obtained with cells from laboratory 1.

The effect of tissue cell concentration can be seen more readily by a comparison of the output factor. The optimal yield was obtained at a total suspension concentration of 40 million cells per 100 ml of suspension with cells obtained from laboratory 2.

TABLE 2. EFFECT OF TISSUE CELL CONCENTRATION OF SUSPENSION CULTURES UPON
THE GROWTH OF RIFT VALLEY FEVER VIRUS

Cell Concentration, 10 ⁶ cells per suspension	Titer, MICID ₅₀		Per Tissue Culture Cell		Output Factor		Per Viral LD ₅₀ of Inoculum	
	Lab 1	Lab 2	Lab 1	Lab 2	Lab 1	Lab 2	Lab 1	Lab 2
20	6.30	6.81	10	32	4	13		
40	6.87	7.45	19	70	8	28		
80	6.56	7.16	5	18	2	8		
120	6.31	7.42	2	22	1	13		
160	6.26	7.16	1	9	>1	4		
200	-a/	7.16	-	7	-	3		

a. Samples gelatinous and could not be resuspended in diluent.

IV. DISCUSSION

Although the source of L cells did not affect the growth of RVF virus in the monolayers, a significant effect was seen in the suspension cultures. With either system, the monolayer or the suspension culture, the optimal input multiplicity is apparently 2.5. Although the difference in input multiplicity varied tenfold there was a complete destruction of the tissue cells by 72 hours in the monolayer systems.

The optimal tissue cell concentration in the suspension system is apparently 40×10^6 cells per 100 ml. Many factors may influence proliferation of the virus. Since the suspension system was not designed with precise controls, the reasons for the optimal growth at this tissue cell concentration cannot be specified. If the suspension study had been conducted in a "controlled" device as described by Daniels et al.⁸ a better understanding of the metabolic effects of viral infection could have been reached. Since cytolytic viruses alter the biosynthetic processes of the tissue system one could then study the changes that occur in the RVF virus infection of tissue cells.

LITERATURE CITED

1. Daubney, R.; Hudson, J.R.; Garnham, P.C. 1931. Enzootic hepatitis or Rift Valley fever, an undescribed virus disease of sheep, cattle and man from East Africa. *J. Pathol. Bacteriol.* 34:545-579.
2. Easterday, R.C.; Murphy, L.C. 1963. The growth of Rift Valley fever virus in cultures of established lines of cells. *Cornell Vet.* 58:3-11.
3. Coackley, W. 1965. Alteration in virulence of Rift Valley fever virus during serial passage in lamb testis cells. *J. Pathol. Bacteriol.* 89:123-131.
4. Tribble, H.R., Jr.; Boyle, J.J. April 1966. Replication of Rift Valley fever virus in L cell suspension cultures, (Technical Manuscript 259). Medical Bacteriology and Virus and Rickettsia Divisions, U. S. Army Biological Laboratories, Fort Detrick, Frederick, Maryland.
5. Kashula, V.R. 1953. The propagation and modification of strains of Rift Valley fever viruses in embryonated eggs and their use as immunizing agents for domestic animals. Doctor of Veterinary Science Thesis, University of Pretoria, Onderstepoort, S. Africa.
6. Daniels, W.F.; Parker, D.A.; Johnson, R.W.; Schneider, L.E. 1965. Controlled pH and oxidation-reduction potential with a new glass tissue culture fermentor. *Biotech. Bioeng.* 7:529-553.
7. Morgan, J.F.; Morton, H.J.; Parker, R.C. 1950. Nutrition of animal cells in tissue culture: I. Studies on synthetic medium. *Proc. Soc. Exp. Biol. Med.* 73:1-15.
8. Reed, L.J.; Muench, H. 1938. A simple method of estimating fifty per cent end points. *Amer. J. Hyg.* 27:493-497.

Unclassified
Security Classification

DOCUMENT CONTROL DATA - R&D		
<i>(Security classification of title, body of abstract and indexing annotation must be entered when the overall report is classified)</i>		
1. ORIGINATING ACTIVITY (Corporate author) Department of the Army Fort Detrick, Frederick, Maryland, 21701		2a. REPORT SECURITY CLASSIFICATION Unclassified
		2b. GROUP
3. REPORT TITLE EFFECT OF INPUT MULTIPLICITY AND TISSUE CELL CONCENTRATION UPON THE GROWTH OF RIFT VALLEY FEVER VIRUS		
4. DESCRIPTIVE NOTES (Type of report and inclusive dates)		
5. AUTHOR(S) (Last name, first name, initial) Orlando, Michael D. DeLauter, Richard D. Riley, Jean M.		
6. REPORT DATE March 1967	7a. TOTAL NO. OF PAGES 12	7b. NO. OF REFS 8
8a. CONTRACT OR GRANT NO. b. PROJECT NO. 1B533601D426 c. d.	9a. ORIGINATOR'S REPORT NUMBER(S) Technical Manuscript 373	
9b. OTHER REPORT NO(S) (Any other numbers that may be assigned this report)		
10. AVAILABILITY/LIMITATION NOTICES Qualified requesters may obtain copies of this publication from DDC. Foreign announcement and dissemination of this publication by DDC is not authorized. Release or announcement to the public is not authorized.		
11. SUPPLEMENTARY NOTE:	12. SPONSORING MILITARY ACTIVITY Department of the Army Fort Detrick, Frederick, Maryland, 21701	
13. ABSTRACT <p>The growth of Rift Valley fever (RVF) virus in L (Earle) cells obtained from two sources and the effect of input multiplicity and tissue cell concentration upon the growth of the virus are described. The titers obtained in the suspension cultures with the cells from two laboratories were significantly different. With both monolayer culture system and suspension culture system, an input multiplicity of 2.5 resulted in the greatest proliferation of virus. Optimal viral yield was obtained in suspension cultures with 40×10^6 tissue cells per 100 ml of suspension.</p>		
14. Key Words *Rift Valley Fever *Growth *Tissue Cultures L Cells Suspension Cultures		

DD FORM 1473
1 JAN 68

Unclassified
Security Classification