

UNCLASSIFIED

AD NUMBER
AD839361
NEW LIMITATION CHANGE
TO Approved for public release, distribution unlimited
FROM Distribution authorized to U.S. Gov't. agencies and their contractors; Critical Technology; JUL 1968. Other requests shall be referred to Department of the Army, Fort Detrick, Attn: Technical Release Branch, Frederick, MD 21701.
AUTHORITY
AMXFD ltr, 9 Feb 1972

THIS PAGE IS UNCLASSIFIED

196688

TRANSLATION NO. 1101

DATE: July 68

DDC AVAILABILITY NOTICE

Reproduction of this publication in whole or in part is prohibited. However, DDC is authorized to reproduce the publication for United States Government purposes.

DDC
RECEIVED
SEP 13 1968
B

STATEMENT #2 UNCLASSIFIED
This document is subject to special export controls and each transmittal to foreign governments or foreign nationals may be made only with prior approval of Dept. of Army, Fort Detrick, ATTN: Technical Release Branch/TID, Frederick, Maryland 21701

DEPARTMENT OF THE ARMY
Fort Detrick
Frederick, Maryland

12

SOMETHING ABOUT THE SYMBIONTS OF MOSQUITOES

Following is the translation of an article by Antonio Verderol Arnal published in Bol. soc. espan. hist. nat. Madrid. Vol. 41 :209-213. 1943. Translation by Bernice MacDonald.

The descriptions of the digestive apparatus of the Culiroids of Le Dantec, Nuttall and Shipley and Szaudinn are classic. We will deal here with the most interesting of these.

After a muscular pharynx is a thin esophagus which dilates at the end to form a stiff chitinous proventricle. The back of this forms a valve with a sphincter which separates it from the middle intestine. On the walls of the proventricle there are microscopic chitinous projections which stem from each cell separately. They strongly resemble those described in Calliphora by Berlesse but they are longer and more slender. Here the esophagus forms three evaginations, two dorsal and one ventral. This one is a large sac which extends underneath the digestive tube as far as to fill a great part of the abdomen attaining its greatest development in the Anopheles. Nuttall and Shipley consider the three to be food reservoirs. Other authors noticing that similar sacs in other diptera occasionally contain food think it very likely that such diverticula serve as storehouses. Szaudinn calls the ventral

diverticulum a sucking stomach (Saugmagen), which will fill up with blood during suction and which would empty itself immediately into the middle intestine. Nuttall and Shipley give it the same meaning, but to Berlesse it is a simple air sac, destined to maintain the shape of the body when the digestive tube is empty and to leave room for it when it distends upon filling up with blood. These two functions do not seem contradictory as we have observed that the mentioned diverticulum appears as a virtual cavity in fasting mosquitoes, while it appears muscular when the animal sucks blood.

During the fasting period the lower part of the proventricle is folded inside the part close to the middle intestine, which dilates giving it a pear-shaped appearance. Some authors mistakenly call this entire area cardias. The name stomach, confirmed by comparative anatomy, would better suit it.

With the ingestion of blood the proventricle disinvaginates and dilates to the shape of a winebag, its cells separated by a gelatinous substance like quitina (?). This gelatine will surround the mass of blood during its passage to the middle intestine and will form a kind of peritrophic membrane although it is of a different origin than that of the rest of the insects.

The wall of the middle intestine, of endothermic

origin is very monotonous. It is made up (a in Pic. VI, fig. B) of prismatic cells with a thick nucleus which in in apical part have a protective sheet stretched vertically, called rhabdorium.(b). For some authors the rhabdorium would also have an absorbent function. These are the cells which separate the digestive enzymes.

In the middle intestine the digestion and absorption of the food takes place, we dont know if through the same cells as Boissezon says these are separated into absorbent and secreting cells to better fulfill these functions. The secreting rhythm is not the same during fast (secretion of hunger) as during contact with food. We, along with Boissezon, also differentiate two kinds of cells for these two secretions.

In the posterior area the so called piloric valve separates the middle intestine from the posterior intestine which again becomes quitinose. on the outside the middle intestine is surrounded by a strong muscular tunic and has a profusion of tracheas.

Forty years ago Schaudinn pointed out the existence of some chainlike fungi in the ventral sac of the Anopheles which until then was supposed to be only an air sac. The function of these fungi

was presumed to manufacture enzymes and at the same time could release a gas in heavy bubbles. Schaudinn supposed that these enzymes are the ones which produce inflammation of the bite. He saw that the gas produced was CO_2 as it clouded barium water. These funguses which he presumed to be ectable were of great importance to the insect. Some symbiontes exist in the eggs. For Schaudinn they would be hereditary residing in internal tissues from which they would pass to the eggs.

More recently, Hertig and Wolbach have found them again in Culex. In the testicles and ovaries of the males and females there are slender little intercellular canes similar to gramnegative Rickettsias. Some of other species have been identified with Schaudinn's fructiferous forms but in others the same result was not obtained.

Therefore, it is not a case of casual infection. These studies prove that they are real specific symbiontes which in all probability intervene in the nutrition of the host.

I have used for my first investigations three species of very abundant hematophages Culicidos: Anopheles, Culex and Aedes (Stegomia), hunting them by the simple means of placing on them, when they are poised, a large diameter test tube. By introducing into the same tube which served to capture

them a strip of paper where they could poise, they can be kept alive for several days.

I effected the fixing "in toto" in formaldehyde deacidified using chalk in the ratio of 10 to 100; I have made the cuts before inclusion in celodene. For dyeing I have used various techniques: especially the first ⁴ variant of Rio-Hortega to the tonocargentic method of Achucarro, and Rio-Hortega's silver carbonate method.

7 Treatment of the whole insect offers the grave inconvenience of defective inclusion. Even in the most favorable cases there always remains a limit marked by the quitine. For this reason the body silhouette of the body breaks away as soon as } rather fine cuts -10 u- are tried. These are necessary for the histophysiologic studies as the cells of the mosquito are very small.

I am now trying another procedure, not lacking in difficulties either. It consists in a previous dissection of the organs (digestive tube, salivary etc.) before fixing and including them. But this can not be done when the digestive tube is full of blood, which is the most interesting case.

The mosquitoes which have just sucked blood have ~~at first sight the~~ ^{a dilated} abdomen, ~~dilated and full of~~ it. ~~It~~ ^{is} can be observed in a sagittal cut. (Pic. VI)

Fig. 1) that the wall of the middle intestine is distended (a) and its cells have lost height. The intestinal content has zones more altered than others where the red blood cells have hemolyzed, and some have broken, showing only membranes. These zones where hemolysis has started are distributed with preference to the back (c'') and in a central mass (c-c') which advances from the cardias.

Examining in detail (pic. VI. fig. B) this zone of hemolysis, there are seen among the red blood cells (f), more or less altered, a profusion of bacillary bacteria, on occasion, lumped together and also forming chains. The hemocytes situated in the vicinity of the groups of bacteria are hemolyzed, and broken, while leucocytes are always unaltered. This fact makes me suppose that the bacteria are there to cause hemolysis. The role of this preparatory step would limit it in digestion, as the real digestive action is the exclusive work of the digestive juices, separated by the cells of the middle intestine, which would surely go to the interior of the mass of blood through the peritrophic gelatine (e).

Boissezon, in his work on the digestion of blood in Culex, says that hemolysis and later crystallization are produced by the action of the

digestive juices. But, he does not mention these bacteria at all which he probably has not seen. As can be observed in pic. VI, fig. A. this accumulation of bacteria penetrates to the interior of the mass of compressed blood and it forges galleries and sponges it with the help of the gases released, and in this way the digestive enzymes which have to act from the periphery.

If, as Nuttall and Shipley suppose, the blood absorbed passes previously through the esophageic diverticula, it seems very likely that it load it self with the bacteria contained in these. In this was the gelatinous ampule surrounds both and constitutes a nutritious bin already prepared for easy digestion.

Considering the Saugmagen of Schaudinn as a bacteria reservoir, that is, of hemolytic enzymes, the proventricle and its annexed diverticula would be the real stomach where the food would be prepared before passing to the middle intestine.

We do not have enough elements to identify these bacteria. In Culex they are diplobacillus sometimes simple and more rarely forming chains (fig. I, a). In Anopheles they are rounded and always form short chains similar to yeasts (fig. I, b). It is necessary to identify them to use adequate bacteriologic methods and if possible obtain cultures.

It gives me great pleasure to make use of this




Fig. 1.—Symbiontes de *Culex* (a) y de *Anopheles* (b) con glóbulos rojos humanos.
(Obj. Zeiss inn. hom. 90X; oc. 20X.)

Fig. 1 Symbiontes of *Culex* (a) and of *Anopheles* (b) with human red blood cells.
(Obj. Zeiss inn. hom. 90X; oc. 20X.)




Fig. A.—Sagittal cut of *Culex* 2 hours after sucking blood. Method: Rio-Ortega's silver carbonate: a, wall of the middle intestine, distended; b, mass of blood not hemolyzed; c c' c'', principal zones hemolysis with abundant bacteria; d d', Malpighi tubes; e, ovule; f, posterior intestine.




Fig. B. Details of the previous figure at c' level. g; wall of middle intestine; cells in different phases of secretion; b, raborium; 6, basal membrane of the epithelium; d, heavy drops of digestive juices; e, surrounding of gelatine with bacteria; f, hematites more or less hemolysed, with abundant compressed bacteria and in chains; g, unaltered leucocytes.

publication to thank Prof. Alvarado for the interest he has had in following my modest investigations.

Laboratory of Comparative Physiology of the Jose Acosta Institute of Natural Sciences.

Bibliography

Nuttall and Shipley

1901. "The structure and biology of Anopheles"
Jour. of Hyg.

LeDanteq

1901. Patologie Exotique. Paris

Schaudinn

1904. Arb. Kais. Gesundheitsamt.

Thompson

1905. Alimentary canal of the mosquito

Jordan

1913. Vergleichenden Physiologie.

Hertig & Wolbach

1928. "Studies on Rickettsia-like microorganism in insects". Jour. of Med. Res.

Hecht

1928. "Über die Sprosspilze der Oesophagusstulpungen von Stech Mücken". Arch. Schiffs u. Tropenhygiene.

Boissezon

1930. "Digestion en Culex". C.R. Soc. Biol. Paris

Buchner

1930. Tier und Pflanze in Symbiose.

Baeza Cuellar

1933. Estudio medico de los Culicidos hematofagos.