

UNCLASSIFIED

AD NUMBER
AD839633
NEW LIMITATION CHANGE
TO Approved for public release, distribution unlimited
FROM Distribution authorized to U.S. Gov't. agencies only; Administrative/Operational Use; 1968. Other requests shall be referred to Army Foreign Service and Technology Center, Washington, DC.
AUTHORITY
USAFSTC ltr, 8 May 1969

THIS PAGE IS UNCLASSIFIED

FSTC-HT-23-417-68

U.S. ARMY FOREIGN SCIENCE AND TECHNOLOGY CENTER

AD 839633



**NITROGENOUS COMPONENTS OF NUTRIENT MEDIUM FOR CULTIVATION
OF CLOSTRIDIUM PERFRINGENS TYPE D
COUNTRY: USSR**

TECHNICAL TRANSLATION

The translation rights for this document have not been obtained. This document is not in the public domain.

Each transmitter of this document outside the agencies of the U. S. Government must have prior approval of the U. S. Army Foreign Science and Technology Center.

Wash. D.C. 10/15

TECHNICAL TRANSLATION

FSTC-HT-23-417-68

NITROGENOUS COMPONENTS OF NUTRIENT MEDIUM FOR
CULTIVATION OF CLOSTRIDIUM PERFRINGENS TYPE D

by
Ye. Sh. Zhuravel'

Source: Veterinariya
No 6, 1968
pp 25-26
USSR

Translated by: George G. Weickhardt
US Army Foreign Science and Technology Center

This translation is a rendition of the original foreign text without any analytical or editorial comment. Statements or theories advocated or implied are those of the source and do not necessarily reflect the position or opinion of the US Army Foreign Science and Technology Center. This translation is published with a minimum of copy editing and graphics preparation in order to expedite the dissemination of information. Requests for additional copies of this document should be addressed to the Defense Documentation Center, Cameron Station, Alexandria, Virginia, ATTN: OSR-2

NITROGENOUS COMPONENTS OF NUTRIENT MEDIUM FOR
CULTIVATION OF CLOSTRIDIUM PERFRINGENS TYPE D

A growing microbe cell is closely related with the nutrient medium in which it is cultivated. From the medium the cell obtains the substances necessary for its development; the cell also releases the by-products of its metabolism to the medium.

We studied the dynamics of the nitrogenous components of the medium in the growth and toxin-formation of Cl. perfringens strains of different toxicity. The results of this research can have significance for the study of the metabolism of a microbe cell and its relation to toxin formation.

MATERIAL AND METHODS: The work was conducted on two strains obtained from the Microbiology and Immunity Laboratory of the All-Union Institute of Experimental Veterinary Medicine (VIEV): typical toxigenic strain 91 and strain 47 which had lost its ability to produce epsilon-toxin, the basic toxin of type D. Both strains were similar in their morphological, cultural, and some of their biochemical properties (lecithinase and hemolytic activity).

The microorganisms were grown on a casein-fungus Vinogradov medium obtained from the N. F. Gamalei Institute of Epidemiology and Microbiology; 50 ml of 18 - 20 hour culture grown on a casein-fungus medium was sown in 2.5 l. of Vinogradov medium. Samples were taken after 1.5, 3, 6, 8, 12, 18, 24, and 48 hours, passed through a cotton filter, and liberated from the microbe mass by centrifuging at 3.5 to 4 thousand rpm. The number of microbes was determined by optical density on a FEK-M instrument.

The lethal properties of the toxin were studied by means of inter-abdominal introduction of a dose of 0.5 ml native and trypsinized toxin into white mice weighing 16 to 18 grams; the lecithinase and hemolytic properties were studied by the method described by F. M. Tyutnikov (1965).

Biochemical investigation of the following indices of the media and filtrates of the cultures of different growth periods were conducted: total and residual nitrogen after settling out of the protein by 21% solution of trichloroacetic acid was determined by the colorimetric method with a Nessler reagent; the protein nitrogen was determined by the difference between the total and residual nitrogen; the amino-nitrogen by the Harting and Maclean method as modified by N. M. Klimov, V. V. Sukhov, and Ye. Sh. Zhuravel' (1966). In all, five series of tests were conducted.

RESULTS OF THE INVESTIGATION: In the analysis of the data received by us, it should be noted that the beginning of toxin-formation coincided with the beginning of cell division (three-hour culture). The most lethal, hemolytic, and lecithinase activity of the culture filtrates was observed after 12 - 24 hours of growth.

In comparison with strain 91, the growth of the bacterial mass in strain 47 was retarded beginning with the eighth hour of culture growth (the number of microbes in strain 47 after 8 hours was 0.85 billion/ml, and in strain 91, 1.04 billion/ml; after 48 hours the respective quantities were 1.33 and 2.24).

We also determined the regularity in variations of particular nitrogen components of the medium in the growth of both strains. Thus in the culturing of both strains for 1.5 to 48 hours, the total nitrogen (10.8--13.01 mg/ml) and protein (3.95--6.18 mg/ml) in the filtrates of the cultures was discovered to be less, while the residual (6.22--6.94 mg/ml) and amino-nitrogen (1.45--3.31 mg/ml) was discovered to be greater than in a sterile medium (the total nitrogen in the medium was 13.3 mg/ml, the residual nitrogen was 6.19, the protein nitrogen was 7.13, and the amino-nitrogen was 1.28 mg/ml).

Some strain differences were noted in the variations of the medium. In the cultivation period of strain 47 from 6 to 18 hours, there was a gradual reduction in the level of protein nitrogen to 3.95 mg/ml; in the same period of strain 91 growth variations in its level were noted (in 6, 12, and 18 hour cultures there was an increase to 6.05--6.18 mg/ml).

The quantity of amino-nitrogen in the initial period of culturing in both strains was decreased (in a three hour culture it reached the lowest level: 1.45--1.48 mg/ml). Then in strain 91 the content of amino-nitrogen sharply increased and was especially pronounced in an eight hour culture (3.31 mg/ml). In the growth of strain 47, the level of amino-nitrogen was stable (1.51--1.81 mg/ml).

Attention is directed to the relatively constant level of residual nitrogen in both strains (6.22--6.94 mg/ml, excluding six hour culture of strain 91, which was 5.94 mg/ml, and eight hour culture of strain 47, which was 7.30 mg/ml); the pH of the medium shifted to the acid side in the growth of both strains (7.7--6.73).

It is necessary to assume that such media variations are related to the proteolytic activity of the microbe cell. The proteolytic ferments of Cl. perfringens type A were investigated by V. A. Blagoveshchenskiy and others (1964). The authors discovered strengthening of the proteolytic activity in the toxigenic strain. Our data about the increase in amino-nitrogen directly confirms this position.

Some of the increase of the total and protein nitrogen in the growth of a toxigenic strain in the period of intensive growth and toxin formation is apparently related to the passage into the medium of toxin of a protein nature.

CONCLUSIONS

1. Toxin formation in Cl. perfringens type D is related to the multiplication of the culture.
2. In culturing of toxigenic and weak-toxigenic strains of Cl. perfringens type D on a semi-synthetic medium based on a hybrid hydrolysate of casein, the level of total and protein nitrogen of the culture filtrates diminished, while the residual and amino-nitrogen increased in comparison with a sterile medium.
3. In the growth of a toxigenic strain there is a sharp increase in the quantity of amino-nitrogen in the medium during the period of intensive growth and toxin formation. In the filtrates of a culture of a weak-toxigenic strain, the level of nitrogen during the whole period of culturing was relatively stable. This regularity is related to the increased proteolytic activity of the toxigenic strain.

UNCLASSIFIED

Security Classification

DOCUMENT CONTROL DATA - R & D

(Security classification of title, body of abstract and indexing annotation must be entered when the overall report is classified)

1. ORIGINATING ACTIVITY (Corporate author) Foreign Science and Technology Center US Army Materiel Command Department of the Army		2a. REPORT SECURITY CLASSIFICATION UNCLASSIFIED	
		2b. GROUP	
3. REPORT TITLE Nitrogenous Components of Nutrient Medium for Cultivation of <u>Clostridium perfringens</u> Type D			
4. DESCRIPTIVE NOTES (Type of report and inclusive date) Translation			
5. AUTHOR(S) (First name, middle initial, last name) Ye. Sh. Zhuravel', All-Union Institute of Experimental Veterinary Medicine			
6. REPORT DATE 1968	7a. TOTAL NO. OF PAGES 3	7b. NO. OF REFS N/A	
8a. CONTRACT OR GRANT NO.	9a. ORIGINATOR'S REPORT NUMBER(S) FSTC-HT-23-417-68		
b. PROJECT NO. c. 8223627 2301	9b. OTHER REPORT NO(S) (Any other numbers that may be assigned to the report) ACSI Control Number (None)		
10. DISTRIBUTION STATEMENT Each transmittal of this document outside the agencies of the U. S. Government must have the prior approval of the US Army Foreign Science and Technology Center.			
11. SUPPLEMENTARY NOTES The translation rights for this document have not been obtained. This document is not in the public domain.		12. SPONSORING MILITARY ACTIVITY US Army Foreign Science and Technology Center	
13. ABSTRACT This article concerns the dynamics of the nitrogenous components of the medium in the growth and toxin-formation of <u>Clostridium perfringens</u> of different toxicity. From their research, the authors made the following conclusions: (1) toxin formation in <u>Cl-perfringens</u> type D is related to the multiplication of the culture; (2) in culturing of toxigenic and weak-toxigenic strains of <u>Cl-perfringens</u> type D on a semi-synthetic medium based on a hybrid hydrolysate of casein, the level of total end protein nitrogen of the culture filtrates diminished, while the residual and amino-nitrogen increased in comparison with a sterile medium; (3) in the growth of a toxigenic strain there is a sharp increase in the quality of amino nitrogen in the medium during the period of intensive growth and toxin formation. In the filtrates of a culture of a weak-toxigenic strain, the level of nitrogen during the whole period of culturing was relatively stable. This regularity is related to the increased proteolytic activity of the toxigenic strain.			

DD FORM 1473

REPLACES DD FORM 1473, 1 JAN 66, WHICH IS OBSOLETE FOR ARMY USE.

UNCLASSIFIED

Security Classification

UNCLASSIFIED

Security Classification

14	LINK A		LINK B		LINK C	
	FILE	WT	ROLE	WT	ROLE	WT
<u>Clostridium perfringens</u> Biological warfare Toxicity						

UNCLASSIFIED

Security Classification

SUPPLEMENTARY

INFORMATION

DISTRIBUTION AND AVAILABILITY CHANGES

<i>IDENTIFICATION</i>	<i>FORMER STATEMENT</i>	<i>NEW STATEMENT</i>	<i>AUTHORITY</i>
AD-839 633L Army Foreign Science and Technology Center, Washington, D. C. Rept. no. FSTC-HT- 23-417-68 1968	USGO: others to Army Foreign Science and Technology Center, Washington, D. C.	No limitation	USAFSTC ltr, 8 May 69