

UNCLASSIFIED

AD NUMBER
AD840850
NEW LIMITATION CHANGE
TO Approved for public release, distribution unlimited
FROM Distribution authorized to U.S. Gov't. agencies and their contractors; Administrative/Operational Use; APR 1967. Other requests shall be referred to DeCommanding Officer, Fort Detrick, Attn: Technical Release Branch/TID, Frederick, MD 21701.
AUTHORITY
SMUFD, D/A ltr, 15 Feb 1972

THIS PAGE IS UNCLASSIFIED

AD840850

TRANSLATION NO. 1471

DATE: 24 Apr (1967)

DDC AVAILABILITY NOTICE

Reproduction of this publication in whole or in part is prohibited. However, DDC is authorized to reproduce the publication for United States Government purposes.

STATEMENT # UNCLASSIFIED

This document is subject to special export controls and each transmittal to foreign governments or foreign nationals may be made only with prior approval of Dept. of Army, Fort Detrick, ATTN: Technical Release Branch/110, Frederick, Maryland 21701

DEPARTMENT OF THE ARMY
Fort Detrick
Frederick, Maryland

1658

THE BINDING OF SOLUBLE ANTIGEN-ANTIBODY COMPLEXES
TO ERYTHROCYTES

Acta Haematologica
Vol 30 No 3 pages 153-158, 1963

F. Gramlich
H.E. Muller

As investigations by Sorokin [1] and Weigle [2] have shown, cellular blood components are essentially concerned in the elimination of antigen-antibody complexes from the circulation. For leucocytes and thrombocytes it has been shown in this connection that they either phagocytize soluble antigen-antibody products developed in the blood plasma or adsorb them to their surface and suffer injuries in the process which bring about cell destruction [3,4].

The present article is intended to make a contribution to the question of what part the erythrocytes are capable of playing in the elimination of soluble antigen-antibody complexes. We started with the suggestion made by Duestberg [5] that a cleansing function in the circulating blood is to be attributed to the erythrocytes on the ground of the multivalent combining power of their large total surface and their consequent adsorption capacity. As earlier articles have been able to show, the adsorption to the surface of the erythrocytes is a quite specific process, as has been demonstrated for heterogenous macromolecular substances such e.g. as bacterial lipopolysaccharides [6] or myxoviruses [7] as well as for homologous plasma proteins [8,9,10,11].

By a technique which has already been applied to the study of the normal plasma protein film on the surface of human erythrocytes [9], we have tested here to what extent an adsorption of soluble antigen-antibody complexes to the surface of rabbit erythrocytes takes place. These soluble immune complexes were always produced in antigen excess. Human albumin and rabbit antibodies served as a model system. In the experimental series sometimes the antigen and sometimes the antibody was marked with iodine-131. For comparison

we used the adsorption quota of the individual components of the immune complex (iodine¹³¹-tagged human albumin and iodine¹³¹-tagged rabbit gamma globulin)

Material

1. Iodine¹³¹-tagged rabbit antibody against human albumin was prepared by the following method: Rabbits were sensitized against human albumin (Behring-Werke, Marburg) with Freund's adjuvant. Blood and serum containing antibody were obtained by heart puncture. The rabbit gamma globulin was precipitated in 1.75 M ammonium sulfate solution, dissolved in Sørensen's phosphate buffer, pH 8.0, and tagged by a slightly modified Pressman and Sternberger [12] method with iodine¹³¹ (Radiochemical Centre, Amersham). From the solution thus treated, at the equivalence point of the nephelometric Heidelberger curve, which is characterized by a turbidity maximum [13,14], a specific albumin-antialbumin-antibody precipitate was precipitated out with the calculated amount of albumin, repeatedly washed with physiological NaCl solution to eliminate in large part any free iodine components, dissolved in 0.01 N HCl, and separated into albumin and antibody in a column with "Sephadex G 100" (height of column 65 cm) [15,16]. In the first fractions the antibody protein, which is present in pure form in the solution, is concentrated, after the albumin always present in small quantities after neutralization with antibodies and precipitate formation has precipitated out. From the last fractions it was possible after neutralization to obtain a soluble antigen-antibody complex, the gamma globulin-antibody content of which was determined from the measured impulse counts of activity measurement. Excess albumin is always obtained with such an antigen-antibody complex for reasons inherent in the method. For purely formal reasons two molecules of albumin were used per molecule of gamma globulin in view of the confidently assumable bivalence of the antibody molecule [7], so that the soluble antigen-antibody complex is made up of 1 molecule of antibody and 2 molecules of antigen.

2. "Very pure" unmarked human albumin was obtained from the Behring-Werke, Marburg.

3. Iodine¹³¹-tagged human albumin came from the Radiochemical Centre, Amersham; it was dialyzed to remove free iodine 24 hours at 4°C against physiological NaCl solution before use.

4. The preparation of soluble antigen-antibody complex was done by adding the quantity of antibody-bearing rabbit serum computed by a Heidelberger curve to the iodine¹³¹-tagged human albumin. Here again we must reckon with free albumin which is not bound in the antigen-antibody complex.

Table I. Specific Surface Activity of an Immune Complex in Comparison with Its Individual Constituents.

1	2	3	4	5	6	7	8
No.	Protein	Equi- librium Amount % FG	Absolute Amount of Protein mg GB	Number of Erythro- cytes $\times 10^{10}$ ZB	Protein on the Erythro- cyte Surface gamma GBE	Molecule on the Ery- throcyte MGB	Factor of Concen- tration P
1	Iodine ¹³¹	0.0749	30	2.0	22.47	9667	
2	Human	0.0800	30	2.0	24.00	10520	
3	Albumin	0.0984	30	1.5	26.52	15153	
Av.						11780	
4	Iodine ¹³¹	0.2230	30	1.5	66.90	38227	3.25
5	Human Albumin	0.3426	30	1.5	102.78	58728	4.99
6	Rabbit	0.2730	30	2.0	81.90	35234	2.99
7	Antibody	0.5733	30	1.5	173.26	99001	8.40
8	Complex	0.6314	30	1.5	189.42	108234	9.19
9	Iodine ¹³¹ Rabbit Globulin Antibody	1.746	16.3	0.9	284.35	121395	
10	Human	4.406	147.8	0.9	1338.24	1472605	12.13
11	Albumin Iodine ¹³¹ Rabbit Antibody Complex	2.808	147.8	0.9	2400.99	938505	7.43

Method

Three ml each of the antigen-antibody complex solutions thus prepared with known immune-complex content (Table I, Column 4), with the antibody in one case and the antigen in the other tagged with iodine¹³¹, and also of pure iodine¹³¹ tagged antibody solutions without albumin and albumin solutions without antibody, were incubated with 2 ml of an erythrocyte suspension from the rabbit for two hours at room temperature and then washed with a Sørensen phosphate buffer solution, pH 7.2, to which physiological NaCl solution had been added in a 1:1 ratio, by the following method: The 5 ml of whole blood solution was mixed with 5 ml of buffer solution, centrifuged for three minutes at 1200 g, and 5 ml of upper stratum piped off in each case. This washing process was repeated ten times. The radioactivities in the various wash waters and in the erythrocyte sediments were measured. For this purpose a scintillation counter with a drilled-out crystal was used, made by Frieseke & Höpfner, Erlangen.

Results

By the equilibrium amount P_G (Table I, Column 3) in % of the original amount of the protein in question is meant the activity and so, the relationship between activity and protein content being known, the absolute quantity of proteins obtained when equal impulse numbers were found in the liquid and the erythrocyte sediment (which also always contained a certain proportion of the liquid). From this in accordance with the scheme used in an earlier article [9] the number of molecules or of immune complexes per individual erythrocyte (M_{EE} , Table I, Column 7) could be computed from the number of erythrocytes in the starting material (Z_E , Table I, Column 5) when the total amount of protein put in (G_B , Table I, Column 4) was known. The concentration factor F is found as the quotient of the number of individual molecules of human albumin or rabbit gamma globulin and the corresponding number of immune complexes which are adsorbed to the surface of the erythrocytes under equal conditions. As is evident from Table I, widely differing M_{EE} numbers were found as between samples 1-3 and 9 and between 4-8 and 10-11. This must be considered as due to the methodologically conditioned fact that in the studies of 1-8 iodine¹³¹-tagged human albumin from the Radiochemical Centre, Amersham, was used, and in samples 9-11 human albumin from the Behring-Werke, Marburg, was used. In the iodine¹³¹-tagged human albumin from Amersham considerable denaturation phenomena were observed which could not be found in the same degree in the corresponding albumin from the Behring-Werke. As control studies showed, the adsorption of proteins is largely conditioned by their provenience. ([Note] Unpublished experiments.) But since only samples 1-3 were compared with 4-8

and 9 with 10-11, the differing behavior plays no part in the present considerations.

Discussion

The results show that both the heterologous albumin as antigen and the homologous gamma globulin as antibody in the form of an immune complex are bound more strongly to the surface of the erythrocytes than the corresponding proteins dissolved in the free state. The soluble antigen-antibody complex formed thus possesses with respect to the erythrocyte surface a greater tendency to adsorption, which is defined as specific surface activity [18], than the individual proteins of which the immune complex is composed. As shown by Table I, it is greater by a factor of 3 to 12 than in the components from which the soluble complex is made up. From this finding the conclusion may be drawn with great probability that the erythrocytes within the circulating blood perform a cleansing function with regard to soluble antigen-antibody complexes formed. Their capacity for a specific adsorptive bond with such substances may be regarded as evidence for their corresponding function. Whether the corpuscles themselves suffer damage in the process and fall prey to an accelerated sequestration, as is obviously the case especially in allergic-hemolytic conditions, must remain an open question.

We thank the German Research Society for its support and Miss E. Buddenhagen and Miss R. Sixt for their valuable assistance.

Summary

The surface of erythrocytes adsorbs soluble antigen-antibody complexes to a greater extent than it does either of the two components of the complex separately. Human albumin and rabbit antihuman-albumin antibody were investigated as a model of a soluble antigen-antibody complex. The specific concentration of the immune complex at the red cell surface suggests that these cells are actively involved in the removal of antigen-antibody complexes from the circulation.

Bibliography

1. Sorkin, E., "The Cellular Uptake of Antigens and Their Intracellular Fate," Mechanisms of Antibody Formation, Publ. House Czechosl. Academy of Sciences, Prague, 1960, page 61.
2. Weigle, W.O., "The Elimination of Heterologous Proteins from the Blood of Animals," Mechanisms of Antibody Formation, Publ. House Czechosl. Academy of Sciences, Prague, 1960, page 53.
3. Miescher, P., "The Experimental Basis of Immunology of the Leucocytes and Thrombocytes," The Immune Hematology

- of the Leucocytes and Thrombocytes," in Miescher and Vorlaender's Immunpathologie in Klinik und Forschung (Immunopathology in Clinic and Research), Thieme, Stuttgart, 1961.
4. Miescher, P. and Straessle, R., "Experimental Studies of the Mechanism of Thrombocyte Damage by Antigen-Antibody Reactions," Vox Sang., Vol 1, 1956, page 83.
 5. Duesberg, R., "The Defensive Function of the Erythrocytes," Schweizerische Medizinische Wochenschrift (Swiss Medical Weekly), Vol 43, 1960, page 1208.
 6. Milderitz, O, Westphal, O., Sievers, K., Kröger, E., Neter, E., and Braun, O.H., "The Fixation of P³²-Tagged Lipopolysaccharide (Endotoxin) from E. coli to Human Erythrocytes," Biochemische Zeitschrift (Biochemical Journal), Vol 330, 1958, page 34.
 7. Burnet, F.M. and Stanley, W.M., "The Viruses," in Anderson's Hemagglutination by Animal Viruses, Academic Press, New York and London, 1959.
 8. Müller, H.E. and Gramlich, F., "Studies of the Natural Serum-Protein Film of the Surface of the Erythrocytes," Acta Haematologica, Vol 29, 1963, page 135.
 9. Müller, H.E. and Gramlich, F., "The Normal Composition of the Plasma Protein Film on the Surface of Normal Human Erythrocytes," Hoppe-Seyler's Zeitschrift für Physiologische Chemie (Hoppe and Seyler's Journal of Physiological Chemistry), gone to press.
 10. Müller, H.E. and Gramlich, F., "Supplementary Studies on the Plasma Protein Film on the Surface of Normal Human Erythrocytes," Die Naturwissenschaften (The Natural Sciences), Vol 50, 1963, page 447.
 11. Hughes-Jones, N.C. and Gardner, B., "The Exchange of I¹³¹-Labeled Protein Between Red Cells and Serum," Biochemical Journal, Vol 83, 1962, page 404.
 12. Pressman, D. and Sternberger, L.A., "The Relative Rate of Iodination of Serum Components and the Effect of Iodination on Antibody Activity," Journal of the American Chemical Society, Vol 72, 1950, page 2226.
 13. Kabat, E.A. and Mayer, M.M., Experimental Immunochemistry, C.C. Thomas, Springfield, Illinois, 1961.
 14. Schultze, H.E. and Schwick, G., "Quantitative Immunological Determination of the Plasma Proteins," Behringwerk-Mitteilungen (Reports from the Behring Works), Vol 35, 1958, page 1.
 15. Gramlich, F., Mohring, D., and Müller, H.E., "The Isolation of Antibodies by Gel Filtration," Die Naturwissenschaften, Vol 49, 1962, page 451.

16. Mohring, D. and Müller, H.E., "Physicochemical Studies of an Isolated Antibody," Hoppe-Seylers Zeitschrift für Physiologische Chemie, gone to press.
17. Müller, H.E. and Gramlich, F., "On the Valence of Precipitating Antigen-Antibody Systems," Zeitschrift für Hygiene (Journal of Hygiene), Vol 148, 1962, page 582.
18. Wolf, K.L., Physik und Chemie der Grenzflächen (Physics and Chemistry of Boundary Surfaces), Springer, Heidelberg, 1957.

Address of the authors: Dr. F. Gramlich and Dr. H.E. Müller,
Institut für Medizinische Klinik
und Poliklinik der Universität,
Langenbeckstrasse 1, Mainz (Germany).

Best Available Copy