UNCLASSIFIED

AD NUMBER

AD843971

NEW LIMITATION CHANGE

TO

Approved for public release, distribution unlimited

FROM

Distribution authorized to U.S. Gov't. agencies and their contractors; Critical Technology; 14 NOV 1966. Other requests shall be referred to Commanding Officer, Fort Detrick, Attn: SMUFD-AE-T, Frederick, MD 21701.

AUTHORITY

SMUFD, D/A ltr, 15 Feb 1972

THIS PAGE IS UNCLASSIFIED

M 843971

18 Carro

TRANSLATION NO. 1915

1977 IN 198 -

STRUCTURE STRUCTURE

The second of the second s

やないなななののの言

÷.

DATE: 14 man 1966

and the second second

DDC AVAILABILITY NOTICE

This document is subject to special export controls and each transmittal to foreign governments or foreign nationals may be made only with prior approval of Commanding Officer, Fort Detrick, ATTN: SMUFD-AE.T, Frederick, Md. 21701.

> DEPARTMENT OF THE ARMY Fort Detrick Frederick, Maryland



Compt. Rend. Acad. Sci. (Paris) 252: 3157-3159 (1961)

CONCENTRATION F1 PROPRIÉTÉS DES ANTICORPS ANTIGACIDE RISONUCLÉIQUE CONCENTRATION AND PROPERTIES OF ANTI-RIBONUCLEIC ACID ANTIBODIES

Ъy

Emanoil Barbu and Jacques Panijel

Summary

It is possible to concentrate and purify anti-RNA antibodies prevent in anti-ribosomal serum. This purification employs precipitation with RNA. These antibodies will also precipitate with certain synthetic p, ribonucleotides.

We have previously demonstrated^{1,2} that antisera against bacterial ribosomes will precipitate not only with the ribosomes used for immunization but also with ribosomes from various other sources. Certain of these antisera will precipitate even with purified ribonucleic acid (RNA)³. Described in this note is a method for the concentration of anti-RNA antibodies starting with anti-ribosomal sera which are incapable of precipitating spontaneously with free RNA. This method is based on the hypothesis that RNA represents the fundamental common antigen among ribosomes of all sources and that it is the only common antigen among ribosomes originally used for preparation of the immune serum and the so-called 'distant' heterologous ribosomes with which a number of the precipitating artibodies in seva react only feebly⁴. Consequently, the 'distant' heterologous ribosomes ought especially to carry down in the specific precipitate the anti-RNA antibodies which having been concentrated in this manner, are further purified.

1915

The method of preparation used is shown as follows:

(A) To 120 ml of goat antiserum for E. coli K_{12} 3000 ribosomes, are added 12 mg of Fusiformis fusiformis riboscres. After sitting for 16 hr at 4°C, the precipitate is collected, which, after washing, contains 45 mg of protein and 7 mg of RNA. This precipitate is suspended in 10 ml of buffer 'A' (0.14 H NaCi in 3.005 M tris(hydroxymethyl) aminomethane buffer, pH 7.5); Setting for 24 hr at 37°C activates the latent ribosomal RNase so that it hydrolyzes the RNA in the precipitated ribosomes; this follows the dissociation of the ribosomes themselves and, consequently, those of the specific precipitate. The insoluble protein of the ribosomes and possibly the antibodies that are bound to them, remain as a precipitate thereas the products of RNA hydrolysis and the anti-RNA antibodics are dissolved; after centrifugation, one recovers in the supermatant 80 %, accessionly more, of the starting protein. The immune globuling are then precipitated with 33 % saturated ammonium suifate and then redissolved in buffer 'A'. After dialysis against buffer, 34,5 mg of antibody is recovered in the supernatant.

(0) The antibody suspension is reacted with 500 µg of RNA. This was prepared from Ehrlich Ascites cells by six successive phenol extractions and reprecipitated with 1.0 M NaCl.

The anti-RNA reaction alions one to obtain a specific prec.pitate containing 4.5 mg protein and 175 µg RNA. The percentages of RNA and protein precipitated are 35 and 13 % respectively. This antibody/antigen ratic is equivalent to 25,

-2-

The following table shows the results obtained with other preparations concentrated under similar conditions:

 $\overline{)}$

0

 \bigcirc

origin and means of precipi- tation of <u>Y-globulin</u>	Y-globulin in mixture	RNA {n mixture	Yuglobulin in precioitate	RNA in precip.	rntibody antigen	
	6r1	61	×	ટર		
goat antiserum for E. coli K ₁₂ . 3000 - productioned of E. coli K ₁₂ .	1406	25	8	24	28	
coccus A23 ribosomes	1400	95 î 10 10	17 26	52	¥8	
goat antiserum for <u>E. coli</u> K ₁₂ 3000 ribosomes, precipitated with <u>Fusiformis fusiformis</u> ribosomes	420 840	55 65	24 20	e n	17	
rabbit antiserum for <u>Proteus vuigaris</u> ribosomes, precipiteted with <u>Strepto-</u> coccus A ₂ } ribosomes.	1500 1500	46 120 240	20 32 22	85 18 18	ማዋወ	

-3-

(C) The purification of anti-RNA antibodies is affected by the following: the specific precipitate (antibody+RNA) obtained in (B) is redissolved in buffer 'A' and treated with RNase ($5\mu g/ml$) for 24 hr at 37°C. The antibodies which go into solution are precipitated twice with 33 % saturated ammonium sulfate (this eliminates significant traces of RNase) and then dialyzed against buffer 'A'. This preparation, which contains 3.5 mg of antibody, gives with 150 μ g of ascites RNA, a specific precipitate which is treated as before. At the end, one obtained 2.35 mg of purified antibody.

Eventhough the final yield is relatively poor, the precipitating ability of these antibodies has been notably improved. In fact, with 1 mg of antibody and 50 μ g of RNA mixed in buffer with 0.01 M MgSO₄ added⁶ one obtains a precipitate containing 318 μ g of protein and 35 μ g of RNA which are 32 and 70 % respectively of the starting quantities. The antibody/antigen ratio is 9.

The study of the nature and properties of anti-RNA antibodies permit one to now speculate on a number of points:

First- the anti-RNA antibodies are entirely γ -globulin. One of the preparations (rabbit serum no. 92 against <u>E. coli</u> K₁₂ 3000 ribosomes, precipitated with <u>Alcaligenes fecalis</u> ribosomes) when subjected to immuno-electrophoresis gave only a single band with goat antiserum against rabbit whole serum; this band corresponded to the one for γ -globulin in whole rabbit serum. When ribosomes were used in the immunoelectrophoresis experiment, again only one band corresponding to γ -globulin was obtained.

Second- it has been demonstrated that anti-RNA antibodies no longer precipitate RNA after it is treated with RNase³. These antibodies are also capable of precipitating with polymerized RNA from diverse sources and will also precipitate with synthetic polynucleotides⁷. This is true moreover not only of antisera that precipitate strongly with RNA but also of the purified and concentrated antibodies. In the case of rabbit antiserum no. 55 against <u>E. coli</u> B ribosomes, which precipitates spontaneously with RNA, 40 to 70 % of the polyadenylic acid added is precipitated. In a similar manner, about 46 % of the polyadenylic acid was precipitated with purified antibody from a goat antiserum against <u>E. coli</u> K₁₂ 3000 ribosomes (precipitated with ribosomes from <u>Streptococcus</u> A₂₃).

Third- It has not been possible, because of the concentrations employed, to recover in a specific precipitate either all of the antigen or all of the antibody. This may be due to the presence of soluble complexes or an inhibitor effect of an RNA analog as is evidenced in the precipitation of heterologous ribosomes³.

These results confirm the existence of anti-RNA antibodies. But one can still question whether RNA represents in itself the antigenic molety or whether it only constitutes a portion of a more complex antigenic molety. Some of our experiments appear to give some credit to the latter hypothesis.

-5-

References

83

 E. Barbu, J. Panijel, Ph. Cayeux, and R. Wahl, Comptes rendus, 249, 1959, p. 338.

-6-

- (2) J. Panijel and E. Barbu, Comptes rendus, 250, 1960, p. 232.
- (3) E. Barbu and J. Panijel, Comptes rendus, 250, 1960, p. 1382.
- (4) E. Barbu, J. Panijel, and G. Quabh, Ann. Inst. Pasteur, 100, 1961, p. 725.
- (5) W. Hoeller and H. Bödtker, Fed. Froc., 20, 1961, p. 357.
- (6) According to Noeller and Bödtker⁵ the degradation of RNA by γ -gloublins, cited by Brown, Ellem, and Colter, can be inhibited by magnesium.

(7) This was furnished to us by Mme. Grunberg-Manago.

Translated by:

William F. Vincent, Ph.D. ILt EmlC 14 November 1966